

Is exposure to pesticides associated with biological aging? A systematic review and meta-analysis

Shanshan Zuo^{1,4}, Vidhya Sasitharan², Gian Luca Di Tanna³, Judith M. Vonk⁴, Maaïke De Vries⁴, Moustafa Sherif⁵, Balázs Ádám⁵, Juan Carlos Rivillas⁶, Valentina Gallo¹

1. University of Groningen, Campus Fryslân, Department of Sustainable Health, Leeuwarden, Netherlands
2. Queen Mary University of London, London, United Kingdom
3. University of Applied Sciences and Arts of Southern Switzerland, Department of Business Economics, Health and Social Care, Lugano, Switzerland
4. University of Groningen, University Medical Center Groningen, Department of Epidemiology and Groningen Research Institute for Asthma and COPD, Groningen, Netherlands.
5. United Arab Emirates University, College of Medicine and Health Sciences, Institute of Public Health, Al Ain, United Arab Emirates
6. Imperial College London, MRC Centre Environment and Health, School of Public Health, Department of Epidemiology and Biostatistics, London, UK

Corresponding author:

Shanshan Zuo
Department of Sustainable Health
Campus Fryslân
University of Groningen
Wirdumerdijk 34
8911 CE Leeuwarden
The Netherlands
s.zuo@rug.nl

Abstract

Objective: Exposure to pesticides is a risk factor for various diseases, yet its association with biological aging remains unclear. We aimed to systematically investigate the relationship between pesticide exposure and biological aging.

Methods: PubMed, Embase and Web of Science were searched from inception to August 2023. Observational studies investigating the association between pesticide exposure and biomarkers of biological aging were included. Three-level random-effect meta-analysis was used to synthesize the data. Risk of bias was assessed by the Newcastle-Ottawa Scale.

Results: Twenty studies evaluating the associations between pesticide exposure and biomarkers of biological aging in 10,368 individuals were included. Sixteen reported telomere length and four reported epigenetic clocks. Meta-analysis showed no statistically significant associations between pesticide exposure and the Hannum clock (pooled $\beta = 0.27$; 95%CI: -0.25, 0.79), or telomere length (pooled Hedges' $g = -0.46$; 95%CI: -1.10, 0.19). However, the opposite direction of effect for the two outcomes showed an indication of possible accelerated biological aging. After removal of influential effect sizes or low-quality studies, shorter telomere length was found in higher-exposed populations.

Conclusion: The existing evidence for associations between pesticide exposure and biological aging is limited due to the scarcity of studies on epigenetic clocks and the substantial heterogeneity across studies examining telomere length. High-quality studies incorporating more biomarkers of biological aging, focusing more on active chemical ingredients of pesticides and accounting for potential confounders are needed to enhance our understanding of the impact of pesticides on biological aging.

Keywords: Pesticides; Aging; Systematic review; Meta-analysis

1. Introduction

Pesticides are defined as a class of chemical compounds used to manage and control pests (Kim et al., 2017). Based on their target, pesticides are categorized mainly into herbicides, insecticides, fungicides and rodenticides, and are widely used globally to protect crops and improve productivity. Most of the pesticides applied contaminate the soil, evaporate into the atmosphere, infiltrate the water flow, and eventually enter the food chain (Sun et al., 2018). Therefore, pesticides are considered ubiquitous in the environment. People may be occupationally exposed to pesticides when farming, applying pesticides or manufacturing them (Figueiredo et al., 2022). In addition, co-habitants of farmers, residents living nearby farmland, and those who apply pesticides in the household can also be exposed (Bloem & Boonstra, 2023; Galdiano et al., 2021; Stanganelli et al., 2020).

Although acute non-intentional poisoning incidents have become infrequent, long-term and low-dose exposure to pesticides has various adverse effects to human health including neurologic diseases (Bemanalizadeh et al., 2022; Yan et al., 2016), cancers (Togawa et al., 2021), birth defects (Enyew et al., 2023) and diabetes (Jia et al., 2023). Moreover, evidence from longitudinal population-based studies indicates that exposure to pesticides is associated with a higher risk of all-cause mortality (Bao et al., 2020; Di et al., 2023).

Aging is characterized by the physiological deterioration of the human body over time (López-Otín et al., 2023). The aging process occurs alongside elevated risks of many chronic diseases and mortality, however there is a substantial between-person variation in the pace at which aging takes place (Fransquet et al., 2019). Biological age has emerged as a new measure for assessing differences in aging rates. It represents the overall aging process within bodies, which can be different from the chronological age. The difference between chronological and biological age can be quantified giving a measure of accelerated aging (Nakamura & Miyao, 2003). In recent years, various biological aging measures have been proposed and utilized in studies, including telomere length, DNA methylation estimator of telomere length (DNAmTL), epigenetic clocks including Horvath, Hannum, PhenoAge, GrimAge, DunedinPoAm and DunedinPACE (Horvath, 2013; Levine et al., 2018; Q. Wang et al., 2023), the Klemera and Doubal method (KDM), homeostatic dysregulation (HD), allostatic load (AL) and the biological health score (BHS) (Mak et al., 2023).

Numerous lifestyle factors, concurrent diseases or morbidities, and environmental exposures including exposure to pesticides were found to be associated with accelerated or decelerated aging (Oblak et al., 2021) and with mortality rates (Fransquet et al., 2019). Several epidemiological studies reported associations between pesticide exposure and biomarkers of biological aging (Cosemans et al., 2022; Lind et al., 2018). However, the evidence available is not consistent and difficult to appraise at glance, given the numerous markers available for measuring biological aging. To date, no attempts were made to capture the association between pesticide exposure and biological aging in a systematic

way. Therefore, the objective of this study is to systematically review and meta-analyse the evidence on the association between exposure to pesticides and biological aging.

2. Methods

This systematic review and meta-analysis was reported in compliance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) reporting guidelines (Page et al., 2021). The protocol was registered in PROSPERO (CRD42023392845).

2.1. Eligibility criteria

Inclusion criteria were applied according to PECO (Population, Exposure, Comparator, Outcome) statement (Table 1). Epidemiological studies using cohort, case-control, and cross-sectional designs and published in English were included. Related systematic reviews were not included but used to scan for references. Ecological studies, methodological papers, and conference abstracts were excluded.

2.2. Search strategy

Three databases (PubMed, EMBASE, Web of Science) were searched for studies published from inception to August 2023. The search strategy was developed for each database, utilizing Emtree in Embase and combining free text and MeSH terms in PubMed for retrieval. All pesticides, categorized by their targets (pesticide, insecticide, herbicide, fumigant, bactericide, and rodenticide), were searched across the databases. Outcomes comprised epigenetic clocks, telomere length, Klemera-Doubal Method, homeostatic dysregulation, allostatic load and biological health score, along with their broad categories including biological aging, accelerated aging and epigenetic age. Detailed search terms were reported in Table A. 2. Reference lists of included papers and relevant reviews were manually scanned for additional publications.

2.3. Study selection

References were de-duplicated in Endnote software X9. All titles and abstracts were screened for relevance then full-text examination was conducted independently by two authors (SZ and VG). Any discrepancies were resolved by discussion. The following information from the included articles was extracted: first author, year of publication, country, study design, sample size, age and gender of participants, type of pesticide(s), exposure setting and assessment, methods for biological aging estimation, statistical model, covariates adjustment, main results including maximally adjusted effect sizes along with standard error (SE) or 95% confidence intervals (CI).

For studies reporting effect sizes between exposed and non-exposed groups, sample size, mean and standard deviation (SD) of the biomarkers were retrieved from each group. When biomarker data were reported as the median and interquartile range (IQR), they were converted to mean and SD (Wan et al., 2014). If results of multiple independent subgroups were reported in one article, they were merged into a single group (Higgins et al., 2019).

If data were only presented in graphs, WebPlotDigitizer (<https://automeris.io/WebPlotDigitizer/>, Version 4.7) was used to extract numerical values. When studies had overlapping populations and information, the one with larger sample size or provided more information (SD, 95%CI or p-value) was selected. Corresponding authors were contacted for missing data.

2.4. Risk of bias assessment

Two reviewers (SZ and VS) independently assessed the risk of bias of included studies by using the Newcastle-Ottawa Scale (NOS) for cohort, case-control studies (https://www.ohri.ca/programs/clinical_epidemiology/oxford.asp), and the adapted version for cross-sectional studies (Herzog et al., 2013). A score of more than 7 was considered high quality, a score of 6–7 was considered moderate quality and a score of less than 6 was considered low quality. The higher the score, the lower the risk of bias. Any disagreements in scoring between two reviewers were resolved by the third reviewer (VG).

2.5. Synthesis of results

From the initial search, telomere length and epigenetic clock estimates were the only two biomarkers of biological aging identified. Synthesis of results encompassed meta-analysis, sensitivity analysis and moderator analysis. For studies that could not be quantitatively analysed in the meta-analysis, the results were summarized in a narrative form.

2.5.1. Meta-analysis

Meta-analysis was performed where comparable data were available in at least two studies. To account for the multiple dependent effect sizes due to more than one pesticide reported within one study, a three-level random-effects model was employed (Harrer et al., 2021). Contrary to the traditional univariate meta-analytic approach, which only considers sampling variance of the extracted effect sizes and variance between studies, the three-level meta-analysis additionally models the variation of effect sizes within studies (Assink & Wibbelink, 2016; Cheung, 2014).

All studies using epigenetic clock as an outcome reported the linear relationship between pesticide exposure as ln-transformed continuous variable and epigenetic clock. In studies which used telomere length as an outcome, various exposure assessment methods led to reporting of different types of effect sizes (i.e., β coefficient, mean difference). The corrected standardized mean difference (Hedges' g) on the raw scale was chosen as the effect size for the meta-analysis, representing the mean difference in telomere length between individuals with higher and lower exposure to pesticides. Positive values indicated longer telomere length in individuals with higher exposure compared to those with lower exposure, while negative values suggested shorter telomere length. The application of Hedges' g allowed the maximisation of inclusion of the results in the meta-analysis; the majority of studies reported telomere length in exposed and unexposed groups. Moreover, effect size transformation methods were available to convert β coefficients from linear regression models with

binary and continuous exposure variables into Hedges' g . Formulas for calculation and transformation were presented in Appendix B.

Heterogeneity was estimated by the I^2 statistic using the `var.comp` function in *demeta* (Cheung, 2014). The total amount of heterogeneity was decomposed into: a) sampling variance at level 1, b) within-study heterogeneity at level 2 (I^2_{within}), and c) between-study heterogeneity at level 3 ($I^2_{between}$). In addition, statistical significance of heterogeneity was tested using Cochran's Q statistic.

2.5.2. Sensitivity analysis

Outlier detection and influence diagnostics were performed to elucidate the source of heterogeneity and test the robustness of the pooled effect estimate. Effect sizes falling below the first quartile or above the third quartile of the interquartile range (IQR) were considered outliers (Gómez Penedo & Flückiger, 2023). Effect sizes were considered influential if their Cook's distance exceeded three times the mean Cook's distance across included studies (Ellis et al., 2023). Sensitivity analysis was carried out by removing outlier effect sizes, influential effect sizes, low-quality studies, and effect sizes with transformations.

2.5.3. Moderator analysis

Additional moderator analysis was performed using omnibus tests to explore potential sources of heterogeneity across the following moderators: chemical class of pesticide (organophosphate or organochlorine), gender (male or female), exposure setting (occupational or environmental) and expression of telomere length (T/S ratio, bp or kb/genome) (Assink & Wibbelink, 2016).

2.5.4. Publication bias

Publication bias were assessed through visual inspection of funnel plots in combination with the Egger's test. All statistical analyses were conducted using the *metafor* and *dmeta* packages of R statistical software (version 4.3.0).

3. Results

3.1. Study selection

The initial search retrieved 2079 records; after 540 duplicates were removed, 1539 records were screened by title and abstract, and 44 articles were selected for full-text eligibility assessment. Among these, 24 articles were excluded (reasons for exclusion were in Fig. 1), yielding a total of 20 eligible articles for the systematic review (Ali et al., 2023; Andreotti et al., 2015; Cosemans et al., 2022; de Oliveira et al., 2019; de Prado-Bert et al., 2021; dos Santos et al., 2022; Duan et al., 2017; Guzzardi et al., 2016; Hoang et al., 2021; Hou et al., 2013; Kahl, da Silva, et al., 2018; Kahl, Dhillon, Fenech, et al., 2018; Kahl, Dhillon, Simon, et al., 2018; Kahl et al., 2016; Karimi et al., 2020; Lind et al., 2018; Lucia et al., 2022; Ock et al., 2020; Saad-Hussein et al., 2019; Shin et al., 2010). Two additional articles were identified through reference lists of selected systematic reviews (Passos et al., 2022) (Fig. 1). Overall, 17 studies were included in meta-analyses while 3 studies were included in the narrative synthesis only.

Despite the overlapping population between (Hou et al., 2013) and (Andreotti et al., 2015), they were treated as two separate studies and included in the systematic review since the size of overlap was small (40/1234 and 40/568).

3.2. Characteristics of the studies

Summary characteristics of the studies were described in Table 2 and Table 3. Twenty studies evaluating the associations between pesticide exposure and two biomarkers of biological aging in 10368 individuals aged between 7 to 92 years (6647 females and 3721 males) were included. Sixteen studies reported telomere length and four studies reported epigenetic clocks. The vast majority were cross-sectional studies, with only 3 studies using a cohort design. The studies were carried out in South America (n = 6), North America (n = 5), Europe (n = 4), Asia (n = 4) and Africa (n = 1). More studies assessed occupational exposures (55%) rather than environmental exposures (40%), while the sample in one study was composed of occupational exposed and non-occupational exposed (Ali et al., 2023). The included studies primarily assessed exposure through the concentration of pesticide metabolites in biological samples (blood/urine). Other approaches included using occupational exposure as a proxy for exposure and self-reported exposure by questionnaires.

3.3. Risk of bias

The quality scores of the selected studies were in the range of 5-9 points (Table A. 5 and Table A. 6). A total of 9 studies were rated high quality, considering adjustments for important confounding variables, and assessing exposure in biological samples or through validated tools. Eight studies were considered as moderate quality, and three were categorized as low quality due to insufficient control of confounding factors or because they relied on occupation as a proxy for exposure, lacking precise or validated tools for exposure assessment.

3.4. Associations between pesticide exposure and epigenetic age

A total of four studies assessed the association between pesticide exposure and epigenetic clocks (Table 2 and Table A. 3). The majority used a cross-sectional design, with only one study having both a longitudinal and a cross-sectional component, investigating prenatal (cohort) and childhood (cross-sectional) pesticide exposure in relation to children's epigenetic age (de Prado-Bert et al., 2021). The total sample size of the included studies was 3672, comprising 2267 males and 1405 females. Three studies focused on middle-aged and elderly individuals, with ages ranging from 45 to 70 years. One study was conducted among children, with an average age of 7 years.

A total of 25 metabolites from 6 chemical categories were reported, with organochlorine and organophosphate pesticides being the most analysed. One study included a wide range of pesticides including chlorophenoxy (Dicamba, 2,4-D), chloroacetanilides (Acetochlor, Metolachlor), Pyridine (Picloram) and Trazines (Atrazine) (Hoang et al., 2021). Most studies focused on more than one epigenetic clock, including Horvath (Hoang et al., 2021; Lucia et al., 2022), Hannum (Hoang et al., 2021; Lind et al., 2018; Lucia et al., 2022), PhenoAge (Hoang et al., 2021; Lucia et al., 2022),

DunedinPoAm (Lucia et al., 2022), GrimAge (Hoang et al., 2021), and Horvath Skin and Blood (de Prado-Bert et al., 2021; Hoang et al., 2021). Among these, the Hannum clock was the most frequently utilized, being employed in three studies.

Three studies reported positive significant associations between metabolites of four pesticides (p,p'-DDE, DDT, TNC, and AMPA) and accelerated epigenetic aging (Hoang et al., 2021; Lind et al., 2018; Lucia et al., 2022). On the contrary, one study reported a negative association between dimethyldithiophosphate (DMDTP) and epigenetic age; they interpreted this as the result of higher intake of fruits and vegetables; however the results did not change substantially once additionally adjusted for fruit intake, vegetable intake or urinary hippurate, a metabolite marker of fruits and vegetables (de Prado-Bert et al., 2021).

Two studies reporting five effect sizes of the association between levels of pesticide metabolites and the Hannum clock were included in the meta-analysis. Results were presented in the forest plot (Fig. 2). Overall, a positive but not significant association was found between pesticide exposure and epigenetic age (pooled $\beta = 0.27$; 95% CI: -0.25, 0.79). 23.83% of overall heterogeneity was attributable to between-study differences ($I^2_{between}$), while 42.57% was attributable to within-study differences (I^2_{within}).

3.5. Associations between pesticide exposure and telomere length

Sixteen studies reported the association between pesticide exposure and telomere length (Table 3). The sample sizes ranged from 80 to 1724, resulting in a total of 6696 participants, including 4380 males and 2316 females. Three studies measured exposure from blood samples (Guzzardi et al., 2016; Karimi et al., 2020; Shin et al., 2010), three from urine samples (Ali et al., 2023; Cosemans et al., 2022; Ock et al., 2020); two studies used questionnaires (Andreotti et al., 2015; Hou et al., 2013), and the remaining eight categorized participants into exposed and non-exposed groups based on their occupation and exposure history (de Oliveira et al., 2019; dos Santos et al., 2022; Duan et al., 2017; Kahl, da Silva, et al., 2018; Kahl, Dhillon, Fenech, et al., 2018; Kahl, Dhillon, Simon, et al., 2018; Kahl et al., 2016; Saad-Hussein et al., 2019).

Nine studies reported results of in total 72 specific pesticides, mainly belonging to the organochlorine and organophosphate classes, followed by carbamate. The remaining seven studies reported on pesticide mixtures. In all studies, telomere length was assessed using the quantitative polymerase chain reaction (qPCR) method. Specifically, telomere length was expressed as the ratio of telomere repeat copy to the relative number of a single copy gene (T/S ratio) in ten studies, as base pairs (bp) in four studies, and as kb/diploid genome in two studies. Overall, the association between pesticide exposure and telomere length was not consistent (Table A. 4): ten studies reported shorter telomeres and two studies reported longer telomeres (Cosemans et al., 2022; Duan et al., 2017), one study found shorter telomeres with diethylphosphate (DE) derived metabolites but not with dimethylphosphate(DM) derived metabolites (Ali et al., 2023). In one study, a negative association was found between 3,5,6-trichloro-2-pyridinol (TCPY) and telomere length, while a positive

association was found for diethyl thiophosphate (DETP) (Ock et al., 2020). Furthermore, one study indicated an inverted U-shaped change in telomere length with increasing pesticide concentration (Shin et al., 2010), and two studies found no overall association between pesticide exposure and telomere length (de Oliveira et al., 2019; dos Santos et al., 2022).

One study was excluded from the meta-analysis because it reported neither standard error, confidence intervals, or p-values. Of the 15 studies included in the primary meta-analysis, 11 studies presented means of telomere length in exposed and unexposed groups. Hedges' *g* was directly calculated from mean difference data. The remaining study results were converted from β coefficients from linear regression models with binary and continuous exposure variables into Hedges' *g*.

Results of the meta-analysis of the association between pesticide exposure and telomere length were shown in the forest plot (Fig. 3). Overall, telomere length in higher-exposed populations was shorter compared to the lower exposed population, but the association fell short of statistical significance (pooled $g = -0.46$; 95%CI: -1.10, 0.19). 99.15 % of the heterogeneity is explained by between-study, 0.39 % and by within-study.

3.5.1. Sensitivity analysis

The sensitivity analysis showed that excluding effect sizes identified as outliers, or through transformation, did not have a strong influence on the results and the pooled effects remained negative in direction.

Notably, significant shorter telomere length in higher exposed population was found after removing the influential effect sizes (pooled $g = -0.28$; 95%CI: -0.54, -0.02) or low-quality studies (pooled $g = -0.29$; 95%CI: -0.54, -0.01) (Table 4).

3.5.2. Moderator analysis

A summary of the moderator results was presented in Table 5. Chemical class of pesticide, gender, exposure setting, or expression of telomere length did not moderate the effect between pesticide exposure and telomere length.

3.5.3. Publication bias

The funnel plot analysis (Fig. A.1) showed the asymmetrical distribution and Egger's test was statistically significant ($P = 0.0051$), suggesting publication bias favoring studies with null-effect sizes.

4. Discussion

This is the first systematic review summarizing the current evidence on the association between pesticide exposure and biological ageing. Using a three-level meta-analysis to quantitatively synthesize data. In our review, pesticide exposure was not statistically associated with either the epigenetic clock or with telomere length. However, the opposite direction of the pooled effects for both outcomes – the positive on epigenetic age and the negative on telomere length, showed a possible indication of biological aging acceleration in relation to pesticide exposure. Importantly,

after removal of influential effect sizes or low-quality studies pesticide exposure was significantly associated with shorter telomere length.

The aging process is a gradual and multifaceted phenomenon characterized by physiological deterioration at the cellular, tissue, and organ levels, resulting in an increased susceptibility to various diseases. Twelve hallmarks of aging are considered to contribute to the aging process: genomic instability, telomere attrition, epigenetic alterations, loss of proteostasis, disabled macroautophagy, deregulated nutrient sensing, mitochondrial dysfunction, cellular senescence, stem cell exhaustion, altered intercellular communication, chronic inflammation and dysbiosis (López-Otín et al., 2013).

In this systematic review, telomere length and epigenetic clocks were the only biomarkers measuring biological aging captured from existing research in relation to exposure to pesticides. Located at the termini of eukaryotic chromosomes, telomeres play a crucial role in preventing degradation during mitosis (Coltell et al., 2023). This protective function is accomplished through the addition of 5'-TTAGGG-3' tandem repeats to the chromosome ends by telomerase, an enzyme predominantly active in germ cells, stem cells, and immortalized cells (Martens & Nawrot, 2016). Previous studies have found that shorter telomere length was associated with higher mortality (Xiong et al., 2023), development of degenerative diseases, and cardiovascular disease (X. Wang et al., 2023). In numerous epidemiological studies, it was employed as a measurement for biological aging and to predict an individual's health condition. However, the variety of approaches used to quantify telomere length sometimes led to varying outcomes, raising concerns about its accuracy (Lin & Yan, 2005). Based on 140 CpGs and validated in leukocytes from several cohorts, Lu et al. developed a DNA methylation estimator of TL (DNAmTL), which outperformed several limitations of the classical techniques in epidemiological studies (Lu, Seeboth, et al., 2019).

The epigenetic clock was developed on large-scale genomewide DNA methylation-based epigenetic analyses using a variety of cell types and tissues and serves as another molecular marker for biological aging (Horvath, 2013). Composed of a subset of cytosinephosphate-guanine sites (CpGs), epigenetic clock-based DNA methylation age can predict chronological age and was associated with mortality and age-related morbidities (Williams et al., 2023). The first generation of epigenetic clocks, represented by Horvath's and Hannum's, exhibited remarkable accuracy in predicting chronological age (Zhou et al., 2022). The second generation comprises PhenoAge and GrimAge. Instead of solely focusing on the correlation with chronological age, the two clocks also incorporated phenotypic or behavioral surrogate markers that reflect age-related health, disease phenotypes, and all-cause mortality (Levine et al., 2018; Lu, Quach, et al., 2019). However, the diverse epigenetic clocks showed varying degrees of associations with risk factors and outcomes, indicating the potential to capture distinct aspects of the aging process (Oblak et al., 2021). The third generation of clocks, represented by the DunedinPACE, have been developed recently. Through using birth cohort data within-individual decline in 19 indicators of organ-system integrity across two decades to model Pace of Aging, these clocks have showed high test reliability and are considered as a complement to the

second generation epigenetic clocks (Belsky et al., n.d.). Moreover, some other markers for measuring biological aging have emerged, such as the Klemera and Doubal method and Biological Health Score (BHS). These approaches measure the aging by integrating biomarkers from various systems of the body and have demonstrated high reliability in predicting biological age and mortality (K. Li et al., 2024). Current recommendations for estimating biological aging propose the necessity of a comprehensive and integrative score that incorporates both molecular biomarkers and physiological functional parameters (Khan et al., 2017).

4.1. Pesticides and epigenetic age

Generally, the meta-analysis showed a positive but nonsignificant association between pesticide exposure and the Hannum clock. However, the results should be interpreted with caution due to the low number of studies included in the meta-analysis, and the inability to incorporate other epigenetic clocks. On one hand, in contrast to the Hannum clock, which was developed to estimate chronological age, the second-generation clocks demonstrated superior performance in predicting age-related clinical phenotypes and all-cause mortality (McCrory et al., 2020). On the other hand, in comparison to the Horvath clock algorithm, designed as a robust multi-tissue age predictor based on DNA methylation at 353 CpG sites, the Hannum clock serves as a blood-based estimator, characterized by DNA methylation at 71 CpG sites (Fransquet et al., 2019). Across all included studies, positive statistical associations with epigenetic clocks were reported for p,p'-DDE, DDT, TNC, and AMPA. The exact biochemical mechanisms linking pesticides to accelerated aging remain uncertain. However, the connection is supported by some epigenome-wide association studies (EWAS) where differential methylation was identified for pesticides. A case-control study found correlations between organophosphate exposure and methylation levels at 70 significant CpGs, with overrepresentation of genes related to GABA-B receptor II and endothelin signaling pathways, which had impact on neurotransmitter release and inflammation, respectively (Paul et al., 2018). Another study in the Netherlands also reported differential methylation related to occupational pesticide exposure, and further found that differential methylation at specific genomic locations induced by pesticides may play a role in airway disease pathogenesis (Plaat et al., 2018). One recent study by (Hoang et al., 2021) found that glyphosate was linked to 24 differentially methylated CpG sites, while 162 CpG sites were discovered in association with eight pesticides (acetochlor, atrazine, dicamba, malathion, metolachlor, mesotrione, picloram, and heptachlor).

4.2. Pesticides and telomere length

A previous systematic review on occupational pesticide exposure and telomere length included a total of six studies which only reported data on exposed and non-exposed groups (Passos et al., 2022). We have now extended this including sixteen papers, including any exposure route, and including more data in the meta-analysis by using Hedges'g. Our results indicated a negative association between pesticide exposure and telomere length, which however did not reach statistical significance. Moderator analysis make no significant change but showed a concordant direction in the pooled

results. The potential mechanisms through which pesticides affect telomere shortening involve mainly oxidative stress. Telomeres are extremely sensitive to damage by oxidative stress because of the high guanine content in certain telomere sequences and the deficiency in the repair of single strand breaks (Honda et al., 2001; Hou et al., 2013). In contrast, some epidemiological studies reported telomere lengthening with adverse health outcomes such as breast cancer (Y. Li & Ma, 2022), gastric cancer (Z. Wang et al., 2018), and lung cancer (Doherty et al., 2018; Tsatsakis et al., 2023). A longer telomere could result in prolonged cell survival, subsequently elevating the probability of accumulating mutations associated with cancer (Aviv et al., 2017).

Substantial heterogeneity between studies was found and this heterogeneity could not be explained after moderator analysis. It may be attributed to systematic differences in characteristics of the study populations as well as to the assessment of exposure. Among the included studies, different pesticide exposure methods were used by the individual studies, such as biological samples, questionnaires and use of occupation as a proxy for exposure. Moreover, despite the wide range of categories of pesticides studied, seven studies concentrated on a broadly defined pesticide category. Pesticides is a heterogeneous group of very different active chemical ingredients, even within their broad application group of insecticides, herbicides and fungicides, which might mix chemicals with no health effects and chemical with a rather significant health effect (Karalexi et al., 2021). In sensitivity analysis, after removing the influential studies or the low-quality studies, statistically significant associations of pesticide exposure with shorter telomere length were found, which highlighted the importance of longitudinal studies with larger sample sizes and proper adjustment for confounders in producing meaningful results. For publication bias, the funnel plot suggested that studies showing a significant negative association between pesticide exposure and telomere length could be missing, and, consequently, our effect size could be underestimated.

4.3. Strengths and limitations

This is the first comprehensive systematic review examining the association between pesticide exposure and biological aging. Considering that the majority of studies reported effects of more than one pesticide, a three-level meta-analytic model was employed to include all effect sizes in our analysis. This approach not only addresses the dependency of effect sizes but also ensures that all available information is preserved, allowing for the attainment of maximum statistical power (Assink & Wibbelink, 2016). Besides, existing effect size transformation methods were applied to homogenize the effect sizes and enhance the interpretability. Finally, we tested the robustness and explored the heterogeneity through sensitivity and moderator meta-analysis.

Nevertheless, several limitations should be noted. Substantial heterogeneity between studies persisted in the meta-analysis of pesticide exposure and telomere length, despite the implementation of moderator analysis, and this might have prevented producing reliable results. Secondly, the predominance of cross-sectional studies included weakened the pooled level of evidence. Thirdly, only two studies were included in the meta-analysis of the epigenetic clock, allowing for effect

estimates being reported as continuous beta coefficients on the same epigenetic clock. Fourthly, a large portion of the studies only reported broad categories of pesticide mixtures rather than the active chemical ingredients, making it difficult to evaluate individual substances. Lastly, only two biomarkers of biological aging were included in this systematic review, restricting our ability to generalize the results to the overall aging process.

5. Conclusion

This systematic review and meta-analysis do not allow a final conclusion on the association between pesticide exposure and biological ageing measured as epigenetic clock or as telomere length. However, both analyses are suggestive of a possible positive association between pesticide exposure and accelerated aging, although this is not always firmly supported by statistically significant pooled effect estimates. Inadequate study quality, high heterogeneity of reported results, publication bias in studies reporting on telomere length, and the small number of studies reporting on epigenetic clocks, are the main constraints limiting our inference. Future studies conducted on large samples coming from high-quality cohort studies, integrate more biomarkers of biological aging and assessing exposure to pesticides through biomarkers with refined adjustments for potential confounders are warranted.

CRedit authorship contribution statement

Shanshan Zuo: Methodology, Software, Formal analysis, Investigation, Writing - Original Draft, Visualization; **Vidhya Sasitharan**: Investigation, Writing - Review & Editing; **Gian Luca Di Tanna**: Writing - Review & Editing; **Judith M. Vonk**: Methodology, Writing - Review & Editing, Supervision; **Maaike De Vries**: Writing - Review & Editing; **Moustafa Sherif**: Writing - Review & Editing; **Balázs Ádám**: Writing - Review & Editing; **Juan Carlos Rivillas**: Writing - Review & Editing; **Valentina Gallo**: Conceptualization, Investigation, Writing - Review & Editing, Supervision.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors

Data availability

Data will be made available on request.

Acknowledgements

Shanshan Zuo is supported by China Scholarship Council (202208620033). The authors would like to thank Lauren B. Ellis at Northeastern University for providing suggestions on data transformation and

synthesis. Similarly, the authors would like to thank the appointed librarian at the University of Groningen, Joost Driesens, who provided guidance on the search terms selection and search engines. Furthermore, the authors appreciate the authors who supplied the necessary data.

References

- Ali, J. H., Abdeen, Z., Azmi, K., Berman, T., Jager, K., Barnett-Itzhaki, Z., & Walter, M. (2023). Influence of exposure to pesticides on telomere length and pregnancy outcome: Diethylphosphates but not Dimethylphosphates are associated with accelerated telomere attrition in a Palestinian cohort. *Ecotoxicology and Environmental Safety*, 256. Embase Medline. <https://doi.org/10.1016/j.ecoenv.2023.114801>
- Andreotti, G., Hoppin, J. A., Hou, L., Koutros, S., Gadalla, S. M., Savage, S. A., Lubin, J., Blair, A., Hoxha, M., Baccarelli, A., Sandler, D., Alavanja, M., & Beane Freeman, L. E. (2015). Pesticide Use and Relative Leukocyte Telomere Length in the Agricultural Health Study. *PloS One*, 10(7), e0133382. <https://doi.org/10.1371/journal.pone.0133382>
- Assink, M., & Wibbelink, C. J. M. (2016). Fitting three-level meta-analytic models in R: A step-by-step tutorial. *The Quantitative Methods for Psychology*, 12(3), 154–174. <https://doi.org/10.20982/tqmp.12.3.p154>
- Aviv, A., Anderson, J. J., & Shay, J. W. (2017). Mutations, Cancer and the Telomere Length Paradox. *Trends in Cancer*, 3(4), 253–258. <https://doi.org/10.1016/j.trecan.2017.02.005>
- Bao, W., Liu, B., Simonsen, D. W., & Lehmler, H.-J. (2020). Association Between Exposure to Pyrethroid Insecticides and Risk of All-Cause and Cause-Specific Mortality in the General US Adult Population. *JAMA Internal Medicine*, 180(3), 367. <https://doi.org/10.1001/jamainternmed.2019.6019>
- Belsky, D. W., Caspi, A., Corcoran, D. L., Sugden, K., Poulton, R., Arseneault, L., Baccarelli, A., Chamarti, K., Gao, X., Hannon, E., Harrington, H. L., Houts, R., Kothari, M., Kwon, D., Mill, J., Schwartz, J., Vokonas, P., Wang, C., Williams, B. S., & Moffitt, T. E. (n.d.). DunedinPACE, a DNA methylation biomarker of the pace of aging. *eLife*, 11, e73420. <https://doi.org/10.7554/eLife.73420>
- Bemanalizadeh, M., Khoshhali, M., Goli, P., Abdollahpour, I., & Kelishadi, R. (2022). Parental Occupational Exposure and Neurodevelopmental Disorders in Offspring: A Systematic Review and Meta-analysis. *Current Environmental Health Reports*, 9(3), 406–422. <https://doi.org/10.1007/s40572-022-00356-6>
- Bloem, B. R., & Boonstra, T. A. (2023). The inadequacy of current pesticide regulations for protecting brain health: The case of glyphosate and Parkinson's disease. *The Lancet Planetary Health*, 7(12), e948–e949. [https://doi.org/10.1016/S2542-5196\(23\)00255-3](https://doi.org/10.1016/S2542-5196(23)00255-3)
- Bubalo, N. M., & Balan, H. M. (2017). Biological age is an integral parameter of the health status of agricultural workers who have suffered acute and chronic pesticide intoxication. *Toxicology Letters*, 280, S103. Embase. <https://doi.org/10.1016/j.toxlet.2017.07.285>
- Cheung, M. W.-L. (2014). Modeling dependent effect sizes with three-level meta-analyses: A structural equation modeling approach. *Psychological Methods*, 19(2), 211–229. <https://doi.org/10.1037/a0032968>
- Coltell, O., Asensio, E. M., Sorlí, J. V., Ortega-Azorín, C., Fernández-Carrión, R., Pascual, E. C., Barragán, R., González, J. I., Estruch, R., Alzate, J. F., Pérez-Fidalgo, A., Portolés, O., Ordovas, J. M., & Corella, D. (2023). Associations between the New DNA-Methylation-Based Telomere Length

Estimator, the Mediterranean Diet and Genetics in a Spanish Population at High Cardiovascular Risk. *Antioxidants*, 12(11), Article 11. <https://doi.org/10.3390/antiox12112004>

Cosemans, C., Van Larebeke, N., Janssen, B. G., Martens, D. S., Baeyens, W., Bruckers, L., Den Hond, E., Coertjens, D., Nelen, V., Schoeters, G., Hoppe, H.-W., Wolfs, E., Smeets, K., Nawrot, T. S., & Plusquin, M. (2022). Glyphosate and AMPA exposure in relation to markers of biological aging in an adult population-based study. *International Journal of Hygiene and Environmental Health*, 240, 113895. <https://doi.org/10.1016/j.ijheh.2021.113895>

de Oliveira, A. F. B., de Souza, M. R., Benedetti, D., Scotti, A. S., Piazza, L. S., Garcia, A. L. H., Dias, J. F., Niekraszewicz, L. A. B., Duarte, A., Bauer, D., Amaral, L., Bassi Branco, C. L., de Melo Reis, É., da Silva, F. R., & da Silva, J. (2019). Investigation of pesticide exposure by genotoxicological, biochemical, genetic polymorphic and in silico analysis. *Ecotoxicology and Environmental Safety*, 179, 135–142. Embase Medline. <https://doi.org/10.1016/j.ecoenv.2019.04.023>

de Prado-Bert, P., Ruiz-Arenas, C., Vives-Usano, M., Andrusaityte, S., Cadiou, S., Carracedo, Á., Casas, M., Chatzi, L., Dadvand, P., González, J. R., Grazuleviciene, R., Gutzkow, K. B., Haug, L. S., Hernandez-Ferrer, C., Keun, H. C., Lepeule, J., Maitre, L., McEachan, R., Nieuwenhuijsen, M. J., ... Bustamante, M. (2021). The early-life exposome and epigenetic age acceleration in children. *Environment International*, 155, 106683. <https://doi.org/10.1016/j.envint.2021.106683>

Di, D., Zhang, R., Zhou, H., Wei, M., Cui, Y., Zhang, J., Yuan, T., Liu, Q., Zhou, T., Liu, J., & Wang, Q. (2023). Exposure to phenols, chlorophenol pesticides, phthalate and PAHs and mortality risk: A prospective study based on 6 rounds of NHANES. *Chemosphere*, 329, 138650. <https://doi.org/10.1016/j.chemosphere.2023.138650>

Doherty, J. A., Grieshober, L., Houck, J. R., Barnett, M. J., De Dieu Tapsoba, J., Thornquist, M. D., Wang, C.-Y., Goodman, G. E., & Chen, C. (2018). Nested case-control study of telomere length and lung cancer risk among heavy smokers in the β -Carotene and Retinol Efficacy Trial. *British Journal of Cancer*, 118(11), 1513–1517. <https://doi.org/10.1038/s41416-018-0075-0>

dos Santos, I. C., da Silva, J. T., Rohr, P., Lengert, A. V. H., de Lima, M. A., Kahl, V. F. S., da Silva, J., Reis, R. M., & Silveira, H. C. S. (2022). Genomic instability evaluation by BMCyt and telomere length in Brazilian family farmers exposed to pesticides. *Mutation Research - Genetic Toxicology and Environmental Mutagenesis*, 878. Embase Medline. <https://doi.org/10.1016/j.mrgentox.2022.503479>

Duan, X., Yang, Y., Wang, S., Feng, X., Wang, T., Wang, P., Liu, S., Li, L., Yao, W., Cui, L., & Wang, W. (2017). Changes in the expression of genes involved in cell cycle regulation and the relative telomere length in the process of canceration induced by omethoate. *Tumor Biology*, 39(7), 1–10. Embase Medline. <https://doi.org/10.1177/1010428317719782>

Ellis, L. B., Molina, K., Robbins, C. R., Freisthler, M., Sgargi, D., Mandrioli, D., & Perry, M. J. (2023). Adult Organophosphate and Carbamate Insecticide Exposure and Sperm Concentration: A Systematic Review and Meta-Analysis of the Epidemiological Evidence. *Environmental Health Perspectives*, 131(11), 116001. <https://doi.org/10.1289/EHP12678>

Enyew, H. D., Bogale, B. G., Hailu, A. B., & Mereta, S. T. (2023). Environmental exposures and adverse pregnancy outcomes in Ethiopia: A systematic review and meta-analysis. *PLOS ONE*, 18(7), e0288240. <https://doi.org/10.1371/journal.pone.0288240>

Figueiredo, D. M., Vermeulen, R. C. H., Jacobs, C., Holterman, H. J., Van De Zande, J. C., Van Den Berg, F., Gooijer, Y. M., Lageschaar, L., Buijtenhuijs, D., Krop, E., Huss, A., & Duyzer, J. (2022). OBOMod—Modelling framework for residents' exposure to pesticides. *Science of The Total Environment*, 825, 153798. <https://doi.org/10.1016/j.scitotenv.2022.153798>

- Fransquet, P. D., Wrigglesworth, J., Woods, R. L., Ernst, M. E., & Ryan, J. (2019). The epigenetic clock as a predictor of disease and mortality risk: A systematic review and meta-analysis. *Clinical Epigenetics*, *11*(1), 62. <https://doi.org/10.1186/s13148-019-0656-7>
- Galdiano, L. L. da S., Baltar, V. T., Polidoro, S., & Gallo, V. (2021). Household pesticide exposure: An online survey and shelf research in the Metropolitan Region of Rio de Janeiro, Brazil. *Cadernos de Saúde Pública*, *37*, e00099420. <https://doi.org/10.1590/0102-311X00099420>
- Gómez Penedo, J. M., & Flückiger, C. (2023). Interpersonal problems as a predictor of outcome in psychotherapy for depressive and anxiety disorders: A multilevel meta-analysis. *Journal of Consulting and Clinical Psychology*, *91*(10), 562–573. <https://doi.org/10.1037/ccp0000828>
- Guzzardi, M. A., Iozzo, P., Salonen, M. K., Kajantie, E., Airaksinen, R., Kiviranta, H., Rantakokko, P., & Eriksson, J. G. (2016). Exposure to persistent organic pollutants predicts telomere length in older age: Results from the Helsinki Birth Cohort Study. *Aging and Disease*, *7*(5), 540–552. Embase. <https://doi.org/10.14336/AD.2016.0209>
- Harrer, M., Cuijpers, P., Furukawa, T. A., & Ebert, D. D. (2021). *Doing Meta-Analysis With R: A Hands-On Guide* (1st ed.). Chapman & Hall/CRC Press. <https://www.routledge.com/Doing-Meta-Analysis-with-R-A-Hands-On-Guide/Harrer-Cuijpers-Furukawa-Ebert/p/book/9780367610074>
- Herzog, R., Álvarez-Pasquin, M. J., Díaz, C., Del Barrio, J. L., Estrada, J. M., & Gil, Á. (2013). Are healthcare workers' intentions to vaccinate related to their knowledge, beliefs and attitudes? A systematic review. *BMC Public Health*, *13*(1), 154. <https://doi.org/10.1186/1471-2458-13-154>
- Hoang, T. T., Qi, C., Paul, K. C., Lee, M., White, J. D., Richards, M., Auerbach, S. S., Long, S., Shrestha, S., Wang, T., Beane Freeman, L. E., Hofmann, J. N., Parks, C., BIOS Consortium, Xu, C.-J., Ritz, B., Koppelman, G. H., & London, S. J. (2021). Epigenome-Wide DNA Methylation and Pesticide Use in the Agricultural Lung Health Study. *Environmental Health Perspectives*, *129*(9), 097008. <https://doi.org/10.1289/EHP8928>
- Honda, S., Hjelmeland, L. M., & Handa, J. T. (2001). Oxidative Stress–Induced Single-Strand Breaks in Chromosomal Telomeres of Human Retinal Pigment Epithelial Cells In Vitro. *Investigative Ophthalmology & Visual Science*, *42*(9), 2139–2144.
- Horvath, S. (2013). DNA methylation age of human tissues and cell types. *Genome Biology*, *14*(10), 3156. <https://doi.org/10.1186/gb-2013-14-10-r115>
- Hou, L., Andreotti, G., Baccarelli, A. A., Savage, S., Hoppin, J. A., Sandler, D. P., Barker, J., Zhu, Z.-Z., Hoxha, M., Dioni, L., Zhang, X., Koutros, S., Freeman, L. E. B., & Alavanja, M. C. (2013). Lifetime pesticide use and telomere shortening among male pesticide applicators in the Agricultural Health Study. *Environmental Health Perspectives*, *121*(8), 919–924. <https://doi.org/10.1289/ehp.1206432>
- Jia, C., Qiu, G., Wang, H., Zhang, S., An, J., Cheng, X., Li, P., Li, W., Zhang, X., Yang, H., Yang, K., Jing, T., Guo, H., Zhang, X., Wu, T., & He, M. (2023). Lipid metabolic links between serum pyrethroid levels and the risk of incident type 2 diabetes: A mediation study in the prospective design. *Journal of Hazardous Materials*, *459*, 132082. <https://doi.org/10.1016/j.jhazmat.2023.132082>
- Kahl, V. F. S., da Silva, F. R., Alves, J. D. S., da Silva, G. F., Picinini, J., Dhillon, V. S., Fenech, M., de Souza, M. R., Dias, J. F., de Souza, C. T., Salvador, M., Branco, C. D. S., Thiesen, F. V., Simon, D., & da Silva, J. (2018). Role of PON1, SOD2, OGG1, XRCC1, and XRCC4 polymorphisms on modulation of DNA damage in workers occupationally exposed to pesticides. *Ecotoxicology and Environmental Safety*, *159*, 164–171. Embase Medline. <https://doi.org/10.1016/j.ecoenv.2018.04.052>

- Kahl, V. F. S., Dhillon, V., Fenech, M., de Souza, M. R., da Silva, F. N., Marroni, N. A. P., Nunes, E. A., Cerchiaro, G., Pedron, T., Batista, B. L., Cappetta, M., Martínez-López, W., Simon, D., & da Silva, J. (2018). Occupational Exposure to Pesticides in Tobacco Fields: The Integrated Evaluation of Nutritional Intake and Susceptibility on Genomic and Epigenetic Instability. *Oxidative Medicine and Cellular Longevity*, 2018, 7017423. <https://doi.org/10.1155/2018/7017423>
- Kahl, V. F. S., Dhillon, V. S., Simon, D., da Silva, F. R., Salvador, M., Branco, C. D. S., Cappetta, M., Martínez-López, W., Thiesen, F. V., Dias, J. F., Souza, C. T. de, Fenech, M., & da Silva, J. (2018). Chronic occupational exposure endured by tobacco farmers from Brazil and association with DNA damage. *Mutagenesis*, 33(2), 119–128. <https://doi.org/10.1093/mutage/gex045>
- Kahl, V. F. S., Simon, D., Salvador, M., Branco, C. dos S., Dias, J. F., da Silva, F. R., de Souza, C. T., & da Silva, J. (2016). Telomere measurement in individuals occupationally exposed to pesticide mixtures in tobacco fields. *Environmental and Molecular Mutagenesis*, 57(1), 74–84. Embase Medline. <https://doi.org/10.1002/em.21984>
- Karalexi, M. A., Tagkas, C. F., Markozannes, G., Tseretopoulou, X., Hernández, A. F., Schüz, J., Halldorsson, T. I., Psaltopoulou, T., Petridou, E. T., Tzoulaki, I., & Ntzani, E. E. (2021). Exposure to pesticides and childhood leukemia risk: A systematic review and meta-analysis. *Environmental Pollution*, 285, 117376. <https://doi.org/10.1016/j.envpol.2021.117376>
- Karimi, B., Nabizadeh, R., & Yunesian, M. (2020). Association Between Leukocyte Telomere Length and Serum Concentrations of PCBs and Organochlorine Pesticides. *Archives of Environmental Contamination and Toxicology*, 79(1), 122–130. Embase Medline. <https://doi.org/10.1007/s00244-020-00732-z>
- Khan, S. S., Singer, B. D., & Vaughan, D. E. (2017). Molecular and physiological manifestations and measurement of aging in humans. *Aging Cell*, 16(4), 624–633. <https://doi.org/10.1111/accel.12601>
- Kim, K.-H., Kabir, E., & Jahan, S. A. (2017). Exposure to pesticides and the associated human health effects. *Science of The Total Environment*, 575, 525–535. <https://doi.org/10.1016/j.scitotenv.2016.09.009>
- Levine, M. E., Lu, A. T., Quach, A., Chen, B. H., Assimes, T. L., Bandinelli, S., Hou, L., Baccarelli, A. A., Stewart, J. D., Li, Y., Whitsel, E. A., Wilson, J. G., Reiner, A. P., Aviv, A., Lohman, K., Liu, Y., Ferrucci, L., & Horvath, S. (2018). An epigenetic biomarker of aging for lifespan and healthspan. *Aging*, 10(4), 573–591. <https://doi.org/10.18632/aging.101414>
- Li, K., Wu, J., Zhou, Q., Zhao, J., Li, Y., Yang, M., Yang, Y., Hu, Y., Xu, J., Zhao, M., & Xu, Q. (2024). The mediating role of accelerated biological aging in the association between blood metals and cognitive function. *Journal of Hazardous Materials*, 462, 132779. <https://doi.org/10.1016/j.jhazmat.2023.132779>
- Li, Y., & Ma, L. (2022). Relationship between telomere length and the prognosis of breast cancer based on estrogen receptor status: A Mendelian randomization study. *Frontiers in Oncology*, 12. <https://www.frontiersin.org/articles/10.3389/fonc.2022.1024772>
- Lin, K.-W., & Yan, J. (2005). The telomere length dynamic and methods of its assessment. *Journal of Cellular and Molecular Medicine*, 9(4), 977–989. <https://doi.org/10.1111/j.1582-4934.2005.tb00395.x>
- Lind, P. M., Salihovic, S., & Lind, L. (2018). High plasma organochlorine pesticide levels are related to increased biological age as calculated by DNA methylation analysis. *Environment International*, 113, 109–113. <https://doi.org/10.1016/j.envint.2018.01.019>
- López-Otín, C., Blasco, M. A., Partridge, L., Serrano, M., & Kroemer, G. (2013). The Hallmarks of Aging. *Cell*, 153(6), 1194–1217. <https://doi.org/10.1016/j.cell.2013.05.039>

- López-Otín, C., Blasco, M. A., Partridge, L., Serrano, M., & Kroemer, G. (2023). Hallmarks of aging: An expanding universe. *Cell*, *186*(2), 243–278. <https://doi.org/10.1016/j.cell.2022.11.001>
- Lu, A. T., Quach, A., Wilson, J. G., Reiner, A. P., Aviv, A., Raj, K., Hou, L., Baccarelli, A. A., Li, Y., Stewart, J. D., Whitsel, E. A., Assimes, T. L., Ferrucci, L., & Horvath, S. (2019). DNA methylation GrimAge strongly predicts lifespan and healthspan. *Aging*, *11*(2), 303–327. <https://doi.org/10.18632/aging.101684>
- Lu, A. T., Seeboth, A., Tsai, P.-C., Sun, D., Quach, A., Reiner, A. P., Kooperberg, C., Ferrucci, L., Hou, L., Baccarelli, A. A., Li, Y., Harris, S. E., Corley, J., Taylor, A., Deary, I. J., Stewart, J. D., Whitsel, E. A., Assimes, T. L., Chen, W., ... Horvath, S. (2019). DNA methylation-based estimator of telomere length. *Aging (Albany NY)*, *11*(16), 5895–5923. <https://doi.org/10.18632/aging.102173>
- Lucia, R. M., Huang, W.-L., Pathak, K. V., McGilvrey, M., David-Dirgo, V., Alvarez, A., Goodman, D., Masunaka, I., Odegaard, A. O., Ziogas, A., Pirrotte, P., Norden-Krichmar, T. M., & Park, H. L. (2022). Association of Glyphosate Exposure with Blood DNA Methylation in a Cross-Sectional Study of Postmenopausal Women. *Environmental Health Perspectives*, *130*(4), 047001. <https://doi.org/10.1289/EHP10174>
- Mak, J. K. L., McMurrin, C. E., Kuja-Halkola, R., Hall, P., Czene, K., Jylhävä, J., & Hägg, S. (2023). Clinical biomarker-based biological aging and risk of cancer in the UK Biobank. *British Journal of Cancer*, *129*(1), Article 1. <https://doi.org/10.1038/s41416-023-02288-w>
- Martens, D. S., & Nawrot, T. S. (2016). Air Pollution Stress and the Aging Phenotype: The Telomere Connection. *Current Environmental Health Reports*, *3*(3), 258–269. <https://doi.org/10.1007/s40572-016-0098-8>
- McCrary, C., Fiorito, G., Hernandez, B., Polidoro, S., O'Halloran, A. M., Hever, A., Ni Cheallaigh, C., Lu, A. T., Horvath, S., Vineis, P., & Kenny, R. A. (2020). GrimAge Outperforms Other Epigenetic Clocks in the Prediction of Age-Related Clinical Phenotypes and All-Cause Mortality. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*, *76*(5), 741–749. <https://doi.org/10.1093/gerona/glaa286>
- Nakamura, E., & Miyao, K. (2003). Further Evaluation of the Basic Nature of the Human Biological Aging Process Based on a Factor Analysis of Age-related Physiological Variables. *The Journals of Gerontology: Series A*, *58*(3), B196–B204. <https://doi.org/10.1093/gerona/58.3.B196>
- Oblak, L., van der Zaag, J., Higgins-Chen, A. T., Levine, M. E., & Boks, M. P. (2021). A systematic review of biological, social and environmental factors associated with epigenetic clock acceleration. *Ageing Research Reviews*, *69*, 101348. <https://doi.org/10.1016/j.arr.2021.101348>
- Ock, J., Kim, J., & Choi, Y.-H. (2020). Organophosphate insecticide exposure and telomere length in U.S. adults. *The Science of the Total Environment*, *709*, 135990. <https://doi.org/10.1016/j.scitotenv.2019.135990>
- Page, M. J., McKenzie, J. E., Bossuyt, P. M., Boutron, I., Hoffmann, T. C., Mulrow, C. D., Shamseer, L., Tetzlaff, J. M., Akl, E. A., Brennan, S. E., Chou, R., Glanville, J., Grimshaw, J. M., Hróbjartsson, A., Lalu, M. M., Li, T., Loder, E. W., Mayo-Wilson, E., McDonald, S., ... Moher, D. (2021). The PRISMA 2020 statement: An updated guideline for reporting systematic reviews. *Systematic Reviews*, *10*(1), 89. <https://doi.org/10.1186/s13643-021-01626-4>
- Passos, J. D. C., Felisbino, K., Laureano, H. A., & Guiloski, I. C. (2022). Occupational exposure to pesticides and its association with telomere length—A systematic review and meta-analysis. *Science of The Total Environment*, *849*, 157715. <https://doi.org/10.1016/j.scitotenv.2022.157715>

- Paul, K. C., Chuang, Y.-H., Cockburn, M., Bronstein, J. M., Horvath, S., & Ritz, B. (2018). Organophosphate pesticide exposure and differential genome-wide DNA methylation. *Science of The Total Environment*, *645*, 1135–1143. <https://doi.org/10.1016/j.scitotenv.2018.07.143>
- Plaat, D. A. van der, Jong, K. de, Vries, M. de, Diemen, C. C. van, Nedeljković, I., Amin, N., Kromhout, H., Consortium, B. I. O. S., Vermeulen, R., Postma, D. S., Duijn, C. M. van, Boezen, H. M., & Vonk, J. M. (2018). Occupational exposure to pesticides is associated with differential DNA methylation. *Occupational and Environmental Medicine*, *75*(6), 427–435. <https://doi.org/10.1136/oemed-2017-104787>
- Saad-Hussein, A., Beshir, S., Taha, M. M., Shahy, E. M., Shaheen, W., Abdel-Shafy, E. A., & Thabet, E. (2019). Early prediction of liver carcinogenicity due to occupational exposure to pesticides. *Mutation Research/Genetic Toxicology and Environmental Mutagenesis*, *838*, 46–53. <https://doi.org/10.1016/j.mrgentox.2018.12.004>
- Shin, J. Y., Choi, Y. Y., Jeon, H. S., Hwang, J. H., Kim, S. A., Kang, J. H., Chang, Y. S., Jacobs, D. R., Park, J. Y., & Lee, D. H. (2010). Low-dose persistent organic pollutants increased telomere length in peripheral leukocytes of healthy Koreans. *Mutagenesis*, *25*(5), 511–516. Embase Medline. <https://doi.org/10.1093/mutage/geq035>
- Stanganelli, I., De Felici, M. b., Mandel, V. d., Caini, S., Raimondi, S., Corso, F., Bellerba, F., Quaglino, P., Sanlorenzo, M., Ribero, S., Medri, M., Farnetani, F., Feliciani, C., Pellacani, G., Gandini, S., & Intergroup, I. the I. M. (2020). The association between pesticide use and cutaneous melanoma: A systematic review and meta-analysis. *Journal of the European Academy of Dermatology and Venereology*, *34*(4), 691–708. <https://doi.org/10.1111/jdv.15964>
- Sun, S., Sidhu, V., Rong, Y., & Zheng, Y. (2018). Pesticide Pollution in Agricultural Soils and Sustainable Remediation Methods: A Review. *Current Pollution Reports*, *4*(3), 240–250. <https://doi.org/10.1007/s40726-018-0092-x>
- Togawa, K., Leon, M. E., Lebailly, P., Beane Freeman, L. E., Nordby, K.-C., Baldi, I., MacFarlane, E., Shin, A., Park, S., Greenlee, R. T., Sigsgaard, T., Basinas, I., Hofmann, J. N., Kjaerheim, K., Douwes, J., Denholm, R., Ferro, G., Sim, M. R., Kromhout, H., & Schüz, J. (2021). Cancer incidence in agricultural workers: Findings from an international consortium of agricultural cohort studies (AGRICOH). *Environment International*, *157*, 106825. <https://doi.org/10.1016/j.envint.2021.106825>
- Tsatsakis, A., Oikonomopoulou, T., Nikolouzakis, T. K., Vakonaki, E., Tzatzarakis, M., Flamourakis, M., Renieri, E., Fragkiadaki, P., Iliaki, E., Bachlitzanaki, M., Karzi, V., Katsikantami, I., Kakridonis, F., Hatzidaki, E., Tolia, M., Svistunov, A. A., Spandidos, D. A., Nikitovic, D., Tsiaoussis, J., & Berdiaki, A. (2023). Role of telomere length in human carcinogenesis (Review). *International Journal of Oncology*, *63*(1), 1–24. <https://doi.org/10.3892/ijo.2023.5526>
- Wan, X., Wang, W., Liu, J., & Tong, T. (2014). Estimating the sample mean and standard deviation from the sample size, median, range and/or interquartile range. *BMC Medical Research Methodology*, *14*(1), 135. <https://doi.org/10.1186/1471-2288-14-135>
- Wang, Q., Hou, T., Wang, Q., He, J., Wang, L., Si, J., & Chen, S. (2023). An evaluation of aging measures: From biomarkers to clocks. *Biogerontology*, *24*(3), 303–328. <https://doi.org/10.1007/s10522-022-09997-4>
- Wang, X., Deng, H., Lin, J., Zhang, K., Ni, J., Li, L., & Fan, G. (2023). Distinct roles of telomerase activity in age-related chronic diseases: An update literature review. *Biomedicine & Pharmacotherapy*, *167*, 115553. <https://doi.org/10.1016/j.biopha.2023.115553>

- Wang, Z., Koh, W.-P., Jin, A., Wang, R., & Yuan, J.-M. (2018). Telomere length and risk of developing gastric adenocarcinoma: The Singapore Chinese Health Study. *Gastric Cancer*, *21*(4), 598–605. <https://doi.org/10.1007/s10120-017-0783-9>
- Williams, A. M., Mandelblatt, J. S., Wang, M., Dong, Q., Armstrong, G. T., Bhakta, N., Brinkman, T. M., Ehrhardt, M. J., Mulrooney, D. A., Gilmore, N., Robison, L. L., Yasui, Y., Small, B. J., Srivastava, D., Hudson, M. M., Ness, K. K., Krull, K. R., & Wang, Z. (2023). Deficit Accumulation Index and Biological Markers of Aging in Survivors of Childhood Cancer. *JAMA Network Open*, *6*(11), e2344015. <https://doi.org/10.1001/jamanetworkopen.2023.44015>
- Xiong, L., Yang, G., Guo, T., Zeng, Z., Liao, T., Li, Y., Li, Y., Chen, F., Yang, S., Kang, L., & Liang, Z. (2023). 17-year follow-up of association between telomere length and all-cause mortality, cardiovascular mortality in individuals with metabolic syndrome: Results from the NHANES database prospective cohort study. *Diabetology & Metabolic Syndrome*, *15*(1), 247. <https://doi.org/10.1186/s13098-023-01206-7>
- Yan, D., Zhang, Y., Liu, L., & Yan, H. (2016). Pesticide exposure and risk of Alzheimer's disease: A systematic review and meta-analysis. *Scientific Reports*, *6*, 32222. <https://doi.org/10.1038/srep32222>
- Zhou, A., Wu, Z., Zaw Phy, A. Z., Torres, D., Vishwanath, S., & Ryan, J. (2022). Epigenetic aging as a biomarker of dementia and related outcomes: A systematic review. *Epigenomics*, *14*(18), 1125–1138. <https://doi.org/10.2217/epi-2022-020>

Tables

Table 1 Population-Exposure-Comparator-Outcome (PECO) statement

PECO Domain	Inclusion criteria	Exclusion criteria
P	General population including children and adults.	Populations with existing diseases or health conditions.
E	Exposure to any type of pesticides, measured via biological markers, self-report, questionnaires, or use of occupation as a proxy for exposure; Environmental or occupational □ setting.	Toxicants not strictly classified as pesticides such as dioxins (and dioxin-like compounds), polychlorinated biphenyls (PCBs), and polychlorinated dibenzofurans (PCDFs); Studies reporting acute exposure events.
C	Reference categories identified as groups with unexposed or less-exposed levels	
O	Biomarkers of biological aging (epigenetic clock, telomere length, Klemra-Doubal method, allostatic load, homeostatic □ dysregulation) reported objectively.	Cellular aging or senescence; DNA methylation analysis without reporting of biological aging biomarkers

Table 2 Characteristics of included studies on pesticide exposure and epigenetic clocks.

Author, year	Country	Study population	N (female %)	Study design	Pesticides	Exposure setting	Exposure assessment	Clocks	Sample, platform	Adjustments
De Prado-Bert, et al., 2021	Multi-countries across Europe: UK, France, Spain, Lithuania, Norway and Greece	Children from HELIX project, six longitudinal population-based birth cohorts across Europe	1173 (45%)	Prenatal exposome: cohort, childhood exposome: cross-sectional	DDE, DDT, HCB, DEP, DETP, DMP, DMTP, DMDTP (only in childhood)	Environmental	Blood	Horvath's Skin and Blood	Blood, 450K	Child's sex, cohort, self-reported maternal education, self-reported ancestry, maternal age at conception
Hoang, et al., 2021	USA	Licensed pesticide applicators from Agricultural Health Study	1170 (0%)	Cross-Sectional	Dicamba, Picloram, Mesotrione, Acetochlor, Metolachlor, Glyphosate, 2,4-D, Atrazine, Malathion, Aldrin, Chlordane, DDT, Dieldrin, Heptachlor, Lindane, Toxaphene	Occupational	Questionnaire	Horvath, Hannum, Levine, Horvath skin and blood, GrimAge	Blood, 450K	Asthma status, age at blood collection, pack-years, smoking, state of residence, cell types
Lind, et al., 2018	Sweden	Community-dwelling elderly from Prospective Study of the Vasculature in Uppsala Seniors (PIVUS)	967 (50%)	Cross-Sectional	p,p'-DDE, HCB, TNC	Environmental	Blood	Hannum	Blood, 450K	Sex, education level, exercise habits, smoking, energy and alcohol consumption and BMI

Lucia, et al., 2022	USA	Postmenopausal women residing in southern California	392 (100%)	Cross-Sectional	Glyphosate, AMPA	Environmental	Urine	Horvath, Hannum, Levine, DunedinPoAm	Blood, 450K	Urinary creatinine concentration, age, race/ethnicity, BMI, smoking status, alcohol consumption, organic eating, diet quality, batch, position on chip and WBC type proportions
---------------------	-----	--	------------	-----------------	------------------	---------------	-------	--------------------------------------	-------------	---

HCB: hexachlorobenzene; TNC: transnonachlor; DMP: dimethylphosphate; DEP: diethylphosphate; DMTP: dimethylthiophosphate; DETP: diethylthiophosphate; DMDTP: dimethyldithiophosphate; DEDTP: diethyldithiophosphate; TCPY: 3,5,6-trichloro-2-pyridinol

Table 3 Characteristics of included studies pesticide exposure and telomere length.

Author, year	Country	Study population	N (female %)	Study design	Pesticides	Exposure setting	Exposure assessment	Telomere length			Adjustments
								Tissue	Method	Expression	
Ali, et al., 2023	Palestine	Pregnant women recruited by consecutive cluster sampling at five hospitals	142 (100%)	Cross-sectional	DMP, DEP, DMTP, DETP, DMDTP, DEDTP	Occupational and environmental	Urine	Blood	qPCR	T/S ratio	Age
Andreotti et al., 2015	USA	Licensed private applicators from AHS	568 (0%)	Cohort	Alachlor, Butylate, Metolachlor, Paraquat, Sethoxydim, 2,4-D, 2,4,5-TP, Aldrin, DDT, Diazinon, Heptachlor, Benomyl, Chlorothalonil, Maneb/Mancozeb, Metalaxyl	Occupational	Questionnaire	Leukocyte	qPCR	T/S ratio	Age at blood draw, the most highly correlated pesticide with telomere length
Cosemans, et al., 2022	Belgium	Sample from Flemish Environment and Health Study	181 (52%)	Cross-sectional	Glyphosate, AMPA	Environmental	Urine	Blood	qPCR	T/S ratio	Sex, age, BMI, smoking status, alcohol, consumption, socioeconomic status, season, and urine specific gravity
De Oliveira, et al., 2019	Brazil	Exposed: occupationally exposed to pesticides used in soybeans; Non-exposed: not occupationally exposed to pesticides or genotoxic substances	148 (0%)	Cross-sectional	Mixture including fungicide, herbicide, insecticide	Occupational	Proxy	Blood	qPCR	bp	None
Dos Santos, et al., 2022	Brazil	Exposed: farm workers; Non-exposed: administrative employees from the Barretos Cancer Hospital	162 (17%)	Cross-sectional	Mixture including herbicides, insecticides and fungicides	Occupational	Proxy	Blood	qPCR	kb/diploid genome	None
Duan, et al., 2017	China	Exposed: recruited from omethoate	295 (35%)	Cross-sectional	Omethoate	Occupational	Proxy	Blood	qPCR	T/S ratio	None

		packaging workshop; Non-exposed: from a company organization in the same city									
Guzzardi, et al., 2016	Finland	Randomly-selected subset of Helsinki Birth Cohort Study	1082 (56%)	Cohort	Oxychlordan, Trans-nonachlor, p, p'-DDE	Environmental	Blood	Blood	qPCR	T/S ratio	Lipids, age or change in age between the two visits, Body Fat % , Diseases
Hou, et al., 2013	USA	Licensed private pesticide applicators (mostly farmers) residing in Iowa and North Carolina from AHS	1234 (0%)	Cohort	Alachlor, Metolachlor, Trifluralin, 2,4-D, DDT, Permethrin (poultry/livestock), Toxaphene, Atrazine, Butylate, Chlorimuron-ethyl, Cyanazine, Dicamba, EPTC, Glyphosate, Imazethapyr, Metribuzin, Paraquat, Pendimethalin, Petroleum oils, 2,4,5 trichlorophenoxy acetic acid, 2,4,5 TP, Aldicarb, Aldrin, Carbaryl, Carbofuran, Chlordane, Chlorpyrifos, Coumaphos, Diazinon, Dichlorvos, Dieldrin, Fonofos, Heptachlor, Lindane, Malathion, Parathion, Permethrin (for crop), Phorate, Terbufos, Aluminum phosphide, Methyl bromide, Ethylene dibromide, Carbon tetrachloride/Carbon disulfide-80/20 mix, Benomyl, Captan, Chlorothalonil, Maneb/Mancozeb, Metalaxyl	Occupational	Questionnaire	Buccal cells	qPCR	T/S ratio	Age at buccal collection, state (Iowa vs, North Carolina), license types (private vs, commercial), regular use of chewing tobacco for ≥ 6 months (yes vs, no), total pesticide exposure days
Kahl, et al.,	Brazil	Exposed:	242 (56%)	Cross-	Mixture including	Occupational	Proxy	Blood	qPCR	bp	None

2018a		agricultural workers; Non-exposed: office and retail employees living in the same region were but not occupationally exposed to any known genotoxic agents		sectional	carbamates, organophosphates, pyrethroids and organochlorines	1						
Kahl, et al., 2018b	Brazil	Exposed: farmers; Nonexposed: indoor office workers living in the same area	80 (53%)	Cross-sectional	Mixture including fungicide, herbicide, insecticide	Occupational	Proxy	Blood	qPCR	bp	None	
Kahl, et al., 2018c	Brazil	Exposed: farmers; Unexposed: not exposed to any known genotoxic agent at their workplace	130 (74%)	Cross-sectional	Mixture	Occupational	Proxy	Blood	qPCR	bp	None	
Kahl, et al., 2016	Brazil	Exposed: self-reported to be tobacco farmers at their "family farms"; Unexposed: subjects living in the same area	124 (52%)	Cross-sectional	Mixture including Glyphosate and flumetralin	Occupational	Proxy	Blood	qPCR	kb/diploid genome	None	
Karimi, et al., 2020	Iran	Random sampling from the male population	300 (0%)	Cross-sectional	Organochlorine pesticides	Environmental	Blood	Blood	qPCR	T/S ratio	Age, education, BMI, smoking, and dietary patterns	
Ock, et al., 2020	USA	Sample from National Health and Nutrition Examination Survey (NHANES)	1724 (53%)	Cross-sectional	TCPY, DMP, DMTP, DMDTP, DEP, DETP, DEDTP	Environmental	Urine	Blood	qPCR	T/S ratio	Age, sex, race/ethnicity, education, pack-years of cigarette smoke, alcohol consumption, physical activity, body mass index, hypertension,	

											diabetes, cardiovascular diseases, cancer, C-reactive protein, and urinary creatinine
Saad-Hussein, et al., 2019	Egypt	Exposed: pesticide sprayers and researchers Non-exposed: same rural areas but not working in pesticide spraying and researcher not exposed	200 (15%)	Cross-sectional	Mixture compounds, mostly OPs (malathionchloropyrifos and dimethoate) and carbamates (carbofuran)	Occupational	Proxy	Blood	qPCR	T/S ratio	None
Shin, et al., 2010	Korea	A community-based health survey	84 (61%)	Cross-sectional	p,p'-DDE, p,p'-DDT, p,p'-DDD, oxychlorane, trans-nonachlor, heptachlor epoxide, β -Hexachlorocyclohexane, Hexachlorobenzene, Mirex	Environmental	Blood	Leukocytes	qPCR	T/S ratio	Age, sex, BMI, cigarette smoking and alcohol drinking

DMP: dimethylphosphate; DEP: diethylphosphate; DMTP: dimethylthiophosphate; DETP: diethylthiophosphate; DMDTP: dimethyldithiophosphate; DEDTP: diethyldithiophosphate; TCPY: 3,5,6-trichloro-2-pyridinol; MA: meta-analysis

Table 4 Sensitivity analysis on the relationship between pesticide exposure and telomere length.

Category	Studies (k)	Effects sizes (n)	Hedges' g	95% CI	% Var (level 2)	% Var (level 3)
Main analysis	15	96	-0.46	[-1.10, 0.19]	0.39	99.15
Outliers removed	8	82	-0.01	[-0.10, 0.09]	21.46	52.17
Influential studies removed	12	93	-0.28	[-0.54, -0.02]	3.10	93.42
Low-quality studies removed	12	93	-0.29	[-0.54, -0.01]	2.94	93.76
Effect sizes with transformation removed	11	77	-0.60	[-1.48, 0.27]	0.37	99.25
Effect sizes with no adjustment of confounders removed	5	33	-0.06	[-0.31, 0.19]	8.25	84.05

Table 5 Moderator analysis on the relationship between pesticide exposure and telomere length.

Category	Studies (k)	Effects sizes (n)	Hedges' g	95% CI	F (df1, df2)	P	% Var (level 2)	% Var (level 3)
Pesticide class					0.10 (1, 47)	0.10	2.49	92.3
Organophosphate	6	27	0.05	[-0.20, 0.3]				
Organochlorine	4	22	-0.02	[-0.28, 0.23]				
Gender					0.75 (1, 82)	0.39	1.97	96.03
Female	5	10	-0.25	[-0.75, 0.25]				
Male	7	74	-0.39	[-0.87, 0.08]				
Exposure setting					0.52 (1, 88)	0.47	0.32	99.30
Occupational	10	71	-0.64	[-1.47, 0.20]				
Environmental	4	19	-0.07	[-1.38, 1.25]				
TL expression					0.04 (2, 93)	0.96	0.34	99.28
T/S ration	9	90	-0.48	[-1.37, 0.42]				
bp	4	4	-0.53	[-1.88, 0.82]				
kb/genome	2	2	-0.23	[-2.14, 1.68]				

Figures




Fig. 1. Flowchart of search process




Fig. 2. Forest plot for meta-analysis of associations between pesticide exposure (ln-transformed) and Hannum epigenetic age




Fig. 3. Forest plot for meta-analysis of associations between pesticide exposure and telomere length