

TITLE PAGE

TITLE: Cortico-Spinal Intermittent Theta Burst Stimulation Propelling Sensorimotor Function Recovery in Complete Spinal Cord Injury: Randomized Control Trial Protocol

Authors: Deeksha Patel¹, Rohit Banerjee¹, Kamran Farooque², Deepak Gupta³, Bhavuk Garg², Nand Kumar⁴, KP Kocchar¹, Suman Jain^{1*}

Affiliation:

¹Department of Physiology, All India Institute of Medical Sciences, Delhi (110049), India

²Department of Orthopaedics, All India Institute of Medical Sciences, Delhi (110049), India

³Department of Neurosurgery, All India Institute of Medical Sciences, Delhi (110049), India

⁴Department of Psychiatry, All India Institute of Medical Sciences, Delhi (110049), India

***Corresponding author-** Suman Jain (Professor)

Email Addresses: sumanjain10@gmail.com (Suman Jain)

Abstract:

Background: Intermittent theta burst stimulation (iTBS) is a non-invasive stimulation technique to induce neuronal and synaptic plasticity. The induced cortical plasticity is imperative in the recovery of motor and sensory functions. Spinal cord injury (SCI) causes damage to neurons and results in sensorimotor dysfunction. The effect of iTBS on recovery of motor and sensory dysfunction in complete SCI (cSCI) is still elusive.

AIM: This study aims to assess the effect of iTBS on cortico-spinal tract integrity, plasticity, and regaining of motor and sensory function in cSCI patients. The rationale behind using an iTBS protocol is to modify and augment the communication between spared neurons of the cortico-spinal tract and strengthen the synaptic transmission, which will improve motor function in underlying muscles.

Method: A total of 48 patients will be recruited and randomly divided into placebo and real stimulation groups. iTBS along with a rehabilitation program will be administered to the placebo and real stimulation groups. Follow-up will be done at 1 month, 2 months, and 3 months after the intervention.

Result: The outcome of the study will be defined by electrophysiological parameters elicited by single and paired-pulse stimulation, ASIA score, pain, activities of daily life, quality of life, anxiety, depression, and biomarkers related to SCI. The results of this study will uncover the effectiveness of iTBS stimulation on (i) recovery of motor and sensory function in cSCI (ii) excitability of cortico-spinal tract (iii) neurological recovery and modulation of pain (iv) cortical reorganization after injury.

Conclusion: Intermittent theta-burst stimulation (iTBS) in conjunction with an individualized rehabilitation program may serve as an integrated strategy to rejuvenate locomotor abilities and improve the overall quality of life for people with complete spinal cord injuries (SCI).

Keywords: Spinal Cord Injury, Intermittent Theta Burst Stimulation, Excitatory-inhibitory circuitry, synaptic plasticity, Physical Rehabilitation, American Spinal Injury Association Impairment Scale.

Key Points:

1. iTBS is a novel neurostimulation technique aimed to restore sensorimotor function after a complete SCI
2. The primary objective of the trial is to evaluate the efficacy of iTBS in promoting sensorimotor function recovery
3. Assessment of the potential impact of iTBS on SCI rehabilitation
4. Understand the underlying mechanism of excitatory-inhibitory circuits associated with SCI
5. Unlock the importance of neuronal plasticity in regaining mobility

Introduction:

Spinal cord injury (SCI) is a neurological condition that causes progressive neurodegeneration. The major symptoms of SCI are paralysis, paraesthesia, pain, spasticity, bladder, bowel, and sexual dysfunction. It largely affects the young age population (21-49 years), thus the psychological impact on a healthy individual to adapt to a paraplegic or quadriplegic condition in their early life is devastating. SCI causes damage to neurons, nerves, and other surrounding cells that send and receive signals from different body parts to the brain via the spinal cord and vice versa(1). Spinal cord injury can be traumatic or non-traumatic. Traumatic spinal cord injury (SCI) is a great challenge for therapeutic management considering its high morbidity. There is complete or incomplete loss of locomotor, sensory, and autonomic functions depending on the severity and level of the lesion(2). The consequences of injury are not just a break in communication between neurons, but a cascade of events that sets up a vicious cycle and leads to widespread neuronal degeneration, cell death, and the formation of glial scars(3). Cellular components such as proteins, phospholipids, neurotransmitters, and metabolites derived from SC neurons and glial cells diffuse from the injury site into the cerebrospinal Fluid (CSF) and blood(4). These biomarkers could serve as good diagnostic markers to predict the severity of the injury. To regain functional connectivity, and attenuate gliosis and secondary injury, activity-dependent strategies have been proposed to be quite effective. The current treatment modalities for SCI include surgery, pain management, and rehabilitation. However, functional recovery by restoring connections of the corticospinal tract after a complete spinal cord injury is challenging.

High-frequency repetitive transcranial magnetic stimulation has decreased neuropathic pain and spasticity in incomplete spinal cord-injured patients(5)(6). A newer form of patterned TMS, intermittent theta burst stimulation (iTBS) is now being studied for the improvement of symptoms in incomplete spinal cord injury(7). There is very limited literature available, showing the effect of either cortical iTBS or trans-spinal TMS in combination with a rehabilitation program to promote repair, regeneration, and recovery in cSCI patients The present study aims to determine the functional outcomes of administering iTBS at the cortex as well as at spinal cord along with intensive rehabilitation program in cSCI patients.

Objectives:

1. To establish an effective site for intermittent theta burst stimulation (iTBS) in complete SCI patients: Motor cortex/ Spinal cord / Motor cortex + Spinal cord
2. To evaluate the efficacy of iTBS along with rehabilitation programs on –
 - a) Recovery of sensory and locomotor functions
 - b) Cortical excitability and plasticity
 - c) Biomarkers associated with spinal cord injury
 - d) Pain, anxiety, depression, and quality of life scores

Material & Methods:

Ethical Approval

Ethical approval has been obtained from the Institutional Ethical Committee of All India Institute of Medical Sciences, Delhi (Project Ref. Id: IECPG/551/7/2022). The study is registered in Clinical Trials Registry- India (CTRI) with reference number CTRI/2022/11/047038. Informed, voluntary, and written consent will be taken and the participants will be given a choice to withdraw from the study at any given point in time. They will be duly informed of the duration of the study along with any potential risks or side effects associated with TMS intervention.

All the experiments will be carried out in the Brain Stimulation and Neuromodulation Laboratory, Department of Physiology, and CARE, dept of Psychiatry, AIIMS, New Delhi.

Study Design

A double-blinded, prospective, randomized, placebo-controlled study will be conducted. The computer-generated sequence will be sealed in sequentially numbered envelopes, which will be opened after the patient is enrolled in the study. Both the patient and investigator will be blinded to the intervention. An experienced therapist will perform the intervention protocol. The study will follow the CONSORT rule (Figure 1).




Figure1. CONSORT diagram

ASIA: American Spinal Cord Injury Association; SCIM: Spinal Cord Injury Independence Measure; VAS: Visual Analogue Scale; STAI: State-trait Anxiety Inventory; QOL: Quality of Life; BDI: Beck depression inventory; MEP: Motor evoked Potential; cSP: Cortical silent period; SICI: Short interval intracortical inhibition; LICI: Long interval intracortical inhibition; IHI: Interhemispheric inhibition; ICF: Intracortical facilitation

Duration of study

The recruitment, intervention, and follow-up phase will be of 2 years, followed by 6 months of data analysis. The recruitment of patients will be started in September 2022 and the trial will be completed by March 2025.

Inclusion-Exclusion Criteria

Adults in the age group of 18 to 60 years with Thoracolumbar Spinal cord Injury having complete motor loss below the level of lesion with ASIA score A will be screened and recruited in the study.

Patients with osteoporotic Fracture, having a history of neurological or orthopedic disease related to the spinal cord, head injury, ferromagnetic metallic implants close to the target stimulation area, pacemaker, cognitive impairment, pregnancy, history of seizures and acute eczema/ bed shore will be excluded from the study.

Confidentiality

The personal privacy of patients will be protected. The results of this study will not disclose any identifying and personal information of the patient without his/her permission.

Experimental Groups:

A total of 48 patients will be randomized into 6 groups using computer-generated random numbers:

1. Group A: Control
2. Group A: Sham (iTBS with placebo coil on motor cortex)
3. Group B: High-frequency rTMS on motor cortex
4. Group C: iTBS on motor cortex
5. Group D: iTBS on the spinal cord
6. Group E: iTBS on the motor cortex and spinal cord

Intervention:

The rTMS/ iTBS will be applied using Neurosoft - Neuro-MS 5 (Neurosoft Ltd®, Ivanovo, Russia), a commercially available transcranial magnetic stimulator equipped with an angulated figure-eight-shaped coil and circular coil will be used for stimulation.

The intervention will be applied over the lower-limb motor area localized in M1 (to stimulate both lower limbs), with the handle of the coil parallel to the interhemispheric midline (pointing occipitally) based on the vertex position of the International 10-20 system of EEG(8,9). For spine stimulation, the circular coil will be placed over the site of injury at the spinal cord.

Standard protocol of iTBS will be used; 3pulse bursts at 50 Hz, repeated at 5 Hz, 2-s train repeated every 10 s for 20 repetitions, for a total of 600 pulses (7). 90% resting motor threshold will be used as the iTBS intensity. rTMS consists of 1600 pulses at 20Hz frequency. 2-s train of rTMS will be repeated every 30 sec for 20 minutes(10). A 90% resting motor threshold will be used as rTMS intensity.

2 weeks postoperatively, rTMS or iTBS will be administered in 2 sessions/day for 5 days, with a total of 10 sessions. The protocol and number of sessions for the sham stimulation group will be the same as that for the actual stimulation group, except the coil will be a placebo. (Figure 2)

Screening and Recruitment	Pre-intervention assessment (3 week)	Post-intervention assessment (4 week)	1 st follow-up (8 week)	2 nd follow-up (12week)	3 rd follow-up (16 week)
Surgical procedure (1 week)	rTMS/iTBS intervention (Twice a day at 2hr intervals for 5 days)				
Rehabilitation training					
	Neurological & Functional parameter	Neurological & Functional parameter	Neurological & Functional parameter	Neurological & Functional parameter	Neurological & Functional parameter
	Electrophysiological & Psychosocial parameter	Electrophysiological & Psychosocial parameter	Electrophysiological & Psychosocial parameter	Electrophysiological & Psychosocial parameter	Electrophysiological & Psychosocial parameter
	Blood sample	Blood sample			Blood sample

Figure 2: Timeline-All the intervention and recording paradigms are organized as a timeline

Methodology:

1. Patients will be recruited from J P N A Trauma Centre, AIIMS Delhi after the vertebral injury has been stabilized. At the first visit, the baseline parameters of the patient will be assessed.
2. During an intervention, the patient will be allocated to one of the interventional groups as per randomization. Motor cortex stimulation will be followed by spinal cord stimulation for 10 sessions twice a day for 5 consecutive days.
3. For cortical TMS intervention, firstly the Resting motor threshold (RMT) will be determined using MEP recorded from Abductor Pollicis Brevis (APB). The figure of eight coils will be used over motor hotspots of APB. The cortical stimulation intensity used to generate MEP in target muscle with amplitude greater than 50 μ V in 5 out of 10 consecutive trials will be considered as RMT. The stimulation intensity for iTBS or rTMS will be kept at 90% of RMT and the coil will be placed at the leg area of the M1 area. In the Sham stimulation group, the protocol will remain the same, but the stimulation coil will be placebo.
4. For spinal TMS intervention, the patient will be lying in a prone position. The figure of eight coil will be held parallel over the vertebral column corresponding to an injured area of the spinal cord. The stimulation protocol will remain similar to cortical stimulation.
5. A trained physiotherapist will be assigned to provide personalized rehabilitation training for 5 days (Figure 3). Thereafter the patient will be guided to perform all the exercises at home and maintain a logbook, share their videos of performance weekly for assessment by the physiotherapist.
6. Blood samples will be collected before intervention, immediately after intervention, and after 3 months of intervention.
- 7.

Customized Exercise protocol for complete SCI patients*		
Upper Limb	Lower Limb	General
<p>Strengthening Exercises: Progressive Resisted Exercises. (using manual resistance, dumbbell, theraband etc.) # Progression based on De-Lorme & Watkins principle at 50-80% RM (3 sets of 20 reps.)</p>	<p>Mobilization exercises- AROM, PROM (10 reps., 2-3 sets) Stretching exercises (15 sec hold, 5 reps, 2-3 sets) Strengthening Exercises (10 reps., 3 sets) Bed mobility exercises e.g. rolling, kneeling; supine to prone, supine to sit and vice versa (10 reps., 3 sets) Physiological standing with orthotic support as per tolerance.</p>	<p>Reaching Exercises (Vector reach exercises). Balance exercises (using gym ball, wobble board) Co-ordination exercises (Peg board ex, Biofeedback) Bladder rehabilitation (5 reps., 10sec hold, 2-3 sets) Sensory re-education (brushing) 15 min, 2 sets/ day</p>

8. **Figure 3.** Outline of physical rehabilitation regime for spinal cord injury patients.

9. *Exercise prescription and progression will be based on the individual's performance and tolerance to baseline exercises.

Post-intervention care

If a patient experiences an adverse event during the trial, the principal investigator will provide treatment and corresponding financial compensation. Patients who are enrolled in the placebo will also receive the conventional rehabilitation program during the study.

Results:

The outcome will be measured at 5-time points: baseline, immediate post-intervention, 1st month, 2nd month and 3rd month of follow-ups. All the assessments will be recorded by the blinded primary investigator to maintain the discrepancy during the intervention (Table 1).

- 1) **Neurological:** The primary outcome will measure the severity of injury with the American Spinal Injury Association (ASIA) Scoring System. The ASIA impairment score (AIS) ranges from complete loss of sensation and movement (AIS = A) to normal neurological function (AIS = E). The ASIA motor score uses a test of the strength of ten key muscles on each side of the body (e.g. elbow flexors, wrist extensors, hip flexors, quadriceps, dorsi flexors). The score ranges from 0 (no contraction) to 5 (normal resistance) through a full range of motion. A total possible score of 50 for the upper extremities (UE) and 50 for the lower extremities (LE) may be obtained (11). The ASIA sensory score involves pinprick and light touch sensation at key points representing each dermatome of the body, scored on a three-point scale (0, 1, and 2). Scores are summed to give a total possible score of 224, where a higher score indicates better sensation than a lower score.
- 2) **Functional:**
 - a) Walking Index for Spinal Cord Injury II (WISCI-II) is a 0–20 level scale, that evaluates the walking activity of a patient based on physical assistance, the need for braces/walker, and other adaptive devices. The levels on the scale are scored from 0 (patient unable to walk) to 20 (patient walking without braces and/or adaptive devices and without any physical assistance for at least 10 m)(12).
 - b) Spinal Cord Independence Measure-III (SCIM-III) includes 19 tasks organized in 3 subclasses based on the patient's general activity: self-care (score 0-20), respiration and sphincter management (score 0-40), and mobility (score 0-40). The overall scores range from 0-100, where a 0 score defines a total dependence of the patient on the caregiver and a score of 100 indicates the complete independence(13).
- 3) **Electrophysiological:**
 - a) **Single-pulse TMS:** A single pulse of TMS at minimum stimulus intensity will be delivered at the motor cortex that elicits a motor-evoked potential (MEP) of $\geq 50\mu V$ at least 5 out of 10 trials in the target muscle and will be recorded as RMT(14). For recording the inhibitory activity, a single TMS is delivered to produce an interruption in ongoing EMG activity during

a tonic contraction followed by the reoccurring EMG activity. The duration of silencing of EMG in response to TMS will be measured as cSP (15). The recruitment curve plotted between the TMS intensity (%) and motor-evoked potential (MEP) sizes.

- b) **Paired-pulse TMS:** Two TMS pulses, conditioning stimulus (CS) and test stimulus (TS) delivered at specific interstimulus intervals result in either facilitation or inhibition. In SICI, MEP elicited by TS is inhibited when preceded by a CS at ~1-5ms intervals. In LICI, the interstimulus interval is kept between 50-200ms, whereas, in ICF, the interval is ~10-30ms (15).

4) **Psychosocial:**

- a) Visual Analogue scale (VAS) measures the perceived intensity of pain on a self-explained scale of 0 to 10, where 0 (no pain) to 10 (severe pain)(16).
- b) Beck depression inventory II is a 21-question multiple-choice self-report inventory to measure the presence and severity of depression. The cut-scores with 0-9 indicating normal, 10-19 indicating mild depression, 20-30 indicating moderate depression, and 31-63 indicating severe depression(17).
- c) State-Trait Anxiety Inventory is a 40-item self-report scale, commonly used to measure anxiety (18).
- d) The WHO-Quality of Life-BREF Scale is developed in the context of the four domains defining the QOL: physical, psychological, social, and environmental. The higher the QOL score the higher the life satisfaction(19).

- 5) **Biochemical Quantification:** The disruption of blood spinal cord barrier (BSCB) and secondary damage following the injury releases several chemokines, growth factors, and neurotransmitters in CSF and systemic circulation which could help predict the severity of injury. The released components such as Myelin Basic Protein (MBP), Interleukins, Phosphorylated neurofilaments, BDNF, Fas-ligand, GABA, and Glutamate may serve as potential biomarkers for remyelination, inflammation, neuronal survival, and excitatory or inhibitory neurotransmitters after SCI (20)(21)(22)(23). Mass spectroscopy and enzyme linked immunosorbent assays will be performed to quantify these biomarkers in plasma and serum samples.

ASSESSMENT	<i>t2</i>	<i>t1</i>	<i>0</i>	<i>T1</i>	<i>T2</i>	<i>T3</i>	<i>T4</i>
<i>Enrolment</i>	*						

<i>Consent</i>	*					
<i>Screening</i>	*					
<i>Demographic</i>	*					
<i>Details</i>						
<i>Vitals</i>	*	*		*	*	*
<i>Medical History</i>	*					
<i>Inclusion</i>	*					
<i>Exclusion</i>	*					
<i>Randomisation</i>		*				
<i>Blinding</i>		*				
<i>Blood sample collection</i>	*			*		*
<i>ASIA-motor</i>		*		*	*	*
<i>ASIA-sensory</i>		*		*	*	*
<i>WSCI</i>		*		*	*	*
<i>MAS</i>		*		*	*	*
<i>SCIM</i>		*		*	*	*
<i>VAS</i>		*		*	*	*
<i>WHOQOL</i>		*		*	*	*
<i>BDI-II</i>		*		*	*	*
<i>STAI</i>		*		*	*	*
<i>TMS pre-therapy questionnaire</i>		*		*	*	*
<i>INTERVENTIONS-</i>			*			
<i>SHAM- rTMS</i>			—————			
<i>SHAM- ITBS</i>			—————			
<i>REAL- rTMS</i>			—————			
<i>REAL- ITBS</i>			—————			
<i>RMT</i>	*	*	*	*	*	*
<i>MEP</i>	*		*	*	*	*
<i>cSP</i>	*		*	*	*	*
<i>SICI</i>	*		*	*	*	*
<i>LICI</i>	*		*	*	*	*

<i>ICF</i>	*	*	*	*	*
<i>UNBLINDING</i>					*
<i>TMS post-therapy questionnaire</i>		*	*	*	*
<i>Rehabilitation Programme</i>					*

Table 1. Protocol for intervention trial and outcomes measure will be recorded at different time intervals.

-t2: Screening before surgery; -t1: Parameters assessment day before intervention; 0: intervention start for consecutive 5 days; T1: Parameters assessed after completion of intervention; T2: follow-up at 1st month; T3: follow-up at 2nd month; T4: follow-up at 3rd month.

Statistical Analysis:

Data will be presented as Mean and standard deviation (SD). The intention-to-treat (ITT) analysis will be the primary analysis. Baseline observation will be carried forward for subjects with missing values. Unless otherwise stated, ITT data will be presented throughout. Per-protocol analysis (all compliant participants) will also be conducted. To see the difference in groups ANOVA analysis followed by multiple comparison tests (Post hoc analysis) will be used. To see the changes at different time points from the baseline, paired t-test or repeated measure ANOVA will be applied. To establish an association between groups and qualitative variables, the Chi-square/fisher exact test will be used. The value of p less than .05 will be considered statistically significant. P value < 0.05 at (95% CI) will be considered statistically significant. Statistical analysis will be done using STATA 14.0 (STATA Corp, Houston, TX, USA) software.

Discussion:

SCI significantly impairs the cortico-spinal integrity and afferent-efferent input/output circuitry. Theoretically, the pyramidal tract is the primary neural pathway that links the cortex and spinal cord to facilitate the movement of distal extremities. The primary purpose of any treatment is to reconstruct the neural circuit immediately after SCI for functional sensory-motor recovery. To facilitate neural circuit reconstruction, it is required to stimulate nerve cell sprouting and regeneration as well as increase the strength of the existing neuronal connections. Previous research demonstrated that individuals with incomplete SCI can benefit from locomotor training to enhance their motor skills. Rehabilitation programs may use learning and relearning mechanisms, uncovering a previously inactive synapse, and forming a new synapse(24). However, the reconstruction of the damaged

neural circuit is quite difficult with locomotor training alone. Studies have shown that the functional effects of exercise along with transcranial magnetic stimulation, can activate spared neural pathways and enhance the possibility of neural reconstruction(25,26).

Transcranial magnetic stimulation (TMS) induces electrical currents in underlying cortical areas, depolarizes neurons, and generates an action potential that modulates the activity of spinal motor neurons and target muscles via the corticospinal tract. Repetitive transcranial magnetic stimulation as well as iTBS is an intervention in various psychiatric and pain conditions. In SCI few studies suggest improvement in locomotor function, spasticity, and pain in incomplete patients(6). Although, the functional mechanism of rTMS or iTBS on sensorimotor recovery in SCI patients is not fully understood, but thought to induce synaptic plasticity via LTP/LTD-like effects (27), thereby can promote functional recovery in SCI patients.

Additionally, spinal cord stimulation can modulate the activity of the local central pattern motor generators, which promote synaptic strengthening (28). In rat models of complete and incomplete SCI a significant attenuation of glial scarring, lesion volume, neurotransmitter imbalance, muscle atrophy, and facilitation of neuronal survival, axonal regeneration, and myogenesis has been shown following whole-body magnetic field exposure(29,30). Therefore, we propose that spinal cord stimulation along with motor cortical stimulation would attenuate secondary damage and promote regeneration even in complete SCI patients, leading to long-term functional recovery.

Various single and paired-pulse paradigms of TMS have been used for the assessment of cortical excitability, plasticity, integrity of the corticospinal tract, and excitatory-inhibitory (E-I) neural circuitry of the motor cortex. We shall use all paired-pulse paradigms (SICI, ICF, and LICI) to objectively assess E-I neural circuitry and single-pulse paradigms (RMT, MEP, recruitment curves, and cSP) for unraveling the excitability of corticospinal circuitry and motor units.

Conclusion:

An intensive individualized rehabilitation regime coupled with iTBS could be a holistic management strategy that can restore locomotor function and quality of life in complete SCI patients. TMS is also a promising tool to evaluate the cortical plasticity and excitability in SCI and thereby understand the mechanism of action of the proposed intervention. characterize the effective connectivity of neural circuits and mechanisms regulating the balance between inhibition and facilitation within the corticospinal pathway.

Conflict of interest disclosure: The authors declare no conflict of interest.

Funding Source: This work was supported by the Department of Biotechnology (DBT) Ministry of Science and Technology, Government of India in 2019 (BT_2052).

Acknowledgment: We want to acknowledge the Department of Biotechnology (DBT) for providing us with funding. We also want to acknowledge Mr. Sanjeev and Mr. Praveen for this technical support during standardization.

References:

1. Medicine I of, Health B on N and B, Injury C on SC. *Spinal Cord Injury: Progress, Promise, and Priorities*. National Academies Press; 2005. 361 p.
2. Alizadeh A, Dyck SM, Karimi-Abdolrezaee S. Traumatic Spinal Cord Injury: An Overview of Pathophysiology, Models and Acute Injury Mechanisms. *Front Neurol*. 2019 Mar 22;10:282.
3. Beattie MS, Farooqui AA, Bresnahan JC. Review of Current Evidence for Apoptosis After Spinal Cord Injury. *J Neurotrauma*. 2000 Oct;17(10):915–25.
4. Ahadi R, Khodagholi F, Daneshi A, Vafaei A, Mafi AA, Jorjani M. Diagnostic Value of Serum Levels of GFAP, pNF-H, and NSE Compared With Clinical Findings in Severity Assessment of Human Traumatic Spinal Cord Injury. *Spine*. 2015 Jul 15;40(14):E823-830.
5. Yılmaz B, Kesikburun S, Yaşar E, Tan AK. The effect of repetitive transcranial magnetic stimulation on refractory neuropathic pain in spinal cord injury. *J Spinal Cord Med*. 2014 Jul 1;37(4):397–400.
6. Benito J, Kumru H, Murillo N, Costa U, Medina J, Tormos J, et al. Motor and Gait Improvement in Patients With Incomplete Spinal Cord Injury Induced by High-Frequency Repetitive Transcranial Magnetic Stimulation. *Top Spinal Cord Inj Rehabil*. 2012 Apr 14;18(2):106–12.
7. Nardone R, Langthaler PB, Orioli A, Höller P, Höller Y, Frey VN, et al. Effects of intermittent theta burst stimulation on spasticity after spinal cord injury. *Restor Neurol Neurosci*. 2017 Jan 1;35(3):287–94.
8. Jetté F, Côté I, Meziane HB, Mercier C. Effect of Single-Session Repetitive Transcranial Magnetic Stimulation Applied Over the Hand Versus Leg Motor Area on Pain After Spinal Cord Injury. *Neurorehabil Neural Repair*. 2013 Sep 1;27(7):636–43.
9. Khedr EM, Farweez HM, Islam H. Therapeutic effect of repetitive transcranial magnetic stimulation on motor function in Parkinson's disease patients. *Eur J Neurol*. 2003;10(5):567–72.
10. Kumru H, Murillo N, Samsó JV, Valls-Sole J, Edwards D, Pelayo R, et al. Reduction of Spasticity With Repetitive Transcranial Magnetic Stimulation in Patients With Spinal Cord Injury. *Neurorehabil Neural Repair*. 2010 Jun;24(5):435–41.
11. Kirshblum SC, Waring W, Biering-Sorensen F, Burns SP, Johansen M, Schmidt-Read M, et al. Reference for the 2011 revision of the international standards for neurological classification of spinal cord injury. *J Spinal Cord Med*. 2011 Nov 1;34(6):547–54.
12. Ditunno JF, Ditunno PL, Graziani V, Scivoletto G, Bernardi M, Castellano V, et al. Walking index for spinal cord injury (WISCI): an international multicenter validity and reliability study. *Spinal Cord*. 2000 Apr;38(4):234–43.
13. Itzkovich M, Shefler H, Front L, Gur-Pollack R, Elkayam K, Bluvshstein V, et al. SCIM III (Spinal Cord Independence Measure version III): reliability of assessment by interview and comparison with assessment by observation. *Spinal Cord*. 2018 Jan;56(1):46–51.

14. Lefaucheur JP, André-Obadia N, Antal A, Ayache SS, Baeken C, Benninger DH, et al. Evidence-based guidelines on the therapeutic use of repetitive transcranial magnetic stimulation (rTMS). *Clin Neurophysiol*. 2014 Nov 1;125(11):2150–206.
15. Arora T, Desai N, Kirshblum S, Chen R. Utility of transcranial magnetic stimulation in the assessment of spinal cord injury: Current status and future directions. *Front Rehabil Sci*. 2022 Oct 5;3:1005111.
16. Jensen MP, Karoly P, Braver S. The measurement of clinical pain intensity: a comparison of six methods. *Pain*. 1986 Oct;27(1):117–26.
17. Kendall PC, Hollon SD, Beck AT, Hammen CL, Ingram RE. Issues and recommendations regarding use of the Beck Depression Inventory. *Cogn Ther Res*. 1987 Jun 1;11(3):289–99.
18. Greene J, Cohen D, Siskowski C, Toyinbo P. The Relationship Between Family Caregiving and the Mental Health of Emerging Young Adult Caregivers. *J Behav Health Serv Res*. 2017 Oct 1;44(4):551–63.
19. Skevington SM, Lotfy M, O’Connell KA. The World Health Organization’s WHOQOL-BREF quality of life assessment: Psychometric properties and results of the international field trial. A Report from the WHOQOL Group. *Qual Life Res*. 2004 Mar 1;13(2):299–310.
20. Albayar AA, Roche A, Swiatkowski P, Antar S, Ouda N, Emara E, et al. Biomarkers in Spinal Cord Injury: Prognostic Insights and Future Potentials. *Front Neurol*. 2019 Jan 29;10:27.
21. Biglari B, Büchler A, Swing T, Child C, Biehl E, Reitzel T, et al. Serum sCD95L concentration in patients with spinal cord injury. *J Int Med Res*. 2015 Apr;43(2):250–6.
22. Löscher W, Fassbender CP, Gram L, Gramer M, Hörstermann D, Zahner B, et al. Determination of GABA and vigabatrin in human plasma by a rapid and simple HPLC method: correlation between clinical response to vigabatrin and increase in plasma GABA. *Epilepsy Res*. 1993 Mar;14(3):245–55.
23. Tanhoffer RA, Yamazaki RK, Nunes EA, Pchevozniki AI, Pchevozniki AM, Nogata C, et al. Glutamine Concentration and Immune Response of Spinal Cord-Injured Rats. *J Spinal Cord Med*. 2007 Jan 1;30(2):140–6.
24. Hubli M, Dietz V. The physiological basis of neurorehabilitation--locomotor training after spinal cord injury. *J Neuroengineering Rehabil*. 2013 Jan 21;10:5.
25. Pulverenti TS, Zaaya M, Grabowski E, Grabowski M, Knikou M. Brain and spinal cord paired stimulation coupled with locomotor training facilitates motor output in human spinal cord injury. *Front Neurol*. 2022;13:1000940.
26. Jo HJ, Richardson MSA, Oudega M, Perez MA. Paired corticospinal-motoneuronal stimulation and exercise after spinal cord injury. *J Spinal Cord Med*. 2021;44(sup1):S23–7.
27. Klomjai W, Katz R, Lackmy-Vallée A. Basic principles of transcranial magnetic stimulation (TMS) and repetitive TMS (rTMS). *Ann Phys Rehabil Med*. 2015 Sep 1;58(4):208–13.
28. Korzhova J, Sinityn D, Chervyakov A, Poydasheva A, Zakharova M, Suponeva N, et al. Transcranial and spinal cord magnetic stimulation in treatment of spasticity: a literature review and meta-analysis. *Eur J Phys Rehabil Med*. 2018 Feb;54(1):75–84.
29. Bhattacharyya S, Dinda A, Vishnubhatla S, Anwar MF, Jain S. A combinatorial approach to modulate microenvironment toward regeneration and repair after spinal cord injury in rats. *Neurosci Lett*. 2021 Jan 10;741:135500.
30. Chakraborty A, Sharma MC, Vishnubhatla S, Jain S. Electromagnetic field stimulation facilitates motor neuron excitability, myogenesis and muscle contractility in spinal cord transected rats. *J Biosci*. 2022;47:78.