

Meta-analysis of Genome wide Association Studies on Childhood ADHD Symptoms and Diagnosis Reveals 17 Novel Loci and 22 Potential Effector Genes

7 Van der Laan, Camiel M.^{1,2}, Ip, Hill F.¹, Schipper, Marijn³, Hottenga, Jouke-Jan¹, Krapohl, Eva
8 M. L.⁴, Brikell, Isabell⁵, Soler Artigas, María^{6,7,8}, Cabana-Domínguez, Judith^{6,7,8}, Llonga
9 Natalia^{6,7,8}, Nolte, Ilja M.⁹, St Pourcain, Beate^{10,11,12}, Bolhuis, Koen¹³, Palviainen, Teemu¹⁴,
10 Zafarmand, Hadi^{15,16}, Gordon, Scott¹⁷, Zayats, Tetyana^{18,19,20}, Aliev, Fazil²¹, Burt, Alexandra S.²²,
11 Wang, Carol A.^{23,24}, Saunders, Gretchen²⁵, Karhunen, Ville^{26,27}, Adkins, Daniel E.^{28,29}, Border,
12 Richard³⁰, Peterson, Roseann E.^{31,32}, Prinz, Joseph A.³³, Thiering, Elisabeth^{34,35}, Vilor-Tejedor,
13 Natàlia³⁶⁻³⁹, Ahluwalia, Tarunveer S.^{40,41}, Allegrini, Andrea⁴, Rimfeld, Kaili⁴, Chen, Qi⁵, Lu, Yi⁵,
14 Martin, Joanna⁵, Bosch, Rosa^{42,6}, Ramos Quiroga, Josep Antoni^{6,43,7,8}, Neumann, Alexander¹³,
15 Ensink, Judith^{44,45,46,47}, Grasby, Katrina¹⁷, Morosoli, José J.^{48,49,50}, Tong, Xiaoran⁵¹, Marrington,
16 Shelby⁵², Scott, James G.^{53,54,55}, Shabalin, Andrey A.²⁹, Corley, Robin⁵⁶, Evans, Luke M.^{56,57},
17 Sugden, Karen^{58,32}, Alemany, Silvia^{6,7,8}, Sass, Lærke³⁹, Vinding, Rebecca³⁹, Ehli, Erik A.⁵⁹,
18 Hagenbeek, Fiona A.^{1,14,45}, Derks, Eske⁶⁰, Larsson, Henrik^{5,61}, Snieder, Harold⁹, Cecil,
19 Charlotte^{13,62,63}, Whipp, Alyce M.¹⁴, Korhonen, Tellervo¹⁴, Vuoksima, Eero¹⁴, Rose, Richard J.⁶⁴,
20 Uitterlinden, André G.^{63,65,66}, Haavik, Jan¹⁸, Harris, Jennifer R.⁶⁷, Helgeland, Øyvind⁶⁸, Johansson,
21 Stefan⁶⁹, Knudsen, Gun Peggy S.⁶⁷, Njolstad, Pal Rasmus⁶⁹, Lu, Qing⁷⁰, Rodriguez, Alina²⁶,
22 Henders, Anjali K.⁷¹, Mamun, Abdullah⁷², Najman, Jakob M.⁵², Brown, Sandy⁷³, Hopfer,
23 Christian⁷⁴, Krauter, Kenneth⁷⁵, Reynolds, Chandra^{76,57}, Smolen, Andrew⁵⁷, Stallings, Michael^{57,77},
24 Wadsworth, Sally⁵⁷, Wall, Tamara⁷³, Eaves, Lindon^{78,32}, Silberg, Judy L.^{78,32}, Miller, Allison⁷⁹,
25 Havdahl, Alexandra^{80,81}, Llop, Sabrina^{82,83}, Lopez-Espinosa, Maria-Jose^{82,83,84}, Bønnelykke,
26 Klaus⁴⁰, Sunyer, Jordi^{85,86,87}, Arseneault, Louise⁸⁸, Standl, Marie³⁴, Heinrich, Joachim^{34,89,90},
27 Boden, Joseph⁹¹, Pearson, John⁹², Horwood, John⁹¹, Kennedy, Martin⁹³, Poulton, Richie⁹⁴, Maes,
28 Hermine H.^{78,32,95}, Hewitt, John^{57,59}, Copeland, William E.⁹⁶, Middeldorp, Christel
29 M.^{44,45,46,50,51,97,47}, Williams, Gail M.⁵², Wray, Naomi^{71,98}, Järvelin, Marjo-Riitta^{26,27,99}, McGue,
30 Matt²⁵, Iacono, William²⁵, Caspi, Avshalom^{59,100,88,101}, Moffitt, Terrie E.^{59,100,88,101}, Whitehouse,
31 Andrew¹⁰², Pennell, Craig E.^{23,24}, Klump, Kelly L.²², Jiang, Chang⁷⁰, Dick, Danielle M.^{103,104,78,105},
32 Reichborn-Kjennerud, Ted^{77,106}, Martin, Nicholas G.¹⁷, Medland, Sarah E.^{17,50,107}, Vrijkotte,
33 Tanja^{108,46,109}, Kaprio, Jaakko^{110,14}, Tiemeier, Henning^{13,111}, Davey Smith, George^{112,10}, Hartman,
34 Catharina A.¹¹³, Oldehinkel, Albertine J.¹¹⁴, Casas, Miquel^{142,43,114,115}, Ribasés, Marta^{6,7,8},
35 Lichtenstein, Paul⁵, Lundström, Sebastian^{116,117}, Plomin, Robert⁴, Bartels, Meike^{1,46}, Nivard,
36 Michel G.¹, Boomsma, Dorret I.^{1,46}

GWAMA OF ADHD SYMPTOMS AND DIAGNOSIS

- 37
38 ¹ Department of Biological Psychology, Vrije Universiteit Amsterdam, Amsterdam, The
39 Netherlands
40 ² Netherlands Institute for the Study of Crime and Law Enforcement
41 ³ Department of Complex Trait Genetics, Center for Neurogenomics and Cognitive Research,
42 Amsterdam Neuroscience, Vrije Universiteit Amsterdam, Amsterdam, The Netherlands
43 ⁴ Institute of Psychiatry, Psychology and Neuroscience, King's College London, United Kingdom
44 ⁵ Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden
45 ⁶ Biomedical Network Research Centre on Mental Health (CIBERSAM), Instituto de Salud Carlos
46 III, Barcelona, Spain
47 ⁷ Department of Mental Health, Hospital Universitari Vall d'Hebron, Barcelona, Spain.
48 ⁸ Psychiatric Genetics Unit, Group of Psychiatry, Mental Health and Addiction, Vall d'Hebron
49 Research Institute (VHIR), Universitat Autònoma de Barcelona, Barcelona, Spain
50 ⁹ Department of Epidemiology, University of Groningen, University Medical Center Groningen,
51 Groningen, The Netherlands
52 ¹⁰ MRC Integrative Epidemiology Unit, University of Bristol, Bristol, UK
53 ¹¹ Max Planck Institute for Psycholinguistics, The Netherlands
54 ¹² Donders Institute for Brain, Cognition and Behaviour, Radboud University, The Netherlands
55 ¹³ Department of Child and Adolescent Psychiatry/Psychology, Erasmus University Medical Center,
56 Rotterdam, Netherlands
57 ¹⁴ Institute for Molecular Medicine FIMM, HiLife, University of Helsinki, Helsinki, Finland
58 ¹⁵ Department of Public Health, Amsterdam Public Health Research Institute, Amsterdam UMC,
59 location Academic Medical Center, University of Amsterdam, Amsterdam, The Netherlands
60 ¹⁶ Department of Clinical Epidemiology, Biostatistics and Bioinformatics, Amsterdam Public Health
61 Research Institute, Amsterdam UMC, location Academic Medical Center, University of
62 Amsterdam, Amsterdam, The Netherlands
63 ¹⁷ QIMR Berghofer Medical Research Institute, Brisbane, Australia
64 ¹⁸ K.G. Jebsen Centre for Neuropsychiatric Disorders, Department of Biomedicine, University of
65 Bergen, Norway
66 ¹⁹ Analytic and Translational Genetics Unit, Department of Medicine, Massachusetts General
67 Hospital and Harvard Medical School, Boston, Massachusetts, USA
68 ²⁰ Stanley Center for Psychiatric Research, Broad Institute of MIT and Harvard, Cambridge,
69 Massachusetts, USA
70 ²¹ Department of Psychiatry, Robert Wood Johnson Medical School, Rutgers University, U.S.A
71 ²² Department of Psychology, Michigan State University, East Lansing, USA
72 ²³ School of Medicine and Public Health. Faculty of Medicine and Health, The University of
73 Newcastle, Newcastle, New South Wales, Australia
74 ²⁴ Hunter Medical Research Institute, Newcastle, New South Wales, Australia
75 ²⁵ Department of Psychology, University of Minnesota, USA
76 ²⁶ MRC Centre for Environment and Health, Department of Epidemiology and Biostatistics, School
77 of Public Health, Imperial College London, London, United Kingdom
78 ²⁷ Faculty of Medicine, University of Oulu, PO Box 8000, FI-90014 Oulun yliopisto, Finland
79 ²⁸ Department of Sociology, College of Social and Behavioral Science, University of Utah
80 ²⁹ Department of Psychiatry, School of Medicine, University of Utah

GWAMA OF ADHD SYMPTOMS AND DIAGNOSIS

- 81 ³⁰ Departments of Neurology and Computer Science, University of California Los Angeles,
82 California, USA
- 83 ³¹ Department of Psychiatry and Behavioral Sciences, State University of New York Downstate
84 Health Sciences University, Brooklyn, NY, USA
- 85 ³² Department of Psychiatry, Virginia Institute for Psychiatric and Behavioral Genetics, Virginia
86 Commonwealth University, Richmond, VA, USA
- 87 ³³ Center for Genomic and Computational Biology, Duke University, Durham, NC, USA
- 88 ³⁴ Institute of Epidemiology, Helmholtz Zentrum München - German Research Center for
89 Environmental Health, Neuherberg, Germany
- 90 ³⁵ Ludwig-Maximilians-University of Munich, Dr. von Hauner Children's Hospital, Division of
91 Metabolic Diseases and Nutritional Medicine, Munich, Germany
- 92 ³⁶ BarcelonaBeta Brain Research Center, Pasqual Maragall Foundation (FPM), Barcelona, Spain
- 93 ³⁷ Centre for Genomic Regulation (CRG), The Barcelona Institute of Science and Technology,
94 Barcelona, Spain
- 95 ³⁸ IMIM (Hospital del Mar Medical Research Institute), Barcelona, Spain
- 96 ³⁹ Department of Clinical Genetics. Erasmus University Medical Center. Rotterdam, Netherlands
- 97 ⁴⁰ COPSAC, Copenhagen Prospective Studies on Asthma in Childhood, Herlev and Gentofte
98 Hospital, University of Copenhagen, Copenhagen, Denmark
- 99 ⁴¹ Steno Diabetes Center Copenhagen, Herlev, Denmark
- 100 ⁴² SJD MIND Schools Program, Hospital Sant Joan de Déu, Institut de Recerca Sant Joan de Déu,
101 Esplugues de Llobregat, Spain
- 102 ⁴³ Department of Psychiatry and Legal Medicine, Universitat Autònoma de Barcelona, Barcelona,
103 Spain
- 104 ⁴⁴ Department of child and adolescent psychiatry and psychology, Amsterdam UMC, Amsterdam
- 105 ⁴⁵ Reproduction and Development research Institute, Amsterdam, the Netherlands
- 106 ⁴⁶ Amsterdam Public Health Research Institute, Amsterdam, the Netherlands
- 107 ⁴⁷ Level, Academic center for Child and Adolescent Psychiatry, Amsterdam, The Netherlands
- 108 ⁴⁸ Department of Clinical, Educational, and Health Psychology, University College London,
109 London, UK
- 110 ⁴⁹ Mental Health and Neuroscience Research Program, QIMR Berghofer Medical Research Institute,
111 Brisbane, Australia
- 112 ⁵⁰ School of Psychology, University of Queensland, Brisbane, Australia
- 113 ⁵¹ Center for Innovation in Public Health, University of Kentucky
- 114 ⁵² School of Public Health, The University of Queensland, Herston, Qld, Australia
- 115 ⁵³ Child Health Research Centre, The University of Queensland, Brisbane, QLD, Australia
- 116 ⁵⁴ Child and Youth Mental Health Service, Children's Health Queensland Hospital and Health
117 Service, Brisbane, QLD, Australia
- 118 ⁵⁵ Queensland Centre for Mental Health Research, QLD, Australia
- 119 ⁵⁶ Institute for Behavioral Genetics, University of Colorado Boulder, Colorado, USA
- 120 ⁵⁷ Department of Ecology and Evolutionary Biology, University of Colorado Boulder, Colorado,
121 USA
- 122 ⁵⁸ Department of Psychology and Neuroscience, Duke University, Durham, N.C., U.S.A.
- 123 ⁵⁹ Avera Institute for Human Genetics, Sioux Falls, South Dakota, USA

GWAMA OF ADHD SYMPTOMS AND DIAGNOSIS

- 124 60 Translational Neurogenomics Laboratory, QIMR Berghofer Medical Research Institute, Brisbane,
125 Queensland, Australia
- 126 61 School of Medical Sciences, Orebro University, Orebro, Sweden
- 127 62 Department of Biomedical Data Sciences, Molecular Epidemiology, Leiden University Medical
128 Center, Leiden, The Netherlands
- 129 63 Department of Epidemiology, Erasmus University Medical Center, Rotterdam, The Netherlands
- 130 64 Department of Psychological and Brain Sciences, Indiana University, Bloomington, IN, USA
- 131 65 Department of Internal Medicine, Erasmus University Medical Center, Rotterdam, The
132 Netherlands
- 133 66 Netherlands Genomics Initiative (NGI)-sponsored Netherlands Consortium for Healthy Aging
134 (NCHA), Leiden, The Netherlands
- 135 67 Centre for Fertility and Health, The Norwegian Institute of Public Health, Oslo, Norway
- 136 68 Department of Genetics and Bioinformatics, Division of Health Data and Digitalization,
137 The Norwegian Institute of Public Health, Bergen, Norway
- 138 69 Department of Clinical Science, University of Bergen, Norway
- 139 70 Department of Biostatistics, University of Florida, Gainesville, USA
- 140 71 Institute for Molecular Bioscience, University of Queensland, QLD, Australia
- 141 72 Institute for Social Science Research, University of Queensland, QLD, Australia
- 142 73 Department of Psychiatry, University of California, San Diego
- 143 74 University of Colorado School of Medicine, Aurora, Colorado, USA
- 144 75 Department of Molecular, Cellular, and Developmental Biology, University of Colorado Boulder,
145 USA
- 146 76 Department of Psychology, University of California Riverside, Riverside, CA, USA
- 147 77 Department of Psychology and Neuroscience, University of Colorado Boulder, Colorado, USA
- 148 78 Department of Human and Molecular Genetics, Virginia Commonwealth University, Richmond,
149 VA, USA
- 150 79 Department of Pathology and Biomedical Science, and Carney Centre for Pharmacogenomics,
151 University of Otago Christchurch, New Zealand
- 152 80 PsychGen Centre for Genetic Epidemiology and Mental Health, Norwegian Institute of Public
153 Health, Oslo, Norway
- 154 81 Nic Waals Institute, Lovisenberg Diaconal Hospital, Oslo, Norway
- 155 82 Epidemiology and Environmental Health Joint Research Unit, FISABIO-Universitat Jaume I-
156 Universitat de València, Valencia (Spain)
- 157 83 Spanish Consortium for Research on Epidemiology and Public Health (CIBERESP), Madrid
158 (Spain)
- 159 84 Department of Nursing, Universitat de València, Valencia, Spain
- 160 85 ISGlobal, Barcelona Institute for Global Health, Barcelona, Spain
- 161 86 Universitat Pompeu Fabra (UPF), Barcelona, Spain
- 162 87 CIBER Epidemiología y Salud Pública (CIBERESP), Spain
- 163 88 Social, Genetic and Developmental Psychiatry Centre, Institute of Psychiatry, Psychology and
164 Neuroscience, King's College London, U.K.
- 165 89 Institute and Outpatient Clinic for Occupational, Social and Environmental Medicine, University
166 of Munich Medical Center, Ludwig-Maximilians-Universität München, Munich, Germany

GWAMA OF ADHD SYMPTOMS AND DIAGNOSIS

- 167 ⁹⁰ Allergy and Lung Health Unit, Melbourne School of Population and Global Health, University of
168 Melbourne, Melbourne, Australia
- 169 ⁹¹ Christchurch Health and Development Study, Department of Psychological Medicine, University
170 of Otago Christchurch, New Zealand
- 171 ⁹² Biostatistics and Computational Biology Unit, Department of Pathology and Biomedical Science,
172 University of Otago Christchurch, New Zealand
- 173 ⁹³ Department of Pathology and Biomedical Science, and Carney Centre for Pharmacogenomics,
174 University of Otago Christchurch, New Zealand
- 175 ⁹⁴ Dunedin Multidisciplinary Health and Development Research Unit, University of Otago,
176 Dunedin, New Zealand
- 177 ⁹⁵ Massey Cancer Center, Virginia Commonwealth University, Richmond, VA, USA
- 178 ⁹⁶ Department of Psychiatry, College of Medicine, University of Vermont
- 179 ⁹⁷ Arkin mental health care, Amsterdam, The Netherlands
- 180 ⁹⁸ Queensland Brain Institute, Institute for Molecular Bioscience, University of Queensland, St
181 Lucia 4072, Australia
- 182 ⁹⁹ Department of Life Sciences, College of Health and Life Sciences, Brunel University London,
183 London, United Kingdom
- 184 ¹⁰⁰ Department of Psychiatry and Behavioral Sciences, Duke University School of Medicine,
185 Durham, N.C., U.S.A.
- 186 ¹⁰¹ Center for Genomic and Computational Biology, Duke University, Durham, N.C., U.S.A.
- 187 ¹⁰² Telethon Kids Institute, University of Western Australia, Nedlands, Western Australia, Australia
- 188 ¹⁰³ Department of Psychiatry, Robert Wood Johnson Medical School, Rutgers University, U.S.A.
- 189 ¹⁰⁴ Department of Psychology, Virginia Commonwealth University, U.S.A.
- 190 ¹⁰⁵ College Behavioral and Emotional Health Institute, Virginia Commonwealth University,
191 Richmond, VA, USA
- 192 ¹⁰⁶ Institute of Clinical Medicine, University of Oslo, Oslo, Norway
- 193 ¹⁰⁷ School of Psychology and Counselling, Queensland University of Technology, Brisbane,
194 Australia
- 195 ¹⁰⁸ Department of Public and Occupational Health, Amsterdam University Medical Center,
196 University of Amsterdam, Amsterdam
- 197 ¹⁰⁹ Amsterdam Reproduction and Development Research Institute, Amsterdam, The Netherlands
- 198 ¹¹⁰ Department of Public Health, Medical Faculty, University of Helsinki, Helsinki, Finland
- 199 ¹¹¹ Department of Social and Behavioral Science, Harvard TH Chan School of Public Health, Boston,
200 USA
- 201 ¹¹² Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK
- 202 ¹¹³ Interdisciplinary Center Psychopathology and Emotion regulation (ICPE), Department of
203 Psychiatry, University of Groningen, University Medical Center Groningen, Groningen, The
204 Netherlands
- 205 ¹¹⁴ Fundació d'Investigació Sant Pau, Barcelona, Spain
- 206 ¹¹⁵ Instituto para el Desarrollo de Terapias Avanzadas en Neurociencias (IDETAN). Barcelona, Spain
- 207 ¹¹⁶ Gillberg Neuropsychiatry Centre, University of Gothenburg, Gothenburg, Sweden
- 208 ¹¹⁷ Centre for Ethics, Law and Mental Health, University of Gothenburg, Gothenburg, Sweden
- 209

210 **Abstract**

211 Attention-deficit/hyperactivity disorder (ADHD) is a heritable neurodevelopmental disorder for
212 which genetic factors explain up to 75% of the variance. In this study, we performed a genome-
213 wide association meta-analysis (GWAMA) of ADHD symptom measures, with an effective sample
214 size of 120,092 (71,733 unique individuals from 28 population-based cohorts, with 288,887
215 quantitative ADHD symptom measures). Next, we meta-analyzed the results with a genome-wide
216 association study (GWAS) of ADHD diagnosis. The GWAMA of ADHD symptoms returned no
217 genome-wide significant variants. However, we estimated strong genetic correlations between our
218 study of quantitative ADHD symptoms and the earlier study of ADHD diagnosis ($r_g = 1.00$, $SE =$
219 0.06). Moderate negative genetic correlations ($r_g < -0.40$) were observed with several cognitive
220 traits. Genetic correlations between ADHD and aggressive behavior and antisocial behavior were
221 around 1. This provides further evidence of the wide pleiotropic effects of genetic variants and the
222 role that genetic variants play in the co-occurrence with (mental) health traits. The GWAMAs of
223 ADHD symptoms and diagnosis identified 2,039 genome-wide significant variants, representing
224 39 independent loci, of which 17 were new. Using a novel fine-mapping and functional annotation
225 method, we identified 22 potential effector genes which implicate several new potential biological
226 processes and pathways that may play a role in ADHD. Our findings support the notion that clinical
227 ADHD is at the extreme end of a continuous liability that is indexed by ADHD symptoms. We
228 show that including ADHD symptom counts in large-scale GWAS can be useful to identify novel
229 genes implicated in ADHD and related symptoms.

230

231 Keywords: ADHD, GWAS, symptoms, diagnosis, genetics, FLAMES

232

233 **Introduction**

234 Attention-deficit/hyperactivity disorder (ADHD) is, for many individuals, a persistent
235 neurodevelopmental disorder (Faraone et al., 2006; Kan et al., 2013). ADHD is characterized by
236 three core symptoms: hyperactivity, impulsivity, and inattention (American Psychiatric
237 Association, 2013). It affects around 5% of children and adolescents and 2.5% of adults worldwide
238 (Faraone et al., 2015). ADHD may be associated with serious consequences for affected
239 individuals, their families, and society at large, with symptoms persisting across multiple settings,
240 i.e. at home, at school, and elsewhere (Caci et al., 2014, 2015). The disorder has a predominantly
241 genetic aetiology, involving both common and rare genetic variants (Faraone et al., 2024). The
242 mean estimated heritability across 37 twin studies of ADHD was 74% (Faraone & Larsson, 2019;
243 Kan et al., 2014; Merwood et al., 2013), with some differences in heritability across ages and raters
244 (Kan et al., 2014; Merwood et al., 2013).

245 In 2019, a genome-wide association meta-analysis (GWAMA) of clinical ADHD, hereafter
246 referred to as ADHD_{DIAG2019}, including data from 20,183 cases and 35,191 controls, identified the
247 first 12 genome-wide significant loci associated with ADHD (Demontis et al., 2019). The study
248 reported that 22% of the variance in ADHD could be explained by all measured single nucleotide
249 polymorphisms (SNPs). They also performed meta-analyses with deCode, 23andMe, and EAGLE
250 (Middeldorp et al., 2016). Four independent loci reached the genome-wide significance threshold
251 ($p < 5 \times 10^{-8}$) in all three meta-analyses. Interestingly, most independent significant loci, 15, were
252 found in the meta-analysis with a quantitative assessment of attention problems in EAGLE,
253 suggesting that quantitative assessments of ADHD can boost the power to identify implicated
254 genetic variants. Ten independent significant loci were found in meta-analyses with deCode and
255 23andMe data. In 2023, Demontis and colleagues presented results from their updated GWAS

GWAMA OF ADHD SYMPTOMS AND DIAGNOSIS

256 meta-analysis of ADHD (ADHD_{DIAG}), combining newly extended data from iPSYCH, deCode,
257 and the PGC, almost doubling the number of cases compared with ADHD_{DIAG2019} (Demontis et al.,
258 2023). The definition of cases was broader, for example by including individuals who used ADHD
259 prescription medication. The study reported 27 independent significant loci and estimated that 14%
260 of the variance in ADHD could be attributed to the included SNPs. The broader definition of
261 ADHD diagnosis resulted in a larger sample and, therefore, more power to detect implicated
262 genetic variants. However, this broader definition also increased the heterogeneity of the
263 phenotype, which may explain the decrease in SNP-heritability (Wang et al., 2023).

264 There is increasing interest and recognition that ADHD symptom counts in non-clinical
265 samples can tap into the same genetic construct as clinically diagnosed ADHD, supporting the
266 notion that ‘clinical ADHD’ is at the extreme end of a continuous measure of ADHD symptoms
267 (Larsson et al., 2012; Levy et al., 1997). This was initially suggested based on multivariate twin
268 studies (Derks et al., 2008). Additionally, the genetic correlation (r_g) between quantitative ADHD
269 symptom counts (Middeldorp et al., 2016) and ADHD_{DIAG2019} (Demontis et al., 2019) was
270 estimated to be 0.97 (SE= 0.21, $p= 2.66 \times 10^{-6}$), suggesting that combining these measures is a
271 viable strategy to increase statistical power in ADHD GWASs. This was additionally supported by
272 the increased number of genome-wide significant loci in the meta-analysis of ADHD_{DIAG2019} and
273 EAGLE, as compared to ADHD_{DIAG2019} alone, and to meta-analyses of ADHD_{DIAG2019}, deCode and
274 23andMe.

275 In this study, we combine information from 28 population-based cohorts in a GWAMA of
276 continuous ADHD symptom scores, leading to a total of 71,733 participants. The measures include
277 repeated assessments (longitudinal data) by multiple raters (maternal, paternal, teachers, and self-
278 assessments) and instruments across ages (range: 2-18 years), for a total of 288,887 measures. We

279 also include retrospective self-report data. We meta-analyze all available data into a cross-
280 rater/cross-age/cross-instrument GWAMA of ADHD symptoms ($ADHD_{SYMP}$), taking into
281 consideration the dependency between multiple assessments within individuals (Ip et al., 2021).
282 Using measures from multiple raters and ages may further increase the power of the analyses due
283 to an increase in the validity of the ADHD symptom measures. Next, we estimate the genetic
284 correlations (r_g) between $ADHD_{SYMP}$ and the meta-analysis of case-control samples (Demontis et
285 al., 2023), and meta-analyze $ADHD_{SYMP}$ with $ADHD_{DIAG}$ ($ADHD_{overall}$). Finally, we performed
286 fine mapping and gene-based tests based on $ADHD_{SYMP}$ and $ADHD_{overall}$, performed several
287 follow-up enrichment and pathway analyses, and estimated genetic correlations between the
288 GWAMA and a set of predefined outcomes from cognitive and externalizing behaviour domains.

289


290

291 **Results**

292 **$ADHD_{SYMP}$ GWAMA**

293 We first meta-analyzed the effect of each SNP across all available univariate GWAS of quantitative
294 ADHD measures. Based on an effective sample size of 120,092, the estimated h_{SNP}^2 of $ADHD_{SYMP}$
295 was 0.04 (SE= 0.01; Z= 8.12). The mean χ^2 statistic was 1.09 along with an LDSC-intercept of
296 1.01 (SE= 0.01), indicating that there was no or very limited inflation in test statistics due to
297 confounding biases such as population stratification. Rather, the GWAMA most likely captured
298 the polygenic nature of childhood ADHD symptoms. The GWAMA of ADHD symptoms did not
299 result in any genome-wide significant SNPs (Figure 1, Supplements Table 11).

GWAMA OF ADHD SYMPTOMS AND DIAGNOSIS



300

301 Figure 1. Manhattan plot: GWAMA of ADHD symptoms.

302


303 **Meta-analysis with ADHD diagnosis GWAS**

304 SNP-heritability estimated with genomicSEM was 0.13 (SE= 0.01) for ADHD_{DIAG}. The estimated
305 genetic correlation between ADHD_{SYMP} and ADHD_{DIAG} was 1.00 (SE= 0.06). The CTI was not
306 significantly different from zero and were subsequently constrained to zero in the following meta-
307 analysis.

308 Because the point estimate of the genetic correlation between ADHD_{SYMP} and ADHD_{DIAG}
309 was not significantly different from 1, we constrained the genetic correlation at unity when pre-
310 adjusting the weights and Z-scores for the meta-analysis of ADHD_{SYMP} and ADHD_{DIAG}. A total of
311 6,571,852 SNPs were included in the meta-analysis. The SNP-heritability of ADHD_{overall} was 0.11
312 (SE= 0.01), with a mean χ^2 statistic of 1.52. The LDSC-intercept and ratio were 1.02 (SE= 0.01)
313 and 0.03 (SE= 0.02), respectively, indicating that approximately 3% of the signal might be due to
314 confounding factors. A Manhattan plot of ADHD_{overall} is shown in Figure 2. 2,039 SNPs reached
315 genome-wide significance ($p < 5 \times 10^{-8}$), of which 644 were also reported in ADHD_{DIAG} and 1,395

GWAMA OF ADHD SYMPTOMS AND DIAGNOSIS

316 were new. The 2,039 SNPs corresponded to 43 independent lead SNPs in 39 independent
317 significant loci, identified with FUMA. Of these 39 loci, 22 were also reported in ADHD_{DIAG} and
318 17 were new. See also Figure 2 and Supplements Tables 12 and 13.



319
320 Figure 2. Manhattan plot: GWAMA of ADHD symptoms and ADHD diagnosis. *Note: orange dots*
321 *reflect lead SNPs.*

322

323

324 **Follow-up analyses**

325 *Fine mapping & gene-based tests*

326 Gene mapping in FUMA mapped the 43 lead SNPS in 39 independent genomic risk loci to 204
327 associated genes (See Supplements Table 14), of which 45 were also reported by Demontis et al
328 (2023). Second, gene-based tests were run in MAGMA (de Leeuw et al., 2015), identifying 64
329 associated genes (see Supplements Table 15), of which 17 were previously reported by Demontis
330 et al. (2023). Third, we ran FLAMES (Schipper et al., 2023), with the aim to get a better
331 understanding of the genes that are causally involved in ADHD. A total of 22 genes had

GWAMA OF ADHD SYMPTOMS AND DIAGNOSIS

332 FLAMES scores larger than 0.05, and were interpreted as potential effector genes (see
333 Supplements Table 16). Fourteen of these effector genes were also tagged by the MAGMA gene-
334 based test. Ten of these potential effector genes were previously reported by Demontis et al.
335 (2023). Four genes were not reported by Demontis et al. (2023) but have previously been linked
336 to ADHD, as listed by GWAS Catalog. Eight potential effector genes were not reported by
337 Demontis et al. (2023) or by any ADHD specific studies listed on GWAS Catalog: *EMCN*,
338 *STK32C*, *PCDH17*, *TCF12*, *PEAK1*, *IGF1R*, *CTNNA2*, and *ABCA12*.

339

340 *Enrichment & tissue specific expression*

341 Gene-set analysis in MAGMA revealed no significant enrichment in any MSigDB v2023
342 gene-sets after correction for multiple testing. MAGMA expression analysis showed significant
343 enrichment of the GWAMA signal in gene-sets differentially expressed in late infancy.
344 Additionally, there was significant enrichment in several brain tissue types, as well as in the
345 pituitary gland (See Supplements Figures 1 and 2).

346 Next, FUMA GENE2FUNC gene-set enrichment analyses of the 204 potential ADHD risk
347 genes mapped by FUMA showed significant enrichment in genes identified in GWAS of ADHD,
348 cognition-related phenotypes, and risk-taking behaviors. The genes were not significantly enriched
349 in any tissue types, or in any of the Brainspan developmental stages of brain samples. Genes were
350 enriched in 29 gene sets that code for transcription factor targets. Zero synapse cellular component
351 terms or biological processes were significantly enriched at 1% FDR (testing terms with at least
352 three matching input genes in SynGO). For a complete overview of all enrichment results and the
353 included gene sets, see Supplemental Figures 3-5, and Supplemental Notes.

354 We repeated the same analyses for the 22 potential effector genes identified by FLAMES.
355 Again, genes were highly enriched for genes identified in GWAS of cognition-related phenotypes,

356 and risk-taking behaviors. The 22 genes were significantly overrepresented in gene sets that are
357 differentially expressed in the frontal cortex, but not in the “general tissue” type brain, or in any of
358 the Brainspan developmental stages. Genes were also enriched in 52 gene sets that code for
359 transcription factor targets, 13 microRNA targets, 4 gene ontology biological processes, 1
360 canonical pathway, and 8 cell type signatures. Zero synapse cellular component terms or biological
361 processes were significantly enriched at 1% FDR (testing terms with at least three matching input
362 genes in SynGO). Nine genes were mapped to SynGO annotations, eight to cellular components
363 and 9 to biological processes. Genes were enriched in integral components of the postsynaptic
364 density membrane ($q= 1.46 \times 10^{-3}$), postsynaptic density ($q= 1.57 \times 10^{-3}$), postsynapse ($q=$
365 5.67×10^{-3}), and synapse ($q= 7.22 \times 10^{-3}$), as well as in postsynaptic modulation of chemical
366 synaptic transmission ($q= 3.28 \times 10^{-5}$), process in the synapse ($q= 3.97 \times 10^{-4}$), and synapse
367 organization ($q= 1.62 \times 10^{-3}$). For a complete overview of all enrichment results and the included
368 gene sets, see Supplemental Figures 6-12, Supplementary Tables 17 and 18, and Supplemental
369 Notes.

370


371 *Genetic correlations*

372 We estimated genetic correlations between $ADHD_{overall}$ and 49 external phenotypes. Results are
373 shown in Figure 4 and Supplementary Table 18. Strong positive genetic correlations were observed
374 between $ADHD_{overall}$ and childhood aggressive behavior ($r_g= 1.13$, $SE= 0.05$) and antisocial
375 behavior ($r_g= 0.97$, $SE= 0.06$). The correlation of 1.13 with childhood aggressive behavior reflects
376 a high genetic correlation that is likely greater than 1 due to sampling variation, correlations
377 estimated by LD score regression in genomicSEM are not bounded between -1 and 1 (Childhood
378 aggression $h^2 Z= 9.03$). Measures of smoking habits ($r_g= 0.46-0.6$, $SE= 0.03$) and number of

GWAMA OF ADHD SYMPTOMS AND DIAGNOSIS

379 children ($r_g = 0.38$, $SE = 0.04$) also showed moderate correlations, as did ratings of overall health
380 ($r_g = -0.59$, $SE = 0.03$), educational attainment ($r_g = -0.55$, $SE = 0.02$), and childhood IQ ($r_g = -0.43$,
381 $SE = 0.06$). In general, $ADHD_{overall}$ showed weak to moderate genetic correlations with
382 psychopathologies, including Major Depressive Disorder ($r_g = 0.57$, $SE = 0.03$) and Autism
383 Spectrum Disorder ($r_g = 0.39$, $SE = 0.04$). Weak negative genetic correlations were found between
384 $ADHD_{overall}$ and alcohol intake frequency ($r_g = -0.28$, $SE = 0.03$), but a weak positive correlation
385 with drinks per week ($r_g = 0.14$, $SE = 0.03$) and a weak positive correlation with cannabis use ($r_g =$
386 0.2 , $SE = 0.03$). Marees and colleagues (Marees et al., 2020) looked into similar genetic correlations
387 for alcohol use, and found evidence to suggest that they are the result of SES influences.
388 $ADHD_{overall}$ was weakly negatively genetically correlated with birth weight ($r_g = -0.1$, $SE = 0.02$),
389 but showed a weak positive genetic correlation with childhood obesity ($r_g = 0.21$, $SE = 0.05$) and a
390 moderate positive genetic correlation with adult BMI ($r_g = 0.3$, $SE = 0.02$).

GWAMA OF ADHD SYMPTOMS AND DIAGNOSIS



391
 392 **Figure 4.** Genetic correlations with external phenotypes. Bars indicate 95% confidence intervals.

393

394

395 Discussion

396 We present a genome-wide association meta-analysis of childhood Attention-Deficit/Hyperactivity

397 Disorder symptoms (ADHD_{SYMP}). A total of 28 cohorts with measures of ADHD symptom counts

GWAMA OF ADHD SYMPTOMS AND DIAGNOSIS

398 took part; contributing data from multiple raters and instruments, across a wide range of ages. We
399 meta-analyzed all continuous measures and combined these results with results from two
400 GWAMAs of ADHD diagnosis ($ADHD_{DIAG}$) by Demontis and colleagues (2019, 2023).

401 We did not identify any genome-wide significant hits for ADHD symptoms specifically.
402 Genetic correlations with ADHD diagnosis was $r_g = 1.00$, $SE = 0.06$. This supports the notion that
403 clinical ADHD is at the extreme end of a continuous genetic liability that is indexed by ADHD
404 symptoms (Larsson et al., 2012; Levy et al., 1997). This was previously suggested based on
405 multivariate twin studies (Derks et al., 2008). The estimated h_{SNP}^2 of $ADHD_{SYMP}$ was 0.04 ($SE =$
406 0.01), which may be considered low compared to the estimated h_{SNP}^2 in Demontis et al., 2019 and
407 2023: 0.22 and 0.14 respectively. This is probably because we introduced heterogeneous
408 measurement error and bias in our phenotyping by including symptom measures from different
409 raters, and at different ages, which could subsequently suppress SNP-heritability.

410 By meta-analyzing GWASs of ADHD symptoms and ADHD diagnosis, we found 2,039
411 genome-wide significant variants in 39 independent loci, of which 17 were new. Demontis et al.
412 found 12 and 27 independent hits in 2019 and 2023, which included data from 23andMe. These
413 23andMe data were excluded from the current meta-analyses. This shows that, because ADHD
414 symptom measures have been widely collected, combining ADHD symptom counts and diagnosis
415 can be effective in identifying implicated genetic variants for ADHD. The estimated h_{SNP}^2 of
416 $ADHD_{overall}$ was 0.11 ($SE = 0.01$), compared to 0.13 ($SE = 0.01$) in $ADHD_{DIAG}$ when excluding
417 23andMe data. This means that by including $ADHD_{SYMP}$ in the $ADHD_{DIAG}$ results, h_{SNP}^2 slightly
418 decreased. We believe this is due to heterogeneous measurement error and bias in the $ADHD_{SYMP}$
419 phenotyping (Wang et al., 2023). The same can be observed when looking at the differences in
420 h_{SNP}^2 between $ADHD_{DIAG}$ from 2019 ($h_{SNP}^2 = 0.22$, including 23andMe), which was strict in its

GWAMA OF ADHD SYMPTOMS AND DIAGNOSIS

421 definition of ADHD cases, and ADHD_{DIAG} from 2023 ($h^2_{SNP} = 0.14$, including 23andMe), which
422 was slightly more lenient in its definition of ADHD cases.

423 MAGMA analyses identified 64 potential ADHD risk genes. These genes were
424 significantly enriched in genes previously identified in GWASs of cognitive phenotypes and risk-
425 taking behaviors. The total GWAS signal was significantly differentially expressed in several brain
426 specific tissue types, general brain tissue types and the pituitary gland, as well as in late infancy
427 Brainspan brain samples. FUMA gene-mapping mapped significant loci to 204 genes. Again, genes
428 were enriched in gene-sets reported by previous GWASs on cognitive behavior, risk-seeking
429 behavior, and brain development. FUMA enrichment analyses further revealed 29 transcription
430 factor targets that may be of interest for ADHD.

431 To get a better understanding of the causal pathways from SNPs to ADHD, we ran
432 FLAMES (Schipper et al., 2023) to identify likely effector genes. FLAMES identified 22 potential
433 effector genes, of which 14 overlapped with the MAGMA genes, 12 were previously reported by
434 Demontis et al. (2023) and 8 were not previously linked to ADHD or not reported on GWAS
435 Catalog. The 22 genes were significantly overrepresented in gene-sets differentially expressed in
436 the frontal cortex, enriched in 4 GO biological processes related to neural and physical
437 development, 52 transcription factor targets, 13 microRNA targets, 8 different cell type signatures,
438 4 synapse cellular components, and 3 synaptic biological processes. In Demontis et al. (2023), the
439 set of potential ADHD risk genes was significantly enriched among genes upregulated during early
440 embryonic brain development, this result was not replicated in our study. A common theme is that
441 implicated genes are enriched in processes that are involved in neural development and functioning.
442 The results provide several new avenues to investigate, which could prove useful in gaining more
443 insights into the etiology of ADHD. The results may provide more potentially useful information
444 for the 22 potential effector genes compared to the 204 genes identified by FUMA positional

GWAMA OF ADHD SYMPTOMS AND DIAGNOSIS

445 mapping, eQTL mapping, and chromatin interaction mapping. We think this difference results from
446 the difference in strategies employed by both methods. FUMA maps every gene for which some
447 functional link is known to exist, whereas FLAMES weighs all these measurements and only
448 prioritizes genes if they are clearly more likely causal genes than the other genes in the locus. Our
449 findings suggests that FLAMES can help identify functional pathways that may remain hidden
450 with other approaches due to a reduction of noise from non-causal genes in the set of prioritized
451 genes, which decreases the power to detect enrichment in functional gene-sets.

452 Next, we examined genetic correlations between $ADHD_{overall}$ and various external
453 phenotypes. Except for anorexia nervosa, $ADHD_{overall}$ showed significant genetic correlations with
454 all examined psychopathological traits. Most striking was the genetic correlation of 1.13 with
455 childhood aggressive behavior, and 0.97 with antisocial behavior. Previous studies reported
456 moderate-to-strong phenotypic correlations across sex-, rater-, age-, and instrument-specific
457 assessment between aggressive behavior and attention problems and hyperactivity (Bartels et al.,
458 2018). We found a negative genetic correlation between ADHD and alcohol intake frequency,
459 and a positive correlation between ADHD and number of drinks per week. Marees et al. (2020)
460 studied similar correlations between alcohol consumption and mental health traits, and found
461 evidence that suggests the different genetic correlations are a result of SES effects. We found a
462 moderate genetic correlation with smoking behaviors, but a small correlation with cannabis use
463 and discrepant findings for alcohol consumption. This was surprising, given the phenotypic
464 associations that have been reported in previous studies (Lee et al., 2011). We found negative
465 correlations with several cognitive traits, such as (childhood) IQ, verbal-numerical reasoning, and
466 educational attainment. Similar to a GWAS of childhood aggression (Ip et al., 2021), genetic
467 correlations with several hormone levels were around zero. Finally, we found a small negative
468 correlation with birthweight, but a weak positive correlation with childhood obesity, and adult

469 BMI. These genetic correlations suggest wide pleiotropic effects of variants involved with ADHD.
470 This is illustrative of the polygenetic nature of most behavioral, cognitive, and (mental) health
471 traits. It also indicates genetic factors play a role in the comorbidity of psychopathological
472 disorders.

473 Combining data collected using different instruments and by different raters helps to
474 increase the sample size, and with that the statistical power of our analyses. This is illustrated by
475 the apparent increase in power to detect genetic variants associated with ADHD in the combination
476 ADHD_{SYMP} and ADHD_{DIAG}, compared to the combination ADHD_{DIAG} and EAGLE ADHD GWAS
477 (Demontis et al., 2019; Middeldorp et al., 2016). However, this approach of including different
478 raters and instruments also has some downsides. Correlations between raters and instruments are
479 not always large, meaning that we introduce heterogeneous measurement error and bias in our
480 phenotyping, which could suppress SNP-heritability. More homogeneous measurement may offer
481 an avenue to higher powered GWAS. In light of increased collaborations between cohorts to
482 expand sample sizes for GWAS, more efforts to standardize the assessment of traits could help
483 further increase the power and specificity of GWAS results. Moreover, downstream analyses, for
484 example using polygenic scores computed with the summary statistics of the current study, will
485 favor cohorts using the same instruments and raters that dominated the discovery sample, in our
486 case the Achenbach System of Empirically Based Assessment (Achenbach et al., 2017).

487 Assessments of ADHD in individuals from non-European ancestry were rare in each of the
488 included cohorts. Because of the low number of assessments, we were forced to exclude non-
489 European individuals from our analyses. We know that results from European ancestry GWASs
490 often also significantly predict differences in non-European ethnic groups, but often effect sizes
491 are diluted towards zero. Regrettably, this means that knowledge generated by these types of
492 studies risks benefit individuals of European ancestry more than from diverse backgrounds. To

493 better understand the etiology of ADHD across individuals and backgrounds, it is important to
494 continue ongoing efforts to increase the inclusivity of GWAS samples.

495 In conclusion, the current study adds novel insight into the genetic etiology of ADHD. By
496 meta-analyzing GWAS results from symptom counts of ADHD in children with a diagnosis of
497 ADHD, we identified novel genome-wide significant loci and genes. The number of genome-wide
498 (significant) genetic variants that are implicated in ADHD provides further insight into the
499 polygenic etiology of ADHD. The 22 potential effector genes that were identified by FLAMES
500 gives insight in several biological processes that may play a causal role in the etiology of ADHD,
501 and provides avenues for further research. The genetic correlations with other phenotypes further
502 indicate the wide pleiotropic effects of genetic variants and the role that genetic variants play in
503 the co-occurrence with (mental) health traits.

504

505

506 **Methods**

507 **Sample & cohorts**

508 Childhood cohorts that collaborate within the ACTION (Aggression in Children: unraveling gene-
509 environment interplay to inform Treatment and InterventiON strategies) consortium (Bartels et al.,
510 2018; Boomsma, 2015; Ip et al., 2021) and EAGLE consortium (Middeldorp et al., 2019) took part
511 in the meta-analysis of ADHD symptom counts (ADHD_{SYMP}). Cohorts assessed ADHD Symptoms
512 in children and adolescents aged 1.5 to 18 years and also included adult retrospective assessments.
513 Each cohort received a standard operation protocol (see: [https://www.action-
514 euproject.eu/content/data-protocols](https://www.action-euproject.eu/content/data-protocols)). Cohorts could contribute one or several univariate genome-
515 wide association studies (GWAS). A separate analysis was performed for every unique

516 combination of rater, instrument, and age, with a minimum of 450 observations per GWAS (so that
517 each GWAS included a maximum of one measure for each individual). Extended information on
518 the cohorts can be found in Supplementary Table 1 and the Supplementary Text. Assessments of
519 individuals of non-European ancestry were limited, and analyses were restricted to individuals of
520 European ancestry. In total, 28 cohorts contributed 154 GWASs, resulting in a total of 288,887
521 observations from 72,483 unique individuals (Supplementary Table 2).

522

523 **Measurement of ADHD symptoms**

524 ADHD symptoms in children and adolescents were rated by parents, teachers, and the individuals
525 themselves. Additionally, retrospective assessments of (pre-) adolescent ADHD symptoms from
526 self-report or maternal report were included in the GWAMA. To maximize sample size, we
527 included measurement of ADHD symptoms across multiple instruments. In total, 22 ADHD
528 symptom assessment instruments were included in the meta-analysis (Supplementary Table 3). The
529 most commonly employed instruments were the Achenbach System of Empirically Based
530 Assessment (Achenbach et al., 2017) and the Strengths and Difficulties Questionnaire (Goodman,
531 2001).

532

533 **Genotyping and quality control**

534 Genotyping was performed within each cohort using common genotyping arrays (see
535 Supplementary Table 4), followed by cohort-specific quality control based on individual- and
536 variant-based call rate, Hardy-Weinberg equilibrium, excessive heterozygosity rates, and minor
537 allele frequency (see Supplementary Table 5). Cohorts removed samples with non-European
538 ancestry. 78.6% of the cohorts imputed their genotypes to 1000 Genomes (1000G) phase 3 version
539 5, while the remaining cohorts used 1000G phase 1 version 3 as the reference set for the imputation

540 (Supplementary Table 6). All genotypes were mapped to build 37 of the Human Genome Reference
541 Consortium assembly (GRCh37).

542
543 **GWA model**
544 Each cohort performed a univariate GWAS where ADHD symptoms were regressed on the SNP
545 genotype, with age, sex, and first five ancestry-based principal components as fixed effects, and, if
546 necessary, cohort-specific covariates (Supplementary Table 7). To correct for dependency between
547 observations within univariate analyses, cohorts with related individuals applied a mixed linear
548 model (Tucker et al., 2015) or a sandwich correction of the standard errors (Minică et al., 2015).

549 GWASs were stratified by (1) rater, (2) instrument, and (3) age, so that observations within
550 a univariate GWAS were independent, with a minimum stratum sample size of 450 observations.
551 In total, summary statistics for 154 univariate GWAS were uploaded. Descriptive statistics for each
552 uploaded GWAS are shown in Supplementary Table 8. Each cohort also supplied information on
553 the degree of sample overlap and phenotypic correlation between their univariate analyses. These
554 statistics allowed us to account for dependency between observations within cohorts.

555
556 **Pre-GWAMA QC**

557 Summary statistics from each GWAS went through quality control (QC) using the EasyQC
558 software package (Winkler et al., 2014). SNPs with a genotyping rate below 95% were removed.
559 We applied variable QC filters on MAF and HWE p -value tailored to the sample size in the specific
560 GWAS. Respective cutoffs of $INFO > 0.6$ and $INFO > 0.7$ were applied to SNPs that were imputed
561 using MACH and IMPUTE (Liu et al., 2015). Reported allele frequencies were compared to the
562 allele frequency in an imputation-matched reference population and variants with an absolute

563 difference in allele frequency larger than 0.2 were removed. Supplementary Table 9 reports the
 564 number of SNPs before and after QC.

565

566 **Meta-analysis of ADHD symptoms**

567 The meta-analysis method is equal to the method described by Ip and colleagues (Ip et al., 2021).
 568 Due to sample overlap between the univariate GWASs, covariance between GWAS test statistics
 569 are a function of sample overlap and a truly shared genetic signal (Bulik-Sullivan, Loh, et al.,
 570 2015). To correct for sample overlap during the meta-analysis, we applied a modified version of
 571 the multivariate meta-analysis approach developed by Baselmans and colleagues (2019), where we
 572 calculated the expected cross-trait-intercept (Bulik-Sullivan, Loh, et al., 2015) based on the
 573 observed sample overlap and phenotypic covariance, as reported by the cohorts. Finally, because
 574 the sum of the number of observations (N_{obs}) was an overestimate of the effective sample size
 575 (N_{eff}), we approximated the effective sample size as proposed by Ip and colleagues (2021) ($N_{eff} =$
 576 $\frac{4}{\left(\frac{1}{N_{cases}} + \frac{1}{N_{controls}}\right)}$). SNPs with $MAF < 0.01$, $N_{eff} < 15,000$, or were observed in only one cohort were
 577 removed from further analyses. SNP-heritability (h_{SNP}^2) was estimated by genomic Structural
 578 Equation Modeling in R (Grotzinger et al., 2019).

579

580 **Meta-analysis with case-control ADHD GWAS**

581 In the next step, we meta-analyzed our ADHD_{SYMP} GWAMA with a GWAS of ADHD diagnosis
 582 (Demontis et al., 2023). In ADHD_{DIAG}, cases are defined as clinically diagnosed with ADHD or
 583 prescribed medication specific for ADHD. ADHD_{DIAG} included data from the Lundbeck
 584 Foundation Initiative for Integrative Psychiatric Research (iPSYCH), the Psychiatric Genomics

585 Consortium (PGC), and deCode. Data were obtained for adults and children, resulting in a total of
 586 38,691 cases and 186,843 controls.

587 For the meta-analysis, we first adjusted the test statistics and sample sizes for ADHD_{SYMP}
 588 and ADHD_{DIAG} as proposed by (Demontis et al., 2019). The lifetime population prevalence of
 589 ADHD was assumed to be 5% (Faraone et al., 2015). SNP-heritability for ADHD_{SYMP}, and
 590 ADHD_{DIAG}, and r_g and cross-trait intercepts (CTI) between ADHD_{SYMP} and ADHD_{DIAG} were
 591 estimated by genomicSEM in R (Grotzinger et al., 2019). We meta-analyzed the results from
 592 ADHD_{SYMP} and ADHD_{DIAG} based on the approach outlined in Baselmans and colleagues
 593 (Baselmans et al., 2019). We specified the effective sample sizes for ADHD_{DIAG} as suggested by
 594 Ip and colleagues (2021): $N_{eff} = \frac{4}{\left(\frac{1}{N_{cases}} + \frac{1}{N_{controls}}\right)}$. SNP-heritability was estimated using LDSC in
 595 genomicSEM (Grotzinger et al., 2019). We assumed no sample overlap between ADHD_{SYMP} and
 596 ADHD_{DIAG}.

597

598 **Follow-up analyses**

599 *Fine mapping & gene-based tests*

600 We used FUMA (Watanabe et al., 2017) to identify independent genome-wide significant loci
 601 and associated genes. LD blocks of independent significant SNPs within 250kb were merged into
 602 a single genomic locus. Protein-coding genes were mapped if they were located within a
 603 maximum distance of 10 kb of an independent significant SNP, or if a credible variant was
 604 annotated to the gene based on eQTL data or chromatin interaction data from human brain (see
 605 Supplemental Text for the included datasets). These are the same settings used by Demontis et al.
 606 (Demontis et al., 2023). Gene-based tests were run in MAGMA (de Leeuw et al., 2015).
 607 MAGMA gene-based tests combine P -values from multiple SNPs inside a gene to obtain a test

608 statistic for each gene (Z_{gene}), while accounting for incomplete linkage disequilibrium between
609 SNPs. To this end, a list of 18,296 genes and their start- and end-positions, and pre-formatted
610 genotypes, based on 1000G phase 3, were obtained from the MAGMA website (see
611 Supplements). We applied a Bonferroni correction for multiple testing at $\alpha = 0.05 / 18,296 =$
612 2.733×10^{-6} .

613 It remains difficult to identify which genes are causally involved in ADHD. Fine-mapped
614 Locus Assessment Model of Effector geneS (FLAMES) was recently developed with the goal of
615 predicting the most likely effector genes from GWAS results. FLAMES is a novel framework that
616 combines SNP-to-gene evidence and convergence-based evidence, outputting a single score per
617 gene from fine-mapped GWAS loci. We performed statistical fine-mapping using FINEMAP
618 version 1.4.1 (Benner et al., 2016), and an LD reference panel of 100,000 unrelated UK biobank
619 participants of European descent. Given that the GWAMA contains cohorts that do not belong to
620 the UK biobank we restricted the maximum number of causal variants in a locus modelled by
621 FINEMAP to 1, to avoid overfitting. We ran FLAMES (version 1.0.0) by inputting pathway naïve
622 PoPS scores (Weeks et al., 2023) for our GWAMA, the FUMA defined loci and corresponding
623 fine-mapped credible sets, resulting in a single FLAMES score per gene. Genes with FLAMES
624 scores above 0.05 were interpreted as potential effector genes, as suggested by the FLAMES
625 authors. For more information on FLAMES, and the included functional annotations, see Schipper
626 et al. (2023). Functional annotation and enrichment analysis was done for a set of genes with
627 FLAMES scores above 0.05.

628

629 *Enrichment & pathway analyses*

630 We performed MAGMA gene-set analyses using the full $\text{ADHD}_{\text{overall}}$ results. Gene property
631 analysis was performed to test relationships between tissue specific gene expression profiles (See

632 Supplementary Notes for an overview) and ADHD-gene associations. Next, genes mapped by
633 FUMA, and the set of potential effector genes identified with FLAMES were used in gene-set
634 enrichment analyses. We ran hypergeometric tests using FUMA genes2func to assess if genes of
635 interest are overrepresented in any of the pre-defined gene sets (See Supplemental Notes for all
636 included gene-sets). We used SynGO (Koopmans et al., 2019) v1.2 (“20231201”) to test for
637 enrichment in genes encoding for proteins involved in synaptic cellular components and biological
638 pathways. The brain expressed background set was used, containing 18,035 unique genes.

639

640 *Genetic correlations*

641 We computed genetic correlations between ADHD_{overall} and 49 preselected traits, including
642 cognition and externalizing behaviors, psychopathologies, anthropometric measures, metabolic,
643 and health outcomes (see Supplementary Table 10). Phenotypes were selected based on established
644 hypotheses or were at least nominally significantly ($P < 0.05$) genetically correlated with
645 ADHD_{DIAG2019} (Demontis et al., 2019). Following Bulik-Sullivan et al. (Bulik-Sullivan, Finucane,
646 et al., 2015) we restricted genetic correlations to external phenotypes for which the Z-scores of the
647 LDSC-based h^2_{SNP} are ≥ 4 .

648

649

650

651

652

653

654

655

656 **Acknowledgements**

657 We very warmly thank all participants, their parents, and teachers for making this study possible.
658 The project was supported by the *Aggression in Children: Unraveling gene-environment*
659 *interplay to inform Treatment and InterventiON strategies* project (ACTION). ACTION received
660 funding from the European Union Seventh Framework Program (FP7/2007-2013) under grant
661 agreement no 602768. Cohort-specific acknowledgements and funding information are included
662 in the Supplementary text.

663

664 **Ethics & Inclusion Statement**

665 This study is the result of a large collaborative effort among multiple clinical and population-
666 based cohorts. Researchers and principal investigators (PIs) representing the individual cohorts
667 were involved in the design and execution of the study. Cohort specific GWAS analyses were
668 performed locally, by local researchers. Local researchers and PIs were included as co-authors in
669 consultation with the PIs of each included cohort. Data collections for the cohorts were approved
670 by local ethics committees. Study approval was obtained from the Central Ethics Committee on
671 Research Involving Human Subjects of the VU University Medical Center, Amsterdam (NTR
672 25th of May 2007 and ACTION 2013/41 and 2014.252), an Institutional Review Board certified
673 by the U.S. Office of Human Research Protections (IRB number IRB00002991 under Federal-
674 wide Assurance- FWA00017598; IRB/institute codes).

675

676 **Competing interests**

677 J.A.R.Q. was on the speakers' bureau and/or acted as consultant for Biogen, Idorsia, Casen-
678 Recordati, Janssen-Cilag, Novartis, Takeda, Bial, Sincrolab, Neuraxpharm, Novartis, BMS,
679 Medice, Rubió, Uriach, Technofarma and Raffo in the last 3 years. He also received travel

GWAMA OF ADHD SYMPTOMS AND DIAGNOSIS

680 awards (air tickets + hotel) for taking part in psychiatric meetings from Idorsia, Janssen-Cilag,
681 Rubió, Takeda, Bial and Medice. The Department of Psychiatry chaired by him received
682 unrestricted educational and research support from the following companies in the last 3 years:
683 Exeltis, Idorsia, Janssen- Cilag, Neuraxpharm, Oryzon, Roche, Probitas and Rubió. M.C. has
684 received fees to give talks for TAKEDA and Laboratorios RUBIO. The authors have no other
685 relevant affiliations or financial involvement with any organization or entity with a financial
686 interest in or financial conflict with the subject matter or materials discussed in the manuscript
687 apart from those disclosed.

688

689 **URLs**

690 GWAS Catalog: <https://www.ebi.ac.uk/gwas/>

691 FUMA: <https://fuma.ctglab.nl>

692 SynGO: <https://www.syngoportal.org>

693 Summary statistics: TBA

694

695 **Code & Summary Statistics**

696 Analysis plan: <https://www.action->

697 [europroject.eu/sites/default/files/Action%20AGG%20AP%20SOP.pdf](https://www.action-europroject.eu/sites/default/files/Action%20AGG%20AP%20SOP.pdf)

698 Code: -link to follow-

699 Summary Statistics: -link to follow-

700

701

702

703

704

705 **References**

- 706 Achenbach, T. M., Ivanova, M. Y., & Rescorla, L. A. (2017). Empirically based assessment and
 707 taxonomy of psychopathology for ages 1½–90+ years: Developmental, multi-informant, and
 708 multicultural findings. *Comprehensive Psychiatry*, 79, 4–18.
 709 <https://doi.org/10.1016/j.comppsy.2017.03.006>
- 710 American Psychiatric Association. (2013). *Diagnostic and Statistical Manual of Mental*
 711 *Disorders, Fifth Edition*.
- 712 Bartels, M., Hendriks, A., Mauri, M., Krapohl, E., Whipp, A., Bolhuis, K., Conde, L. C.,
 713 Luningham, J., Fung Ip, H., Hagenbeek, F., Roetman, P., Gatej, R., Lamers, A., Nivard, M.,
 714 van Dongen, J., Lu, Y., Middeldorp, C., van Beijsterveldt, T., Vermeiren, R., ... Boomsma,
 715 D. I. (2018). Childhood aggression and the co-occurrence of behavioural and emotional
 716 problems: results across ages 3–16 years from multiple raters in six cohorts in the EU-
 717 ACTION project. *European Child & Adolescent Psychiatry*, 27(9), 1105–1121.
 718 <https://doi.org/10.1007/s00787-018-1169-1>
- 719 Baselmans, B. M. L., Jansen, R., Ip, H. F., van Dongen, J., Abdellaoui, A., van de Weijer, M. P.,
 720 Bao, Y., Smart, M., Kumari, M., Willemsen, G., Hottenga, J.-J., Boomsma, D. I., de Geus,
 721 E. J. C., Nivard, M. G., & Bartels, M. (2019). Multivariate genome-wide analyses of the
 722 well-being spectrum. *Nature Genetics*, 51(3), 445–451. [https://doi.org/10.1038/s41588-018-](https://doi.org/10.1038/s41588-018-0320-8)
 723 [0320-8](https://doi.org/10.1038/s41588-018-0320-8)
- 724 Benner, C., Spencer, C. C. A., Havulinna, A. S., Salomaa, V., Ripatti, S., & Pirinen, M. (2016).
 725 FINEMAP: efficient variable selection using summary data from genome-wide association
 726 studies. *Bioinformatics*, 32(10), 1493–1501. <https://doi.org/10.1093/bioinformatics/btw018>
- 727 Boomsma, D. I. (2015). Aggression in children: unravelling the interplay of genes and
 728 environment through (epi)genetics and metabolomics. *Journal of Pediatric and Neonatal*
 729 *Individualized Medicine*, 4(2), e040251. <https://doi.org/https://doi.org/10.7363/040251>
- 730 Bulik-Sullivan, B. K., Finucane, H. K., Anttila, V., Gusev, A., Day, F. R., Loh, P.-R., Duncan,
 731 L., Perry, J. R. B., Patterson, N., Robinson, E. B., Daly, M. J., Price, A. L., & Neale, B. M.
 732 (2015). An atlas of genetic correlations across human diseases and traits. *Nature Genetics*,
 733 47(11), 1236–1241. <https://doi.org/10.1038/ng.3406>
- 734 Bulik-Sullivan, B. K., Loh, P.-R., Finucane, H. K., Ripke, S., Yang, J., Patterson, N., Daly, M. J.,
 735 Price, A. L., & Neale, B. M. (2015). LD Score regression distinguishes confounding from
 736 polygenicity in genome-wide association studies. *Nature Genetics*, 47(3), 291–295.
 737 <https://doi.org/10.1038/ng.3211>
- 738 Caci, H., Asherson, P., Donfrancesco, R., Faraone, S. V., Hervas, A., Fitzgerald, M., & Döpfner,
 739 M. (2015). Daily life impairments associated with childhood/adolescent attention-
 740 deficit/hyperactivity disorder as recalled by adults: results from the European Lifetime
 741 Impairment Survey. *CNS Spectrums*, 20(2), 112–121.
 742 <https://doi.org/10.1017/S1092852914000078>
- 743 Caci, H., Doepfner, M., Asherson, P., Donfrancesco, R., Faraone, S. V., Hervas, A., & Fitzgerald,
 744 M. (2014). Daily life impairments associated with self-reported childhood/adolescent
 745 attention-deficit/hyperactivity disorder and experiences of diagnosis and treatment: Results

- 746 from the European Lifetime Impairment Survey. *European Psychiatry*, 29(5), 316–323.
 747 <https://doi.org/10.1016/j.eurpsy.2013.10.007>
- 748 de Leeuw, C. A., Mooij, J. M., Heskes, T., & Posthuma, D. (2015). MAGMA: Generalized Gene-
 749 Set Analysis of GWAS Data. *PLOS Computational Biology*, 11(4), e1004219.
 750 <https://doi.org/10.1371/journal.pcbi.1004219>
- 751 Demontis, D., Walters, G. B., Athanasiadis, G., Walters, R., Therrien, K., Nielsen, T. T.,
 752 Farajzadeh, L., Voloudakis, G., Bendl, J., Zeng, B., Zhang, W., Grove, J., Als, T. D., Duan,
 753 J., Satterstrom, F. K., Bybjerg-Grauholm, J., Bækved-Hansen, M., Gudmundsson, O. O.,
 754 Magnusson, S. H., ... Børglum, A. D. (2023). Genome-wide analyses of ADHD identify 27
 755 risk loci, refine the genetic architecture and implicate several cognitive domains. *Nature*
 756 *Genetics*, 55(2), 198–208. <https://doi.org/10.1038/s41588-022-01285-8>
- 757 Demontis, D., Walters, R. K., Martin, J., Mattheisen, M., Als, T. D., Agerbo, E., Baldursson, G.,
 758 Belliveau, R., Bybjerg-Grauholm, J., Bækvad-Hansen, M., Cerrato, F., Chambert, K.,
 759 Churchhouse, C., Dumont, A., Eriksson, N., Gandal, M., Goldstein, J. I., Grasby, K. L.,
 760 Grove, J., ... Neale, B. M. (2019). Discovery of the first genome-wide significant risk loci
 761 for attention deficit/hyperactivity disorder. *Nature Genetics*, 51(1), 63–75.
 762 <https://doi.org/10.1038/s41588-018-0269-7>
- 763 Derks, E. M., Hudziak, J. J., Dolan, C. V., van Beijsterveldt, T. C. E. M., Verhulst, F. C., &
 764 Boomsma, D. I. (2008). Genetic and Environmental Influences on the Relation Between
 765 Attention Problems and Attention Deficit Hyperactivity Disorder. *Behavior Genetics*, 38(1),
 766 11–23. <https://doi.org/10.1007/s10519-007-9178-8>
- 767 Faraone, S. V., Asherson, P., Banaschewski, T., Biederman, J., Buitelaar, J. K., Ramos-Quiroga,
 768 J. A., Rohde, L. A., Sonuga-Barke, E. J. S., Tannock, R., & Franke, B. (2015). Attention-
 769 deficit/hyperactivity disorder. *Nature Reviews Disease Primers*, 1(1), 15020.
 770 <https://doi.org/10.1038/nrdp.2015.20>
- 771 Faraone, S. V., Bellgrove, M. A., Brikell, I., Cortese, S., Hartman, C. A., Hollis, C., Newcorn, J.
 772 H., Philipsen, A., Polanczyk, G. V., Rubia, K., Sibley, M. H., & Buitelaar, J. K. (2024).
 773 Attention-deficit/hyperactivity disorder. *Nature Reviews Disease Primers*, 10(1), 11.
 774 <https://doi.org/10.1038/s41572-024-00495-0>
- 775 Faraone, S. V., Biederman, J., & Mick, E. (2006). The age-dependent decline of attention deficit
 776 hyperactivity disorder: a meta-analysis of follow-up studies. *Psychological Medicine*, 36(2),
 777 159–165. <https://doi.org/10.1017/S003329170500471X>
- 778 Faraone, S. V., & Larsson, H. (2019). Genetics of attention deficit hyperactivity disorder.
 779 *Molecular Psychiatry*, 24(4), 562–575. <https://doi.org/10.1038/s41380-018-0070-0>
- 780 Goodman, R. (2001). Psychometric Properties of the Strengths and Difficulties Questionnaire.
 781 *Journal of the American Academy of Child & Adolescent Psychiatry*, 40(11), 1337–1345.
 782 <https://doi.org/10.1097/00004583-200111000-00015>
- 783 Grotzinger, A. D., Rhemtulla, M., de Vlaming, R., Ritchie, S. J., Mallard, T. T., Hill, W. D., Ip,
 784 H. F., Marioni, R. E., McIntosh, A. M., Deary, I. J., Koellinger, P. D., Harden, K. P., Nivard,
 785 M. G., & Tucker-Drob, E. M. (2019). Genomic structural equation modelling provides
 786 insights into the multivariate genetic architecture of complex traits. *Nature Human*
 787 *Behaviour*, 3(5), 513–525. <https://doi.org/10.1038/s41562-019-0566-x>
- 788 Ip, H. F., van der Laan, C. M., Krapohl, E. M. L., Brikell, I., Sánchez-Mora, C., Nolte, I. M., St
 789 Pourcain, B., Bolhuis, K., Palviainen, T., Zafarmand, H., Colodro-Conde, L., Gordon, S.,
 790 Zayats, T., Aliev, F., Jiang, C., Wang, C. A., Saunders, G., Karhunen, V., Hammerschlag, A.
 791 R., ... Boomsma, D. I. (2021). Genetic association study of childhood aggression across

- 792 raters, instruments, and age. *Translational Psychiatry*, *11*(1), 413.
 793 <https://doi.org/10.1038/s41398-021-01480-x>
- 794 Kan, K.-J., Dolan, C. V., Nivard, M. G., Middeldorp, C. M., van Beijsterveldt, C. E. M.,
 795 Willemsen, G., & Boomsma, D. I. (2013). Genetic and Environmental Stability in Attention
 796 Problems Across the Lifespan: Evidence From the Netherlands Twin Register. *Journal of*
 797 *the American Academy of Child & Adolescent Psychiatry*, *52*(1), 12–25.
 798 <https://doi.org/10.1016/j.jaac.2012.10.009>
- 799 Kan, K.-J., van Beijsterveldt, C. E. M., Bartels, M., & Boomsma, D. I. (2014). Assessing Genetic
 800 Influences on Behavior: Informant and Context Dependency as Illustrated by the Analysis of
 801 Attention Problems. *Behavior Genetics*, *44*(4), 326–336. [https://doi.org/10.1007/s10519-](https://doi.org/10.1007/s10519-014-9657-7)
 802 [014-9657-7](https://doi.org/10.1007/s10519-014-9657-7)
- 803 Koopmans, F., van Nierop, P., Andres-Alonso, M., Byrnes, A., Cijssouw, T., Coba, M. P.,
 804 Cornelisse, L. N., Farrell, R. J., Goldschmidt, H. L., Howrigan, D. P., Hussain, N. K., Imig,
 805 C., de Jong, A. P. H., Jung, H., Kohansalnodehi, M., Kramarz, B., Lipstein, N., Lovering, R.
 806 C., MacGillavry, H., ... Verhage, M. (2019). SynGO: An Evidence-Based, Expert-Curated
 807 Knowledge Base for the Synapse. *Neuron*, *103*(2), 217–234.e4.
 808 <https://doi.org/10.1016/j.neuron.2019.05.002>
- 809 Larsson, H., Anckarsater, H., Råstam, M., Chang, Z., & Lichtenstein, P. (2012). Childhood
 810 attention-deficit hyperactivity disorder as an extreme of a continuous trait: a quantitative
 811 genetic study of 8,500 twin pairs. *Journal of Child Psychology and Psychiatry*, *53*(1), 73–
 812 80. <https://doi.org/10.1111/j.1469-7610.2011.02467.x>
- 813 Lee, S. S., Humphreys, K. L., Flory, K., Liu, R., & Glass, K. (2011). Prospective association of
 814 childhood attention-deficit/hyperactivity disorder (ADHD) and substance use and
 815 abuse/dependence: A meta-analytic review. *Clinical Psychology Review*, *31*(3), 328–341.
 816 <https://doi.org/10.1016/j.cpr.2011.01.006>
- 817 Levy, F., Hay, D. A., McStephen, M., Wood, C., & Waldman, I. (1997). Attention-Deficit
 818 Hyperactivity Disorder: A Category or a Continuum? Genetic Analysis of a Large-Scale
 819 Twin Study. *Journal of the American Academy of Child & Adolescent Psychiatry*, *36*(6),
 820 737–744. <https://doi.org/10.1097/00004583-199706000-00009>
- 821 Liu, Q., Cirulli, E. T., Han, Y., Yao, S., Liu, S., & Zhu, Q. (2015). Systematic assessment of
 822 imputation performance using the 1000 Genomes reference panels. *Briefings in*
 823 *Bioinformatics*, *16*(4), 549–562. <https://doi.org/10.1093/bib/bbu035>
- 824 Marees, A. T., Smit, D. J. A., Ong, J.-S., MacGregor, S., An, J., Denys, D., Vorspan, F., van den
 825 Brink, W., & Derks, E. M. (2020). Potential influence of socioeconomic status on genetic
 826 correlations between alcohol consumption measures and mental health. *Psychological*
 827 *Medicine*, *50*(3), 484–498. <https://doi.org/10.1017/S0033291719000357>
- 828 Merwood, A., Greven, C. U., Price, T. S., Rijdsdijk, F., Kuntsi, J., McLoughlin, G., Larsson, H., &
 829 Asherson, P. J. (2013). Different heritabilities but shared etiological influences for parent,
 830 teacher and self-ratings of ADHD symptoms: an adolescent twin study. *Psychological*
 831 *Medicine*, *43*(9), 1973–1984. <https://doi.org/10.1017/S0033291712002978>
- 832 Middeldorp, C. M., Felix, J. F., Mahajan, A., & McCarthy, M. I. (2019). The Early Growth
 833 Genetics (EGG) and EARly Genetics and Lifecourse Epidemiology (EAGLE) consortia:
 834 design, results and future prospects. *European Journal of Epidemiology*, *34*(3), 279–300.
 835 <https://doi.org/10.1007/s10654-019-00502-9>
- 836 Middeldorp, C. M., Hammerschlag, A. R., Ouwens, K. G., Groen-Blokhuis, M. M., St. Pourcain,
 837 B., Greven, C. U., Pappa, I., Tiesler, C. M. T., Ang, W., Nolte, I. M., Vilor-Tejedor, N.,
 838 Bacelis, J., Ebejer, J. L., Zhao, H., Davies, G. E., Ehli, E. A., Evans, D. M., Fedko, I. O.,

GWAMA OF ADHD SYMPTOMS AND DIAGNOSIS

- 839 Guxens, M., ... Boomsma, D. I. (2016). A Genome-Wide Association Meta-Analysis of
840 Attention-Deficit/Hyperactivity Disorder Symptoms in Population-Based Pediatric Cohorts.
841 *Journal of the American Academy of Child & Adolescent Psychiatry*, 55(10), 896-905.e6.
842 <https://doi.org/10.1016/j.jaac.2016.05.025>
- 843 Minică, C. C., Dolan, C. V., Kampert, M. M. D., Boomsma, D. I., & Vink, J. M. (2015).
844 Sandwich corrected standard errors in family-based genome-wide association studies.
845 *European Journal of Human Genetics*, 23(3), 388–394. <https://doi.org/10.1038/ejhg.2014.94>
- 846 Schipper, M., de Leeuw, C. A., Maciel, B. A. P. C., Wightman, D. P., Hubers, N., Boomsma, D.
847 I., O'Donovan, M. C., & Posthuma, D. (2023). A framework for accurate prediction of
848 effector genes in genetic loci for complex traits. *MedRxiv*.
- 849 Tucker, G., Loh, P.-R., MacLeod, I. M., Hayes, B. J., Goddard, M. E., Berger, B., & Price, A. L.
850 (2015). Two-Variance-Component Model Improves Genetic Prediction in Family Datasets.
851 *The American Journal of Human Genetics*, 97(5), 677–690.
852 <https://doi.org/10.1016/j.ajhg.2015.10.002>
- 853 Wang, X., Walker, A., Revez, J. A., Ni, G., Adams, M. J., McIntosh, A. M., Visscher, P. M.,
854 Wray, N. R., Wray, N. R., Ripke, S., Mattheisen, M., Trzaskowski, M., Byrne, E. M.,
855 Abdellaoui, A., Adams, M. J., Agerbo, E., Air, T. M., Andlauer, T. F. M., Bacanu, S.-A., ...
856 Sullivan, P. F. (2023). Polygenic risk prediction: why and when out-of-sample prediction R²
857 can exceed SNP-based heritability. *The American Journal of Human Genetics*, 110(7),
858 1207–1215. <https://doi.org/10.1016/j.ajhg.2023.06.006>
- 859 Watanabe, K., Taskesen, E., van Bochoven, A., & Posthuma, D. (2017). Functional mapping and
860 annotation of genetic associations with FUMA. *Nature Communications*, 8(1), 1826.
861 <https://doi.org/10.1038/s41467-017-01261-5>
- 862 Weeks, E. M., Ulirsch, J. C., Cheng, N. Y., Trippe, B. L., Fine, R. S., Miao, J., Patwardhan, T.
863 A., Kanai, M., Nasser, J., Fulco, C. P., Tashman, K. C., Aguet, F., Li, T., Ordovas-
864 Montanes, J., Smillie, C. S., Biton, M., Shalek, A. K., Ananthakrishnan, A. N., Xavier, R. J.,
865 ... Finucane, H. K. (2023). Leveraging polygenic enrichments of gene features to predict
866 genes underlying complex traits and diseases. *Nature Genetics*, 55(8), 1267–1276.
867 <https://doi.org/10.1038/s41588-023-01443-6>
868