

# Genetic variants associated with longevity in long-living Indians

---

Sandhya Kiran Pemmasani, Shakthiraju R G, Suraj V, Raunaq Bhattacharya, Chetan Patel, Anil Kumar Gupta, Anuradha Acharya

Mapmygenome India Limited

## ABSTRACT

Genetic factors play a significant role in determining an individual's longevity. The present study was aimed at identifying the genetic variants associated with longevity in Indian population. Long living individuals (LLIs), aged 85+, were compared with younger controls, aged 18-49 years, using data from GenomeDB, a genetic database of Indians living in India. An in-house developed custom chip, having variants associated with various cancers, cardiovascular, neurological, gastro-intestinal, metabolic and auto-immune disorders, was used to generate genotype data. Logistic regression analysis with top five genetic principal components as covariates resulted in 11 variants to be significantly associated with longevity at a p-value threshold of  $5 \times 10^{-4}$ . Alleles associated with slower heart rate (rs365990, MYH6), lower risk of osteoporosis and short body height (rs2982570, ESR1), decreased risk of schizophrenia (rs1339227, RIMS1-KCNQ5) and decreased risk of anxiety and neuroticism (rs391957, HSPA5) were found to have higher frequency in LLIs. Alleles associated with increased risk of atrial fibrillation (rs3903239, GORAB-PRRX1) and biliary disorders (rs2002042, ABCC2) were found have lower frequency. The G allele of rs2802292 from FOXO3A gene, associated with longevity in Japanese, German and French centenarians, was also found to be significant in this population ( $P=0.027$ ). Pathway enrichment analysis revealed that the genes involved in oxidative stress, apoptosis, DNA damage repair, glucose metabolism and energy metabolism were significantly involved in increasing the longevity. Results of our study demonstrate the genetic basis of healthy aging and longevity in the population.

## INTRODUCTION

Longevity and healthy ageing are complex phenomena influenced by various factors, like socioeconomic status, nutrition, lifestyle, physical activity, gender and genetics. Among them, genetics plays a major role, and its influence is estimated to be around 25-40 percent [1]. Studies indicate that offspring of long-lived individuals tend to lead a healthy and longer life compared to general population [2]. This could be due to inheritance of genetic variants associated with healthy lipid profiles, enhanced insulin sensitivity, slower cognitive decline and lower incidence of age-related diseases, like Alzheimer's and cardiovascular diseases [3-7]. There are also a certain category of variants that are identified to be prevalent in long-living individuals through multiple genome-wide association studies (GWAS), candidate studies and meta-analyses related to longevity. They are the variants from Forkhead Box O3A (FOXO3A) and Apolipoprotein E (APOE) genes that are part of pathways associated with cell apoptosis, metabolism and oxidative stress [8-10]. Variants that decrease the length of telomeres also influence an individual's lifespan [11-12].

Genetic associations observed in one population might not reflect in other populations due to variable frequencies of effect allele across populations. In developed countries or high-income countries, reaching 100 years of age is easier compared to low-income countries, and hence selection pressure is less. This leads to relatively lesser frequencies of effect alleles [13]. Also, age thresholds to define "cases" or "long-living individuals" change as per the population under study. Studies done on isolated populations (Blue Zones) show that the environmental factors work in synergy with genetics in contributing to exceptional longevity in those populations [14]. So, for any study done on genetics of longevity, it is necessary to understand longevity of the individuals in that population, define the age thresholds and take cases and controls from the same environment.

According to the latest estimates of Longitudinal Aging Study in India (LASI - 2019), India has world's second-largest number of older people, aged 60 and above [15]. It is further estimated that the number will double by 2050. With improved access to vaccines, antibiotics and clean water, and increased awareness on healthy diets and physical activity, life expectancy at birth has improved from 62.1 years in 2000 to 70.8 years in 2019 [16]. Despite this, only 0.4% of the population reaches the age of 85 years. Understanding the genetics of these long living individuals (LLIs), of age 85+, in India is still a challenge due to lack of genetic research on the population. It is necessary to conduct genome-wide studies by selecting cases and controls from the individuals living in India. Identifying genetic variants associated with longevity, and in turn inferring factors that influence longevity helps in recommending strategies for healthy aging at population-wide and an individual level.

In the present study, we have identified variants that have significantly higher frequency in long-living Indians, compared to younger controls, using GenomagaDB of Mapmygenome [17]. Through literature mining on the identified variants we have inferred the factors that help in healthy aging. To our knowledge this is the first genetic study to investigate variants associated with longevity in Indian population.

## MATERIALS AND METHODS

### Study Data

Participants in the current study were taken from GenomegaDB of Mapmygenome, which is a genotype-phenotype database of Indians living in India. Cases were defined as participants with self-reported age of greater than 85 years at the time of sample collection. Controls were the participants with 18-49 years of age at the time of sample collection. A total of 133 cases and 1155 controls were considered for the present study, to investigate the variants associated with longevity in Indian population. Genotyping was done using a custom chip developed in-house at Mapmygenome. It has 8,768 variants associated with various phenotypes, like cancers, cardiovascular, neurological, gastrointestinal, metabolic and autoimmune disorders. Written informed consent was taken from each participant. Research was carried out in compliance with the Helsinki declaration and the procedures had been approved by internal bio-safety committee at Mapmygenome.

### Data pre-processing

Variants with genotype call rate <90%, minor allele frequency (MAF) <5% and are not in Hardy-Weinberg Equilibrium (HWE) in controls ( $P_{HWE} \leq 0.0001$ ) were removed from further analysis. Also, the participants with genotype call rate <95% were removed from the analysis. In the case of relatives, only one participant, preferably the elder one was retained in the analysis. Participants who were genetic outliers, defined as being more than three standard deviations away from the mean of top five genetic principal components, were excluded from the analysis. R [18] and PLINK [19] software were used for all the data pre-processing.

### Association analysis

The association of each variant with longevity was assessed using additive logistic regression with Firth's bias reduction available in PLINK software. Top five genetic principal components were added as covariates in the model, to correct for population stratification. Variants with  $p$ -value  $\leq 5 \times 10^{-4}$  were considered to be significantly associated with longevity. Manhattan plots were generated using *ggplot* package of R v4.2. Gene information was obtained through NCBI's E-utilities [20].

### Pathway enrichment analysis

Genes corresponding to the significant SNPs were mapped to KEGG (Kyoto Encyclopedia of Genes and Genomes) pathway maps to understand the possible functional processes associated with longevity. DAVID Functional Annotation Tool v2024q1 [21-22] and ClueGO v2.5.10 [23] were used to perform pathway enrichment analysis and to draw interacting networks.

## RESULTS

Table 1 gives information on characteristics of participants included in the analysis. Mean age of cases was found to be 93.4 years, and mean age of controls was found to be 36.4 years. There was equal representation of participants from both the genders. Out of 8,768 variants, 6,300 autosomal SNPs passed quality control filters, and were finally considered in the analysis.

**Table 1 Characteristics of study participants**

	Cases	Controls
Number of participants	133	1155
Age, mean (SD)	93.4 (6.0)	36.4 (7.7)
Male, n (%)	65 (49%)	647 (56%)

Association analysis using additive logistic regression along with top five genetic principal components resulted in 11 variants to be significantly associated with longevity, at a p-value threshold of  $5 \times 10^{-4}$ . Table 2 gives information on minor allele, major allele, minor allele frequency (MAF), p-value and odds ratio obtained for these variants. Figure 1 shows Manhattan plot of associations. 492 variants that have suggestive association at a nominal p-value of 0.05 are given in Supplementary Table 1.

**Table 2 Variants significantly associated with longevity in Indian population**

ID	Chr	Position	Minor allele	Major allele	MAF in Cases	MAF in Controls	P	OR	95% CI	Gene/Nearest Gene
rs12083278	1	31151182	G	C	0.149	0.084	3.125E-04	2.119	1.409 - 3.188	
rs1877455	1	114556471	T	C	0.297	0.433	1.489E-04	0.588	0.447 - 0.774	TRIM33-BCAS2
rs3903239	1	170600176	G	A	0.128	0.232	2.500E-04	0.498	0.343 - 0.723	GORAB-PRRX1
rs9363918	6	68432116	T	G	0.624	0.484	4.700E-05	1.716	1.323 - 2.226	LINC02549-ADGRB3
rs1339227	6	72445999	T	C	0.493	0.375	3.602E-04	1.604	1.237 - 2.079	RIMS1-KCNQ5
rs2982570	6	151692613	T	C	0.290	0.195	3.882E-04	1.692	1.265 - 2.262	ESR1
rs10970985	9	32453121	C	G	0.267	0.398	5.984E-05	0.56	0.422 - 0.743	ACO1
rs391957	9	125241745	T	C	0.417	0.311	4.831E-04	1.586	1.224 - 2.055	HSPA5
rs2002042	10	99828174	T	C	0.105	0.197	4.309E-04	0.487	0.326 - 0.727	ABCC2
rs2762051	13	50261579	T	C	0.150	0.071	3.220E-05	2.222	1.525 - 3.237	DLEU1
rs365990	14	23392602	G	A	0.365	0.263	3.965E-04	1.625	1.242 - 2.126	MYH6

ID: rsID of variant; Chr: Chromosome number; Position: GRCh38 position; MAF in Cases: Frequency of minor allele in Cases; MAF in Controls: Frequency of minor allele in Controls; P: P-value calculated from logistic regression after adjusting for top five principal components; OR: Odds Ratio for minor allele; 95% CI: 95% Confidence Interval of Odds Ratio;




Figure 1 Manhattan plot of association analysis results

Statistical significance of the association, as measured by the negative logarithm of the p-value, is shown on y-axis. Larger values correspond to smaller p-values. X-axis represents the chromosomal locations. Red horizontal dashed line indicates the p-value threshold of  $5 \times 10^{-4}$ . Significant variants are highlighted in red, with their information in rectangular green boxes.

Literature mining on significantly associated SNPs revealed that they are implicated in various diseases, indirectly affecting the lifespan. As can be seen in Table 1, G allele of rs3903239, known to be associated with atrial fibrillation [24-25], decreases the chances of surviving to older age by 0.498 times (95% CI: 0.343 - 0.723). Its frequency is very low in LLIs compared to younger controls. Significant association was also observed for rs1339227, an intergenic SNP between RIMS1 and KCNQ5 genes. Several studies indicated that the C allele of this SNP is associated with schizophrenia [26-27]. Carriers of the other allele, T allele, at this locus have 1.604 times higher chances of surviving up to the age of 85 (95% CI: 1.237 - 2.079). The rs2982570, from ESR1 gene that is involved in growth and metabolism, is known to be associated with the risk of osteoporosis [28]. The non-effect allele, T allele, at this locus has an odds ratio of 1.692 (95%CI: 1.265 - 2.261) for longevity compared to the risk allele. The rs10970985, from ACO1 gene that is involved in controlling the iron levels in the cells, is found to have lower frequency of C allele in LLIs. The C allele at this locus decreases the survival to older age by 0.56 times (95% CI: 0.422 - 0.743). Another variant, rs391957 from HSPA5 gene, which is associated with decreased anxiety and neuroticism [29-30], was found to have high frequency in LLIs, with an odds ratio of 1.586 (95%CI: 1.224 - 2.055) for longer survival. Rs2002042 from ABCC2 gene, which is known to be associated with non-alcoholic fatty liver disease [31], has low frequency of risk allele in LLIs compared to controls. Presence of risk allele (T allele) decreases the chances of survival till 85+ by 0.486 times (95%CI: 0.326 - 0.727). Significant association with longevity was also observed for rs365990 from MYH6 gene that is involved in cardiac muscle contraction. G allele at this locus is known to be associated with prolonged PR interval

and slower heart rate [32-33]. Carriers of G allele have 1.625 times higher chances of surviving to older age (95%CI: 1.242 - 2.126) than the carriers of A allele. The additional four SNPs in Table 1, rs12083278 [C] associated with severity of COVID-19 [34], rs1877455 [T] associated with autism along with other SNPs as haplotype [35], rs9363918 [T] associated with pancreatic cancer [36], and rs2762051 [T] associated with celiac disease [37], have limited literature support for their associations, and further research studies are required to understand their role in increasing the lifespan.

Previous studies showed multiple SNPs from FOXO3A gene to be associated with longer lifespan. In particular, the G-allele of rs2802292 showed significant presence in Japanese, German and French centenarians [38-40]. In our study, this SNP reached nominal significance with a p-value of 0.027 and an OR of 1.34. Other SNPs from FOXO3A (rs9400239, rs2153960, rs4946935 and rs4946936) did not show statistical significance, but the direction of association is consistent with other studies. Figure 2 shows the results of association analysis on SNPs from FOXO3A gene.




Figure 2 Association analysis on SNPs from FOXO3A gene

Pathway enrichment analysis was performed on 492 variants that have suggestive association with longevity (p-value < 0.05). 84 variants located in intergenic regions did not map to any genes, and were excluded from the analysis. Significantly enriched KEGG pathways identified through DAVID Functional Analysis tool, with p-value threshold of 0.05, are given in Supplementary Table 2. Figure 3 shows the most significant pathways identified through ClueGO (P-value <  $5 \times 10^{-4}$ ). Significant pathways identified by both the tools were longevity regulating pathway, FoxO signaling pathway, adipocytokine signaling pathway and non-alcoholic fatty liver disease pathway. Some of the other pathways identified by the tools included insulin resistance, cholesterol metabolism, inflammatory bowel disease and PI3K-Akt signaling pathway.





Figure 3 Pathway analysis through ClueGO. A) Barograph showing the % of genes mapped to each pathway term. Colors indicate the grouping of pathways done based on Kappa score B) Network of pathways grouped along with Longevity regulating pathway.

## DISCUSSION

In the present study, we analyzed genotype data of long-living Indians, aged 85+, to identify genetic polymorphisms associated with longevity in the population. By mapping the identified variants to genes, pathways and their functions, and by doing thorough literature search we inferred the factors that influence longevity and healthy aging. The alleles that increase the risk of atrial fibrillation, biliary disorders, schizophrenia, anxiety and neuroticism were in low frequency in long-living individuals compared to younger controls. Alleles associated with slower heart rate, lower risk of bone fractures, short body height were in high frequency.

Atrial fibrillation is one of the risk factors that reduce the life expectancy. It contributes to complications such as stroke and heart failure. Several genetic research studies indicated the association of rs3903239, an SNP in the upstream of PRRX1 gene, with atrial fibrillation. PRRX1 (Paired related homeobox 1) regulates muscle creatine kinase and controls the growth and development of mesodermal muscles such as the heart. SNPs in the upstream of PRRX1 may reduce the expression of the gene, and shorten the atrial action potential duration [41]. Ellinor et al [24] conducted a GWAS meta-analysis in individuals of European ancestry, and found G allele of rs3903239 to be significantly associated with atrial fibrillation. The association was also observed in Japanese and Korean population [25]. In the present study we observed a lower frequency of G allele in LLIs, indicating the lower risk of cardiovascular disorders, and increased longevity.

The T allele of rs1339227, which decreases the risk of schizophrenia, is present in higher frequency in LLIs. Schizophrenia Working Group of the Psychiatric Genomics Consortium identified significant association of rs1339227 with schizophrenia [26]. The T allele was found to decrease the risk of disease with an odds ratio of 0.942. Yao et al [27] found a decreased effect of T allele in the association of schizophrenia. Higher frequency of T allele in LLIs suggests protective association of this SNP in increasing the lifespan.

Estrogen receptor 1 gene (ESR1) encodes an estrogen receptor (ER-alpha) that plays a major role in bone metabolism and maintenance of the skeletal system. Several genome-wide association studies have found variations in this gene to be associated with osteoporosis and risk of fractures via low bone mineral density (BMD). Estrogen-replacement therapy in post-menopausal women is known to prevent bone loss and osteoporosis. Trajanoska et al [42] found the C allele of rs2982570 from ESR1 to be associated with increased risk of fractures and decreased femoral neck BMD and lumbar spine BMD. In the present study, we observed low frequency of C-allele in LLIs, indicating lower risk of skeletal and bone related problems. The other allele (T-allele), which is also found to be associated with decreased body height [43], is in high frequency in LLIs. Multiple research studies found a negative correlation between height and longevity. Shorter people tend to have higher resistance to chronic diseases, especially in middle age [44].

Anemia is also one of the major health concerns in older adults. It is associated with cognitive deficits and reduced physical performance. Several genes and genetic variants are known to affect iron levels, hemoglobin levels and anemia. ACO1 is one such gene that encodes a bifunctional protein which controls the levels of iron within the cells. We observed that the variant in this gene, rs10970985, has lower frequency of C allele in LLIs compared to controls. Further research studies are required to understand the role of this variant.

HSPA5 gene (heat shock protein family A (Hsp70) member 5) encodes immunoglobulin protein that is an essential component of folding/unfolding processes and regulation of Ca<sup>2+</sup> homeostasis in the endoplasmic reticulum (ER) [45]. Disturbed homeostasis leads to ER stress, causing inflammation, metabolic disorders and neurodegenerative diseases. It is also associated with anxiety disorders such as depression and post-traumatic stress disorder (PTSD) [46]. Research studies found T-allele of rs391957 from HSPA5 gene to be a protective factor for anxiety and neuroticism [29-30]. In the present investigation we observed the frequency of T-allele to be higher in LLIs compared to younger controls, indicating that lesser anxiety leads to increased lifespan.

ABCC2 (ATP binding cassette subfamily C member 2) gene encodes a protein called multidrug resistance protein 2 (MRP2) which is involved in metabolism and clearance of certain drugs from organs and



tissues. It is also known to play a major role in biliary transport. Mutations in ABCC2 gene were observed in patients with Dubin-Johnson syndrome, an autosomal recessive disorder characterized by conjugated hyperbilirubinemia. T allele of rs2002042 from ABCC2 gene was found to be associated with intrahepatic cholestasis of pregnancy and nonalcoholic fatty liver disease [47,31]. In our study, we observed lower frequency of T allele in LLIs compared to controls, indicating undisturbed biliary transport, good enterohepatic circulation of bile acids and healthy gut leading to longer lifespan.

Carriers of G allele at rs365990 have 1.725 times higher chances of surviving to older age. Previous studies indicate that G allele of rs365990 is associated with decrease in heart rate [32-33]. Low resting heart rate results in increased lifespan due to reduced stress on arterial cells, reduced stiffening of aorta and reduced cardiovascular abnormalities [48-49]. Faster heart rate is related to high metabolic rate, development of free radicals, oxidative stress and faster aging. Studies also indicate that heart rate lowering through medication or selective sinus node inhibition increases the lifespan.

Our pathway enrichment analysis confirmed that the molecular processes associated or pathways with longevity are significantly different in LLIs. Pathways with the genes (through SNPs) involved in oxidative stress, apoptosis, DNA damage repair, glucose metabolism and energy metabolism are differentially regulated in LLIs. IRS2 and PRKAA1 genes that are related to regulation of insulin secretion and glucose levels are common across most of the significant pathways. Several previous studies indicated the role of PI3K-Akt signaling pathway in ageing [50, 51]. In addition to that, our study revealed pathways related to non-alcoholic fatty liver disease (NFLD), inflammatory bowel disease (IBD) and inflammation to be associated with ageing. Screening the variants associated with these disorders early on in life helps in recommending preventive measures, and in promoting healthy ageing.

Major limitation in our study is the sample size. Increasing the sample size will increase the statistical power of the study, and may identify additional variants that increase our understanding of factors affecting longevity. Also, gender-specific analysis is not feasible due to small sample size. But our results can provide some indication on variants and factors associated with longevity in the population, and help in further follow-up studies with larger sample sizes. Additional information on clinical and lifestyle variables, such as BMI, cholesterol levels, diet patterns, physical activity, smoking and alcohol consumption help in identifying the variants associated with interaction of those variables with longevity.

Allele frequencies of the longevity-associated SNPs vary among populations studied under 1000 Genomes project (Supplementary Table 3). Major alleles in one population were observed to be minor alleles in other populations. For example, G allele of rs3903239 associated with atrial fibrillation is a minor allele in SAS, but major allele in EAS (East Asians). Similarly, alleles associated with longevity from the SNPs of FOXO3 gene have in general higher frequency in AFR and SAS populations. This indicates the need for replication studies in different populations to confirm the associations in each population.

In conclusion, our study identified genetic variants associated with longevity in Indian population that is underrepresented in genetic research. Clinical and lifestyle factors inferred through the identified variants can guide in recommending strategies for healthy aging and longevity at population-wide and an individual level.

## REFERENCES

1. Bin-Jumah MN, Nadeem MS, Gilani SJ, et al. Genes and Longevity of Lifespan. *Int J Mol Sci*. 2022;23(3):1499. Published 2022 Jan 28. doi:10.3390/ijms23031499
2. Gudmundsson H, Gudbjartsson DF, Frigge M, Gulcher JR, Stefánsson K. Inheritance of human longevity in Iceland. *Eur J Hum Genet*. 2000;8(10):743-749. doi:10.1038/sj.ejhg.5200527
3. Pilling LC, Kuo CL, Sicinski K, et al. Human longevity: 25 genetic loci associated in 389,166 UK biobank participants. *Aging (Albany NY)*. 2017;9(12):2504-2520. doi:10.18632/aging.101334
4. Vaarhorst AA, Beekman M, Suchiman EH, et al. Lipid metabolism in long-lived families: the Leiden Longevity Study. *Age (Dordr)*. 2011;33(2):219-227. doi:10.1007/s11357-010-9172-6
5. Wijsman CA, Rosing MP, Streefland TC, et al. Familial longevity is marked by enhanced insulin sensitivity. *Aging Cell*. 2011;10(1):114-121. doi:10.1111/j.1474-9726.2010.00650.x
6. Dutta A, Henley W, Robine JM, et al. Aging children of long-lived parents experience slower cognitive decline. *Alzheimers Dement*. 2014;10(5 Suppl):S315-S322. doi:10.1016/j.jalz.2013.07.002
7. Ash, Arlene S et al. "Are Members of Long-Lived Families Healthier Than Their Equally Long-Lived Peers? Evidence From the Long Life Family Study." *The journals of gerontology. Series A, Biological sciences and medical sciences* vol. 70,8 (2015): 971-6. doi:10.1093/gerona/glv015
8. Willcox, Bradley J et al. "FOXO3A genotype is strongly associated with human longevity." *Proceedings of the National Academy of Sciences of the United States of America* vol. 105,37 (2008): 13987-92. doi:10.1073/pnas.0801030105
9. APOE
10. Deelen, J., Evans, D.S., Arking, D.E. et al. A meta-analysis of genome-wide association studies identifies multiple longevity genes. *Nat Commun* 10, 3669 (2019). <https://doi.org/10.1038/s41467-019-11558-2>
11. Mohamadkhani, Ashraf et al. "Significant SNPs Related to Telomere Length and Hepatocellular Carcinoma Risk in Chronic Hepatitis B Carriers." *Asian Pacific journal of cancer prevention : APJCP* vol. 19,3 585-590. 27 Mar. 2018, doi:10.22034/APJCP.2018.19.3.585
12. Lili, Ma et al. "Genetic variations associated with telomere length affect the risk of gastric carcinoma." *Medicine* vol. 99,23 (2020): e20551. doi:10.1097/MD.00000000000020551
13. Nygaard M, Lindahl-Jacobsen R, Soerensen M, et al. Birth cohort differences in the prevalence of longevity-associated variants in APOE and FOXO3A in Danish long-lived individuals. *Exp Gerontol*. 2014;57:41-46. doi:10.1016/j.exger.2014.04.018
14. Poulain M, Herm A, Errigo A, et al. Specific features of the oldest old from the Longevity Blue Zones in Ikaria and Sardinia. *Mech Ageing Dev*. 2021;198:111543. doi:10.1016/j.mad.2021.111543
15. Bloom, D.E., Sekher, T.V. & Lee, J. Longitudinal Aging Study in India (LASI): new data resources for addressing aging in India. *Nat Aging* 1, 1070–1072 (2021). <https://doi.org/10.1038/s43587-021-00155-y>
16. <https://data.who.int/countries/356>
17. <https://mapmygenome.in/>
18. R Core Team (2022). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.
19. Purcell S, Neale B, Todd-Brown K, et al. PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet*. 2007;81(3):559-575. doi:10.1086/519795
20. Kans J. Entrez Direct: E-utilities on the Unix Command Line. 2013 Apr 23 [Updated 2024 Jan 25]. In: Entrez Programming Utilities Help [Internet]. Bethesda (MD): National Center for Biotechnology Information (US); 2010-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK179288/>

21. B.T. Sherman, M. Hao, J. Qiu, X. Jiao, M.W. Baseler, H.C. Lane, T. Imamichi and W. Chang. DAVID: a web server for functional enrichment analysis and functional annotation of gene lists (2021 update). *Nucleic Acids Research*. 23 March 2022. doi:10.1093/nar/gkac194.[PubMed]
22. Huang DW, Sherman BT, Lempicki RA. Systematic and integrative analysis of large gene lists using DAVID Bioinformatics Resources. *Nature Protoc*. 2009;4(1):44-57. [PubMed]
23. Bindea G, Mlecnik B, Hackl H, et al. ClueGO: a Cytoscape plug-in to decipher functionally grouped gene ontology and pathway annotation networks. *Bioinformatics*. 2009;25(8):1091-1093. doi:10.1093/bioinformatics/btp101
24. Ellinor PT, Lunetta KL, Albert CM, et al. Meta-analysis identifies six new susceptibility loci for atrial fibrillation. *Nat Genet*. 2012;44(6):670-675. Published 2012 Apr 29. doi:10.1038/ng.2261
25. Lee JY, Kim TH, Yang PS, et al. Korean atrial fibrillation network genome-wide association study for early-onset atrial fibrillation identifies novel susceptibility loci. *Eur Heart J*. 2017 Sep 7;38(34):2586-2594. doi: 10.1093/eurheartj/ehx213.
26. Schizophrenia Working Group of the Psychiatric Genomics Consortium. Biological insights from 108 schizophrenia-associated genetic loci. *Nature*. 2014 Jul 24;511(7510):421-7. doi: 10.1038/nature13595. Epub 2014 Jul 22.
27. Yao X, Glessner JT, Li J, et al. Integrative analysis of genome-wide association studies identifies novel loci associated with neuropsychiatric disorders. *Transl Psychiatry*. 2021 Jan 21;11(1):69. doi: 10.1038/s41398-020-01195-5.
28. Koromani F, Trajanoska K, Rivadeneira F, Oei L. Recent Advances in the Genetics of Fractures in Osteoporosis. *Front Endocrinol (Lausanne)*. 2019;10:337. Published 2019 Jun 4. doi:10.3389/fendo.2019.00337
29. Thorp, J.G., Campos, A.I., Grotzinger, A.D. et al. Symptom-level modelling unravels the shared genetic architecture of anxiety and depression. *Nat Hum Behav* 5, 1432–1442 (2021). <https://doi.org/10.1038/s41562-021-01094-9>
30. Nagel M, Speed D, van der Sluis S, Østergaard SD. Genome-wide association study of the sensitivity to environmental stress and adversity neuroticism cluster. *Acta Psychiatrica Scandinavica*. 2020 May;141(5):476-478. DOI: 10.1111/acps.13155. PMID: 31972866.
31. Sookoian S, Castaño G, Gianotti TF, Gemma C, Pirola CJ. Polymorphisms of MRP2 (ABCC2) are associated with susceptibility to nonalcoholic fatty liver disease. *J Nutr Biochem*. 2009 Oct;20(10):765-70. doi: 10.1016/j.jnutbio.2008.07.005. Epub 2008 Oct 15. PMID: 18926681.
32. Eijgelsheim, Mark et al. Genome-wide association analysis identifies multiple loci related to resting heart rate. *Human molecular genetics vol. 19,19 (2010): 3885-94*. doi:10.1093/hmg/ddq303
33. Holm H, Gudbjartsson DF, Sulem P, et al. A rare variant in MYH6 is associated with high risk of sick sinus syndrome. *Nat Genet*. 2011;43(4):316-320. Published 2011 Mar 6. doi:10.1038/ng.781
34. Dite GS, Murphy NM, Allman R. An integrated clinical and genetic model for predicting risk of severe COVID-19: A population-based case-control study. *PLoS One*. 2021 Feb 16;16(2):e0247205. doi: 10.1371/journal.pone.0247205. PMID: 33592063; PMCID: PMC7886160.
35. Xia K, Guo H, Hu Z, et al. Common genetic variants on 1p13.2 associate with risk of autism. *Mol Psychiatry*. 2014;19(11):1212-1219. doi:10.1038/mp.2013.146
36. Wu C, Miao X, Huang L, et al. Genome-wide association study identifies five loci associated with susceptibility to pancreatic cancer in Chinese populations. *Nat Genet*. 2011;44(1):62-66. Published 2011 Dec 11. doi:10.1038/ng.1020
37. Dubois PC, Trynka G, Franke L, et al. Multiple common variants for celiac disease influencing immune gene expression. *Nat Genet*. 2010 Apr;42(4):295-302. doi: 10.1038/ng.543. Epub 2010 Feb 28. Erratum in: *Nat Genet*.2010 May;42(5):465.
38. Willcox BJ, Donlon TA, He Q, et al. FOXO3A genotype is strongly associated with human longevity. *Proc Natl Acad Sci U S A*. 2008;105(37):13987-13992. doi:10.1073/pnas.0801030105

39. Flachsbart F, Caliebe A, Kleindorp R, et al. Association of FOXO3A variation with human longevity confirmed in German centenarians. *Proc Natl Acad Sci U S A*. 2009;106(8):2700-2705. doi:10.1073/pnas.0809594106
40. Sun, L., Hu, C., Zheng, C. et al. FOXO3 variants are beneficial for longevity in Southern Chinese living in the Red River Basin: A case-control study and meta-analysis. *Sci Rep* 5, 9852 (2015). <https://doi.org/10.1038/srep09852>
41. Tucker NR, Dolmatova EV, Lin H, et al. Diminished PRRX1 Expression Is Associated With Increased Risk of Atrial Fibrillation and Shortening of the Cardiac Action Potential. *Circ Cardiovasc Genet*. 2017 Oct;10(5):e001902. doi: 10.1161/CIRCGENETICS.117.001902.
42. Trajanoska K, Morris JA, Oei L, et al. Assessment of the genetic and clinical determinants of fracture risk: genome wide association and mendelian randomisation study. *BMJ (Clinical Research ed.)*. 2018 Aug;362:k3225. DOI: 10.1136/bmj.k3225. PMID: 30158200; PMCID: PMC6113773.
43. Yengo, Loïc et al. A saturated map of common genetic variants associated with human height. *Nature* vol. 610,7933 (2022): 704-712. doi:10.1038/s41586-022-05275-y
44. Chmielewski PP. The association between body height and longevity: evidence from a national population sample. *Folia Morphol (Warsz)*. Published online January 27, 2023. doi:10.5603/FM.a2023.0005
45. Jie Wang, Jessica Lee, David Liem, Peipei Ping (2017) HSPA5 Gene encoding Hsp70 chaperone BiP in the endoplasmic reticulum, *Gene*, Volume 618, 2017, Pages 14-23, ISSN 0378-1119, <https://doi.org/10.1016/j.gene.2017.03.005>.
46. Mao, J., Hu, Y., Ruan, L., Ji, Y., & Lou, Z. (2019). Role of endoplasmic reticulum stress in depression (Review). *Molecular Medicine Reports*, 20, 4774-4780. <https://doi.org/10.3892/mmr.2019.10789>
47. Ref7: Sookoian S, Castano G, Burgueno A, Gianotti TF, Pirola CJ. Association of the multidrug-resistance-associated protein gene (ABCC2) variants with intrahepatic cholestasis of pregnancy. *J Hepatol*. 2008;48:125–132.
48. Boudoulas KD, Borer JS, Boudoulas H. Heart Rate, Life Expectancy and the Cardiovascular System: Therapeutic Considerations. *Cardiology*. 2015;132(4):199-212. doi:10.1159/000435947
49. Jensen MT. Resting heart rate and relation to disease and longevity: past, present and future. *Scand J Clin Lab Invest*. 2019;79(1-2):108-116. doi:10.1080/00365513.2019.1566567
50. McIntyre RL, Liu YJ, Hu M, et al. Pharmaceutical and nutraceutical activation of FOXO3 for healthy longevity. *Ageing Res Rev*. 2022 Jun;78:101621. doi: 10.1016/j.arr.2022.101621. Epub 2022 Apr 11. PMID: 35421606.
51. Long HZ, Cheng Y, Zhou ZW, et al. PI3K/AKT Signal Pathway: A Target of Natural Products in the Prevention and Treatment of Alzheimer's Disease and Parkinson's Disease. *Front Pharmacol*. 2021 Apr 15;12:648636. doi: 10.3389/fphar.2021.648636. PMID: 33935751; PMCID: PMC8082498.