

Rossell: COVID-19 mental health Australia

**An overview of current mental health in the general population of Australia during the
COVID-19 pandemic: Results from the COLLATE project**

Susan L Rossell^{1,2}, Erica Neill^{1,2,3#}, Andrea Phillipou^{1,2,3,4#}, Eric J Tan^{1,2#}, Wei Lin Toh^{1#}, Tamsyn E
Van Rheenen^{1,5#}, Denny Meyer¹

All middle authors contributed equally, hence are included in alphabetical order

¹*Centre for Mental Health, Faculty of Health, Arts & Design, Swinburne University of Technology,
Melbourne, VIC, Australia*

²*Department of Mental Health, St Vincent's Hospital, Melbourne, VIC, Australia*

³*Department of Psychiatry, University of Melbourne, Melbourne, VIC, Australia*

⁴*Department of Mental Health, Austin Hospital, Melbourne, VIC, Australia*

⁵*Melbourne Neuropsychiatry Centre, Department of Psychiatry, University of Melbourne &
Melbourne Health, Melbourne, VIC, Australia*

Word count: 3681

Abstract word count: 291 words

References: 20

Correspondence: Professor Susan Rossell, Centre for Mental Health, Faculty of Health, Arts &
Design, Swinburne University of Technology, PO Box 218, Hawthorn VIC 3122, Australia;
telephone: +613 9214 8173; email: srossell@swin.edu.au

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

Rossell: COVID-19 mental health Australia

Abstract

The novel coronavirus disease (COVID-19) poses significant mental health challenges globally; however, to date, there is limited community level data. This study reports on the first wave of data from the COLLATE project (COvid-19 and you: mentaL heaLth in AusTralia now survEy), an ongoing study aimed at understanding the impact of the COVID-19 pandemic on the mental health and well-being of Australians. This paper addresses prevailing primary concerns related to the COVID-19 pandemic, current levels of negative emotions and risk factors predicting these negative emotions. On April 1st to 4th 2020, 5158 adult members of the Australian general public completed an online survey. Participants ranked their top ten current primary concerns about COVID-19, and completed standardized measures to ascertain levels of negative emotions (specifically, depression, anxiety and stress). Socio-demographic information was also collected and used in the assessment of risk factors. The top three primary concerns were all related to the health and well-being of family and loved ones. As expected, levels of negative emotion were exceptionally high. Modelling of predictors of negative emotions established several risk factors related to demographic variables, personal vulnerabilities, financial stresses, and social distancing experiences; particularly being young, being female, or having a mental illness diagnosis. The data provides important characterization of the current mental health of Australians during the COVID-19 pandemic. Critically, it appears that specific groups in the Australian community may need special attention to ensure their mental health is protected during these difficult times. The data further suggests the need for immediate action to combat high levels of psychological distress, along with the exacerbation of mental health conditions, in relation to the COVID-19 pandemic in Australia. These results may provide some direction for international researchers hoping to characterize similar issues in other countries.

Keywords: Coronavirus, COVID-19, mental health, general population, concerns, negative emotions

Rossell: COVID-19 mental health Australia

Introduction

The novel coronavirus disease (COVID-19) emerged in China in late 2019 and has spread rapidly across the globe. It is a contagious viral infection presenting with respiratory, neurological, gastrointestinal, and cardiac symptoms that range in severity from non-symptomatic through to causing death (Mao et al., 2020). The first case in Australia reported symptoms on January 13th 2020 (2019-nCoV National Incident Room Surveillance Team, 2020). This was followed by an exponential increase in infections, and unfortunately, deaths (the first Australian death occurred on February 24th). Since COVID-19 was declared a pandemic by the World Health Organization (WHO) on March 11th 2020 (WHO, 2020), the world has been engulfed in an unprecedented global crisis characterized by threatened or actual healthcare system collapse, job losses, and a failing global economy. In Australia, the crisis has been compounded by the implementation of government-regulated restrictions to contain the virus affecting social liberties.

In light of this, COVID-19 poses a significant mental health challenge to the Australian population, both now and in the long term. However, at the time of conducting this project there was no community level data in relation to the mental health implications of the COVID-19 pandemic in Australia, and only one published study from another Western country. In that study of 1310 adults in Spain, heightened negative emotions were associated with being female, being younger, and having negative self-perceptions (Losada-Baltar et al., 2020). Four studies on the psychological impacts of COVID-19 have also emerged from China. One study analyzed Weibo (Chinese social media platform) posts from 17,865 active users using online ecological recognition based on machine-learning predictive models (Li et al., 2020). The results showed that negative emotions increased (e.g., anxiety, depression and indignation), while positive emotions (e.g., happiness) and life satisfaction decreased over a two-week period from January 13th to January 26th 2020. Two other studies compared the psychological status of medical and non-medical (administration) health workers, illustrating increased insomnia, fear, anxiety, depression, somatization, and obsessive-

Rossell: COVID-19 mental health Australia

compulsive symptoms in medical health workers (Lu et al., 2020, Zhang et al., 2020). Living in rural areas, being female, and being in contact with COVID-19-positive patients were reported as risk factors for negative emotions. Finally, Wang et al. (Wang et al., 2020), using an online survey and snowball sampling in the general population between January 31st and February 2nd 2020, reported that ~50% of the 1210 respondents rated the psychological impact of COVID-19 as moderate-to-severe, with 33% reporting moderate-to-severe anxiety. Student status, being female and poor self-rated health were reported as risk factors for negative emotions.

In Australia, several health and economic measures had been implemented by March 31st 2020, in an attempt to control the spread of COVID-19 and stave off economic recession (e.g. \$130b towards keeping Australians employed). While a recent funding announcement of \$1.1b to boost digital mental health services is welcome, if we are to adequately manage this COVID-19 mental health crisis, there is a time-critical need to empirically characterize the current psychological impacts of the pandemic on the Australian population. This is particularly relevant given the Australian Government's current implementation of 'social distancing', a key transmission-prevention measure that describes the maintenance of minimum physical space between oneself and those outside of one's home. Social distancing restrictions, which limit one's out-of-home movements unless absolutely essential, have been found to increase social isolation and loneliness (Zhang et al., 2020), alcohol abuse (Wu et al., 2008), and domestic violence (Galea et al., 2020). This could translate to widespread fear, anxiety, and depression in general society, particularly exacerbated in persons with existing mental health conditions who have an increased susceptibility to the adverse impacts of stress (Duan and Zhu, 2020).

Here we report on the first wave of data collected from the COLLATE project (COvid-19 and you: mental health in Australia now survey), an ongoing study aimed at understanding the impact of the COVID-19 pandemic on the mental health and wellbeing of Australians. The COLLATE project

Rossell: COVID-19 mental health Australia

(described below) focuses on identifying the current concerns, emotional experiences and risk factors for adverse COVID-19-related mental health outcomes in people currently living in Australia. In our initial analysis of wave 1 data, we focused on characterizing the primary concerns of respondents related to the current COVID-19 pandemic as of April 1st to 4th 2020. Levels of negative emotion (depression, anxiety, and stress) were examined and compared with existing Australian population norms; and were modelled as an outcome to identify possible risks factors related to demographic variables, personal vulnerabilities, financial stresses, and social distancing experiences.

Methods

This study received ethics approval from Swinburne University Human Ethics Review Committee (approval number: 20202917-4107) and complied with the Declaration of Helsinki.

Study Design and Population

On April 1st 2020, adult members of the Australian general public (aged 18+) were invited to participate in an anonymous ~15-20mins online survey, completed at their convenience (i.e. the inclusion criteria to participate were being aged 18+ years and currently residing in Australia). Participants were informed that 16 surveys would be issued over the course of the project. These would be active for 72 hours per month, from 9am on the 1st to 8:59am on the 4th (Australian Eastern Standard Time), occurring monthly for the first year and then annually for the subsequent four years (Tan et al., 2020). Participants were informed that they could complete as many or as few surveys as they wanted, with surveys from the same respondent being linked by a personalized pseudonym (thus a subsample would provide us with longitudinal data, with the remaining data cross-sectional snapshots over the 16 surveys).

Rossell: COVID-19 mental health Australia

After online consent, participants completed the survey which covered three broad topics: a) current concerns, b) current emotional experiences, and c) socio-demographics/risk factors. Items from previously validated surveys were incorporated where possible, in line with good practice in survey creation (Thayer-Hart, Dykema, Elver, Schaeffer, & Stevenson, 2010). Relevant existing scales and measures were included if they had good reliability/validity. Demographic items were included based on examinations of other large-scale Australian surveys (including the Household, Income and Labor Dynamics in Australia (HILDA) Survey, the National Drug Strategy Household Survey and the Domestic and Family Violence Survey, the Australian Bureau of Statistics National Health Survey) additional items were created where necessary to ensure that all areas of interest were covered. In terms of item structure, many of the demographic questions were multiple choice or check box options. For more exploratory items, open ended questions with text boxes for responses were provided. As noted, the data described here relate to survey round 1: April 2020, and only the measures addressing our aims for this manuscript are described below.

Invitations to complete the survey were placed on digital university and community noticeboards and social media (e.g. Facebook, LinkedIn, Instagram, and Twitter) as well as participant registries held within the Centre for Mental Health at Swinburne University, which included participants with identified mental health conditions. Exponential non-discriminative snowball sampling was used, with all participants asked to pass the invitation onto their networks.

Measures

Primary concerns: Participants were asked to identify and rank their top 10 current concerns (out of 23) relating to the COVID-19 pandemic, with 1 being their greatest concern (see Table 2 for the full list of concerns).

Negative emotions: The Depression Anxiety Stress Scale (DASS-21) was used. It is a 21-item self-report measure yielding three subscales – depression, anxiety, and stress – each containing seven

Rossell: COVID-19 mental health Australia

items (Lovibond and Lovibond, 1995). Individual items are scored on a four-point Likert scale (0 to 3). DASS-21 raw scores were doubled to render them comparable to full-length DASS scores (42 items).

Risk factors: Measurement of risk factors were divided into four categories: *Demographics*

including: age, gender, education, living situation, geographical location/state, whether born in Australia, ethnicity, and religion; *Personal vulnerabilities* including: being someone at increased

mortality risk (e.g. immune-compromised, >60years), having lived experience of mental illness,

being a carer of someone with a mental illness or special needs, and being a healthcare professional

or ‘essential’ worker; *Financial stresses:* fortnightly take-home pay, cash savings, mortgage

repayments/rent, self-employment, job loss, and occupation; *Social distancing experiences:*

perceived positives of the situation, perception of government restrictions on mental health,

perception of social distancing measures duration, and working from home.

Statistical Analyses

Data were analyzed in *SPSS v26.0*. For all analyses, weights were used to adjust for imbalances in the sample based on the Australian Bureau of Statistics (ABS) population data for age, gender and geographical location/state (ABS, 2016). In all, there were 12 categories for age (18-19; 20-24; 25-29; 30-34; 35-39; 40-44; 45-49; 50-54; 55-59; 60-64; 65-70; 70+), two categories for gender (male; female) and four categories for state (Victoria; New South Wales; Queensland; Australian Capital Territory + Northern Territory + Western Australia + South Australia + Tasmania). By March 31st 2020 23:59 Australian Eastern Standard Time (AEST) there were 4,707 confirmed cases of COVID-19 in Australia, with 18 deaths. The majority of confirmed cases were in the states of New South Wales (n=2,182), Queensland (n=743) and Victoria (n=917), with the other states and territories reporting a total of 865 cases together. Given confirmed case numbers, we stratified by state by examining these three states independently from the other states and territories, which were combined.

Rossell: COVID-19 mental health Australia

Primary concerns: To characterize the top ten primary concerns, the number of respondents endorsing each concern was obtained and mean rankings were computed for the ten most commonly selected options. Rankings of zero were assigned to options not endorsed by a participant, and rankings of 1 to 10 were computed for endorsed concerns, with 10 for the option of greatest concern. In this case, the weights developed used the joint distributions of the three weighting variables ensuring that the sum of weights assigned was 5545 (i.e. the total number of respondents to this question who also provided age, gender and geographical location/state data).

Negative emotions: Depression, anxiety, and stress subscales and total DASS scores were compared to Australian population norms (Lovibond and Lovibond, 1995) using *t*-tests. Additionally, the percentage of participants (weighted and non-weighted) were calculated across the four defined severity levels (normal, mild, moderate, severe/extremely severe) for the three negative emotions. Respondents who failed to complete more than 10% of the DASS items were removed from the analysis. Remaining missing items were imputed using the EM algorithm as Little's MCAR test showed items were missing completely at random. For this analysis, the sum of weights and sample size were equal to 5158.

Risk factors: Using a transformed (SQRT) total DASS, the relationships between negative emotions and the four domains (*demographics, personal vulnerabilities, financial stresses and social distancing experiences*) were explored using general linear model analyses.

Results

Sample description

8014 participants started the survey, with $n=5545$ respondents (~30% attrition) completing the primary concerns ranking question and providing demographic data. For the negative emotion analyses $n=5158$ respondents completed the DASS. Demographic data is displayed in Table 1. The sample was biased in favour of females aged 25-45 living in the state of Victoria, making the use of post-stratification weighting essential in subsequent analyses.

Rossell: COVID-19 mental health Australia

Table 1: Sociodemographic description of the sample

Demographic Variables		Primary Concerns Analyses		Negative Emotion and Risk Factors Analyses	
		N	%	N	%
Age (years)	18-19	82	1.5	77	1.5
	20-24	436	7.9	397	7.7
	25-29	927	16.7	852	16.5
	30-34	894	16.1	806	15.6
	35-39	785	14.2	731	14.2
	40-44	630	11.4	592	11.5
	45-49	497	9.0	472	9.2
	50-54	405	7.3	383	7.4
	55-59	266	4.8	257	5.0
	60-64	288	5.2	274	5.3
	65-70	183	3.3	177	3.4
	70+	152	2.7	140	2.7
	Gender	Male	958	17.3	896
Female		4483	80.8	4172	80.9
Other		104	1.9	90	1.7
Highest Education	Postgraduate	1744	31.8	1642	31.8
	Undergraduate	2275	41.5	2152	41.7
	Diploma/ Certificate	956	17.4	889	17.2
	High School	511	9.3	475	9.2
Living situation	Single person living alone	693	12.8	664	12.9
	Non-related adults sharing a home	454	8.4	428	8.3
	Couple living with no children	1384	25.6	1318	25.6
	Couple with dependent children living at home	1597	29.6	1523	29.6
	Single parent with dependent children living at home	267	4.9	260	5.0
	Single person living with extended family (with or without children)	354	6.6	331	6.4
	Couple living with extended family (with or without children)	221	4.1	214	4.2
Other	434	8.0	415	8.1	
Geographical location / state	ACT, SA, WA, TAS, NT	665	12.0	621	12.0
	NSW	1028	18.5	955	18.5
	VIC	3489	62.9	3239	62.8
	QLD	363	6.5	343	6.6
Born in Australia	Yes	4211	75.9	3917	78.5
	No	1148	20.7	1070	21.5

Legend: Australian Capital Territory – ACT; South Australia – SA; Western Australia – WA; Tasmania – TAS; Northern Territory – NT; New South Wales – NSW; Victoria - VIC; Queensland – QLD.

Primary concerns: The primary concern data, with the percentage of respondents rating their top 10 concerns, is presented in Table 2. Mean rankings were ordered from 1 to 23 in declining order of importance (mean and standard deviations calculated for the rankings of each concern).

Rossell: COVID-19 mental health Australia

“Implications for health and wellbeing of family loved ones” was the most commonly endorsed concern, however, it ranked 3rd in terms of mean rankings. “Loved one dying of coronavirus” and “Loved one catching coronavirus” were the next most commonly endorsed primary concerns, and had the highest mean rankings.

Table 2: Rank data illustrating the ten primary concerns endorsed by Australians (N= 5545)

Concern [^]	Order Mean Ranks	Mean Rank#	SD Rank#	% Top 10 Rankings	Order Top 10 Ranks
Loved one dying from COVID-19	1	6.41	4.09	77.75	2
Loved one catching coronavirus	2	5.40	3.91	73.13	3
Implications for health and wellbeing of family/loved ones	3	5.27	3.20	82.36	1
Implications for health and wellbeing of society	4	3.48	3.19	67.57	4
Catching COVID-19 myself	5	3.43	3.62	58.95	7
Implications for health and wellbeing of self	6	3.32	3.28	61.61	6
Dying of COVID-19 myself	7	2.98	3.79	47.56	11
Social isolation and social distancing~	8	2.98	3.22	57.05	8
Australian economy	9	2.83	2.94	62.68	5
Risk of unemployment or reduced employment	10	2.63	3.52	44.35	13
Personal finances	11	2.21	3.07	45.89	12
Access to appropriate medical care	12	2.11	2.75	47.60	10
Availability of food and medicines	13	2.10	2.63	51.20	9
World economy	14	1.75	2.61	43.33	14
Balancing work & caring for children/dependents	15	1.25	2.76	20.98	17
The rapidly changing landscape	16	1.24	2.35	29.60	15
Travel restrictions	17	1.21	2.30	28.87	16
Government communication of key messages	18	0.91	2.09	20.76	18
Media coverage of the pandemic	19	0.69	1.83	17.92	19
Domestic violence	20	0.66	1.80	15.53	21
Adapting to working from home (e.g. IT/connectivity issues)	21	0.64	1.80	15.61	20
Others	22	0.46	1.76	9.51	22
Not being able to attend regular place of worship	23	0.34	1.45	6.92	23


Legend: [^] Analyses weighted to adjust for imbalance in the sample of respondents based on Australian Bureau of Statistics (ABS)(ABS, 2016) population data for age, gender and geographical location (State). # Rankings of zero were assigned to options not endorsed by a participant and, for endorsed concerns, rankings of 1 to 10 were computed, with 10 for the option of greatest concern. ~ On 31st March 2020, Australia was at Stage 3 activity restrictions, with Australians told to stay at home except for four reasons - food and essential supplies, medical attention, exercise, and work and study if cannot do so remotely. The first COLLATE survey was launched the following day, 1st

Rossell: COVID-19 mental health Australia

April 2020. Black text = Top ten concerns in rank order, Grey text = Remaining thirteen concerns in rank order.

Negative emotions: Figure 1 (also see Supplementary Table 1) shows mean values for the DASS scores compared to Australian norms. In all cases, they were significantly greater than the norms ($p < .001$); with people self-identifying as having a mental health diagnosis (MH DX) scoring 5-5.5-fold higher than those without such a diagnosis, who themselves scored 3 times higher than normative levels. Table 3 presents the score distributions across the four severity levels (normal, mild, moderate, severe/extremely severe); 21-35% of the population demonstrated moderate-to-extremely severe depression, anxiety and stress.

Figure 1: Comparison of weighted sample results with existing Australian norms for the DASS-21



Legend: Norms in Australia from Lovibond & Lovibond (Lovibond and Lovibond, 1995), MH = Mental Health

Rossell: COVID-19 mental health Australia

Table 3: DASS scores across the severity levels in the current Australian data from COLLATE in comparison to the Chinese data from Wang et al. (Wang et al., 2020)

		Depression			Anxiety			Stress		
		Aus W	Aus NW	China	Aus W	Aus NW	China	Aus W	Aus NW	China
Normal	N	2796	2521	843	3480	3058	770	2328	1870	821
	%	54.2%	48.9%	69.7%	67.5%	59.3%	63.6%	45.1%	36.3%	67.9%
Mild	N	752	810	167	402	439	91	1730	1825	292
	%	14.6%	15.7%	13.8%	7.8%	8.5%	7.5%	33.5%	35.4%	24.1%
Moderate	N	922	1040	148	713	892	247	725	929	66
	%	17.9%	20.2%	12.2%	13.8%	17.3%	20.4%	14.1%	18.0%	5.5%
Severe/Extremely Severe	N	687	787	52	562	770	102	375	534	31
	%	13.3%	15.3%	4.3%	10.9%	14.9%	8.4%	7.3%	10.4%	2.6%

Legend: Aus W = Australia weighted data, Aus NW = Australia non-weighted data; Depression: normal (score: 0-6), mild (score: 10-12), moderate (score: 13-20), severe/extremely severe (score: 21-42); Anxiety: normal (score: 0-6), mild (score: 7-9), moderate (score: 10-14), severe/extremely severe (score: 15-42) and Stress: normal (score: 0-10), mild (score: 11-18), moderate (score: 19-26), severe/extremely severe (score: 27-42). Scores as per Lovibond & Lovibond (Lovibond and Lovibond, 1995), it should be noted that DASS is not a categorical measure of clinical diagnosis, for clinical purposes it can be helpful to have ‘labels’ to characterize degree of severity relative to the population.

Risk factors: Demographics – 18.7% of the variation in negative emotions was explained by demographic factors (Supplementary Table 2). Lower levels of negative emotions were demonstrated by: people aged 30-34 and 70+, males, people with higher levels of education, couples (with or without children), and non-Australian born residents. The two states with the most COVID-19 cases (i.e. New South Wales and Victoria) showed lower negative emotions than the other states and territories.

Personal vulnerabilities - Adding personal vulnerabilities to the initial model explained an additional 10.5% of the variation in negative emotions (Supplementary Table 3). People with a higher mortality risk, lived experience of mental illness, carer responsibilities for someone with a

Rossell: COVID-19 mental health Australia

mental illness or special needs, as well as people in “essential” occupations, all had higher levels of negative emotions.

Financial stresses - Adding finance-related variables to the model explained an additional 5.2% of the variation (Supplementary Table 4). Individuals with higher fortnightly incomes and cash savings demonstrated lower levels of negative emotions. Higher levels of negative emotions were experienced by people under financial stress to meet mortgage and rental payments as well as those expecting to lose their jobs. Additionally, highest negative emotions were present for the unemployed, closely followed by homemakers, volunteers, or retired people.

Social distancing experiences - An additional 8.4% of the variation was explained by adding social distancing variables (Supplementary Table 5). Generally speaking, higher negative emotions were recorded for those who found they now had more free time. This included those who had more down-time, more time to spend communicating with family, more time to do jobs around the house and for those who identified no positive influences in the current situation. However, negative emotions were lower for those who found they now had more time for hobbies. Negative emotions were higher for those who reported that the government restrictions were adversely impacting their mental health, and for those who thought that the current restrictions might continue for more than 12 months. Finally, negative emotions were higher for those not working from home.

Risk factor summary – The four domains explained 42.8% of the variance in negative emotions (summarized in Table 4). Important predictors for high negative emotions (i.e. $\eta^2 \geq 0.010$) were being young (18-24), being female, being single, living in states with lower COVID-19 cases (QLD, ACT, SA, WAS, TAS, NT), being at higher risk of mortality and having a lived experience of mental illness. The perceived negative effect of government restrictions on mental health was also highly associated with negative emotions, demonstrating the largest effect size, $\eta^2 = 0.102$. Having sizeable cash savings, owning one’s own home and predicting a short duration of the current situation were protective factors against experiencing negative emotions.

Rossell: COVID-19 mental health Australia

Table 4: R-Square values and most important predictor variables for negative emotions

Predictor Domains	Increase in R-Square	R-Square	Most important predictors
Demographics	18.7%	18.7%	Age ($\eta^2=0.099$) Gender ($\eta^2=0.027$) Living situation ($\eta^2=0.011$) State ($\eta^2=0.016$)
Personal vulnerabilities	10.5%	29.2%	Higher mortality risk ($\eta^2=0.020$) Lived experience mental health illness ($\eta^2=0.095$)
Financial stresses	5.2%	34.4%	Home rental stress ($\eta^2=0.012$) Level of cash savings ($\eta^2=0.012$)
Social distancing experiences	8.2%	42.8%	Effects of government restrictions ($\eta^2=0.102$) Expected duration of current situation ($\eta^2=0.010$)

Legend: All selected predictors with partial effect sizes $\eta^2 \geq 0.010$

Discussion

The first wave of data from the COLLATE project provides an important characterization of the current mental health of Australians during the COVID-19 pandemic. The top three primary concerns among the general public were all related to the health and well-being of family and loved ones, specifically loved ones catching or dying from COVID-19. As expected, levels of negative emotions (depression, anxiety and stress) were exceptionally high, approximately three times greater than existing population norms in those with no pre-existing mental health conditions. Of concern was the finding that those with a pre-existing mental health condition demonstrated negative emotions 5 to 5.5 times greater than population norms. When the current Australian DASS data was compared with Chinese data (Wang et al., 2020), two differences emerged. First, the mean total DASS score from China of 20.16 (SD 20.42) was lower than that of Australia, even for individuals not reporting a mental health condition (18-24years: 33.56 (SD 25.49) and 25years+: 22.29 (SD 16.90)). Second, more Australians were classified as having moderate-to-extremely severe negative emotions (see Table 3). These apparent cross-cultural differences will need to be further investigated with a specifically designed comparison study, with differences in social norms and civil liberties between the two countries being possible influences. Nonetheless, both the

Rossell: COVID-19 mental health Australia

Australian and Chinese data speak to the elevation of negative emotions in the general population during the COVID-19 pandemic.

In our data, modelling predictors of negative emotions established several risks factors related to demographic variables, personal vulnerabilities, financial stresses, and social distancing experiences. This included young people (18-24 years) as well as those that are approaching middle-age (35-50 years). Given that the number of young people experiencing mental health conditions has been rising over the last decade in Australia (Carlisle et al., 2019), and internationally (Miron et al., 2019), this current data of such high levels of negative emotions in young people (up to age 24 here) is of particular concern. Increased negative emotions in our middle-aged respondents were associated with increased childcare duties and/or financial stresses that are specific to the immediate situation. This speaks to the importance of monitoring negative emotions in both young people and the middle-aged group longitudinally, that is, in the short and long term.

Being female was another significant risk factor for high levels of negative emotions (Lu et al., 2020, Wang et al., 2020, Zhang et al., 2020). While possible reasons for this remain to be determined, we speculate this could relate to juggling work and increased childcare duties, heightened risk of being in a domestic violence situation, as well as the higher risks of job loss and/or higher likelihood of being an ‘essential’ worker. Those under financial strain and those who were unemployed are also at increased risk of psychological distress. Thus, methods for targeting this ‘financial strain’ population to offer them more affordable options for mental health support will be important.

Finally, those with pre-existing mental illness are of specific concern (Phillipou et al., 2020; Neill et al, 2020; Van Rheenen et al, 2020), and existing mental health services will likely need increased

Rossell: COVID-19 mental health Australia

support to meet the rising needs of these consumers. It is notable that since this survey was conducted, the Australian government has announced a significant boost in funding to support mental health (\$76m over the next two years) with initiatives including dedicated websites and phone lines to support people experiencing stress and anxiety from prevailing COVID-19 related pressures, as well as a public information campaign. The current findings strongly underlie the need for such initiatives to be more targeted to specific groups.

Two other findings in our data warrant discussion. First, respondents from the two states with the highest number of COVID-19 cases, New South Wales and Victoria, were found to have lower negative emotions. While this was unexpected, further examination of risk factors established there were a number of protective personal vulnerabilities and financial stresses for persons living in these two states. That is, these states encompassed a lower percentage of respondents with lived experience of a mental illness and a greater percentage of respondents with financial stability.

Another important, albeit unsurprising, finding from our data was that individuals who perceived that current government restrictions were very negatively impacting their mental health also had the most pronounced negative emotions. In the context of respondents' primary concerns with the health and well-being of family members, this finding reinforces that government restrictions, such as social distancing, may be better framed in public messaging as necessary for protecting loved ones and ourselves from contracting the virus. Such a refocusing on positive outcomes may provide individuals with a sense of agency that tempers the powerlessness of being given a legal mandate to stay home.

A limitation of the study was the snowballing approach to survey recruitment; this resulted in a non-representative sample of the Australian population, which included some respondents with known mental health diagnosis. To address this, weightings were used based on ABS data (ABS, 2016) to statistically correct for any bias. However, even with statistical weighting, it is difficult to account

Rossell: COVID-19 mental health Australia

for specific subgroups, for example those without access to the internet. Furthermore, despite >8000 participants starting the survey, only ~5500 had useable datasets due to considerable attrition (30%), which is albeit typical of online research. This data provides a snapshot of mental health and well-being of Australians in April 2020 in relation to COVID-19; to do so we compared current negative emotions to existing Australian norms. It is possible differences in sampling factors related to the current sample data and existing norms may explain some of the differences rather than COVID-19 itself. However, given the magnitude of our findings in terms of elevated negative emotions in the general community such sampling differences are unlikely to explain the large variance between current and norm data. We also did not assess the mental health of school-aged children and adolescents in the current survey, the majority are whom are being home-schooled and not permitted face-to-face contact with their peers in Australia at present. School-age children may thus be particularly vulnerable to stress and anxiety, and a critical avenue for future work will be to examine this group specifically.

Conclusion

The data collected from the COLLATE project will provide a reference for healthcare professionals in terms of current mental health needs in Australia, in addition to guiding policymakers in making accurate provisions within mental health services and actionable policies. The findings are predicted to be applicable across other nations with similar healthcare systems and government management of the COVID-19 pandemic. Important findings from the data are that: a) people with existing mental health conditions have very high levels of negative emotions (as per (Duan and Zhu, 2020)), in addition, there are b) high levels of psychological distress in the general community, with some individuals particularly vulnerable (as per (Wang et al., 2020)). The fluid nature of the situation throughout this pandemic makes our continuing and longitudinal comparisons of the mental health effects of government restrictions and lock-down duration forecasts a priority for future study.

Rossell: COVID-19 mental health Australia

Overall, this data has made it clear that increased mental health support will be of paramount importance as the world faces the consequences of the COVID-19 pandemic.

Rossell: COVID-19 mental health Australia

References

- 2019-NCOV NATIONAL INCIDENT ROOM SURVEILLANCE TEAM 2020. 2019-nCoV acute respiratory disease, Australia Epidemiology Report 1: Reporting week 26 January – 1 February 2020. *Communicable Diseases Intelligence*, 44.
- ABS 2016. QuickStats. Available at: <https://www.abs.gov.au/websitedbs/D3310114.nsf/Home/2016%20QuickStats>.
- CARLISLE, E., FILDES, J., HALL, S., PERRENS, B., PERDRIAU, A. & PLUMMER, J. 2019. Youth Survey Report 2019. *sydney: NSW: Mission Australia*.
- DUAN, L. & ZHU, G. 2020. Psychological interventions for people affected by the COVID-19 epidemic. *Lancet Psychiatry*, 7, 300-302.
- GALEA, S., MERCHANT, R. M. & LURIE, N. 2020. The Mental Health Consequences of COVID-19 and Physical Distancing: The Need for Prevention and Early Intervention. *JAMA Intern Med*.
- LI, S., WANG, Y., XUE, J., ZHAO, N. & ZHU, T. 2020. The Impact of COVID-19 Epidemic Declaration on Psychological Consequences: A Study on Active Weibo Users. *Int J Environ Res Public Health*, 17, E2032.
- LOSADA-BALTAR, A., JIMENEZ-GONZALO, L., GALLEGO-ALBERTO, L., PEDROSO-CHAPARRO, M. D. S., FERNANDES-PIRES, J. & MARQUEZ-GONZALEZ, M. 2020. "We're staying at home". Association of self-perceptions of aging, personal and family resources and loneliness with psychological distress during the lock-down period of COVID-19. *J Gerontol B Psychol Sci Soc Sci*. in press.
- LOVIBOND, S. H. & LOVIBOND, P. F. 1995. Manual for the depression anxiety stress scales. *Sydney: Psychology Foundation Monograph*.
- LU, W., WANG, H., LIN, Y. & LI, L. 2020. Psychological status of medical workforce during the COVID-19 pandemic: A cross-sectional study. *Psychiatry Res*, 288, 112936.
- MAO, L., JIN, H., WANG, M., HU, Y., CHEN, S., HE, Q., CHANG, J., HONG, C., ZHOU, Y., WANG, D., MIAO, X., LI, Y. & HU, B. 2020. Neurologic Manifestations of Hospitalized Patients With Coronavirus Disease 2019 in Wuhan, China. *JAMA Neurol*.
- MIRON, O., YU, K. H., WILF-MIRON, R. & KOHANE, I. S. 2019. Suicide Rates Among Adolescents and Young Adults in the United States, 2000-2017. *JAMA*, 321, 2362-2364.
- NEILL, E., MEYER, D., TOH, W.L., VAN RHEENEN, T.E. PHILLIPOU, A., TAN, E.J., ROSSELL, S.L. 2020. Alcohol Use in Australia during the Early Days of the COVID-19 Pandemic: Initial results from the COLLATE project. *Psychiatry and Clinical Neurosciences*, doi: 10.1111/pcn.13099.
- PHILLIPOU, A., MEYER, D., NEILL, E., TAN, E., TOH, W.L., VAN RHEENEN, T., ROSSELL, S.L. 2020. Eating and exercise behaviours in eating disorders and the general population during the COVID-19 pandemic in Australia: initial results from the COLLATE project. *International Journal of Eating Disorders*, doi: 10.1002/eat.23317.
- TAN, E., MEYER, D., NEILL, E., PHILLIPOU, A., TOH, W., VAN RHEENEN, T. & ROSSELL, S. 2020. Considerations for assessing the impact of the COVID-19 pandemic on mental health in Australia. *Australian and New Zealand Journal of Psychiatry*. in press.
- THAYER-HART, N., DYKEMA, J., ELVER, K., SCHAEFFER, N.C., STEVENSON, J. 2010. Survey fundamentals: A guide to designing and implementing surveys. Madison: Board of Regents, University of Wisconsin Sytem.
- VAN RHEENEN, T.E., MEYER, D., NEILL, E., PHILLIPOU, A., TAN, E.J., TOH, W.L., ROSSELL, S.L. 2020. Mental health status of individuals with a mood-disorder during the COVID-19 Pandemic in Australia: Initial Results from the COLLATE Project. *Journal of Affective Disorders*, 275:69-77. doi: 10.1016/j.jad.2020.06.037.
- WANG, C., PAN, R., WAN, X., TAN, Y., XU, L., HO, C. S. & HO, R. C. 2020. Immediate Psychological Responses and Associated Factors during the Initial Stage of the 2019

Rossell: COVID-19 mental health Australia

Coronavirus Disease (COVID-19) Epidemic among the General Population in China. *Int J Environ Res Public Health*, 17, 1729.

WHO. 2020. *WHO Director-General's opening remarks at the media briefing on COVID-19 - 11 March 2020* [Online]. Available: <https://www.who.int/dg/speeches/detail/who-director-general-s-opening-remarks-at-the-media-briefing-on-covid-19---11-march-2020> [Accessed 14/04/2020].

WU, P., LIU, X., FANG, Y., FAN, B., FULLER, C. J., GUAN, Z., YAO, Z., KONG, J., LU, J. & LITVAK, I. J. 2008. Alcohol abuse/dependence symptoms among hospital employees exposed to a SARS outbreak. *Alcohol*, 43, 706-12.

ZHANG, W. R., WANG, K., YIN, L., ZHAO, W. F., XUE, Q., PENG, M., MIN, B. Q., TIAN, Q., LENG, H. X., DU, J. L., CHANG, H., YANG, Y., LI, W., SHANGGUAN, F. F., YAN, T. Y., DONG, H. Q., HAN, Y., WANG, Y. P., COSCI, F. & WANG, H. X. 2020. Mental Health and Psychosocial Problems of Medical Health Workers during the COVID-19 Epidemic in China. *Psychother Psychosom*, 1-9.

Rossell: COVID-19 mental health Australia

Acknowledgements

SLR holds a Senior National Health and Medical Research Council (NHMRC) Fellowship (GNT1154651), and EJT (GNT1142424) and TVR (GNT1088785) hold Early Career NHMRC Fellowships. AP (GNT1159953) and WLT (GNT1161609) are supported by NHMRC New Investigator Project Grants. The authors would also like to thank all the participants who took the time to participate in this study.

Author Contributions

SLR conceived the project and the design. All authors SLR, DM, EN, AP, EJT, WLT and TVR finalized the design, constructed the survey, obtained ethics, engaged in data collection and interpretation of findings. DM completed all the data analyses in consultation with the other authors. All authors prepared the manuscript and agreed to its final form.

Declaration of conflicting interests

The authors have declared no potential conflicts of interest with respect to the research, authorship, and publication of this article.

Funding

This project received no funding.

Rossell: COVID-19 mental health Australia

Supplementary Material

Supplementary Table 1: Comprehensive comparison of weighted sample results with existing Australian norms for the DASS-21

	Australian Norms 18-24 years(Lovibond and Lovibond, 1995)	18-24 years No MH DX Nwt=410	18-24 years Previous MH DX Nwt=201	Australian Norms 25-90 years(Lovibond and Lovibond, 1995)	25-90 years No MH DX Nwt=3148	25-90 years Previous MH DX Nwt=1399
Depression	M 3.96 SD 4.52 Md 3 Range 0-20	12.57 10.65 10 0-42	20.83 11.57 20 0-42	2.21 3.60 1 0-19	7.64 6.98 6 0-40	14.48 10.07 12 0-42
Anxiety	M 2.76 SD 3.52 Md 2 Range 0-15	7.34 7.86 4 0-36	15.93 10.28 16 0-40	1.48 2.60 0 0-17	4.15 5.17 2 0-42	8.70 7.86 6 0-42
Stress	M 4.78 SD 4.71 Md 3 Range 0-21	13.65 9.60 12 0-42	21.37 8.89 22 0-42	3.79 4.10 3 0-21	10.50 7.49 10 0-42	16.80 8.90 16 0-42
Total DASS	M 11.51 SD 11.51 Md 8.5 Range 0-54	33.56 25.49 28 0-120	58.12 25.52 60 0-124	7.48 9.18 4 0-52	22.29 16.90 18 0-124	39.98 23.33 36 0-124

Legend: Norms in Australia from Lovibond & Lovibond (Lovibond and Lovibond, 1995), Nwt = weighted sample size, M= mean, SD = standard deviation, Md = Median, MH = mental health, DX = diagnosis

Rossell: COVID-19 mental health Australia