

Treatment with ACE-inhibitors is associated with less severe disease with SARS-Covid-19 infection in a multi-site UK acute Hospital Trust

Authors

Daniel M Bean^{1,2+}, Zeljko Kraljevic¹, Thomas Searle¹, Rebecca Bendayan^{1,4}, Andrew Pickles¹, Amos Folarin^{1,2,3,7}, Lukasz Roguski^{2,3,7}, Kawsar Noor^{2,3,7}, Anthony Shek⁸, Kevin O'Gallagher^{5,6}, Rosita Zakeri^{5,6}, Ajay M Shah^{5,6}, James TH Teo^{5,8+*}, Richard JB Dobson^{1,2,3,4,7+*}

*joint author

+corresponding authors: Dr Dan Bean (daniel.bean@kcl.ac.uk); Dr James Teo

(jamesteo@nhs.net; +44(0) 20 7828 0800); Prof Richard Dobson (richard.j.dobson@kcl.ac.uk; +44(0) 20 7848 0473)

1. Department of Biostatistics and Health Informatics, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, U.K. 2. Health Data Research UK London, University College London, London, U.K. 3. Institute of Health Informatics, University College London, London, U.K. 4. NIHR Biomedical Research Centre at South London and Maudsley NHS Foundation Trust and King's College London, London, U.K. 5. Kings College Hospital NHS Foundation Trust, London, U.K. 6. School of Cardiovascular Medicine & Sciences, King's College London British Heart Foundation Centre of Excellence, London SE5 9NU, U.K. 7. NIHR Biomedical Research Centre at University College London Hospitals NHS Foundation Trust, London. 8. Dept of Clinical Neuroscience, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London

Abstract:

Background: The SARS-Cov2 virus binds to the ACE2 receptor for cell entry. It has been suggested that ACE-inhibitors, which are commonly used in patients with hypertension or diabetes and which raise ACE2 levels, may increase the risk of severe COVID-19 infection.

Methods: We evaluated this hypothesis in an early cohort of 205 acute inpatients with COVID-19 at King's College Hospital and Princess Royal University Hospital, London, UK with the primary endpoint being death or transfer to a critical care unit for organ support within 7-days of symptom onset.

Findings: 53 patients out of 205 patients reached the primary endpoint. Contrary to the hypothesis, treatment with ACE-inhibitors was associated with a reduced risk of rapidly deteriorating severe disease. There was a lower rate of death or transfer to a critical care unit within 7 days in patients on an ACE-inhibitor OR 0.29 (CI 0.10-0.75, $p < 0.01$), adjusting for age, gender, comorbidities (hypertension, diabetes mellitus, ischaemic heart disease and heart failure).

Interpretation: Although a small sample size, we do not see evidence for ACE-inhibitors increasing the short-term severity of COVID-19 disease and patients on treatment with ACE-inhibitors should continue these drugs during their COVID-19 illness. A potential beneficial effect needs to be explored as more data becomes available.

Introduction

Early data from China during the SARS-Cov2 pandemic suggest that patients with hypertension or diabetes have an increased risk of severe COVID-19 disease.¹ It has been hypothesized that treatment with ACE-inhibitors (ACEi) or angiotensin receptor blockers (ARB) in such patients may increase the expression of ACE2, the receptor for SARS-Cov2 binding.

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

and entry into cells.^{1,2} However, reduced activation of the renin angiotensin system (RAS) and/or increased levels of ACE2 may be protective during severe lung injury.³ The effect of ACEi and ARB during infection with SARS-CoV-2 is therefore controversial but needs urgent clarification.⁴⁻⁷

We tested for association between treatment with ACEi or ARB and disease severity in the first 205 patients with COVID-19 disease admitted to a multi-site acute NHS Trust in the United Kingdom (King's College Hospital NHS Foundation Trust). We used an informatics pipeline to allow rapid evaluation of disease concepts during the rapidly evolving pandemic.

Methods:

This project operated under London South East Research Ethics Committee (reference 18/LO/2048) approval granted to the King's Electronic Records Research Interface (KERRI); specific work on COVID-19 research was reviewed with expert patient input on a virtual committee with Caldicott Guardian oversight.

Study Design:

The study cohort was defined as all inpatients testing positive for SARS-Cov2 by RT-PCR at King's College Hospital and Princess Royal University Hospital from 1 March to 22nd March 2020. Only patients symptomatic and requiring inpatient admission were included. The primary endpoint was defined as death or admission to a critical care unit for organ-support within 7 days of symptoms onset (symptoms defined as fever, cough, dyspnoea, myalgia, chest pain or delirium). Patients were stratified according to drug exposure to ACEi or ARB within 7 days before symptoms or during inpatient treatment (prior to an endpoint being reached). Specifically, we divided the total cohort into patients with prescription and/or mention in any text record of (1) ACEi (Ramipril, Perindopril, Lisinopril, Enalapril, Captopril, Quinapril, Imidapril, Fosinopril, Trandolapril) and (2) ARB (Candesartan, Irbesartan, Losartan, Olmesartan, Telmisartan, Valsartan) up to 7days before or after symptom onset. We considered patients whose medication was withheld during admission as positive for drug exposure. Patients with no medication orders or clinical documents created in the study period (because never admitted) were excluded. One newborn was also excluded. The primary endpoint was manually verified by a clinician review of the electronic health record.

Data Processing:

The data (demographic, emergency department letters, discharge summaries, clinical notes, radiology reports, medication orders, lab results) was retrieved and analyzed in near real-time from the structured and unstructured components of the electronic health record (EHR) using a variety of natural language processing (NLP) informatics tools belonging to the CogStack ecosystem,⁸ namely DrugPipeline,⁹ MedCAT¹⁰ and MedCATTrainer.¹¹ The CogStack NLP pipeline captures negation, synonyms, and acronyms for medical SNOMED-CT concepts as well as surrounding linguistic context using deep learning and long short-term memory networks. DrugPipeline was used to annotate medications and MedCAT produced unsupervised annotations for all SNOMED-CT concepts under parent terms Clinical Finding, Disorder, Organism, and Event with disambiguation, pre-trained on MIMIC-III.¹² Further supervised training improved detection of annotations and meta-annotations such as experiencer (is the concept annotated experienced by the patient or other), negation (is the

concept annotated negated or not) and temporality (is the concept annotated in the past or present) with MedCATTrainer. Meta-annotations for hypothetical and experienter were merged into Irrelevant meaning that any concept annotated as either hypothetical or where the experienter was not the patient was annotated as irrelevant. Performance of the MedCAT NLP pipeline for disorders mentioned in the text was evaluated on 138 documents by 4 annotators (TS, ZK, DB, AS) and F1, precision and recall recorded. Additional full case review for correct subsequent diagnosis assignment was performed by 3 clinicians (JT, KOG, RZ) for key comorbidities: Diabetes Mellitus, Hypertension, Heart Failure and Ischaemic Heart Disease. Performance of DrugPipeline has previously been described.⁹ All detected drug mentions were manually reviewed to exclude false positives (e.g. allergy).

Statistical Analysis:

In order to investigate the association between ACEi and disease severity measured as critical care admission or death, we performed a series of logistic regressions applying Firth's correction.¹³ This procedure has shown to be robust for low prevalence events and low-dimensional settings.^{14,15} In a first step, we explored independently the association for ACEi (Baseline model). In a second step, we adjusted the model for age and sex (Model 1). Then, we additionally adjusted for hypertension (Model 2), and additionally adjusted by other comorbidities diabetes and ischemic heart disease or heart failure (Model 3). We also explored the independent association for hypertension following the same modelling approach. Sensitivity analyses were performed i) using exact logistic regression with models adjusted variable by variable; ii) requiring at least two detections of medication for positive exposure; iii) using only structured data on in-hospital medication orders; iv) ignoring our 7 day window for medications; v) testing sensitivity to unmeasured confounders.

Role of the funding source:

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Results:

Our total cohort consists of 205 confirmed positive symptomatic inpatients aged 63±20 (SD) years and 52% males (Table 1). Baseline characteristics are 51·2% with hypertension, 30·2% with diabetes and 14·6% with ischaemic heart disease or heart failure. The percentage of patients that have a positive mention of a certain disorder in each of the two groups (Dead or Critical Care, Other) derived via the NLP for medical concept annotations with F1 > 80% and more than 10 annotated mentions are shown in Figure 1 (performance shown in Figure 2). All NLP-detected positive mentions of hypertension, diabetes, ischaemic heart disease or heart failure were manually reviewed at a patient level and false positive rates calculated 1·9%, 3·2%, 31%, 0% respectively.

Of the 205 patients, 53 patients died or required critical care support within 7 days of symptoms and 152 patients did not. The inclusion criteria of only patients needing admission is likely why this critical outcome figure is relatively high (25·9%) compared to fatality rate in population studies¹⁶ but is comparable to hospital case series.¹⁷ 14% (5/37) patients with exposure to an ACE-inhibitor died or required critical care support compared to 29% (48/168) for patients without such exposure.

Findings from unadjusted logistic regression models indicated that individuals on ACEi had lower likelihood of severe disease (OR 0.42 (CI 0.14-1.00), $p=0.058$). These associations were only partially attenuated when adjustments for gender and age were included (Model 1 in Table 2). Furthermore, these associations remained significant and were only partially attenuated when the model was additionally adjusted for hypertension (Model 2 in Table 2) and further for other comorbidities diabetes and ischaemic heart disease or heart failure (Model 3 in Table 2). Odds ratios and p-values for all variables in each model are shown in Supplementary Table 1. Males were found to have a higher likelihood of severe disease in Model 3 (OR 2.00 (CI 1.00-4.00), $p=0.037$).

We also examined the independent association between hypertension and disease severity. Our results showed that individuals diagnosed with hypertension had a similar likelihood of developing severe disease as those that were not diagnosed with hypertension, either in unadjusted models (OR 1.60 (CI 0.88-3.10); $p=0.12$) or models adjusted for age and gender (OR 1.80 (CI 0.83-3.80); $p=0.14$).

We did not run the regression analysis on the ARB group as there are only 9 patients in our cohort. We intend to carry out this analysis as our cohort grows.

Sensitivity analyses showed similar results when compared with the results from exact logistic regression analyses with univariate adjustment. We also compared our results to those using the penalised regression model and criteria for ACEi exposure that were either more strict (requiring multiple mentions or using only medications ordered in hospital) and less strict (including any detection of ACEi outside our 7 day window). In all cases we found that estimates of the impact of ACEi exposure were consistently in the same direction as those in Table 2 but were not significant.

We assessed the robustness to unmeasured confounders of the fully adjusted estimate of ACEi protective effect using the e-value approach.¹⁸ An e-value of 3.32 for the point estimate indicated that it was robust to plausible residual confounding that might explain away the estimate. However, the e-value of 1.58 for the upper confidence of 0.75 showed that in the presence of residual confounding of a level often considered plausible, this study could not fully exclude the possibility of small harmful effects.

Discussion:

This study suggests that ACE-inhibitors do not increase the severity of COVID-19 disease as hypothesised⁴⁻⁷ but may reduce severity. This holds true even after adjusting for conditions where ACEi may be used (hypertension, diabetes mellitus, ischaemic heart disease and heart failure). No meaningful comment can be made about ARB effect given the low prevalence of their use in this cohort, although ARB have a different mechanism of action compared to ACEi.

This study used an NLP approach to perform very rapid analysis of high volume, unstructured real world clinical data. This however introduces the possibility of missing circumlocutory mentions of disease, symptoms or medications. We have mitigated against this by manually validating annotations in a subset of records and also verified ACEi and ARB annotations

against inpatient electronic prescription data. Moreover, we have performed sensitivity analyses to test the impact of different criteria to define the ACEi exposed cohort on our results, finding that although not significant the OR remained <1.0 for ACEi exposure in all analyses. The lack of significance in the more strict analyses is likely due to the loss of power as some detections of ACEi medication are excluded. For the less strict analysis, the lack of significance may be due to noise introduced (e.g. prescription halted before the study period). The NLP output in the less strict analysis is also not manually reviewed and is highly likely to contain some irrelevant mentions e.g. previous allergic reaction. While the findings of robustness to bias due to unmeasured confounding increased our confidence, the need for replication in a larger sample remains.

One limitation with this study is the relatively small sample taken from a single UK centre over a short follow-up. Although we have used statistical procedures to provide robust results with our current sample, as numbers increase further updates to the analysis will be required to better understand our findings and confirm the directionality of these associations. Our group will provide regular updates during the pandemic to the analysis at the link <https://cogstack.org/cogstack-kch-covid-19-analyses/> including longer follow-up and pooled analysis with other organisations. Whether these results also apply to infection severity in the non-hospital setting or to different global populations requires further study.

A tentative favourable association of ACE-inhibitors with less severe early outcomes is suggested by this study. A putative mechanism could be reduced RAS activation in patients on ACE-inhibitors, which is considered protective in Acute Respiratory Distress Syndrome, ARDS.³ Furthermore, elevation of ACE2 also reduces RAS activation and is protective in acute lung injury,¹⁹ including in ARDS of SARS1 infection.²⁰

In summary, based on these early results and the absence of any evidence suggesting harm, patients on treatment with ACE-inhibitors should continue these drugs during their COVID-19 illness as per current guidelines.²¹ Active research is merited on whether ACE inhibition or enhancement of ACE2 may have a therapeutic role in severe COVID-19 disease.

Acknowledgements:

DMB is funded by a UKRI Innovation Fellowship as part of Health Data Research UK MR/S00310X/1 (<https://www.hdruk.ac.uk>). RB is funded in part by grant MR/R016372/1 for the King's College London MRC Skills Development Fellowship programme funded by the UK Medical Research Council (MRC, <https://mrc.ukri.org>) and by grant IS-BRC-1215-20018 for the National Institute for Health Research (NIHR, <https://www.nihr.ac.uk>) Biomedical Research Centre at South London and Maudsley NHS Foundation Trust and King's College London. AF is supported by grant IS-BRC-1215-20018 for the National Institute for Health Research (NIHR, <https://www.nihr.ac.uk>) Biomedical Research Centre at South London and Maudsley NHS Foundation Trust and King's College London and The National Institute for Health Research University College London Hospitals Biomedical Research Centre. RJBD is supported by: 1. Health Data Research UK, which is funded by the UK Medical Research Council, Engineering and Physical Sciences Research Council, Economic and Social Research Council, Department of Health and Social Care (England), Chief Scientist Office of the Scottish Government Health and Social Care Directorates, Health and Social Care

Research and Development Division (Welsh Government), Public Health Agency (Northern Ireland), British Heart Foundation and Wellcome Trust. 2. The BigData@Heart Consortium, funded by the Innovative Medicines Initiative-2 Joint Undertaking under grant agreement No. 116074. This Joint Undertaking receives support from the European Union's Horizon 2020 research and innovation programme and EFPIA; it is chaired by DE Grobbee and SD Anker, partnering with 20 academic and industry partners and ESC. 3. The National Institute for Health Research University College London Hospitals Biomedical Research Centre. 4. National Institute for Health Research (NIHR) Biomedical Research Centre at South London and Maudsley NHS Foundation Trust and King's College London. KO'G is supported by an MRC Clinical Training Fellowship. RZ is supported by a King's Prize Fellowship. AS is supported by a King's Medical Research Trust studentship. AMS is supported by the British Heart Foundation (CH/1999001/11735), the National Institute for Health Research (NIHR) Biomedical Research Centre at Guy's & St Thomas' NHS Foundation Trust and King's College London (IS-BRC-1215-20006), and the Fondation Leducq. AP is partially supported by NIHR NF-SI-0617-10120. This work was supported by the National Institute for Health Research (NIHR) University College London Hospitals (UCLH) Biomedical Research Centre (BRC) Clinical and Research Informatics Unit (CRIU), NIHR Health Informatics Collaborative (HIC), and by awards establishing the Institute of Health Informatics at University College London (UCL). This work was also supported by Health Data Research UK, which is funded by the UK Medical Research Council, Engineering and Physical Sciences Research Council, Economic and Social Research Council, Department of Health and Social Care (England), Chief Scientist Office of the Scottish Government Health and Social Care Directorates, Health and Social Care Research and Development Division (Welsh Government), Public Health Agency (Northern Ireland), British Heart Foundation and the Wellcome Trust. This paper represents independent research part funded by the National Institute for Health Research (NIHR) Biomedical Research Centre at South London and Maudsley NHS Foundation Trust and King's College London. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health and Social Care. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. We would also like to thank all the clinicians managing the patients, the patient experts of the KERRI committee, Professor Irene Higginson, Professor Alastair Baker, Professor Jules Wendon, Dan Persson and Damian Lewsley for their support.

Declaration of Interests:

JTHT received research support and funding from InnovateUK, Bristol-Myers-Squibb, iRhythm Technologies, and holds shares <£5,000 in Glaxo Smithkline and Biogen.

Author Contributions:

JTHT, RJBD, DMB, ZK, TS, AF conceived the study design
 DMB, ZK, AS, TS, JTHT, LR, KN performed data processing and software development
 KOG, RZ, JTHT performed data validation
 DMB, AP, RB performed statistical analysis
 JTHT, DMB, ZK, TS, AF, RJBD, AMS performed critical review
 JTHT, RJBD, DMB, AMS, ZK, TS, AF, DMB, AP, RB wrote the manuscript

References:

1. Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* [Internet]. 2020 Feb 15;395(10223):497–506. Available from: [http://dx.doi.org/10.1016/S0140-6736\(20\)30183-5](http://dx.doi.org/10.1016/S0140-6736(20)30183-5)
2. Hoffmann M, Kleine-Weber H, Schroeder S, Krüger N, Herrler T, Erichsen S, et al. SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. *Cell* [Internet]. 2020 Mar 4; Available from: <http://dx.doi.org/10.1016/j.cell.2020.02.052>
3. World leading geneticist and immunologist Josef Penninger on potential COVID-19 treatment [Internet]. Youtube; 2020 [cited 2020 Mar 28]. Available from: <https://www.youtube.com/watch?v=jAW6VBWTiAA>
4. Di Guardo G. SARS-CoV-2, hypertension and ACE-inhibitors. 2020; Available from: <https://research.unite.it/handle/11575/107297>
5. Sommerstein R, Gräni C. Rapid response: re: preventing a covid-19 pandemic: ACE inhibitors as a potential risk factor for fatal Covid-19. *BMJ*. 2020;368:m810.
6. Fang L, Karakiulakis G, Roth M. Are patients with hypertension and diabetes mellitus at increased risk for COVID-19 infection? *Lancet Respir Med* [Internet]. 2020 Mar 11; Available from: [http://dx.doi.org/10.1016/S2213-2600\(20\)30116-8](http://dx.doi.org/10.1016/S2213-2600(20)30116-8)
7. Kuster GM, Pfister O, Burkard T, Zhou Q, Twerenbold R, Haaf P, et al. SARS-CoV2: should inhibitors of the renin–angiotensin system be withdrawn in patients with COVID-19? *Eur Heart J* [Internet]. 2020 Mar 20 [cited 2020 Mar 28]; Available from: <https://academic.oup.com/eurheartj/advance-article-abstract/doi/10.1093/eurheartj/ehaa235/5810479>
8. Jackson R, Kartoglu I, Stringer C, Gorrell G, Roberts A, Song X, et al. CogStack - experiences of deploying integrated information retrieval and extraction services in a large National Health Service Foundation Trust hospital. *BMC Med Inform Decis Mak* [Internet]. 2018 Jun 25;18(1):47. Available from: <http://dx.doi.org/10.1186/s12911-018-0623-9>
9. Bean DM, Teo J, Wu H, Oliveira R, Patel R, Bendayan R, et al. Semantic computational analysis of anticoagulation use in atrial fibrillation from real world data. *PLoS One* [Internet]. 2019 Nov 25;14(11):e0225625. Available from: <http://dx.doi.org/10.1371/journal.pone.0225625>
10. Kraljevic Z, Bean D, Mascio A, Roguski L, Folarin A, Roberts A, et al. MedCAT -- Medical Concept Annotation Tool [Internet]. arXiv [cs.CL]. 2019. Available from: <http://arxiv.org/abs/1912.10166>
11. Searle T, Kraljevic Z, Bendayan R, Bean D, Dobson R. MedCATTrainer: A Biomedical Free Text Annotation Interface with Active Learning and Research Use Case Specific Customisation. In: Proceedings of the 2019 Conference on Empirical Methods in Natural Language Processing and the 9th International Joint Conference on Natural Language Processing (EMNLP-IJCNLP): System Demonstrations [Internet]. Stroudsburg, PA, USA: Association for Computational Linguistics; 2019. p. 139–44. Available from: <https://www.aclweb.org/anthology/D19-3024>

12. Johnson AEW, Pollard TJ, Shen L, Lehman L-WH, Feng M, Ghassemi M, et al. MIMIC-III, a freely accessible critical care database. *Sci Data* [Internet]. 2016 May 24;3:160035. Available from: <http://dx.doi.org/10.1038/sdata.2016.35>
13. Firth D. Bias reduction of maximum likelihood estimates. *Biometrika* [Internet]. 1993 Mar 1 [cited 2020 Mar 28];80(1):27–38. Available from: <https://academic.oup.com/biomet/article-abstract/80/1/27/228364>
14. Wang X. Firth logistic regression for rare variant association tests. *Front Genet* [Internet]. 2014 Jun 19;5:187. Available from: <http://dx.doi.org/10.3389/fgene.2014.00187>
15. Doerken S, Avalos M, Lagarde E, Schumacher M. Penalized logistic regression with low prevalence exposures beyond high dimensional settings. *PLoS One* [Internet]. 2019 May 20;14(5):e0217057. Available from: <http://dx.doi.org/10.1371/journal.pone.0217057>
16. Onder G, Rezza G, Brusaferro S. Case-Fatality Rate and Characteristics of Patients Dying in Relation to COVID-19 in Italy. *JAMA* [Internet]. 2020 Mar 23; Available from: <http://dx.doi.org/10.1001/jama.2020.4683>
17. Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* [Internet]. 2020 Mar 11; Available from: [http://dx.doi.org/10.1016/S0140-6736\(20\)30566-3](http://dx.doi.org/10.1016/S0140-6736(20)30566-3)
18. Linden A, Mathur MB, VanderWeele TJ. Conducting sensitivity analysis for unmeasured confounding in observational studies using E-values: The evalua package. *Stata J* [Internet]. 2020 Mar 1;20(1):162–75. Available from: <https://doi.org/10.1177/1536867X20909696>
19. Imai Y, Kuba K, Rao S, Huan Y, Guo F, Guan B, et al. Angiotensin-converting enzyme 2 protects from severe acute lung failure. *Nature* [Internet]. 2005 Jul 7;436(7047):112–6. Available from: <http://dx.doi.org/10.1038/nature03712>
20. Kuba K, Imai Y, Rao S, Gao H, Guo F, Guan B, et al. A crucial role of angiotensin converting enzyme 2 (ACE2) in SARS coronavirus–induced lung injury. *Nat Med* [Internet]. 2005 Jul 10 [cited 2020 Mar 28];11(8):875–9. Available from: <https://www.nature.com/articles/nm1267>
21. Simone GD. Position Statement of the ESC Council on Hypertension on ACE-Inhibitors and Angiotensin Receptor Blockers [Internet]. 2020 [cited 2020 Mar 28]. Available from: [https://www.escardio.org/Councils/Council-on-Hypertension-\(CHT\)/News/position-statement-of-the-esc-council-on-hypertension-on-ace-inhibitors-and-ang](https://www.escardio.org/Councils/Council-on-Hypertension-(CHT)/News/position-statement-of-the-esc-council-on-hypertension-on-ace-inhibitors-and-ang)




Figure 1. The percentage of patients that have a positive mention of a disorder in each of the two groups (Dead or Critical Care, Other). Dead or Critical Care - patients that have died or that are in the Intensive Treatment Unit; and Other - patients that are neither dead nor in ITU at day 7. All diseases were extracted from free-text using Cogstack and MedCAT. Only medical concept annotations with $F1 > 80\%$ and more than 10 annotated samples are shown.




Figure 2. Performance of the CogStack and MedCAT NLP pipeline in detecting disease mentions within the electronic health record text. Precision (P), Recall (R) and F1 (harmonic mean of precision and recall). Only medical concept annotations with F1 > 80% and more than 10 annotated samples are shown.

	Overall	On ACEi	Not on ACEi	On ARB	Not on ARB	Primary endpoint reached	Primary endpoint not reached
N	205	37	168	9	196	53	152
Age	62.95 (19.94)	74.51 (13.41)	60.4 (20.27)	65.0 (13.53)	62.85 (20.21)	64.91 (17.4)	62.26 (20.77)
Male	106 (51.7%)	22 (59.5%)	84 (50.0%)	3 (33.3%)	103 (52.6%)	33 (62.3%)	73 (48.0%)
On ACEi	37 (18.0%)	37 (100.0%)	0 (0.0%)	0 (0.0%)	37 (18.9%)	5 (9.4%)	32 (21.1%)
On ARB	9 (4.4%)	0 (0.0%)	9 (5.4%)	9 (100.0%)	0 (0.0%)	4 (7.5%)	5 (3.3%)
Hypertension	105 (51.2%)	31 (83.8%)	74 (44.0%)	7 (77.8%)	98 (50.0%)	32 (60.4%)	73 (48.0%)
Diabetes mellitus	62 (30.2%)	17 (45.9%)	45 (26.8%)	6 (66.7%)	56 (28.6%)	20 (37.7%)	42 (27.6%)
Ischaemic heart disease or heart failure	30 (14.6%)	11 (29.7%)	19 (11.3%)	2 (22.2%)	28 (14.3%)	6 (11.3%)	24 (15.8%)
Primary Endpoint	53 (25.9%)	5 (13.5%)	48 (28.6%)	4 (44.4%)	49 (25.0%)	53 (100.0%)	0 (0.0%)

Table 1. Characteristics of study cohort. All patients are positive for COVID-19. All variables are shown as N (% of column) except age which is mean (SD). ACEi = Angiotensin converting enzyme inhibitor; ARB = Angiotensin 2 Receptor Blocker.

Model	Adjustments	OR (95% CI) on ACEi vs Not on ACEi	p
Baseline	-	0.42 (0.14-1.00)	0.058
Model 1	Age, sex	0.34 (0.11-0.86)	0.02
Model 2	Age, sex, hypertension	0.28 (0.09-0.73)	<0.01
Model 3	Age, sex, hypertension, diabetes mellitus, ischaemic heart disease, heart failure	0.29 (0.10-0.75)	<0.01

Table 2. Summary of odds ratios for ACE inhibitor drug exposure and primary endpoint. Odds ratios and p-values calculated from logistic regressions applying Firth's correction (Firth, 1993). Baseline no adjustments; Model 1 adjusted for age and gender; Model 2 additionally adjusted for hypertension; Model 3 additionally adjusted for hypertension, diabetes mellitus, ischaemic heart disease or heart failure. ACEi = Angiotensin converting enzyme inhibitor. OR = Odds ratio.

Model	Variable	OR (95% CI)	P-value
Baseline	On ACEi	0.42 (0.14-1.00)	0.058
Model 1	On ACEi	0.34 (0.11-0.86)	0.022
Model 1	Age per 10 years	1.10 (0.94-1.30)	0.21
Model 1	Male	1.80 (0.95-3.40)	0.074
Model 2	On ACEi	0.28 (0.09-0.73)	<0.01
Model 2	hypertension	2.20 (1.00-4.90)	0.047
Model 2	Age per 10 years	1.00 (0.82-1.20)	0.98
Model 2	Male	2.00 (1.00-3.90)	0.038
Model 3	On ACEi	0.29 (0.10-0.75)	<0.01
Model 3	Age per 10 years	1.00 (0.85-1.30)	0.72
Model 3	Male	2.00 (1.00-4.00)	0.037
Model 3	diabetes	1.50 (0.71-3.00)	0.29
Model 3	hypertension	2.10 (0.90-4.80)	0.085
Model 3	ischaemic heart disease or heart failure	0.53 (0.18-1.40)	0.22
Hypertension unadjusted	hypertension	1.60 (0.88-3.10)	0.12
Hypertension adjusted	hypertension	1.80 (0.83-3.80)	0.14

Hypertension adjusted	Age per 10 years	0.98 (0.81-1.20)	0.84
Hypertension adjusted	Male	1.80 (0.97-3.50)	0.064

Supplementary Table 1. Odds ratios and p-values for all variables and primary endpoint. Odds ratios and p-values calculated from logistic regressions applying Firth's correction¹. ACEi = Angiotensin converting enzyme inhibitor. OR = Odds ratio.

References

1. Firth D. Bias reduction of maximum likelihood estimates. Biometrika [Internet]. 1993 Mar 1 [cited 2020 Mar 28];**80**(1):27–38. Available from: <https://academic.oup.com/biomet/article-abstract/80/1/27/228364>