

Syndromic surveillance insights from a symptom assessment app before and during COVID-19 measures in Germany and the United Kingdom: results from repeated cross-sectional analyses

Alicia Mehl^{1*}, MSc; Francois Bergey^{1*}, MSc; Caoimhe Cawley¹, PhD; Andreas Gilsdorf¹, MD, Dr. med.

*Equal contribution

Correspondence to
Alicia Mehl; science@ada.com

1 – Ada Health GmbH, Karl-Liebknecht-Str. 1, 10178 Berlin, Germany

Abstract

Background: Unprecedented lockdown measures have been introduced in countries across the world to mitigate the spread and consequences of COVID-19. While attention has focused on the effects of these measures on epidemiological indicators relating directly to the infection, there is increased recognition of their broader health implications. However, assessing these implications in real time is a challenge, due to limitations of existing syndromic surveillance data and tools.

Objective: To explore the added value of mobile phone app-based symptom assessment tools as real time health insight providers to inform public health policy makers.

Methods: A comparative and descriptive analysis of the proportion of all self-reported symptoms entered by users during an Ada assessment in Germany and the United Kingdom (UK) was conducted between two periods: before and after the implementation of “Phase One” COVID-19 measures. Additional analyses were performed to explore the association between symptom trends and seasonality, and symptom trends and weather. Differences in the proportion of unique symptoms between the periods were analysed using Pearson’s Chi-squared test and reported as Log₂ Fold Changes (Log₂ FC).

Results: Between 48,300-54,900 symptomatic users reported 140,500-170,400 symptoms during the Baseline and Measures periods in Germany. Between 34,200-37,400 symptomatic users in the UK reported 112,100-131,900 symptoms during the Baseline and Measures periods. The majority of symptomatic users were female (Germany 68,600/103,200, 66.52%; UK 51,200/71,600, 72.74%). The majority (Germany 68,500/100,000, 68.45%; UK 50,900/68,800, 73.91%) were aged between 10 and 29 years, and about a quarter (Germany 26,200/100,000, 26.15%; UK 14,900/68,800, 21.65%) were between 30-59 years. 103 symptoms were reported either more or less frequently (with statistically significant differences) during the Measures as compared to the Baseline period, and 34 of these were found in both countries. The following mental health symptoms (Log₂ FC, *P*-value) were reported less often during the Measures period: *inability to manage constant stress and demands at work* (-1.07, *P*<.001), *memory difficulty* (-0.56, *P*<.001), *depressed mood* (-0.42, *P*<.001), and *impaired concentration* (-0.46, *P*<.001). *Diminished sense of taste* (2.26, *P*<.001) and *hyposmia* (2.20, *P*<.001) were reported more frequently during the Measures period. None of the 34

symptoms were found to be different between the same dates in 2019. Fourteen of the 34 symptoms had statistically significant associations with weather variables.

Conclusions: Symptom assessment apps have an important role to play in facilitating improved understanding of the implications of public health policies such as COVID-19 lockdown measures. Not only do they provide the means to complement and cross-validate hypotheses based on data collected through more traditional channels, they can also generate novel insights through a real-time syndromic surveillance system.

Keywords: COVID-19; symptom assessment apps; symptom checker app; syndromic surveillance; health effects of COVID-19 measures; COVID-19 lockdown; digital public health, COVID-19 measures

Introduction

Background

Since its emergence at the end of 2019, COVID-19 has had an enormous and wide-ranging impact, with millions of confirmed cases and hundreds of thousands of deaths reported worldwide [1]. Governments have introduced a series of measures ranging from work and school closures to social distancing and lockdowns to mitigate the spread and consequences of infection. The measures have been unprecedented on many levels: in how disruptive they are to daily life, by the proportion of populations affected, in the duration of implementation, and by their global reach. As a result, the daily lives of millions have been upended.

The extent of the counter-measures to the pandemic raises questions as to their impact on public and individual health. While much of the initial focus in the medical and scientific community has been on understanding the virus and infection itself [2], as well as on the effects of various policy measures on epidemiological indicators relating directly to COVID-19 [3,4], there has also been increased recognition of the broader health effects [5].

Some of the broader implications concern the direct and immediate consequences of lockdown. For instance, as people are distressed due to social isolation and the economic fallout of the crises, an upsurge in the incidence and severity of mental health problems has been predicted [5–11]. Increased handwashing (a primary recommendation to reduce transmission of COVID-19) is expected to result in increased skin irritation and dermatitis [12,13]. Other implications are related to more indirect factors such as delays and cancellations to surgeries and non-urgent treatments for cancer and other patients [14–16], interruptions in drug and commodities supply chains [17,18], and drops in vaccination rates for vaccine-preventable diseases [19,20]. The consequences of these interruptions to medical services during this pandemic will likely create a higher morbidity, but the impact may not be visible for several years to come [17].

Assessing the health implications of lockdown policies in real-time is a challenge. Apart from the time lag, evidence cited in support of purported health effects is often anecdotal or based on surveys conducted among medical professionals. The latter provide important

insights into changes in health symptoms identified at the point of contact, but may not account for the effects of social distancing guidelines on health-seeking behavior. Traditional syndromic surveillance data [21] provides valuable information for a defined set of indicators over time, but is unable to distinguish whether the trends reflect changes in disease incidence or in the uptake of health services. More significant efforts towards data collection among the general public are currently being undertaken [22]. However, these are missing a Baseline prior to the emergence of COVID-19 that would facilitate a clearer understanding of the impact. In general, surveys can provide snapshots of a highly dynamic situation, but tend to be restricted in scope, meaning that changes in health beyond pre-defined indicators risk being overlooked.

In the past years, a growing number of studies have shown the benefits of mobile phone app-based symptom assessment tools for improved health outcomes, for example, in lowering the barrier to seek help [23–29]. If the literature on symptom assessment tools has primarily focused on individual health benefits, e.g. early detection of rare diseases [30], recent research has also demonstrated their potential contribution to public health. Specifically with regard to COVID-19, symptom tracker apps have helped flag anosmia (loss of the sense of smell) as a potentially relevant symptom of infection [31,32], as well as to challenge common understandings of presenting symptoms of COVID-19 [33].

Goal of this study

In the present study, we add to this body of literature by exploring the added value of symptom assessment tools for the analysis of broader health implications resulting from public health interventions such as lockdown measures. The study is based on self-reported symptom data from Ada, a digital symptom assessment app which uses a probabilistic reasoning engine that collects demographic information, symptoms, and medical history to suggest possible conditions and then guide individuals to the most appropriate care. The Ada app is available in seven languages and has over ten million users. It is described in Gilbert, et al in further detail [34]. Using an inductive approach and focusing on the immediate impact of COVID-19 control measures, we compared symptom data reported by users in Germany and the UK before and during the first phase of COVID-19 lockdown measures to identify changes and continuities in the incidence of self-reported symptoms.

The aim of this study was to explore the potential of the Ada symptom assessment app to generate real-time health insights to inform public health policy makers.

Methods

Study focus

This analysis is a comparative descriptive study of self-reported symptoms entered by users in an Ada assessment completed during the time of the “Phase One” COVID-19 interventions (the Measures period) compared to a Baseline period in Germany and the United Kingdom. The Measures periods began in each country when all five major non-pharmaceutical interventions described in Flaxman et al [4] were implemented (specifically: school closure ordered, case-based measures (strong recommendation of self-isolation when showing COVID-19-like symptoms), public events banned, social

distancing encouraged, lockdown decreed) and lasted until April 22nd, 2020, when data were extracted. The end of the Baseline period was defined as the day before any of the five interventions were implemented in the respective country, and the length of the Baseline period was equal to the Measures period in the same country. The period between the Baseline and Measures periods is excluded from the analysis due to the partial implementation of “Phase One” COVID-19 measures. This is an exploratory analysis of all symptoms (not only COVID-related) reported by symptomatic users during the defined periods. In this study, we consider a symptomatic user as one who completed at least one assessment during the periods of analysis, either themselves or by someone on their behalf (i.e. a legal guardian if under 16 years). We considered trends if in both countries, the proportion that a specific symptom was reported (out of all reported symptoms) was significantly different between the two periods.

Additional analyses

Ad-hoc analyses were later performed to test identified trends for symptoms against selected potential confounders, such as seasonality and weather. To explore the potential impact of seasonality on trends, the results were compared to the same analysis conducted for the same dates in 2019. To explore the impact of weather on trends, associations between the monthly proportion of reported symptoms and weather variables (average monthly temperature, monthly precipitation (mm), and monthly hours of sunshine) were investigated during the period January 2019 - March 2020. The weather analysis was restricted to Germany.

Participants

All assessments completed by Ada users in Germany and the UK during the Baseline and Measures period were included in the analysis. We analysed pseudonymized health data for public health purposes, according to the European General Data Protection Regulation (GDPR), and users are duly informed of such use of their data (information available at any time in Ada’s Privacy Policy). Additionally, users maintain their right to object to such processing for reasons arising from their particular situation, as required by the GDPR.

Variables

An Ada assessment consists of different parts: 1) the user enters an unlimited number of symptoms, 2) the user is asked about other potential symptoms they could have, and 3) an assessment result is provided with conditions that could potentially explain the reported symptoms and adequate triage. This analysis only includes symptoms that are self-reported by a user in the first part of the assessment: that is, responses to the initial question “*Let’s start with the symptom that’s troubling you the most*”, followed by “*Do you have any other symptoms?*”. Upon entering free text, the user is then given a range of medically-curated options (based on linguistic relevance) to select from.

The variable of interest $S_{i,k}$, representing the proportion a symptom i is self-reported during the period k is defined as

$$S_{i,k} = \frac{\sum_{j \text{ in } \{users\}} s_{i,j,k}}{\sum_{j \text{ in } \{users\}} x_{j,k}},$$

where

$s_{i,j,k}$ equals 1 if the user j self-reported the symptom i during the period k , and equals 0 otherwise.

$x_{j,k}$ equals 1 if the user j has finished one assessment during the period k , and equals 0 otherwise.

The age variable was grouped using the following categories (years): 0-9, 10-19, 20-29, 30-39, 40-59 and 60+.

Weather data (average monthly temperature (°C), monthly precipitation (mm) and monthly hours of sunshine) for Germany were extracted from the Deutscher Wetter Dienst database for the period January 2019 until March 2020 [35].

For ease of reporting and to aid interpretation, symptoms which were reported in significantly different proportions during the Baseline and Measures periods were grouped using the International Classification of Diseases version 2019 (ICD-10) of the World Health Organization [36]. ICD-10 R sub-groups, named “Symptoms, signs and abnormal clinical and laboratory symptoms, not elsewhere classified” were used when possible. Similar categories were grouped together later, as presented in (Table 1).

Table 1. List of groups based on the ICD-10 classification

		ICD-10 classification
Group name		
	Circulatory and respiratory systems	R00- R09, J
	Digestive system and abdomen	R10 – R19
	Skin and subcutaneous tissue	R20 – R23, L
	Musculoskeletal	M
	Genitourinary system	R30 – R39, N
	Cognition, perception, emotional state and behaviour	R40 – R46, F

	Speech and voice	R47 – R49
	General symptoms and signs	R50 – R69
	Nervous system	G
	Eye and adnexa	H

Bias

As Ada’s medical model and databases are continuously updated, we defined the Baseline period to be as close as possible to the Measures period to limit the impact of these changes on the data. All modeled symptoms that were added, deleted, significantly modified, or significantly affected by the modification of any other symptom from the first day of the Baseline period until the end of the Measures period were removed from this study.

Statistical methods

Sex and age groups of symptomatic users were reported as percentages and tested for differences between the periods with Pearson’s Chi-squared test. Differences in the proportion of symptoms between the periods were reported as Log2 Fold Changes and were analyzed with Pearson’s Chi-squared test. A Log2 Fold Change of 0.5 means that the proportion of that reported symptom was 1.41 times as large during the Measures compared to the Baseline. A Log2 Fold Change of 1 is interpreted as being twice as large during the Measures compared to the Baseline, and a Log2 Fold Change of 2 is four times as large. Conversely, a Log2 Fold Change of -1 means that the proportion of the reported symptom was twice as large during the Baseline compared to the Measures period. In general, Log2 Fold Change calculations are helpful in understanding relative differences in the proportions of users reporting each symptom between the two periods, but do not reflect how common reporting of that symptom was overall. Associations between weather variables and the proportion of symptoms were tested based on the Spearman’s Rank correlation coefficient.

When required, *P*-values were adjusted for multiple testing using the false discovery rate method. *P*-values less than or equal to 0.05 were considered statistically significant. Statistical analyses and figures were executed using R (version 3.6.1).

The analysis was done using exact numbers, but results representing user numbers are presented rounded to the closest hundred, to ensure a fully anonymised presentation of the results.

Results

An overview of the Baseline and Measures periods in Germany and the UK (numbers of Ada users, numbers of symptoms reported) are shown in (Table 2).

Table 2. Key parameters

	Germany		United Kingdom	
	Baseline	Measures	Baseline	Measures
Dates of period (dd.mm.yy)	03.02.20 – 05.03.20	22.03.20 – 22.04.20	11.02.20 – 11.03.20	24.03.20 – 22.04.20
Number of days per period	32	32	30	30
Number of all users (N _a)	467,000	483,100	488,800	501,300
Number of symptomatic users (N _a)	54,900	48,300	37,400	34,200
Number of reported symptoms by symptomatic users (N _a)	170,400	140,500	131,900	112,100
Number of symptoms reported at least once in Baseline or Measures period (N _a)	1,328		1,294	

^a rounded to the nearest hundred

Demographic characteristics of users are shown in (Table 3). During both the Baseline and Measures periods, the majority (Germany Baseline: 36,300/54,900, 66.19%; Germany Measures: 32,300/48,300, 66.90%; UK Baseline: 26,600/37,400, 71.17%; UK Measures: 24,600/34,200, 71.94%) of symptomatic users in both countries were female. The majority were aged between 10 and 29 years (Germany Baseline: 37,000/53,200, 69.51%; Germany Measures: 31,400/46,800, 67.13%; UK Baseline: 27,200/35,800, 75.76%; UK Measures: 23,700/32,900, 71.89%). Those aged between 30-59 years represented roughly a quarter (Germany Baseline: 13,300/53,200, 24.94%; Germany Measures: 12,900/46,800, 27.53%; UK Baseline: 7,100/35,800, 19.92%; UK Measures: 7,700/32,900, 23.54%) of symptomatic users. The number of symptomatic users in the Baseline (54,900 Germany, 37,400 UK) was slightly higher than in the Measures period (48,300 Germany, 34,200 UK).

Table 3. Demographic characteristics of the study population

		Germany					United Kingdom				
		Baseline		Measures		P-value ^c	Baseline		Measures		P-value ^c
		N ^a	%	N ^a	%		N ^a	%	N ^a	%	
Sex											
	female	36,300	66.2	32,300	66.9	0.02	26,600	72.2	24,600	72.9	0.02
Age^b (years)											
	0-9	1,300	2.4	700	1.6	<.001	900	2.4	700	2.0	<.001
	10-19	17,100	32.1	13,700	29.2	<.001	16,200	45.1	13,600	41.2	<.001
	20-29	20,000	37.5	17,800	37.9	0.28	11,000	30.7	10,100	30.7	0.95
	30-39	6,400	12.0	6,000	12.9	<.001	3,100	8.7	3,300	9.9	<.001
	40-59	6,900	13.0	6,800	14.6	<.001	4,000	11.2	4,500	13.6	<.001
	60+	1,700	3.2	1,800	3.9	<.001	700	2.1	900	2.7	<.001

^a rounded to the nearest hundred

^b Users who did not report a birth year were excluded from the analysis

^c Pearson's chi-squared test for differences between Baseline and Measures


Twenty-one symptoms were excluded from the analysis as they had been added to or deleted from the medical model during either the Baseline or Measures period. Three additional symptoms were excluded from the analysis as there were significant changes in the associated terms (text entered by users to match the description of a symptom to the term used in the model), which could affect the number of times a symptom is reported. A list of these symptoms is included in Table A of the Supplementary Materials.

Main analyses

Out of 1,328 and 1,294 symptoms investigated respectively in Germany and in the United Kingdom, 103 symptoms were reported either more or less frequently, in either country, during the Measures as compared to the Baseline period. The complete results can be found in (Tables B and C) of the Supplementary Materials.

Figure 2 presents the 34 symptoms that showed a statistically significant difference in both countries. 24 symptoms were reported less often and 10 were reported more often in the Measures period in comparison to the Baseline period.

Figure 2. Relative difference (Log2 Fold Change values) in the proportions of Ada users' reported symptoms with statistically significant differences between the Baseline and COVID-19 Measures periods in Germany and the UK



Out of the 34 significant symptoms, all skin and tissue-related symptoms (*pruritic nasal cavity*, *spot on the upper extremity*, *dry skin of the hands*, and *pruritic eyes*) were reported more frequently during the Measures period. In contrast, all speech and voice symptoms (*aphonia* and *hoarseness*), all eye and adnexa symptoms (*odynophonia*, *inability to clear the ear with changing barometric pressure*, *aural fullness*, and *otalgia*), all digestive system and abdomen symptoms (*odynophagia*, *diarrhea*, and *nausea*) and all musculoskeletal and nervous systems symptoms (*excessive daytime sleepiness* and *myalgia*) were reported less frequently during the Measures period.

In the cognition, perception, emotional state and behaviour symptoms group, two perception symptoms (*diminished sense of taste* and *hyposmia*), were reported more

frequently during the Measures period whereas the mental health symptoms (*inability to manage constant stress and demands at work, memory difficulty, depressed mood, and impaired concentration*) were reported less frequently during the Measures period. Out of the circulatory and respiratory symptoms, three were reported more frequently during the Measures period (*throat clearing, dyspnea, and chest pain*) and four were reported less frequently (*nasal discharge, sinus pain, sore throat, and cough*). One general symptom was reported more frequently (*lowered temperature of the foot*) and three (*malaise, chills, and fatigue*) were reported less frequently during the Measures period.

Additional analyses

Out of the 34 symptoms found to be different between the Baseline and Measures period in both Germany and the UK in 2020, none were found to be different between the same periods in 2019 in both countries. However, looking at the countries separately, in Germany, the data shows that eight of the 34 significant symptoms were also reported less frequently during the Measures period in 2019: *cough, chills, nasal discharge, myalgia, sore throat, malaise, fatigue* and *sinus pain*. The complete results can be found in Tables D and E of the Supplementary Materials.

The monthly weather reports in February 2020 (which corresponds to most of the Baseline period) differed in comparison to March 2020 (which corresponds to most of the Measures period) in Germany. Average temperature increased by 5.1°C (from 5.3°C to 10.4°C), monthly precipitation decreased by 107.8mm (from 124.1mm to 16.3mm) and the number of hours of sunshine per month increased by 228.5h (from 63.9h to 292.4h). Fourteen of the 34 significant symptoms had statistically significant associations with weather. Increased temperature was positively associated with *spot on the upper extremity* and negatively associated with *chest pain, lowered temperature of the foot, odynophagia, malaise, myalgia, cough, otalgia, chills, vomiting, sinus pain, and nasal congestion*. Increased hours of sunshine was positively associated with *pruritic eyes, spot on the upper extremity, and pruritic nasal cavity* and negatively associated with *lowered temperature of the foot* and *sinus pain*. The complete results can be found in Table F & Figure G of the Supplementary Materials.

Discussion

Principal Results

The results presented above show significant differences in the frequency and proportion of self-reported symptoms in Ada assessments before and after the implementation of measures aimed at reducing the transmission of COVID-19. Importantly, the same differences were found in both Germany and the UK, despite the divergent trajectories of these countries in the lead-up to the implementation of lockdown policies [37], as well as other national differences such as those relating to health systems [38]. Furthermore, these differences were not found during the same time periods in 2019, suggesting that the lockdown measures could have contributed to the results.

Many of the observed differences were to be expected. The reduced frequency of reported respiratory symptoms (*nasal discharge, sore throat, cough, sinus pain, nasal congestion, hoarseness, odynophonia, aphonia*) and influenza-like illness (i.e. *malaise, fatigue, chills, myalgia*) following the measures is understandable as the cold and flu season has also waned exceptionally rapidly during this period [39], likely facilitated by reduced contact resulting from lockdown. To support this interpretation, *cough, chills, nasal discharge, myalgia, sore throat, malaise, fatigue* and *sinus pain* were also reported less frequently during the Measures periods in 2019 in Germany, suggesting that seasonal changes are reflected in the data. The increased reporting of dry hands following the measures is consistent with more frequent handwashing during the Measures period, as expected by dermatologists [12,13]. Increased reports of *pruritic eyes* and *pruritic nasal cavity* following the measures could be a consequence of seasonal hay fever (known to be worse during the Spring compared to Winter months due to increased pollen in the air [40]), as these symptoms were found to be associated with increased sunshine (presumably when people spend more time outdoors).

The reduction of gastrointestinal symptoms could be associated with the closing of preschool/ day care settings (as they are known to contribute to the spread of these diseases [41]), restaurant closures, or due to improved hand hygiene. The decrease of ear problems could be a result of the extremely reduced air travel with much less people experiencing pressure adjustment problems called “airplane ear” [42], or related to the end of the cold and flu season. In these cases more research would be needed to explore these hypotheses.

We have also observed an increase in the reporting of *hyposmia, diminished sense of taste*, and *dyspnea* during the Measures period. These are less frequent but also typical COVID-19 symptoms and were increasingly recognized in the general public [31,43,44]. They were not found to be associated with seasonality or weather. This increase could be related to COVID-19 infections, or an artefact resulting from increased awareness of these symptoms due to media coverage.

A more surprising result is that *depressed mood, inability to manage constant stress and demands at work, impaired concentration, memory difficulty*, and *excessive daytime sleepiness* were reported in a lower proportion during the Measures period. This is not only contrary to conventional wisdom, but also runs counter to what was observed during previous infectious disease outbreaks, such as SARS, MERS, and Ebola [45,46] as well as to the literature reporting on the effects of COVID-19 on Mental Health [6–11,47,48]. Despite the fact that during the Measures period the temperature was warmer and there was more sunshine than usual, the analysis on weather data for Germany did not show a significant impact on the changed mental health symptoms. Our findings are supported by a growing body of evidence from ongoing studies [49] and recently published research [50], as well as anecdotal evidence reported in various mainstream media reports [51–53] which suggest that at least in the short-term the mental health effects of the COVID-19 measures may not be as negative as expected. One factor may be that the stress of everyday life and work/study is reduced during lockdown and home office/schooling. In addition, the reduction of *excessive daytime sleepiness* may be due to

the fact that people may sleep better or more during lockdown as there are less available places to go out and socialize in the evenings. Ongoing studies into the mental health effects of COVID-19 and its countermeasures may shed further light on these questions.

Limitations

It is important to interpret the study results taking into consideration the characteristics of the study population and the normal use case for Ada. Due to the specific age and sex distribution of users (predominantly young and female), the results may not be generalisable to other population groups. Furthermore, this analysis was limited to Germany and the UK (due to sufficient user numbers), and represents a two month snapshot of reported symptoms. Users who know they have a disease might not use Ada if their symptoms deteriorate, as the cause is already known. In addition, this is an analysis of patient-reported symptoms which are not validated. The impact of user acquisition strategies is not known. However, the similarity in trends observed across two countries (and for respiratory symptoms, over the same period of time in 2019) adds weight to our findings.

Despite these limitations, the analysis presented in this paper has a number of unique strengths. First, the analysis was conducted using a large, existing dataset that updates in real-time and covers over 1,400 unique self-reported symptoms since November 2016, allowing the monitoring of changes in trends over time. Second, the data is user-driven as a user self-reports their symptoms during an assessment on their own initiative. This allows for identification of changes that would not be detected in traditional studies focusing on specific and/or predefined areas. Third, the large number of symptoms presented in the results that are consistent with expectations, the observation of seasonal differences, and that the results were observed in both Germany and the UK is an indication of the reliability of Ada's data, demonstrated especially by the detected increase in the less frequent but typical COVID-19 symptoms *hyposmia* and *diminished sense of taste*. While the clinical soundness of Ada's model at the level of individual diagnostics has already been demonstrated in other studies [54,55], the presented results build confidence that the data collected through the Ada app can also detect health changes in a population in real time.

Future research

Future research can build on these strengths focusing on reasons for some of the detected changes and by expanding the analysis to more countries. Of particular interest are countries from the Global South and low- and middle-income countries, given the comparative paucity of up-to-date health data in these countries and the differentials in the burden of disease. In addition, investigating changes in trends over time as the implementation of the COVID-19 measures adapts according to the reality in different countries (i.e. as individuals return to work) will offer meaningful insights into the effects of policy changes.

Conclusions

Our findings suggest that symptom assessment tools might have a role to play in improving understanding of the implications of public health measures. In this analysis we have shown an innovative use of an existing dataset that enables policy makers to

inform and monitor public health measures with a real time, low-resource syndromic surveillance system that is relevant both during the COVID-19 pandemic but also in the future.

Acknowledgements

Author Statement AM & FB contributed to the planning (study conception, protocol development) which was revised by CC and AG. AM & FB contributed to the conduct (data collection and data analysis). AM, FB, CC, & AG contributed to the reporting (report writing). All the authors contributed to commenting on drafts of the report. AM is the guarantor for this work. The corresponding author attests that all listed authors meet authorship criteria and that no others meeting the criteria have been omitted.

Disclosures AM, FB, CC, & AG are employees or company directors of Ada Health GmbH.

Funding This study was funded by Ada Health.

Disclaimer None

Acknowledgements Stephen Gilbert (Director of Clinical Evaluation, Ada Health GmbH), Vincent Jörres (Communications & Public Affairs Lead, Ada Health GmbH), Vanessa Lemarié (Director of Business Development & Program Owner of Rare Disease Initiative in Business Development), Nicola Vona (Director of Reasoning Technology in Product), and Paul Wicks reviewed the manuscript and provided feedback.

References

1. COVID-19 Map - Johns Hopkins Coronavirus Resource Center [Internet]. [cited 2020 May 11]. Available from: <https://coronavirus.jhu.edu/map.html>
2. EPPI-Mapper [Internet]. [cited 2020 May 13]. Available from: http://eppi.ioe.ac.uk/COVID19_MAP/covid_map_v10.html
3. Prem K, Liu Y, Russell TW, Kucharski AJ, Eggo RM, Davies N, Jit M, Klepac P, Flasche S, Clifford S, Pearson CAB, Munday JD, Abbott S, Gibbs H, Rosello A, Quilty BJ, Jombart T, Sun F, Diamond C, Gimma A, van Zandvoort K, Funk S, Jarvis CI, Edmunds WJ, Bosse NI, Hellewell J. The effect of control strategies to reduce social mixing on outcomes of the COVID-19 epidemic in Wuhan, China: a modelling study. *Lancet Public Health* [Internet] 2020 May [cited 2020 May 11];5(5):e261–e270. [doi: 10.1016/S2468-2667(20)30073-6]
4. Flaxman S, Mishra S, Gandy A, Unwin H, Coupland H, Mellan T, Zhu H, Berah T, Eaton J, Perez Guzman P, Schmit N, Cilloni L, Ainslie K, Baguelin M, Blake I, Boonyasiri A, Boyd O, Cattarino L, Ciavarella C, Cooper L, Cucunuba Perez Z, Cuomo-Dannenburg G, Dighe A, Djaafara A, Dorigatti I, Van Elsland S, Fitzjohn R, Fu H, Gaythorpe K, Geidelberg L, Grassly N, Green W, Hallett T, Hamlet A, Hinsley W, Jeffrey B, Jorgensen D, Knock E, Laydon D, Nedjati Gilani G, Nouvellet P, Parag K, Siveroni I, Thompson H, Verity R, Volz E, Walters C, Wang H, Wang Y, Watson O, Winskill P, Xi X, Whittaker C, Walker P, Ghani A, Donnelly C, Riley S, Okell L, Vollmer M, Ferguson N, Bhatt S. Report 13: Estimating the number of infections and the impact of non-pharmaceutical interventions on COVID-19 in 11 European countries [Internet]. Imperial College London; 2020 Mar. [doi: 10.25561/77731]

5. Douglas M, Katikireddi SV, Taulbut M, McKee M, McCartney G. Mitigating the wider health effects of covid-19 pandemic response. *BMJ* [Internet] 2020 Apr 27 [cited 2020 May 11];m1557. [doi: 10.1136/bmj.m1557]
6. United Nations Policy Brief: COVID-19 and the Need for Action on Mental Health [Internet]. 2020 May p. 17. Available from: https://www.un.org/sites/un2.un.org/files/un_policy_brief-covid_and_mental_health_final.pdf
7. Torales J, O'Higgins M, Castaldelli-Maia JM, Ventriglio A. The outbreak of COVID-19 coronavirus and its impact on global mental health. *Int J Soc Psychiatry* [Internet] 2020 Mar 31 [cited 2020 May 20];002076402091521. [doi: 10.1177/0020764020915212]
8. Shigemura J, Ursano RJ, Morganstein JC, Kurosawa M, Benedek DM. Public responses to the novel 2019 coronavirus (2019-nCoV) in Japan: Mental health consequences and target populations. *Psychiatry Clin Neurosci* [Internet] 2020 Apr [cited 2020 May 20];74(4):281–282. [doi: 10.1111/pcn.12988]
9. Moccia L, Janiri D, Pepe M, Dattoli L, Molinaro M, De Martin V, Chieffo D, Janiri L, Fiorillo A, Sani G, Di Nicola M. Affective temperament, attachment style, and the psychological impact of the COVID-19 outbreak: an early report on the Italian general population. *Brain Behav Immun* [Internet] 2020 Apr [cited 2020 May 20];S0889159120305869. [doi: 10.1016/j.bbi.2020.04.048]
10. Wang C, Pan R, Wan X, Tan Y, Xu L, Ho CS, Ho RC. Immediate Psychological Responses and Associated Factors during the Initial Stage of the 2019 Coronavirus Disease (COVID-19) Epidemic among the General Population in China. *Int J Environ Res Public Health* [Internet] 2020 Mar 6 [cited 2020 May 20];17(5):1729. [doi: 10.3390/ijerph17051729]
11. Galea S, Merchant RM, Lurie N. The Mental Health Consequences of COVID-19 and Physical Distancing: The Need for Prevention and Early Intervention. *JAMA Intern Med* [Internet] 2020 Apr 10 [cited 2020 May 20]; [doi: 10.1001/jamainternmed.2020.1562]
12. Beiu C, Mihai M, Popa L, Cima L, Popescu MN. Frequent Hand Washing for COVID-19 Prevention Can Cause Hand Dermatitis: Management Tips. *Cureus* [Internet] 2020 Apr 2 [cited 2020 May 19]; [doi: 10.7759/cureus.7506]
13. MacGibeny MA, Wassef C. Preventing adverse cutaneous reactions from amplified hygiene practices during the COVID-19 pandemic: how dermatologists can help through anticipatory guidance. *Arch Dermatol Res* [Internet] 2020 May 9 [cited 2020 May 19]; [doi: 10.1007/s00403-020-02086-x]
14. The Lancet Oncology. COVID-19: global consequences for oncology. *Lancet Oncol* [Internet] 2020 Apr [cited 2020 May 13];21(4):467. [doi: 10.1016/S1470-2045(20)30175-3]
15. Tan K-K, Lau J. Cessation of cancer screening: An unseen cost of the COVID-19 pandemic? *Eur J Surg Oncol* [Internet] 2020 May [cited 2020 May 19];S0748798320304571. [doi: 10.1016/j.ejso.2020.05.004]
16. Laupacis A. Working together to contain and manage COVID-19. *Can Med Assoc J* [Internet] 2020 Mar 30 [cited 2020 May 19];192(13):E340–E341. [doi: 10.1503/cmaj.200428]

17. Hogan A, Jewell B, Sherrard-Smith E, Vesga J, Watson O, Whittaker C, Hamlet A, Smith J, Ainslie K, Baguelin M, Bhatt S, Boonyasiri A, Brazeau N, Cattarino L, Charles G, Cooper L, Coupland H, Cuomo-Dannenburg G, Dighe A, Djaafara A, Donnelly C, Dorigatti I, Eaton J, Van Elsland S, Fitzjohn R, Fu H, Gaythorpe K, Green W, Haw D, Hayes S, Hinsley W, Imai N, Knock E, Laydon D, Lees J, Mangal T, Mellan T, Mishra S, Nedjati Gilani G, Nouvellet P, Okell L, Ower A, Parag K, Pickles M, Stopard I, Thompson H, Unwin H, Verity R, Vollmer M, Walters C, Wang H, Wang Y, Whittles L, Winskill P, Xi X, Ferguson N, Churcher T, Arinaminpathy N, Ghani A, Walker P, Hallett T. Report 19: The potential impact of the COVID-19 epidemic on HIV, TB and malaria in low- and middle-income countries [Internet]. Imperial College London; 2020 May. [doi: 10.25561/78670]
18. Riley, Sully, Ahmed, Biddlecom. Estimates of the Potential Impact of the COVID-19 Pandemic on Sexual and Reproductive Health In Low- and Middle-Income Countries. *Int Perspect Sex Reprod Health* [Internet] 2020 [cited 2020 May 19];46:73. [doi: 10.1363/46e9020]
19. More than 117 million children at risk of missing out on measles vaccines, as COVID-19 surges [Internet]. Measles Rubella Initiat. [cited 2020 May 11]. Available from: <https://measlesrubellainitiative.org/measles-news/more-than-117-million-children-at-risk-of-missing-out-on-measles-vaccines-as-covid-19-surges/>
20. Hoffman J. Vaccine Rates Drop Dangerously as Parents Avoid Doctor's Visits. *N Y Times* [Internet] 2020 Apr 23 [cited 2020 May 11]; Available from: <https://www.nytimes.com/2020/04/23/health/coronavirus-measles-vaccines.html>
21. PHE Real-time Syndromic Surveillance Team. Emergency Department Syndromic Surveillance System: England [Internet]. Public Health England; 2020 Apr. Available from: https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/882234/EDSSSBulletin2020wk17.pdf
22. Longitudinal studies | COVID-MINDS [Internet]. [cited 2020 May 11]. Available from: <https://www.covidminds.org/longitudinalstudies>
23. Laranjo L, Dunn AG, Tong HL, Kocaballi AB, Chen J, Bashir R, Surian D, Gallego B, Magrabi F, Lau AYS, Coiera E. Conversational agents in healthcare: a systematic review. *J Am Med Inform Assoc* [Internet] 2018 Sep 1 [cited 2020 May 11];25(9):1248–1258. [doi: 10.1093/jamia/ocy072]
24. Lucas GM, Gratch J, King A, Morency L-P. It's only a computer: Virtual humans increase willingness to disclose. *Comput Hum Behav* [Internet] 2014 Aug 1 [cited 2020 May 11];37:94–100. [doi: 10.1016/j.chb.2014.04.043]
25. Miner AS, Milstein A, Schueller S, Hegde R, Mangurian C, Linos E. Smartphone-Based Conversational Agents and Responses to Questions about Mental Health, Interpersonal Violence, and Physical Health. *JAMA Intern Med* [Internet] 2016 May 1 [cited 2020 May 11];176(5):619–625. PMID:26974260
26. Steinhubl SR, Topol EJ. Now we're talking: bringing a voice to digital medicine. *The Lancet* [Internet] 2018 Aug [cited 2020 May 11];392(10148):627. [doi: 10.1016/S0140-6736(18)31803-8]
27. Nobles AL, Leas EC, Caputi TL, Zhu S-H, Strathdee SA, Ayers JW. Responses to addiction help-seeking from Alexa, Siri, Google Assistant, Cortana, and Bixby

- intelligent virtual assistants. *Npj Digit Med* [Internet] 2020 Dec [cited 2020 May 11];3(1):11. [doi: 10.1038/s41746-019-0215-9]
28. Kocaballi AB, Quiroz JC, Rezazadegan D, Berkovsky S, Magrabi F, Coiera E, Laranjo L. Responses of Conversational Agents to Health and Lifestyle Prompts: Investigation of Appropriateness and Presentation Structures. *J Med Internet Res* [Internet] 2020 Feb 9 [cited 2020 May 11];22(2):e15823. [doi: 10.2196/15823]
 29. Bickmore TW, Trinh H, Olafsson S, O’Leary TK, Asadi R, Rickles NM, Cruz R. Patient and Consumer Safety Risks When Using Conversational Assistants for Medical Information: An Observational Study of Siri, Alexa, and Google Assistant. *J Med Internet Res* [Internet] 2018 Sep 4 [cited 2020 May 11];20(9):e11510. [doi: 10.2196/11510]
 30. Ronicke S, Hirsch MC, Türk E, Larionov K, Tientcheu D, Wagner AD. Can a decision support system accelerate rare disease diagnosis? Evaluating the potential impact of Ada DX in a retrospective study. *Orphanet J Rare Dis* [Internet] 2019 Dec [cited 2019 Aug 14];14(1):69. [doi: 10.1186/s13023-019-1040-6]
 31. Menni C, Valdes AM, Freidin MB, Sudre CH, Nguyen LH, Drew DA, Ganesh S, Varsavsky T, Cardoso MJ, El-Sayed Moustafa JS, Visconti A, Hysi P, Bowyer RCE, Mangino M, Falchi M, Wolf J, Ourselin S, Chan AT, Steves CJ, Spector TD. Real-time tracking of self-reported symptoms to predict potential COVID-19. *Nat Med* [Internet] 2020 May 11 [cited 2020 May 19]; [doi: 10.1038/s41591-020-0916-2]
 32. Denis F, Galmiche S, Dinh A, Fontanet A, Scherpereel A, Benezit F, Lescure F-X. Epidemiological Observations on the Association Between Anosmia and COVID-19 Infection: Analysis of Data From a Self-Assessment Web Application. *J Med Internet Res* [Internet] 2020 [cited 2020 Jun 16];22(6):e19855. [doi: 10.2196/19855]
 33. Drew DA, Nguyen LH, Steves CJ, Menni C, Freydin M, Varsavsky T, Sudre CH, Cardoso MJ, Ourselin S, Wolf J, Spector TD, Chan AT, COPE Consortium§. Rapid implementation of mobile technology for real-time epidemiology of COVID-19. *Science* [Internet] 2020 May 5 [cited 2020 May 19];eabc0473. [doi: 10.1126/science.abc0473]
 34. Gilbert S, Mehl A, Baluch A, Cawley C, Challiner J, Fraser H, Millen E, Multmeier J, Pick F, Richter C, Tuerk E, Upadhyay S, Virani V, Vona N, Wicks P, Novorol C. Original research: How accurate are digital symptom assessment apps for suggesting conditions and urgency advice?: a clinical vignettes comparison to GPs [Internet]. *Health Systems and Quality Improvement*; 2020 May. [doi: 10.1101/2020.05.07.20093872]
 35. Wetter und Klima - Deutscher Wetterdienst - Klimaüberwachung - Deutschland - Zeitreihen und Trends [Internet]. [cited 2020 May 12]. Available from: <https://www.dwd.de/DE/leistungen/zeitreihen/zeitreihen.html?nn=480164#buehne>
Top
 36. ICD-10 Version:2019 [Internet]. [cited 2020 May 12]. Available from: <https://icd.who.int/browse10/2019/en>
 37. Whose coronavirus strategy worked best? Scientists hunt most effective policies [Internet]. [cited 2020 Jun 3]. Available from: <https://www.nature.com/articles/d41586-020-01248-1>

38. Papanicolas I, Mossialos E, Gundersen A, Woskie L, Jha AK. Performance of UK National Health Service compared with other high income countries: observational study. *BMJ* [Internet] 2019 Nov 27 [cited 2020 Jun 3];l6326. [doi: 10.1136/bmj.l6326]
39. Koch-Institut R. *Epidemiologisches Bulletin* 16/2020. 2020;11.
40. Information NC for B, Pike USNL of M 8600 R, MD B, Usa 20894. Allergies: Overview [Internet]. *Inf Internet*. Institute for Quality and Efficiency in Health Care (IQWiG); 2020 [cited 2020 Jun 3]. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK447112/>
41. Hebbelstrup Jensen B, Jokelainen P, Nielsen ACY, Franck KT, Rejkjær Holm D, Schønning K, Petersen AM, Krogfelt KA. Children Attending Day Care Centers are a Year-round Reservoir of Gastrointestinal Viruses. *Sci Rep* [Internet] 2019 Dec [cited 2020 May 21];9(1):3286. [doi: 10.1038/s41598-019-40077-9]
42. Bhattacharya S, Singh A, Rillera Marzo R, 1 Department of Community Medicine, Himalayan Institute of Medical Sciences, Dehradun, India, 2 Department of Community Medicine, School of Public Health, Postgraduate Institute of Medical Education and Research, Chandigarh, India, 3 Department of Public Health, Asia Metropolitan University, Johor Bahru, Malaysia. “Airplane ear”—A neglected yet preventable problem. *AIMS Public Health* [Internet] 2019 [cited 2020 May 21];6(3):320–325. [doi: 10.3934/publichealth.2019.3.320]
43. Bénézit F, Le Turnier P, Declerck C, Paillé C, Revest M, Dubée V, Tattevin P, Arvieux C, Baldeyrou M, Chapplain J-M, Comacle P, Patrat-Delon S, Maillard A, Poinot M, Pronier C, Souala F, Thibault V, Abgueguen P, Cormier H, Delbos V, de la Chapelle M, Ducancelle A, Mahieu R, Rabier V, Rehaiem S, Vandamme Y, Biron C, Brochon J, Boutoille D, Chauveau M, Deschanvres C, Gaborit BJ, Jenvrin J, Lecomte R, Lefebvre M, Raffi F. Utility of hyposmia and hypogeusia for the diagnosis of COVID-19. *Lancet Infect Dis* [Internet] 2020 Apr [cited 2020 May 20];S1473309920302978. [doi: 10.1016/S1473-3099(20)30297-8]
44. Lao WP, Imam SA, Nguyen SA. Anosmia, hyposmia, and dysgeusia as indicators for positive SARS-CoV-2 infection. *World J Otorhinolaryngol - Head Neck Surg* [Internet] 2020 Apr [cited 2020 May 20];S2095881120300330. [doi: 10.1016/j.wjorl.2020.04.001]
45. Sommer IE, Bakker PR. What can psychiatrists learn from SARS and MERS outbreaks? *Lancet Psychiatry* [Internet] 2020 May [cited 2020 May 20];S2215036620302194. [doi: 10.1016/S2215-0366(20)30219-4]
46. Van Bortel T, Basnayake A, Wurie F, Jambai M, Koroma AS, Muana AT, Hann K, Eaton J, Martin S, Nellums LB. Psychosocial effects of an Ebola outbreak at individual, community and international levels. *Bull World Health Organ* [Internet] 2016 Mar 1 [cited 2020 May 20];94(3):210–214. [doi: 10.2471/BLT.15.158543]
47. Holmes EA, O’Connor RC, Perry VH, Tracey I, Wessely S, Arseneault L, Ballard C, Christensen H, Cohen Silver R, Everall I, Ford T, John A, Kabir T, King K, Madan I, Michie S, Przybylski AK, Shafran R, Sweeney A, Worthman CM, Yardley L, Cowan K, Cope C, Hotopf M, Bullmore E. Multidisciplinary research priorities for the COVID-19 pandemic: a call for action for mental health science. *Lancet Psychiatry* [Internet] 2020 Apr [cited 2020 May 11];S2215036620301681. [doi: 10.1016/S2215-0366(20)30168-1]

48. Agteren J van, Bartholomaeus J, Fassnacht DB, Iasiello M, Ali K, Lo L, Kyrios M. Using Internet-Based Psychological Measurement to Capture the Deteriorating Community Mental Health Profile During COVID-19: Observational Study. *JMIR Ment Health* [Internet] 2020 [cited 2020 Jun 16];7(6):e20696. [doi: 10.2196/20696]
49. HOME | CovidSocialStudy [Internet]. [cited 2020 May 20]. Available from: <https://www.covidsocialstudy.org/>
50. Jacobson NC, Lekkas D, Price G, Heinz MV, Song M, O'Malley AJ, Barr PJ. Flattening the Mental Health Curve: COVID-19 Stay-at-Home Orders Are Associated With Alterations in Mental Health Search Behavior in the United States. *JMIR Ment Health* [Internet] 2020 [cited 2020 Jun 16];7(6):e19347. [doi: 10.2196/19347]
51. Farr M. Australians are as happy as before coronavirus lockdown, Guardian Essential poll finds. *The Guardian* [Internet] 2020 Apr 13 [cited 2020 May 21]; Available from: <https://www.theguardian.com/australia-news/2020/apr/14/australians-are-as-happy-as-before-coronavirus-lockdown-guardian-essential-poll-finds>
52. Vu V. Wie geht es Ihnen heute: Geht es Ihnen seit Corona besser? *Zeit* [Internet] Hamburg; 2020 Apr 29 [cited 2020 May 21]; Available from: <https://www.zeit.de/gesellschaft/zeitgeschehen/2020-04/wie-geht-es-ihnen-heute-warum-besser>
53. Jarral F. The lockdown paradox: why some people's anxiety is improving during the crisis | Farrah Jarral. *The Guardian* [Internet] 2020 Apr 29 [cited 2020 May 21]; Available from: <https://www.theguardian.com/commentisfree/2020/apr/29/coronavirus-lockdown-anxiety-mental-health>
54. Jungmann SM, Klan T, Kuhn S, Jungmann F. Accuracy of a Chatbot (Ada) in the Diagnosis of Mental Disorders: Comparative Case Study With Lay and Expert Users. *JMIR Form Res* [Internet] 2019 Oct 29 [cited 2019 Dec 16];3(4):e13863. [doi: 10.2196/13863]
55. Nateqi J, Lin S, Krobath H, Gruarin S, Lutz T, Dvorak T, Gruschina A, Ortner R. Vom Symptom zur Diagnose – Tauglichkeit von Symptom-Checkern: Update aus Sicht der HNO. *HNO* [Internet] 2019 May [cited 2020 May 21];67(5):334–342. [doi: 10.1007/s00106-019-0666-y]

Abbreviations

°C: degrees Celsius

COVID-19: Coronavirus disease 2019

GDPR: General Data Protection Regulation

ICD-10: International Classification of Diseases version 2019

Log2 FC: Log2 Fold Change

mm: millimeters

P-value: probability value

UK: United Kingdom