

1 Body mass index multiple regression formula testable by
2 eight Bradford Hill causality criteria: Worldwide
3 ecological cohort data analysis to inform dietary guidance

4

5

6 David K Cundiff (0000-0002-3206-9665), independent researcher,^{1 3} Chunyi Wu (0000-0002-
7 2186-3433), Research Epidemiologist/Statistician^{2 3}

8 ¹ Long Beach, California, USA

9 ² Area Specialist Lead in Epidemiology and Statistics, Michigan Medicine, Ann Arbor,
10 Michigan, USA

11 ³ Volunteer collaborators with the Institute of Health Metrics and Evaluation, Seattle,
12 Washington, USA

13 Correspondence to: David K Cundiff davidkcundiff@gmail.com Phone: 01-562-438-8805

14 Word count: abstract=399, text=3603

15

16 **Abstract**

17 **Objectives** (1) Derive a worldwide body mass index (BMI) multiple regression formula (BMI
18 formula), (2) Quantify the percent weights attributable to dietary and other risk factors from this
19 BMI formula, and (3) Test relevant dietary guidelines and diets using the BMI formula generated
20 BMI estimates

21 **Design** BMI and risk factor cohort data from 1990-2017 from the Institute of Health Metrics and
22 Evaluation (IHME) were formatted and population-weighted. We empirically explored the
23 univariate and multiple regression correlations of risk factors with BMI to maximize BMI
24 formula functionality.

25 **Setting** Worldwide

26 **Ecological cohort data** Over 12,000 Global Burden of Disease (GBD) risk factor surveys of
27 people 15-69 years old from 195 countries analysed and synthesised into representative cohort
28 risk factor values by staff and volunteer collaborators with the IHME

29 **Main outcome measures** Performance of a worldwide BMI formula when tested with eight
30 Bradford Hill causality criteria, each scored with 0-5 scale: 0=negative to 5=very strong support
31 (40 possible score)

32 **Results** In the BMI formula derived, all foods were expressed in kilocalories/day (kcal/day), and
33 all risk factor coefficients were adjusted to equate with their percent weight impacts on BMI.
34 BMI formula =0.29*processed meat +11.87*red meat +0.02*fish +3.74*milk +11.09*poultry
35 +5.46*eggs +1.95*alcohol +6.29*sugary beverages +0.29*corn +3.53*potatoes +2.71 *
36 saturated fatty acids+ 0.64*polyunsaturated fatty acids+0.06*trans fatty acids-0.37*fruit -
37 0.52*vegetables -0.03*nuts and seeds -0.12 *whole grains -0.47*legumes-8.26*rice -18*sweet
38 potatoes -9.93*physical activity (METs/week) +5.78*total kcals/day available-4.13*child

39 underweight+0.92*discontinued breast feeding. BMI increasing foods have + signs in the BMI
40 formula and BMI decreasing foods have - signs. Bradford Hill causality criteria test scores (0-5):
41 (1) strength=5, (2) consistency=5, (3) dose-response=5, (4) temporality=5, (5) analogy=4, (6)
42 plausibility=5, (7) experimentation=5 and (8) coherence=5. Total=39/40. Based on the USA
43 BMI trend data, the mean adult BMI in 1990 was 25.45, is projected to be 28.13 in 2020, and
44 31.81 in 2050. BMI formula based mean BMI estimates included the following: Dietary
45 Guidelines for Americans=26.34, USA with 50% reduction of BMI increasing food intake
46 =23.34, USA with 25% reduction of BMI increasing food intake and increasing physical activity
47 by running at 6 mph for 1 hour/day on average=23.67, vegetarian diet=22.54, low carbohydrate,
48 high fat diet=31.76.

49 **Conclusions** Eight relevant Bradford Hill causality criteria strongly supported that the BMI
50 formula derived was causally related to mean BMIs of worldwide cohorts, subgroups, and
51 individual risk factor patterns. The statistical analysis methodology introduced could inform
52 individual, clinical, and public health strategies regarding overweight/obesity
53 prevention/treatment and other health outcomes.

54

55 **Introduction**

56 The scientific validity of the Dietary Guidelines for Americans for 2015-2020 was challenged by
57 Journalist Nina Teicholz in the *BMJ*.¹ The Center for Science in the Public Interest called for the
58 *BMJ* to retract the article. The peer reviewers selected to adjudicate the far reaching dispute
59 concluded that, “Teicholz’s criticisms of the methods used by Dietary Guidelines for Americans
60 Committee are within the realm of scientific debate.”² Regarding scientific study of treatments of
61 overweight/obesity, *JAMA* published a representative randomised trial comparing a low fat diet
62 with a low carbohydrate diet. It reported no difference in the very modest weight loss achieved at
63 one year.³ This suggests that current short term dietary intervention trials (< 5 years) are
64 meaningless in determining the long-term relationship of diet (> 20 years) with
65 overweight/obesity.

66
67 Influential Stanford University meta researcher, Dr. John Ioannidis, called for radical reform of
68 all nutritional epidemiology methodologies used to influence food/agricultural policies and to
69 produce dietary guidelines for clinicians and the public.⁴ Currently, no methodology for relating
70 body mass index (BMI) long-term or BMI change/year to food intake and physical activity has
71 been generally accepted as rigorous, replicable, and scientifically valid. This study analysing
72 worldwide data for dietary and other risk factors for BMI attempts to answer Dr. Ioannidis’ call
73 for a more rigorous and reliable nutritional epidemiology methodology to base public health
74 policies and individual dietary guidance.

75
76 The first objective of this analysis was to derive a multiple regression risk factor formula using
77 worldwide ecological data on BMIs of male and female cohorts (dependent variable) and dietary

78 and other risk factors (independent variables). Secondly, we used Bradford Hill causality criteria
79 to test the BMI formula containing worldwide diet and other risk factors. Satisfying Bradford
80 Hill causality criteria is considered validating in epidemiological research.⁵ Third we wanted to
81 apply the BMI formula to assorted diet and other risk factor patterns to derive mean BMI
82 estimates associated with long-term (>20 years) following of those diet and other risk factor
83 patterns. We only hypothesised that the objectives were achievable.

84

85

86

87 **Methods**

88 As volunteer collaborators with the Institute of Health Metrics and Evaluation (IHME), we
89 received raw Global Burden of Disease (GBD) ecological data (≈ 1.4 Gigabytes) on mean BMIs
90 of male and female cohorts 15-49 years old and 50-69 years old from each year 1990-2017 from
91 195 countries and 365 subnational locations ($n=1\,120$ cohorts). We also utilised GBD data on
92 exposures to 32 risk factors and covariates potentially related to BMI. IHME dietary covariate
93 data originally came from Food and Agriculture Organization surveys of animal and plant food
94 commodities available percapita—as opposed to consumed per capita—in countries worldwide.⁶
95 Food risk factors came as g/day consumed percapita. Other variables were utilised as part of
96 deriving the BMI formula and testing the BMI formula according to the Bradford Hill causality
97 criteria: physical activity (METs/day), kilocalories per day available percapita (kcal/day, a
98 covariate), severe child underweight (2 SD below the mean weight for age), and discontinued
99 breast feeding before 6 months. Other available variables tested included socio-demographic
100 index, LDL-cholesterol, fasting plasma glucose and, systolic blood pressure. Supplementary
101 Table 1 lists the relevant GBD risk factors, covariates, and other available variables with
102 definitions of those risk factor exposures.⁷
103
104 GBD worldwide citations of over 12,000 surveys constituting ecological data inputs for this
105 analysis are in Appendix 1.^{6,8} The main characteristics of IHME GBD data sources for BMI and
106 all risk factor values have been published by IHME GBD data researchers and discussed
107 elsewhere.⁹⁻¹¹ These included detailed descriptions of categories of input data, potentially
108 important biases, and methodologies of analysis. We did not clean or pre-process any of the
109 GBD data. GBD cohort risk factor and BMI data from the IHME had no missing records.

110 Because this was a post hoc analysis and the GBD data for the study came from IHME, no ethics
111 committee approval or institutional review board review was needed for this statistical analysis.

112 The raw data for this analysis may be obtained by volunteer researchers collaborating with
113 IHME.¹²

114
115 IHME made available GBD BMI, risk factor, and covariate exposure data for each year from
116 1990-2017 for male and female cohorts from ages 15-49 years old and from 50-69 years old. To
117 maximally utilise the available data, we averaged the values for ages 15-49 years old together
118 with 50-69 years old for BMI and for each risk factor exposure for each male and female cohort
119 for each year. Finally, for each male and female cohort, data from all 28 years (1990-2017) on
120 mean BMI and on each of the risk factor exposures were averaged using the computer software
121 program R.

122
123 To weigh the country and subnational data according to population, internet searches (mostly
124 Wikipedia) yielded the most recent population estimates for countries and subnational states,
125 provinces, and regions. World population data from the World Bank and the Organisation for
126 Economic Co-operation and Development could not be used because they did not include all 195
127 countries or any subnational data.

128
129 Using the above described formatted dataset of risk factors, covariates, and BMIs, a software
130 program in R generated a population-weighted analysis dataset. Each male or female cohort in
131 the population-weighted analysis dataset represented approximately 1 million people (range: <
132 100,000 to 1.5 million). The analysis dataset had n=7886 cohorts (rows of data), half male and

133 half female, representative of over seven billion people. For example, India with about 1.234
134 billion people had 617 rows of the same data for males and 617 rows for females. Maldives, with
135 about 445,000 people, had a single row of data for males and another for females. Without
136 population-weighting the data, cohorts in India and Maldives each would have had one row in
137 the analysis dataset, invalidating the analysis results.

138
139 This report follows the STrengthening the Reporting of OBservational studies in Epidemiology
140 (STROBE) guidelines for reporting global health estimates.¹³

141
142 Supplementary Table 2 details how omega-3 fatty acid g/day was converted to fish g/day using
143 data on the omega-3 fatty acid content of frequently eaten fish from the National Institutes of
144 Health Office of Dietary Supplements (USA).¹⁴ As shown in Supplementary Table 3, we
145 converted all of the animal and plant food data, including alcohol and sugary beverage
146 consumption, from g/day to kcal/day. For the g to kcal conversions, we used the Nutritionix track
147 app,¹⁵ which tracks types and quantities of foods consumed. Saturated fatty acids risk factor (0-1
148 portion of the entire diet) was not available with GBD data from 2017, so GBD saturated fatty
149 acids risk factor data from 2016 was used. Polyunsaturated fatty acid and trans fatty acid GBD
150 risk factor data from 2017 (0-1 portion of the entire diet) was also utilised. These fatty acid data
151 were converted to kcal/day by multiplying by the kcal/day available for each cohort.

152

153 **Statistical methods**

154 To determine the strengths of the risk factor correlations with mean BMIs of population
155 weighted worldwide cohorts (7886 cohorts) or subgroups of cohorts (e.g., continents, socio-

156 demographic quartiles, etc.), we utilised Pearson correlation coefficients: r , 95% confidence
157 intervals (95% CIs), and P values.

158

159 In this first of its kind, post hoc, exploratory analysis; the methodology was determined as we
160 proceeded by experimenting with strategies to optimise the functioning of the BMI formula
161 derived. We sought to derive a BMI multiple regression formula from worldwide data with the
162 following characteristics:

163

- 164 1. including as many as possible of the available animal and plant food variables,
- 165 2. including physical activity and other plausibly informative available ecological variables,
166 and
- 167 3. combining dietary variables if appropriate to utilise more variables and improve BMI
168 formula strength (i.e., R^2 with BMI) and consistency at predicting subgroup mean BMIs.

169

170 Appendix 2 further details the statistical methodology and explains the use of Bradford Hill
171 causality criteria to assess whether the worldwide BMI was causally related to the risk factors in
172 the BMI formula. Briefly, we tested the BMI formula output with the Bradford Hill causality
173 criteria (1) strength, (2) consistency, (3) dose response, (4) temporality, (5) analogy, (6)
174 plausibility, (7) experimentation, and (8) coherence. For each criterion, we used a 0-5 scale to
175 assess the magnitude of support of the BMI formula output being causally related to the BMIs
176 worldwide (0=no support of causality to 5=very strong support of causality).

177

178 In determining the variables to include and exclude in worldwide BMI formula, we set the
179 statistical threshold for a variable to enter and to remain in the formula at $P < 0.25$. We used
180 SAS and SAS Studio statistical software 9.4 (SAS Institute, Cary, NC) for the data analysis.

181 **Results**

182 Table 1 shows the basic statistics and univariate correlations of mean BMI with dietary and other
183 risk factors and with the BMI formula derived from worldwide cohorts. Table 2 shows the
184 univariate correlations of each of the 24 BMI formula risk factors with each other. All six animal
185 foods (processed meat, red meat, fish, milk, poultry, and eggs), alcohol, sugary beverages, corn
186 availability, potato availability, saturated fatty acids, polyunsaturated fatty acids, and trans fatty
187 acids all positively correlated with BMI. Table 2 shows that corn availability (kcal/day percapita,
188 a covariate) correlated moderately strongly with sugary beverages ($r=0.419$, 95% CI 0.400 to
189 0.437, $P<0.0001$), suggesting that high fructose corn syrup may account for the positive
190 correlation with BMI. Potato availability (kcal/day percapita, a covariate), which positively
191 correlated with BMI, included $\geq 50\%$ highly processed potato products worldwide according to
192 the International Potato Center.¹⁶ Four starchy plant foods (whole grains, legumes, rice, and
193 sweet potatoes) all negatively correlated with BMI.

194

195 The strong positive correlations of fruits, vegetables, and nuts and seeds with BMI were
196 unexpected since nutritional literature broadly supports these foods as non-obesogenic. It
197 suggested possible multicollinearity (when an independent variable is highly correlated with one
198 or more known or unknown other independent variables). Supplementary Table 4 demonstrates
199 the multicollinearity of fruits, vegetables, and nuts and seeds with BMI increasing foods:

200

- 201 • BMI increasing foods (animal foods + alcohol + sugary beverages + corn + potatoes) as a
202 group were very strongly correlated with BMI ($r= 0.597$, 95% CI 0.583 to 0.611, $P<$
203 0.0001).

- 204 • Fruits, vegetables, nuts and seeds grouped together were strongly positively correlated
205 with BMI ($r= 0.655$, 95% CI 0.642 to 0.677, $P< 0.0001$) but also with BMI increasing
206 foods ($r= 0.323$, 95% CI 0.304 to 0.343, $P< 0.0001$).

207 Supplementary Table 4 also shows that the top sociodemographic index (SDI) cohort quartile's
208 mean BMI was about 3.6 BMI units higher than the three lowest SDI quartiles (24.54 versus
209 20.88), meaning that the more developed countries had much higher mean BMIs. BMI increasing
210 foods in the top SDI quartile exceeded intake in the three lowest quartiles by 58% (905/kcal/day /
211 571 kcal/day=1.58). Fruits, vegetables, and nuts and seeds in the top SDI quartile exceeded
212 intake in the three lower quartiles by 1.9 to 1 (199/kcal/day / 106 kcal/day=1.88). Because of the
213 multicollinearities of fruits, vegetables and nuts and seeds with the 10 BMI increasing foods, we
214 grouped fruits, vegetables and nuts and seeds with BMI decreasing foods (including whole
215 grains, legumes, rice, and sweet potatoes) in a combination variable used to derive the BMI
216 formula, gives the coefficients of risk factors equated to their percent weights (shown in
217 Supplementary Table 5):

218
219 BMI formula =0.29*processed meat +11.87*red meat +0.02*fish +3.74*milk
220 +11.09*poultry +5.46*eggs +1.95*alcohol +6.29*sugary beverages +0.29*corn
221 +3.53*potatoes +2.71 * saturated fatty acids+ 0.64*polyunsaturated fatty
222 acids+0.06*trans fatty acids-0.37*fruit -0.52*vegetables -0.03*nuts and seeds -
223 0.12 *whole grains -0.47*legumes-8.26*rice-0.18*sweet potatoes -9.93*physical
224 activity (METs/week) +5.78*total kcals/day available-4.13*child
225 underweight+0.92*discontinued breast feeding. Percent weights of risk factors
226 were equated to risk factor coefficients. Total percent weights=78.64. (BMI

227 formula $R^2=0.7864$, $r=0.887$, 95% CI 0.882 to 0.891, $P<0.0001$, + signs mean
228 BMI increasing and - signs mean BMI decreasing).

229
230 As detailed in the methods and Appendix 2, the standardised BMI formula coefficients of the
231 dietary and other risk factors were all adjusted to equate to the percent weights of risk factors in
232 the BMI formula.

233
234 As shown in Supplementary Table 5, the 13 BMI increasing foods and seven BMI decreasing
235 foods underwent three adjustments: (1) multiplied times their mean kcal/day values (Column C),
236 (2) multiplied times their R^2 values in univariate correlation with BMI (Column E), and (3)
237 multiplied times the ratio of the mean kcal/day of the second, third, and fourth sociodemographic
238 index (SDI) quartiles divided by the mean kcal/day of the first SDI quartile (Column G). Table 3
239 demonstrates that these adjustments improved the consistency of the BMI formula as measured
240 by the closeness of fit of the multiple regression derived BMI formula outputs with the mean
241 BMIs of 37 subgroups of the worldwide data.

242

243 **BMI formula output analysed by Bradford Hill causality criteria**

244 Eight Bradford Hill causality criteria tested the functionality of the BMI formula. With the
245 methodology detailed in Appendix 2, we assessed the relationship of the worldwide BMI
246 formula output versus the mean BMI as follows:

- 247
- 248 **1.** Strength=5—The correlation coefficient of the BMI formula regressed with BMI: $r=$
249 0.887 (95% CI: 0.882 to 0.891), $P<0.0001$.

- 250 **2.** Consistency score=5— Table 3 shows the average absolute differences between subgroup
251 mean BMIs and BMI formula estimates of the BMI formula compared with two other
252 variations of the BMI formula. The BMI formula with fruits, vegetables, and nuts and
253 seeds grouped with BMI decreasing foods as negatively correlated with BMI had a
254 slightly closer fit than a comparison BMI formula with fruits, vegetables, and nuts and
255 seeds grouped with BMI increasing foods positively correlated with BMI (average BMI -
256 BMI formula estimate (absolute difference): 0.249 versus 0.255 BMI units). Without the
257 three adjustments to each of the food variables, the average BMI - BMI formula estimate
258 absolute difference was 0.368 BMI units.
- 259 **3.** Dose-response (Biological gradient) score=5—Table 3 shows that the mean of the BMI
260 absolute differences between the BMI formula estimates and mean BMIs and in the four
261 dose-response quartiles was 0.302 BMI units.
- 262 **4.** Temporality score=5—a multiple regression with the worldwide BMI trend 1990-2007
263 (dependent variable) versus the risk factor trends (independent variables) generated a
264 BMI trend formula versus risk factor BMI trends: $r = 0.615$, (95% CI: 0.601 to 0.628),
265 $P < 0.0001$). Fifteen out of 24 risk factor trends contributed to the BMI trend formula. The
266 derivation of standardised BMI trend versus risk factors trends formula with sign
267 concordant coefficients adjusted to equal the percent weights of the significant risk
268 factors shown in Supplementary Table 7 was as follows:
- 269 BMI trend formula= $4.84 * \text{Red meat} + 0.63 * \text{Milk} + 4.61 * \text{Poultry} + 2.62 * \text{Eggs} + 3.91$
270 $* \text{Alcohol} + 2.69 * \text{Sugary beverages} + 0.40 * \text{corn} + 3.60 * \text{Potatoes} + 3.90 * \text{PUFA} -$
271 $0.65 * \text{Legumes} - 3.08 * \text{Rice} - 0.80 * \text{Sweet potatoes} + 6.06 * \text{kcal available}$ (total
272 percent weight 37.79, $R^2 = 0.3779$).
- 273

- 274 **5.** Analogy score=4—Systolic blood pressure had 15/24 risk factors concordant in
275 sign with BMI. SBP correlated weakly with the BMI formula ($r=0.024$, 95% CI
276 0.002 to 0.046, $P=0.0361$). The low density lipoprotein cholesterol (LDL-C) and
277 the fasting plasma glucose (FPG) formulas had 23/24 and 21/24 risk factors
278 concordant in sign with the BMI formula, respectively. The LDL-C and the FPG
279 were both strongly correlated with BMI ($r=0.757$, 95% CI 0.747 to 0.766, $P <$
280 0.0001 and $r=0.565$, 95% CI 0.550 to 0.580, $P < 0.0001$, respectively). They
281 were also strongly correlated with the BMI formula ($r=0.757$, 95% CI 0.747 to
282 0.766, $P < 0.0001$ and $r=0.581$, 95% CI 0.566 to 0.595), $P < 0.0001$, respectively.
- 283 **6.** Plausibility: Score=5—Based on systematic medical literature reviews, physical activity
284 inversely correlated with BMI¹⁷ and BMI directly correlated with intakes of sugar,¹⁸
285 alcohol,¹⁹ and animal foods.²⁰ The relationship of adult BMI with early childhood severe
286 underweight has not been reported worldwide. Since people in poor countries have less
287 animal foods and alcohol and more physical exercise than in developed countries, it is
288 plausible that childhood severe underweight positively correlates with lower BMI in
289 adulthood.
- 290 **7.** Experiment: Score=5—Table 4 demonstrates cross validation of the BMI formula with a
291 very high degree of reproducibility in the percent weight distributions from 20 randomly
292 selected subgroups of the dataset ($n=100$ for each, which are shown in Supplementary
293 Table 7).
- 294 **8.** Coherence: Score=5—As evidenced by the near perfect score 34/35 on the first seven
295 criteria, the Bradford Hill causality criteria overall were strongly supportive that the

296 worldwide BMI formula accurately modeled the 24 risk factors that led to increased and
297 decreased BMI—total causality criteria score=39/40.

298
299 Relative to other countries, the USA had one of the world’s highest levels of kcal/day of BMI
300 increasing foods (exclusive of fatty acids), 1015 kcal/day and below average consumption of
301 BMI decreasing foods, 326 kcal/day, corresponding to an average adult BMI=26.66. Based on
302 the USA BMI trend data, the mean adult BMI in 1990 was 25.45 and is projected to be 28.13 in
303 2020, and 30.81 in 2050. Table 5 shows BMI formula estimates for various relevant patterns of
304 diet and/or other BMI formula risk factors. Following the Dietary Guidelines for Americans
305 2015 -2020 would result in a mean adult BMI of 26.34, which is almost the mean USA adult
306 BMI from 1990 - 2017 (26.66). A major increase in exercise on a regular basis alone would
307 decrease the BMI formula estimated mean long term BMI. In addition to the current mean level
308 of physical activity (3853 METs/week), running two hours/day on average at 6 miles per hour
309 (8400 METs/week)²¹ would reduce the estimated mean BMI by 2.68 BMI units to 23.98.

310 **Discussion**

311 The eight Bradford Hill criteria used in this analysis of long term worldwide data all support that
312 the BMI formula is causally related to adult BMI.

313

314 Short of extreme and unrealistic for most people increases in exercise to control weight, dietary
315 change from the USA current dietary patterns or even the USA dietary guidelines recommended
316 diet would be required to reach the normal range for BMI ($BMI \geq 18.5$ and $BMI < 25$). For
317 example, shifting 50% of kcal/day from BMI increasing foods to BMI decreasing foods equates
318 to a BMI formula estimate=23.26 BMI units over the long term. While following a low
319 carbohydrate, high fat diet has been demonstrated to cause modest short term weight loss in
320 obese people,²² the BMI formula projects that the long-term effect (> 20 years) would be obesity
321 (projected mean BMI=31.76).

322

323 The perhaps counterintuitive findings that fruits, vegetables, and nuts and seeds were positively
324 correlated with BMI ($r=0.655$, 95% CI 0.642 to 0.677, $P < 0.0001$) may be largely explained by
325 the multicollinearity between fruits, vegetables, and nuts and seeds and BMI increasing foods
326 ($r=0.323$, 95% CI 0.304 to 0.343, $P < 0.0001$, Table 2). The high cost of fruits, vegetables, and
327 nuts and seeds may account for much of this multicollinearity both worldwide and within
328 wealthy countries that have high economic inequality levels like the USA. A systematic review
329 of the literature on food cost relative to nutrient quality found that the median costs of starches
330 (€0.14/100 kcal) was quite low relative to fruits and vegetables (€0.82/100 kcal), meat/eggs/fish
331 (€0.64/100 kcal), fresh dairy (€0.32/100 kcal), and nuts (€0.25/100 kcal).²³

332

333 The findings in this study, particularly those in Table 5, should influence government food policy
334 decisions. The Supplemental Nutrition Assistance Program (SNAP—formerly Food Stamps)
335 spends an estimated 22.6% of its \$73 billion/year budget²⁶ on payments to low-income
336 Americans for “sweetened beverages, prepared desserts, salty snacks, candy, and sugar.”²⁷
337 Additionally, the US Department of Agriculture (USDA) subsidises crops that go primarily for
338 animal feed or that are processed into sugars while not subsidizing fruits and vegetables.²⁴ While
339 the USDA recognises the relatively low intake of fruits and vegetables in the USA and sponsors
340 a publicity campaign to increase fruits and vegetables,²⁵ USDA expenditures should promote
341 reduced prices of BMI decreasing foods and increased prices of BMI increasing foods.

342
343 The strong correlation of discontinuation of breastfeeding with BMI ($r=0.802$, 95% CI 0.794 to
344 0.810, $P<0.0001$) was in accord with a meta-analysis of breast feeding related to subsequent
345 childhood and adult BMI by Jeanne Stolzer.²⁶ However, the estimated benefit in reducing adult
346 BMI of breastfeeding in this study was very modest (0.19 BMI unit, see Table 5). Breastfeeding
347 for at least six months is recommended by the American Academy of Pediatrics^{27,28} and the
348 National Institute of Clinical Excellence.²⁹

349

350 **Limitations**

351 The GBD data on animal and plant foods were not comprehensive and comprised only 1199
352 kcal/day on average. Subnational data were available on only four countries. Because the data
353 formatting and statistical methodology were new, this was necessarily a post hoc analysis and no
354 pre-analysis protocol was possible. As detailed in the Foresight Report on obesity,³⁰ obesity is
355 affected by a complex system of interacting factors besides diet, physical activity, and breast

356 feeding. So genes,³¹ gut microbiome,³² ultra processing of food,^{33,34} and other influences on BMI
357 were outside of the purview of this analysis.

358

359 **Generalisability**

360 Given the strength and consistency of the relationship of the BMI formula in modeling BMI
361 (SDI quartiles, continents, and subnationals, etc., see Tables 3 and 5), the findings should be
362 generalisable to people all over the world.

363 **Conclusion**

364 Eight Bradford Hill causality criteria strongly supported that the worldwide obesity epidemic is
365 causally related to the 24 risk factors in proportion to their coefficients in the BMI formula. The
366 findings in this study should be considered by health policymakers drafting dietary guidelines for
367 healthy weight management. While this study deals only with dietary and other risk factors for
368 BMI (overweight/obesity), the methodology introduced could easily apply to estimating percent
369 weights of multiple dietary and other risk factors that pertain to dozens of non communicable
370 diseases, for which the IHME have GBD data.

371

372

373

374
375

Table 1. BMI risk factor basic statistics and Bradford Hill criterion strength

BMI and risk factors: n=7886	Mean	Std Dev	Mini- mum	Maxi- mum	r	95% CI	P	R²
BMI kg/M2	21.79	2.31	17.95	29.39				
BMI formula	21.79	2.31	18.15	30.32	0.887	0.882 to 0.891	<.0001	0.791
Processed kcal/day	5.49	9.94	0.20	68.8	0.604	0.590 to 0.618	<.0001	0.365
Red meat kcal/day	50.70	45.45	3.21	235.9	0.662	0.649 to 0.674	<.0001	0.438
Fish kcal/day	10.01	36.43	0.40	370.4	0.103	0.081 to 0.124	<.0001	0.011
Milk kcal/day	25.37	27.39	1.06	146.8	0.684	0.672 to 0.696	<.0001	0.468
Poultry kcal/day avail	45.06	51.03	1.06	411.9	0.813	0.805 to 0.820	<.0001	0.661
Eggs kcal/day avail	19.47	14.76	0.79	69.6	0.688	0.676 to 0.699	<.0001	0.473
Alcohol kcal/day	81.71	58.06	4.25	430	0.166	0.145 to 0.188	<.0001	0.028
Sugary beverages kcal/day	297.65	152.33	73	1472	0.119	0.097 to 0.141	<.0001	0.014
Corn kcal/day avail	34.67	48.16	0.16	305.2	0.073	0.051 to 0.095	<.0001	0.005
Potatoes kcal/day avail	84.16	74.43	3.07	533.9	0.211	0.189 to 0.232	<.0001	0.044
Saturated FA % kcal available	191.27	64.49	70.79	481.1	0.710	0.699 to 0.721	<.0001	0.504
Polyunsaturated fatty acids kcal %	82.24	74.98	2.93	381.31	0.737	0.727 to 0.747	<.0001	0.543
Trans fatty acids % kcal	13.4	13.82	1.99	77.76	0.488	0.471 to 0.504	<.0001	0.238
Fruits kcal/day	40.39	22.59	3.58	161.4	0.622	0.609 to 0.636	<.0001	0.387
Vegetables kcal/day	80.14	43.36	9.48	304.2	0.520	0.503 to 0.536	<.0001	0.270
Nuts and seeds kcal/day	8.51	8.44	0.05	103.0	0.486	0.469 to 0.503	<.0001	0.237
Whole grains kcal/day	55.65	30.85	1.14	235.1	-0.201	-0.222 to -0.179	<.0001	0.040
Legumes kcal/day	51.66	32.16	0.51	194.7	-0.384	-0.403 to -0.365	<.0001	0.148
Rice kcal/day avail	141.23	116.38	1.42	461.8	-0.561	-0.576 to -0.545	<.0001	0.314
Sweet potatoes kcal/day avail	22.67	35.88	0.02	364.7	-0.152	-0.173 to -0.130	<.0001	0.023
Physical activity METs	4709	1367	1609	7669	-0.445	-0.462 to -0.427	<.0001	0.198
Kcal available/day	2579	421	1579	3898	0.843	0.837 to 0.850	<.0001	0.711
Child/infant underweight	0.19	0.17	0.00	0.53	-0.794	-0.802 to -0.786	<.0001	0.631
Discontinued breast feeding< 6 months	0.12	0.06	0.02	0.24	0.802	0.794 to 0.810	<.0001	0.644
Sodium g/day	4.45	2.33	1.33	9.21	-0.019	-0.041 to -0.003	0.099	0.000
Calcium g/day	0.3	0.18	0.08	1.04	0.771	0.762 to 0.779	<.0001	0.594
Dietary fiber g/day	9.22	3.15	2.72	22.68	0.310	0.290 to 0.330	<.0001	0.096
Sugar available kcal/day	447.1	283.8	48.3	1249	0.423	0.405 to 0.441	<.0001	0.179
Fasting plasma glucose mmol/dl	4.31	0.35	3.32	5.58	0.565	0.550 to 0.580	<.0001	0.32
Systolic BP mm/Hg	133.9	4.33	123.4	147.9	0.087	0.065 to 0.109	<.0001	0.008
LDL cholesterol mmol/dl	2.36	0.4	1.27	3.25	0.757	0.747 to 0.766	<.0001	0.573
Socio-demographic index	0.54	0.17	0.11	0.9	0.738	0.728 to 0.748	<.0001	0.544

376
377

378 **Table 2. Correlations between the risk factors that are in Table 1**

Variables (foods in kcal/day)	Process ed meat	Red meat	Fish	Milk	Poultry	Eggs	Alcohol
Processed meat	1.000						
Red meat	0.656	1.000					
Fish	0.281	0.118	1.000				
Milk	0.763	0.682	0.109	1.000			
Poultry	0.675	0.670	0.162	0.684	1.000		
Eggs	0.574	0.786	0.392	0.532	0.633	1.000	
Alcohol	0.592	0.478	0.292	0.402	0.372	0.352	1.000
Sugary beverages	-0.118	0.037	-0.163	0.059	0.126	0.014	-0.098
Corn	-0.052	-0.144	-0.018	-0.108	0.008	-0.100	-0.108
Potatoes	0.160	0.124	-0.058	0.202	0.104	0.032	-0.007
Saturated fatty acids	0.633	0.709	0.039	0.772	0.709	0.601	0.304
Poly unsaturated fatty acids	0.711	0.699	0.162	0.691	0.844	0.635	0.400
Trans fatty acids	0.479	0.203	-0.006	0.445	0.465	0.217	0.174
Fruits	0.391	0.380	0.099	0.499	0.613	0.374	0.075
Vegetables	0.308	0.392	0.271	0.362	0.360	0.519	0.209
Nuts and seeds	0.520	0.399	0.073	0.590	0.519	0.344	0.253
Whole grains	-0.022	-0.111	0.091	-0.250	-0.023	-0.076	0.065
Legumes	-0.199	-0.404	0.061	-0.197	-0.222	-0.418	0.133
Rice	-0.396	-0.357	-0.071	-0.568	-0.396	-0.277	-0.139
Sweet potatoes	-0.197	0.000	-0.075	-0.395	-0.242	0.045	-0.068
Physical activity METs	-0.215	-0.072	-0.163	-0.431	-0.397	-0.119	0.097
Kcal /day available	0.602	0.649	0.098	0.775	0.754	0.644	0.231
Child/infant 2SD underweight	-0.416	-0.737	-0.189	-0.409	-0.618	-0.804	-0.141
Discontinued breast feeding <6 mo	0.583	0.812	0.219	0.639	0.727	0.847	0.307

379
380
381
382

Table 2. Correlations between the risk factors that are in Table 1 (continued)

Variables (foods in kcal/day)	Sugary beverages	Corn	Potatoes	Fruits	Vegetables	Nuts and seeds	Whole grains
Corn	0.419	1.000					
Potatoes	-0.026	0.158	1.000				
Fruits	0.189	0.100	0.243	1.000			
Vegetables	-0.173	-0.203	-0.044	0.394	1.000		
Nuts and seeds	-0.154	-0.114	0.188	0.586	0.556	1.000	
Whole grains	0.204	0.428	-0.058	0.012	-0.346	-0.208	1.000
Legumes	0.199	0.269	0.018	-0.032	-0.160	-0.028	0.100
Rice	-0.142	-0.248	-0.378	-0.325	-0.237	-0.359	0.547
Sweet potatoes	-0.215	-0.027	0.238	-0.100	-0.099	-0.079	-0.039
Physical activity METs	-0.099	0.028	0.021	-0.393	-0.212	-0.306	0.268
Kcal /day available	0.114	-0.044	0.182	0.640	0.615	0.601	-0.187
Child/infant 2SD underweight	-0.060	-0.081	-0.216	-0.497	-0.457	-0.300	0.089
Discontinued breast feeding <6 mo	0.133	-0.064	0.132	0.564	0.480	0.373	-0.125

383
384

385 **Table 2. Correlations between the risk factors that are in Table 1 (continued)**

Variables (foods in kcal/day)	Legumes	Rice	Sweet potatoes	Physical activity	Kcal	Child/infant under-weight
Rice	-0.047	1.000				
Sweet potatoes	-0.053	0.072	1.000			
Physical activity METs	-0.066	0.388	0.328	1.000		
Kcal /day available	-0.250	-0.429	-0.257	-0.500	1.000	
Child/infant 2SD underweight	0.507	0.377	-0.164	0.138	-0.621	1.000
Discontinued breast feeding <6 mo	-0.406	-0.419	-0.062	-0.227	0.727	-0.836

386
387

388 **Table 3. Bradford Hill criteria consistency and dose response: Mean BMI and BMI**
 389 **formula output differences in 37 subgroups**
 390

Subsets of worldwide data on risk factors related to BMI	n	Mean BMI	BMI formula output with mean and SD equated with mean BMI	BMI and BMI formula [‡] output difference	BMI and BMI formula [¶] output difference	BMI and BMI formula [†] output difference
Socio-demographic index quartile 1	1966	24.54	24.75	0.210	0.243	0.289
SDI quartile 2	2096	21.82	21.69	0.130	0.210	0.402
SDI quartile 3	2220	20.40	20.30	0.099	0.104	0.107
SDI quartile 4	1604	20.31	20.31	0.007	0.079	0.012
Africa	1682	21.67	21.25	0.411	0.009	0.519
Asia	4188	20.47	20.46	0.013	0.065	0.058
Europe	880	24.31	24.66	0.343	0.388	0.234
North America	592	25.97	26.00	0.032	0.079	0.312
Oceania	54	24.43	24.66	0.229	0.277	0.206
South America	474	23.48	24.24	0.761	0.667	0.463
Four countries with subnational data	730	25.18	25.45	0.267	0.300	0.964
Quartile 1 BMI increasing foods	2007	24.14	24.70	0.565	0.572	0.654
Q2	1834	21.96	21.86	0.096	0.103	0.274
Q3	2050	20.60	20.31	0.293	0.281	0.205
Q4	1995	20.49	20.28	0.212	0.226	0.224
Quartile 1 BMI decreasing foods	1475	20.61	20.16	0.454	0.570	0.197
Q2 BMI decreasing foods	2275	21.12	21.03	0.092	0.114	0.114
Q3 BMI decreasing foods	2211	22.48	22.74	0.258	0.294	0.239
Q4 BMI decreasing foods	1925	22.69	22.82	0.125	0.199	0.112
Quartile 1 physical activity	2371	20.48	20.56	0.081	0.089	0.019
Q2 physical activity	1573	21.66	21.13	0.533	0.598	0.554
Q3 physical activity	2018	21.37	21.55	0.175	0.198	0.308
Q4 physical activity	1924	23.95	24.07	0.118	0.136	0.125
Male	3943	21.51	21.91	0.406	0.401	0.368
Female	3943	22.07	21.65	0.423	0.419	0.382
Random controlled trial #1	100	21.92	21.96	0.038	0.054	0.061
RCT #2	100	22.00	21.76	0.236	0.226	0.198
RCT #3	100	21.67	21.78	0.109	0.111	0.149
RCT #4	100	21.89	21.81	0.074	0.069	0.064
UK	66	24.99	25.13	0.132	0.132	0.393
USA	376	26.66	26.74	0.082	0.132	0.380
Japan	158	21.89	22.87	0.978	0.964	2.888
Mexico	130	24.99	24.99	0.006	0.062	0.606
Quartile 1 Dose-response: BMI formula versus BMI	1862	24.82	25.33	0.509	0.538	0.663
Q2 Dose-response	1802	22.44	22.06	0.379	0.350	0.539
Q3 Dose-response	1773	21.10	20.84	0.266	0.034	0.201
Q4 Dose-response	2449	19.51	19.56	0.057	0.158	0.128
Mean of BMI - BMI formula absolute values				0.249	0.255	0.368

Mean of BMI-BMI formula dose response absolute values				0.302	0.284	0.493
--	--	--	--	-------	-------	-------

391
392 ‡In combination variables grouped by BMI increasing foods and BMI decreasing foods, each food was multiplied
393 times (1) its kcal/day (2) its coefficient of determination (R^2) with BMI, and (3) its ratio of kcal/day in SDI quartiles
394 2-4 /SDI quartile 1. Fruits, vegetables, and nuts/seeds grouped with plants that negatively correlate with BMI.
395 ¶ As with ‡ but with fruits, vegetables, and nuts/seeds grouped with plants that positively correlate with BMI
396 † As with § but with food risk factor unadjusted but with food risk factors unadjusted
397
398

399 **Table 4. Cross validation: summary percent weight data for 20 RCT derived BMI**
 400 **formulas↓**
 401

Risk factors	Mean	Std Dev	Minimum	Maximum
Processed meat kcal/day	0.26	0.09	0.11	0.43
Red meat kcal/day	9.82	3.05	5.41	14.38
Fish kcal/day	0.18	0.51	0.01	1.97
Milk kcal/day	3.13	1.22	1.38	4.88
Poultry kcal/day available	9.10	2.88	5.35	15.53
Eggs kcal/day available	4.66	1.09	2.50	7.73
Alcohol kcal/day	1.78	2.08	0.01	7.92
Sugary beverages kcal/day	7.24	1.51	0.01	14.62
Corn kcal/day available	0.80	0.70	0.00	2.00
potatoes kcal/day available	2.79	1.94	0.01	7.81
Saturated fatty acids kcal/day	3.09	2.09	0.90	8.18
Poly unsaturated fatty acids kcal/day	0.71	0.38	0.18	1.50
Trans fatty acids kcal/day	0.09	0.05	0.03	0.18
Fruits kcal/day	0.46	0.21	0.12	0.88
Vegetables kcal/day	0.63	0.30	0.23	1.31
Nuts and seeds kcal/day	0.03	0.02	0.01	0.07
Whole grains kcal/day	0.13	0.02	0.00	0.39
Legumes kcal/day	0.56	3.51	0.09	1.54
Rice kcal/day available	8.89	2.65	3.40	13.19
Sweet potatoes kcal/day available	0.32	0.32	0.02	1.13
Physical activity METs	9.76	4.09	4.24	13.29
Kcal available/day	7.89	3.23	3.68	13.56
Child/infant 2SD underweight	5.41	4.51	2.49	8.29
Discontinued breast feeding <6 mo	1.19	0.84	0.20	2.96
Total percent weights	78.90	2.88	74.65	83.04

402
 403 ↓ See Supplementary Table 7 for the data on the 20 RCTs each with n=100

404

405

406 **Table 5. BMI formula estimates with common dietary and other risk factor scenarios^β**
 407

Testing common dietary and other risk factor scenarios with standardised BMI formula estimates	BMI kg/M ²	BMI formula estimate
World (n=7886 cohorts)	21.79	21.71
United Kingdom (n=66 cohorts)	24.99	25.09
USA (n=376 cohorts)	26.66	26.67
Following USA Dietary Guidelines 2015-2020 recommendations		26.34
USA with 25% reduction of BMI increasing food intake [†]		25.04
USA with 50% reduction of BMI increasing food intake [†]		23.34
USA mean physical activity plus 2 hours/day running at 6 mph		23.98
USA mean physical activity plus 1 hour/day running at 6 mph and 25% reduction of BMI increasing food intake [†]		23.67
USA with no red or processed meat [†]		24.81
USA with no sugary beverage intake [†]		26.31
USA vegetarian (no meat, poultry, fish) [†]		22.54
USA vegan (no meat, poultry, fish, dairy or eggs) [†]		21.11
USA breast feeding continued at least 6 months in all		26.68
USA breast feeding discontinued before 6 months in all		26.87
USA no occurrence of child severe underweight		26.73
USA all children with severe underweight (theoretically)		25.54
EAT-Lancet diet		22.47
Low Carbohydrate Mediterranean Diet		31.76

408
 409 ^β BMI formula estimates based on 28 years of following dietary and/or risk factor patterns
 410 [†] Kcal/day isocalorically shifted to the 7 BMI decreasing foods in the BMI formula, distributed
 411 equally
 412

413 **Acknowledgments**

414• **Authors' contributions:** DKC acts as guarantor; conceived and designed the study, acquired
415 and analysed the data, interpreted the study findings, drafted the manuscript, critically reviewed
416 and edited the manuscript and tables, and approved the final version for publication.

417
418 CW designed software programs in R to format and population weight the data,
419 aided with the SAS statistical analysis, critically reviewed the manuscript, and
420 approved the final version for publication.

421
422 The corresponding author attests that all listed authors meet authorship criteria and that no others
423 meeting the criteria have been omitted.

424
425 **Copyright:** The retain the copyright to the paper

426
427 The Patient and Public Involvement

428 • When and how were patients/public first involved in the research?

429 IHME acquired, catalogued and extracted information from over 12,000 surveys from
430 government and non government agencies in order to compile the GBD database.

431 • How were the research question(s) developed and informed by their priorities,
432 experience, and preferences?

433 The diverse surveys contributing to the GBD database had many different priorities and
434 intentions unrelated to this post hoc analysis. The current analysis research questions were

435 developed to give researchers, policymakers, and the public the methodological tool to quantify
436 the BMI impacts of dietary and other risk factor patterns.

437 ● How were patients/public involved in

438 (a) the design and conduct of the study?

439 The way patients/public were involved in the collection of the surveys has not been
440 systematically studied and reported on. Patients/public are the intended beneficiaries of
441 this analysis of the GBD data.

442 (b) choice of outcome measures?

443 The outcome measures chosen all related to human health and to adult BMI.

444 (c) recruitment to the study?

445 The recruitment methods varied by the survey.

446 ● How were (or will) patients/ public be involved in choosing the methods and agreeing
447 plans for dissemination of the study results to participants and linked communities?

448 IHME and all the health surveyors that contributed will make these decisions. This analysis of
449 the data will be available to all by open access.

450

451● **Competing interests statement:** Both authors have completed the ICMJE uniform disclosure
452 form at www.icmje.org/coi_disclosure.pdf and declare: no support from any organisation for the
453 submitted work; no financial relationships with any organisations that might have an interest in
454 the submitted work in the previous three years; no other relationships or activities that could
455 appear to have influenced the submitted work.

456

457 **Contributors:** Martin Sebera, from the Department of Kinesiology, Faculty of Sports Studies

458 Masaryk University, Czech Republic, critiqued statistical aspects of the manuscript and provided
459 useful input. Pavel Grasgruber, from Masaryk University, Czech Republic, provided suggestions
460 after reviewing the manuscript.

461
462 **Transparency declaration:** The manuscript is an honest, accurate, and transparent account of
463 the study being reported; that no important aspects of the study have been omitted; and that any
464 discrepancies from the study as planned have been explained.

465
466 **Ethics:** Studies based solely on data from IHME GBD database do not need approval from any
467 bioethics committee.

468
469• **Funding:** This research received no specific grant from any funding agency in the public,
470 commercial or not-for-profit sectors. The Bill and Melinda Gates Foundation funded the
471 acquisition of the data for this analysis by the IHME. The data were provided to the authors as
472 volunteer collaborators with IHME.

473•
474
475 **Details of the role of the study sponsors:** While IHME GBD faculty and staff by virtue of
476 Gates Foundation grants provided the raw data for this analysis, they did not vet the analysis or
477 sponsor the manuscript.

478
479 **Statement of independence of researchers from funders:** The researchers have received no
480 funding. Gates Foundation funded IHME to collect and analyse the GBD data.

481 **Dissemination declaration:** Dissemination of this manuscript to the participants of the more
482 than 12,000 surveys is not possible individually, but the information will become in the public
483 domain.

484
485 **Disclosures:** We thank Scott Glenn and Brent Bell from IHME who supplied us with the GBD
486 risk factor exposure data for the risk factors and for BMI data.

487•
488 **Ethics:** Studies were based solely on data from the IHME GBD database and do not need
489 approval from any bioethics committee.

490
491 **Participant informed consents:** Not applicable.

492
493 **Author access to data:** As volunteer collaborators with the Institute of Health Metrics and
494 Evaluation, we received about 1.4 gigabytes of raw data on BMI and 32 relevant risk factors for
495 BMI and other health outcomes.

496
497 **Data sharing statement:** The raw, unformatted data used in this analysis is now out of date. The
498 2019 GBD data on all the variables in this analysis may be obtained from the IHME by volunteer
499 collaborating researchers. Our data formatting software code in R and SAS and our formatted
500 database are available on request to researchers.

501
502 **Protocol, submitted as a supplementary file:** Not applicable.

503

504 **STROBE checklist:** Submitted.

505

506 **Patient consent:** Not applicable.

507

508 **Clinical trial registration:** Not applicable.

509 **References**

- 510 1. Teicholz N. **The scientific report guiding the US dietary guidelines: is it scientific?**
511 *BMJ*. 2015; **351**. <http://www.bmj.com/bmj/351/bmj.h4962.full.pdf>
- 512 2. **Outcome of Post-publication review of article by Nina Teicholz.** *BMJ*. 2015; **351**:
513 h4962. <https://www.bmj.com/content/351/bmj.h4962/rr-48>
- 514 3. Gardner CD, Trepanowski JF, Del Gobbo LC, et al. **Effect of Low-Fat vs Low-**
515 **Carbohydrate Diet on 12-Month Weight Loss in Overweight Adults and the**
516 **Association With Genotype Pattern or Insulin Secretion: The DIETFITS**
517 **Randomized Clinical Trial.** *JAMA*. 2018; **319** (7): 667-679.
518 <https://jamanetwork.com/journals/jama/fullarticle/2673150>
- 519 4. Ioannidis JA. **The challenge of reforming nutritional epidemiologic research** *JAMA*
520 2018; **320** (10): 969-70. <http://dx.doi.org/10.1001/jama.2018.11025>
- 521 5. Hill AB. **THE ENVIRONMENT AND DISEASE: ASSOCIATION OR**
522 **CAUSATION?** *Proceedings of the Royal Society of Medicine*. 1965; **58** (5): 295-300.
523 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1898525/>
- 524 6. Global Burden of Disease Study 2017 (GBD 2017) Data Input Sources Tool. Seattle,
525 Washington: Institute of Health Metrics and Evaluation Accessed August 10, 2019.
526 <http://ghdx.healthdata.org/gbd-2017/data-input-sources>
- 527 7. **GBD 2017 Risk Factor Collaborators. Global, regional, and national comparative**
528 **risk assessment of 84 behavioural, environmental and occupational, and metabolic**
529 **risks or clusters of risks for 195 countries and territories, 1990-2017: a systematic**
530 **analysis for the Global Burden of Disease Study 2017: Table of risk factor**
531 **definitions.** *The Lancet*. 2018; **392** (10159): 1923-94.

- 532 [https://www.thelancet.com/action/showFullTableHTML?isHtml=true&tableId=tbl1&pii=](https://www.thelancet.com/action/showFullTableHTML?isHtml=true&tableId=tbl1&pii=S0140-6736%2818%2932225-6)
533 [S0140-6736%2818%2932225-6](https://www.thelancet.com/action/showFullTableHTML?isHtml=true&tableId=tbl1&pii=S0140-6736%2818%2932225-6)
- 534 8. PROTOCOL FOR THE GLOBAL BURDEN OF DISEASES, INJURIES, AND RISK
535 FACTORS STUDY (GBD) Version 3.0; Issue 26. Seattle, Washington: Institute for
536 Health Metrics and Evaluation Accessed August 2, 2019.
537 http://www.healthdata.org/sites/default/files/files/Projects/GBD/GBD_Protocol.pdf
- 538 9. Naghavi M, Abajobir AA, Abbafati C, et al. **Global, regional, and national age-sex**
539 **specific mortality for 264 causes of death, 1980-2016: a systematic analysis for the**
540 **Global Burden of Disease Study 2016.** *The Lancet.* 2017; **390** (10100): 1151-1210.
541 [https://doi.org/10.1016/S0140-6736\(17\)32152-9](https://doi.org/10.1016/S0140-6736(17)32152-9)
- 542 10. Flaxman AD, Lee YY, Vos T, et al. An Integrative Metaregression Framework for
543 Descriptive Epidemiology. Seattle, WA: University of Washington Press, 2015.
- 544 11. GBD 2017 Risk Factor Collaborators. **Global, regional, and national comparative risk**
545 **assessment of 84 behavioural, environmental and occupational, and metabolic risks**
546 **or clusters of risks for 195 countries and territories, 1990-2017: a systematic**
547 **analysis for the Global Burden of Disease Study 2017.** . *The Lancet.* 2018; **392**
548 (10159): 1923-94. [https://doi.org/10.1016/S0140-6736\(18\)32225-6](https://doi.org/10.1016/S0140-6736(18)32225-6)
- 549 12. Call for Collaborators. Seattle, Washington: Institute for Health Metrics and Evaluation
550 at the University of Washington 2019; Accessed April 25, 2019.
551 <http://www.healthdata.org/gbd/call-for-collaborators>
- 552 13. Stevens GA, Alkema L, Black PRE, et al. **Guidelines for Accurate and Transparent**
553 **Health Estimates Reporting: the GATHER statement.** *The Lancet.* 2016; **388**
554 (10062): e19-e23. [https://doi.org/10.1016/S0140-6736\(16\)30388-9](https://doi.org/10.1016/S0140-6736(16)30388-9)

- 555 14. Omega 3 Fatty Acids: Fact Sheet for Health Professionals. The Office of Dietary
556 Supplements 2018; Accessed September 1, 2018.
557 <https://ods.od.nih.gov/factsheets/Omega3FattyAcids-HealthProfessional/>
- 558 15. Nutritionix Track App. Syndigo LLC Accessed April 25, 2019.
559 <https://www.nutritionix.com/>
- 560 16. POTATO PROCESSING AND USES. Lima, Peru: International Potato Center
561 Accessed May 28, 2020. <https://cipotato.org/potato/potato-processing-uses/>
- 562 17. Ruotsalainen H, Kyngäs H, Tammelin T, Kääriäinen M. **Systematic review of physical**
563 **activity and exercise interventions on body mass indices, subsequent physical**
564 **activity and psychological symptoms in overweight and obese adolescents.** *Journal of*
565 *Advanced Nursing.* 2015; **71** (11): 2461-2477.
566 <https://onlinelibrary.wiley.com/doi/abs/10.1111/jan.12696>
- 567 18. Te Morenga L, Mallard S, Mann J. **Dietary sugars and body weight: systematic review**
568 **and meta-analyses of randomised controlled trials and cohort studies.** *BMJ* 2013;
569 **346:** e7492. <https://www.bmj.com/content/bmj/346/bmj.e7492.full.pdf>
- 570 19. Traversy G, Chaput J-P. **Alcohol Consumption and Obesity: An Update.** *Current*
571 *obesity reports.* 2015; **4** (1): 122-130.
572 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4338356/>
- 573 20. Turner-McGrievy G, Mandes T, Crimarco A. **A plant-based diet for overweight and**
574 **obesity prevention and treatment.** *Journal of geriatric cardiology : JGC.* 2017; **14** (5):
575 369-374. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5466943/>
- 576 21. AINSWORTH BE, HASKELL WL, HERRMANN SD, et al. **Compendium of Physical**
577 **Activities: A Second Update of Codes and MET Values.** *Medicine & Science in Sports*

- 578 & *Exercise*. 2011 **43** (8): 1575-1581. [https://journals.lww.com/acsm-](https://journals.lww.com/acsm-msse/Fulltext/2011/08000/2011_Compndium_of_Physical_Activities__A_Second.25.aspx)
579 [msse/Fulltext/2011/08000/2011_Compndium_of_Physical_Activities__A_Second.25.as](https://journals.lww.com/acsm-msse/Fulltext/2011/08000/2011_Compndium_of_Physical_Activities__A_Second.25.aspx)
580 px
- 581 22. Ebbeling CB, Feldman HA, Klein GL, et al. **Effects of a low carbohydrate diet on**
582 **energy expenditure during weight loss maintenance: randomized trial.** *BMJ*. 2018;
583 **363**: k4583.
- 584 23. Darmon N, Drewnowski A. **Contribution of food prices and diet cost to**
585 **socioeconomic disparities in diet quality and health: a systematic review and**
586 **analysis.** *Nutrition reviews*. 2015; **73** (10): 643-660.
587 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4586446/>
- 588 24. Wakamo B. We Subsidize the Wrong Kind of Agriculture. Inequality.org Accessed.
589 <https://inequality.org/great-divide/subsidize-wrong-kind-agriculture/>
- 590 25. Tagtow A. Fruits and Veggies Now Chock Full of Marketing Power. Washington, DC
591 2017; Accessed June 24, 2020. [https://www.usda.gov/media/blog/2015/05/07/fruits-and-](https://www.usda.gov/media/blog/2015/05/07/fruits-and-veggies-now-chock-full-marketing-power)
592 [veggies-now-chock-full-marketing-power](https://www.usda.gov/media/blog/2015/05/07/fruits-and-veggies-now-chock-full-marketing-power)
- 593 26. Stolzer JM. **Breastfeeding and obesity: a meta-analysis.** *Open Journal of Preventive*
594 *Medicine*. 2011; **1** (3): 88-93.
595 <http://www.scirp.org/journal/paperinformation.aspx?paperid=8509>
- 596 27. AAP Reaffirms Breastfeeding Guidelines. The American Academy of Pediatrics 2012;
597 Accessed August 10, 2019. [https://www.aap.org/en-us/about-the-aap/aap-press-](https://www.aap.org/en-us/about-the-aap/aap-press-room/Pages/AAP-Reaffirms-Breastfeeding-Guidelines.aspx)
598 [room/Pages/AAP-Reaffirms-Breastfeeding-Guidelines.aspx](https://www.aap.org/en-us/about-the-aap/aap-press-room/Pages/AAP-Reaffirms-Breastfeeding-Guidelines.aspx)
- 599 28. **Breastfeeding and the Use of Human Milk.** *Pediatrics*. 2012; **129** (3): e827-e841.
600 doi:10.1542/peds.2011-3552

- 601 29. Quality statement 5: Breastfeeding. National Institute of Clinical Excellence 2013;
602 Accessed August 10, 2019. [https://www.nice.org.uk/guidance/qs37/chapter/Quality-](https://www.nice.org.uk/guidance/qs37/chapter/Quality-statement-5-Breastfeeding)
603 [statement-5-Breastfeeding](https://www.nice.org.uk/guidance/qs37/chapter/Quality-statement-5-Breastfeeding)
- 604 30. FORESIGHT--Tackling Obesities: Future Choices -Follow Up Actions. London,
605 England: Government Office for Science (UK) October 2007; Accessed August 5, 2019.
606 [https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/295885/07-1526-obesity-action-plan.pdf)
607 [data/file/295885/07-1526-obesity-action-plan.pdf](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/295885/07-1526-obesity-action-plan.pdf)
- 608 31. Mărginean CO, Mărginean C, Meliț LE. **New Insights Regarding Genetic Aspects of**
609 **Childhood Obesity: A Minireview.** *Frontiers in pediatrics.* 2018; **6**: 271-271.
610 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6180186/>
- 611 32. Castaner O, Goday A, Park Y-M, et al. **The Gut Microbiome Profile in Obesity: A**
612 **Systematic Review.** *International journal of endocrinology.* 2018; **2018**: 4095789-
613 4095789. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5933040/>
- 614 33. Monteiro CA, Moubarac J-C, Levy RB, Canella DS, Louzada MLdC, Cannon G.
615 **Household availability of ultra-processed foods and obesity in nineteen European**
616 **countries.** *Public Health Nutrition.* 2017; **21** (1): 18-26.
617 [https://www.cambridge.org/core/article/household-availability-of-ultraprocessed-foods-](https://www.cambridge.org/core/article/household-availability-of-ultraprocessed-foods-and-obesity-in-nineteen-european-countries/D63EF7095E8EFE72BD825AFC2F331149)
618 [and-obesity-in-nineteen-european-countries/D63EF7095E8EFE72BD825AFC2F331149](https://www.cambridge.org/core/article/household-availability-of-ultraprocessed-foods-and-obesity-in-nineteen-european-countries/D63EF7095E8EFE72BD825AFC2F331149)
- 619 34. Monteiro CA, Cannon G, Levy RB, et al. **Ultra-processed foods: what they are and**
620 **how to identify them.** *Public Health Nutrition.* 2019; **22** (5): 936-941.
621 [https://www.cambridge.org/core/article/ultraprocessed-foods-what-they-are-and-how-to-](https://www.cambridge.org/core/article/ultraprocessed-foods-what-they-are-and-how-to-identify-them/E6D744D714B1FF09D5BCA3E74D53A185)
622 [identify-them/E6D744D714B1FF09D5BCA3E74D53A185](https://www.cambridge.org/core/article/ultraprocessed-foods-what-they-are-and-how-to-identify-them/E6D744D714B1FF09D5BCA3E74D53A185)
- 623

624 Supplementary Table 1. Definitions of IHME GBD risk factors and covariates related to BMI

Variables	Definition
Alcohol	Any alcohol consumption (g/day)
Body-mass index	Body mass index (BMI) (kg/m ²)—the dependent variable of interest
Child underweight	Proportion of children - 3 SD to - 2 SD of the WHO 2006 standard weight-for-age curve (0-1)
Corn	Corn availability percapita (g/day), a covariate
Discontinued breast feeding	Proportion of children aged 6-23 months who do not receive any breast milk
Eggs	Eggs availability percapita (g/day) a covariate
Fasting plasma glucose	Fasting plasma glucose (mmol/L)
Fish	This variable expressed in g/day was derived by determining the weight of fish in g corresponding to 1 g of omega-3 fatty acids (eicosapentaenoic acid and docosahexaenoic acid) by averaging the fish g per 1 g of omega-3 fatty acids 20 species of fish= 117.04 g/day fish/1 g/day omega-3 fatty acids (Supplementary Table 2)
Fruits	Consumption of fruits (includes fresh, frozen, cooked, canned, or dried fruit but excludes fruit juices and salted or pickled fruits) (g/day)
Kilocalories available /day	The mean number of kilocalories percapita available per day to people in each location (kcal/day available), a covariate
LDL cholesterol	Serum low-density lipoprotein cholesterol (mmol/L)
Legumes	Consumption of beans, lentils, pulses (g/day)
Milk	Consumption of milk including non-fat, low-fat, and full-fat milk but excluding soy milk and other plant derivatives (g/day)
Nuts and seeds	Consumption of nuts and seeds (g/day)
Physical activity	Average weekly physical activity at work, home, transport-related and recreational measured by MET min per week. Less than 3000 METs per week constitutes low physical activity.
Poultry	Poultry availability percapita (g/day), a covariate
Potatoes	Potatoes availability percapita (g/day), a covariate
Processed meat	Consumption of any processed meat (includes meat preserved by smoking, curing, salting, or addition of chemical preservatives, including bacon, salami, sausages, or deli or luncheon meats like ham, turkey, and pastrami) (g/day)
Red meat	Consumption of red meat (includes beef, pork, lamb, and goat but excludes poultry, fish, eggs, and all processed meats) (g/day)
Rice	Rice availability percapita (g/day), a covariate
Seafood omega-3 fatty acids	Seafood omega-3 fatty acids (eicosapentaenoic acid and docosahexaenoic acid) in tablet or fish form (g/day)
Socio-demographic index	SDI is a composite indicator of development status that was originally constructed for GBD 2015, and is derived from components that correlate strongly with health outcomes. It is the geometric mean for indices of the total fertility rate among women younger than 25 years, mean education for those aged 15 years or older, and lag-distributed income per capita. The resulting metric ranges from 0 to 1, with higher values corresponding to higher levels of development.
Sugar-sweetened beverages	Consumption of any beverage with ≥50 calories of sugar per one-cup serving, including carbonated beverages, sodas, energy drinks, fruit drinks but excluding 100% fruit and vegetable juices (g/day)
Sweet potatoes	Sweet potato availability percapita (g/day), a covariate
Systolic blood pressure	Systolic blood pressure (mm Hg)
Total sugar	Total sugar availability percapita (g/day), a covariate

Vegetables	Consumption of frozen, cooked, canned, or dried vegetables (including legumes but excluding salted or pickled, juices, nuts and seeds, and starchy vegetables such as potatoes or corn) (g/day)
Whole grains	Consumption of whole grains (bran, germ, and endosperm in their natural proportions) from breakfast cereals, bread, rice, pasta, biscuits, muffins, tortillas, pancakes, and others (g/day)

625

626

627 **Supplementary Table 2. Omega-3 Fatty Acid g to fish g calculation¶¶**

Fish	DHA g/3 ounce fish	EPA g/3 ounce fish	Omega-3 Fatty Acids (DHA + EPA) g/3 ounce fish mean	Fish 3 ounces = 85.02 g	Fish (g) per omega-3 Fatty Acids (g)=columns E/F
Salmon Atlantic farmed	1.24	0.59			
Salmon Atlantic wild	1.22	0.35			
Herring Atlantic	0.94	0.77			
Sardines canned in tomato sauce drained	0.74	0.45			
Mackerel Atlantic	0.59	0.43			
Salmon pink canned drained	0.63	0.28			
Trout rainbow wild	0.44	0.40			
Oysters eastern wild	0.23	0.30			
Sea bass	0.47	0.18			
Shrimp	0.12	0.12			
Lobster	0.07	0.10			
Tuna light canned in water drained	0.17	0.02			
Tilapia	0.11				
Scallops	0.09	0.06			
Cod Pacific	0.1	0.04			
Tuna yellowfin	0.09	0.01			
Mean DHA and EPA Omega-3 Fatty Acids g/3 ounce fish	0.4531	0.2733			
Calculations total Omega-3 FA g to fish g			0.7264	85.02	117.043

628 ¶¶ Data on omega-3 fatty acid content of varieties of fish came from the National Institutes of
 629 Health Office of Dietary Supplements (USA)
 630

631 **Supplementary Table 3. Calculations of kcal/day from g/day of animal and plant foods¶**

Foods	Food sub-categories	kcal/serving	g/serving	kcal/g
Milk (2% fat)		122	244	0.5
Fish		218	170	1.28
Eggs		72	50	1.44
Poultry		187	85	2.91
Red meat		247	85	2.91
Processed meat				
	Salami	222	59	3.76
	Pastrami	104	71	1.46
	Ring baloney	86	28	3.07
	Pepperoni	94	100	0.94
Average processed meat		126.5	64.5	1.96
Fruits		97	162	0.60
Vegetables		59	91	0.65
Legumes		249	179	1.39
Nuts		172	28	6.14
Seeds				
	Flax seeds	55	10	5.5
	Chia seeds	58	12	4.83
	Fennel seeds	34.5	10	3.45
	Hemp seeds	55.3	10	5.53
Average of seeds		50.7	10.5	4.83
Average of nuts and seeds		111.4	19.25	5.78
Corn		99	103	0.96
Potatoes		161	173	0.93
Sweet potatoes		115	151	0.76
Rice		205	158	1.3
Whole grains		120	52	2.31

632 ¶ Source: NutritionIX app

633

634 **Supplementary Table 4. Multicollinearity of fruits, vegetables, and nuts and seeds**

High BMI risk factors (n=7886 cohorts)	Mean Kcal/day	BMI kg/M ² r	95% CI	P	BMI increasing r †	95% CI	P
BMI kg/m ²	21.79				0.724	0.713 to 0.734	<0.0001
BMI increasing r	654	0.597	0.583 to 0.611	<0.0001			
Fruits + vegetables + nuts and seeds	129	0.655	0.642 to 0.677	<0.0001	0.323	0.304 to 0.343	<0.0001
Fruits	40.4	0.622	0.609 to 0.636	<0.0001	0.498	0.482 to 0.515	<0.0001
Vegetables	80.1	0.520	0.503 to 0.536	<0.0001	0.150	0.128 to 0.171	<0.0001
Nuts and seeds	8.51	0.486	0.469 to 0.503	<0.0001	0.279	0.258 to 0.299	<0.0001
Top socio-demographic index (SDI) quartile n=1996	Kcal/day	BMI kg/M ² r	95% CI	P	BMI increasing r †	95% CI	P
BMI kg/m ²	24.54				0.364	0.325 to 0.402	<0.0001
BMI increasing	905	0.364	0.325 to 0.402	<0.0001			
Fruits + vegetables + nuts and seeds	199	0.104	0.060 to 0.148	<0.0001	-0.306	0.346 to 0.266	<0.0001
Fruits	61.1	0.296	0.255 to 0.336	<0.0001	-0.022	-0.007 to -0.095	0.32
Vegetables	120	-0.064	-0.108 to -0.020	0.0043	-0.388	-0.425 to -0.350	<0.0001
Nuts and seeds	17.5	0.314	0.273 to 0.353	0.0018	-0.118	-0.161 to -0.074	<0.0001
Bottom three SDI quartiles (n=5920 cohorts)	Kcal/day	BMI kg/M ² r	95% CI	P	BMI increasing r †	95% CI	P
BMI kg/m ²	20.88				0.364	0.342 to 0.386	<0.0001
BMI increasing	571	0.364	0.342 to 0.386	<0.0001			
Fruits + vegetables + nuts and seeds	106	0.536	0.517 to 0.554	<0.0001	0.108	0.083 to 0.133	<0.0001
Fruits	33.5	0.466	0.446 to 0.486	<0.0001	0.429	0.408 to 0.450	<0.0001
Vegetables	66.8	0.395	0.374 to 0.417	0.0043	-0.103	-0.128 to -0.077	0.0186
Nuts and seeds	5.52	-0.040	-0.066 to -0.015	0.0018	-0.072	-0.098 to -0.047	<0.0001

635
 636 † BMI increasing foods include all six animal products, alcohol, sugary beverages, corn (high
 637 fructose corn syrup included), and potatoes (highly processed potato products included).
 638 Saturated fatty acids, PUFAs, and TFAs also correlate positively with BMI. However, for the
 639 purposes of this table, they were not included with BMI increasing foods, because animal foods
 640 contain most of the fatty acids.
 641

642 **Supplementary Table 5. Derivation of the BMI formula (variables all standardized)**

A	B	C	D	E	F	G	H	I	J	K
Variables in BMI formula		Mean		R2 with BMI		SDI Q 2-4 / SDI Q 1		Foods and physical activity multiple regression derived parameter estimates		Foods and physical activity synthesis parameter estimates= Columns C*E*G*I absolute values
Processed meat kcal/day	*	5.486	*	0.365	*	0.099	*	0.02972	=	0.00586
Red meat kcal/day	*	50.70	*	0.438	*	0.364	*	0.02972	=	0.24003
Fish kcal/day	*	10.02	*	0.011	*	0.106	*	0.02972	=	0.00033
Milk kcal/day	*	25.37	*	0.468	*	0.214	*	0.02972	=	0.07553
Poultry kcal/day availability	*	45.06	*	0.661	*	0.253	*	0.02972	=	0.22416
Eggs kcal/day availability	*	19.47	*	0.473	*	0.402	*	0.02972	=	0.11011
Alcohol kcal/day	*	81.71	*	0.028	*	0.587	*	0.02972	=	0.03948
Sugary beverages kcal/day	*	297.7	*	0.014	*	1.013	*	0.02972	=	0.12722
Corn kcal/day availability	*	34.67	*	0.005	*	1.081	*	0.02972	=	0.00590
Potatoes kcal/day availability	*	84.16	*	0.044	*	0.644	*	0.02972	=	0.07137
Saturated fatty acids	*	191.3	*	0.504	*	0.653	*	0.0008705	=	0.05481
Polyunsaturated fatty acids	*	82.24	*	0.543	*	0.334	*	0.0008705	=	0.01297
Trans fatty acids	*	13.39	*	0.238	*	0.471	*	0.0008705	=	0.00131
Fruits kcal/day	*	40.39	*	0.387	*	0.549	*	0.0008705	=	0.00747
Vegetables kcal/day	*	80.14	*	0.270	*	0.555	*	0.0008705	=	0.01046
Nuts and seeds kcal/day	*	8.511	*	0.237	*	0.315	*	0.0008705	=	0.00055
Whole grains kcal/day	*	55.65	*	0.040	*	1.248	*	0.0008705	=	0.00243
Legumes kcal/day	*	51.65	*	0.148	*	1.432	*	0.0008705	=	0.00951
Rice kcal/day availability	*	141.2	*	0.314	*	4.320	*	0.0008705	=	0.16697
Sweet potatoes kcal/day availability	*	22.67	*	0.023	*	7.977	*	0.0008705	=	0.00362
Physical activity METs								0.20087	=	0.20087
Total Kcal/day availability										
Child/infant underweight										
Discontinued breast feed <6 months										
Sums										1.37098
Sums										1.37098
BMI formula R2										0.73530
BMI formula R2 / sum of parameter estimates										0.53633

643
644
645
646

647

Supplementary Table 5 Continued

A	L	M	N	O	P	Q	R	S
Variables in BMI formula	Adjust column K μ	Add other risk factors	Adjust column M by multiplying by M28 = 0.1211	Com-bine columns L and N		Check result with statistics software SAS Studio		Final BMI formula percent weights: Column O * 0.9208 (O30) *100
Processed meat kcal/day	0.00315			0.00315	+	Processed meat	*	0.29
Red meat kcal/day	0.12874			0.12874	+	Red meat	*	11.87
Fish kcal/day	0.00018			0.00018	+	Fish	*	0.02
Milk kcal/day	0.04051			0.04051	+	Milk	*	3.74
Poultry kcal/day	0.12022			0.12022	+	Poultry	*	11.09
Eggs kcal/day	0.05906			0.05906	+	Eggs	*	5.45
Alcohol kcal/day	0.02117			0.02117	+	Alcohol	*	1.95
Sugary beverages kcal/day	0.06823			0.06823	+	Sugary beverages	*	6.29
Corn kcal/day	0.00317			0.00317	+	Corn	*	0.29
Potatoes kcal/day	0.03828			0.03828	+	Potatoes	*	3.53
Saturated fatty acids	0.02940			0.02940	+	SFA	*	2.71
Polyunsaturated fatty acids	0.00696			0.00696	+	PUFA	*	0.64
Trans fatty acids	0.00070			0.00070	+	TFA	*	0.06
Fruits kcal/day	0.00401			0.00401	-	Fruits	*	0.37
Vegetables kcal/day	0.00561			0.00561	-	Vegetables	*	0.52
Nuts and seeds kcal/day	0.00030			0.00030	-	Nuts and seeds	*	0.03
Whole grains kcal/day	0.00130			0.00130	-	Whole grains	*	0.12
Legumes kcal/day	0.00510			0.00510	-	Legumes	*	0.47
Rice kcal/day	0.08955			0.08955	-	Rice	*	8.26
Sweet potatoes kcal/day	0.00194			0.00194	-	Sweet Potatoes	*	0.18
Physical activity	0.10773			0.10773	-	Physical activity	*	9.93
Total Kcal/day		0.5408	0.063	0.06270	+	Total kcal	*	5.78
Child/infant underweight		0.3865	0.045	0.04481	-	Child underweight	*	4.13
Discontinued breast feed <6 months		0.0861	0.010	0.00999	+	Discontinued breast feeding	*	0.92
Sums	0.7353	1.0133	0.118	0.85280				78.64
Sums		1.0133		0.85280				
BMI formula R2		0.1175		0.78643				0.79
BMI formula R2 / sum of parameter estimates		0.1160		0.92218				100.00

648 μ Adjust column K by K28 = 0.5314 (foods and physical activity BMI formula R² (K27) / parameter estimate sums (K26)

649

650

651 **Supplementary Table 6. Spreadsheet derivation of BMI trend formula**

A	B	C	D	E	F	G	H	I
BMI trend risk factor trends		Mean		Risk factor with BMI17fts R2		Parameter estimates from multiple regression		Column C * Column E * Column G= preliminary parameter estimates
Processed meat kcal/day trend	*	5.486	*	0.0001	*		=	0.000
Red meat kcal/day trend	*	50.695	*	0.1650	*	0.01645	=	0.138
Fish kcal/day trend	*	10.015	*	0.0251	*		=	0.000
Milk kcal/day trend	*	25.372	*	0.0425	*	0.01645	=	0.018
Poultry kcal/day trend	*	45.059	*	0.1768	*	0.01645	=	0.131
Eggs kcal/day trend	*	19.466	*	0.2322	*	0.01645	=	0.074
Alcohol kcal/day trend	*	81.713	*	0.0826	*	0.01645	=	0.111
Sugary beverages kcal/day trend	*	297.648	*	0.0156	*	0.01645	=	0.077
Corn kcal/day available trend	*	34.669	*	0.0198	*	0.01645	=	0.011
Potatoes available kcal/day available trend	*	84.160	*	0.0739	*	0.01645	=	0.102
Saturated fatty acids trend	*	191.27	*					
Polyunsaturated fatty acids trend	*	82.24	*	0.0820		0.01645		0.111
Trans fatty acids trend	*	13.4	*					
Fruits kcal/day trend	*	40.388	*	0.2301	*			0.000
Vegetables kcal/day trend	*	80.141	*	0.1559	*			0.000
Nuts and seeds kcal/day trend	*	8.510	*	0.0152	*			0.000
Whole grains kcal/day trend	*	55.647	*	0.0107	*		=	0.000
Legumes kcal/day trend	*	51.657	*	0.0887	*	0.00404	=	0.019
Rice kcal/day available trend	*	141.230	*	0.1536	*	0.00404	=	0.088
Sweet potatoes available kcal/day trend	*	22.667	*	0.2487	*	0.00404	=	0.023
Physical activity mets trend								
Kcal available/day trend					*	0.17225	=	0.17225
Child/infant 2SD underweight trend								
Discontinued breast feeding <6 mo trend								
Sums								1.07414
Sum								1.07414
R²								0.37790
Trend R² / trend BMI formula parameter estimates sum								0.35182

652
653
654

655

Spreadsheet derivation of BMI trend formula continued

A	I	J	K	L	M	N
BMI trend risk factor trends	Column C * Column E * Column G= preliminary parameter estimates	Column I adjusted to equate to parameter estimates sum to R2 of the BMI trend formula		Final BMI trend formula to check percent weights with SAS Studio		Column J * 100 to derive percent weights of risk factors
Processed meat kcal/day trend	0.000			pmeat17KCts		
Red meat kcal/day trend	0.138	0.048	+	rmeat17KCts	*	4.841
Fish kcal/day trend	0.000	0.000		fish17KCts		
Milk kcal/day trend	0.018	0.006	+	milk17KCts	*	0.624
Poultry kcal/day trend	0.131	0.046	+	poultry16KCts	*	4.610
Eggs kcal/day trend	0.074	0.026	+	eggs16KCts	*	2.616
Alcohol kcal/day trend	0.111	0.039	+	Alcohol17KCts	*	3.908
Sugary beverages kcal/day trend	0.077	0.027	+	Sugarb17KCts	*	2.692
Corn kcal/day available trend	0.011	0.004	+	corn16KCts	*	0.397
Potatoes available kcal/day available trend	0.102	0.036	+	potatoes16KCts	*	3.600
Saturated fatty acids trend		0.000				
Polyunsaturated fatty acids trend	0.111	0.039	+	PUFA17KCts	*	3.904
Trans fatty acids trend		0.000				
Fruits kcal/day trend	0.000	0.000		fruits17KCts		
Vegetables kcal/day trend	0.000	0.000		Vegetables17KCts		
Nuts and seeds kcal/day trend	0.000	0.000		nutsseeds17KCts		
Whole grains kcal/day trend	0.000	0.000		wgrains17KCts		
Legumes kcal/day trend	0.019	0.007	-	legumes17KCts	*	0.651
Rice kcal/day available trend	0.088	0.031	-	rice16KCts	*	3.084
Sweet potatoes available kcal/day trend	0.023	0.008	-	swtpot16KCts	*	0.801
Physical activity mets trend		0.000		PAMETs17ts		
Kcal available/day trend	0.172	0.061	+	KCal2016ts	*	6.060
Child/infant 2SD underweight trend				Childunwt17ts		
Discontinued breast feeding <6 mo trend				discbreastF17ts		
Sums	1.074	0.378				37.79
Sum	1.074					0.37790
R²	0.378					100.00
Trend R² / trend BMI formula parameter estimates sum	0.352					

656

657

658 **Supplementary Table 7. Cross validation experiment with 20 randomised controlled trials**
 659 **of n=100**

Risk factors	RCT1 Percent Weights	RCT2 Percent Weights	RCT3 Percent Weights	RCT4 Percent Weights	RCT5 Percent Weights
Processed meat kcal/day	0.39	0.22	0.14	0.22	0.11
Red meat kcal/day	13.27	8.57	6.41	7.71	5.41
Fish kcal/day	0.00	0.01	0.01	0.02	0.01
Milk kcal/day	5.96	3.58	1.38	2.72	1.42
Poultry kcal/day available	13.66	8.22	5.35	8.90	5.53
Eggs kcal/day available	5.43	3.72	2.50	4.35	3.72
Alcohol kcal/day	0.56	1.58	0.30	1.76	0.93
Sugary beverages kcal/day	0.01	8.49	13.37	9.67	11.67
Corn kcal/day available	0.01	0.16	0.19	1.96	1.04
potatoes kcal/day available	2.17	0.83	4.55	2.36	2.60
Saturated fatty acids kcal/day	1.79	7.70	3.12	1.44	3.71
Polyunsaturated fatty acids kcal/day	0.51	1.58	0.88	0.48	0.86
Trans fatty acids kcal/day	0.05	0.20	0.08	0.07	0.07
Fruits kcal/day	0.29	0.73	0.56	0.28	0.79
Vegetables kcal/day	0.45	1.05	0.59	0.63	0.89
Nuts and seeds kcal/day	0.03	0.06	0.03	0.02	0.05
Whole grains kcal/day	0.18	0.26	0.04	0.06	0.01
Legumes kcal/day	0.46	1.10	0.65	0.24	0.47
Rice kcal/day available	3.40	12.76	9.29	7.81	8.14
Sweet potatoes kcal/day available	0.05	0.74	0.18	0.53	0.06
Physical activity METs	13.08	13.10	7.90	9.28	7.11
Kcal available/day	6.52	3.72	10.22	8.31	14.90
Child/infant 2SD underweight	4.86	2.49	7.62	6.03	10.51
Discontinued breast feeding <6 mo	1.52	0.52	0.42	0.45	2.34
Total percent weights in BMI formula	74.65	81.39	75.77	75.28	82.35

660 **Supplementary Table 7. continued**
 661

Risk factors	RCT6 Percent weights	RCT7 Percent weights	RCT8 Percent weights	RCT9 Percent weights	RCT10 Percent weights
Processed meat kcal/day	0.27	0.34	0.26	0.39	0.33
Red meat kcal/day	8.14	11.88	8.69	13.49	14.38
Fish kcal/day	0.01	0.04	0.02	0.04	0.03
Milk kcal/day	2.89	3.98	1.89	3.47	4.88
Poultry kcal/day available	6.69	11.11	5.89	7.89	15.53
Eggs kcal/day available	4.41	5.84	4.14	6.21	7.18
Alcohol kcal/day	0.11	4.39	0.62	7.92	1.26
Sugary beverages kcal/day	10.44	4.11	14.62	0.34	1.79
Corn kcal/day available	0.01	0.00	0.82	1.49	0.01
potatoes kcal/day available	1.82	1.97	1.80	2.15	5.32
Saturated fatty acids kcal/day	8.18	1.87	1.97	3.01	2.42

Polyunsaturated fatty acids kcal/day	1.22	0.41	0.54	0.56	0.45
Trans fatty acids kcal/day	0.13	0.07	0.07	0.06	0.04
Fruits kcal/day	0.71	0.33	0.36	0.43	0.35
Vegetables kcal/day	1.31	0.37	0.41	0.77	0.41
Nuts and seeds kcal/day	0.07	0.02	0.03	0.06	0.03
Whole grains kcal/day	0.30	0.16	0.00	0.22	0.08
Legumes kcal/day	1.54	0.39	0.47	0.57	0.46
Rice kcal/day available	11.87	11.51	8.24	8.93	3.53
Sweet potatoes kcal/day available	0.61	0.03	0.79	0.32	0.10
Physical activity METs	4.24	11.48	11.91	9.73	11.59
Kcal available/day	7.47	4.25	6.31	7.51	3.68
Child/infant 2SD underweight	5.93	4.09	4.88	4.10	4.02
Discontinued breast feeding <6 mo	1.92	1.47	2.96	1.65	0.62
Total percent weights in BMI formula	74.65	81.39	75.77	75.28	82.35

662
663
664

Supplementary Table 7. continued

Risk factors	RCT11 Percent weights	RCT12 Percent weights	RCT13 Percent weights	RCT14 Percent weights	RCT15 Percent weights
Processed meat kcal/day	0.13	0.18	0.28	0.35	0.16
Red meat kcal/day	6.31	6.46	12.56	11.78	6.90
Fish kcal/day	0.01	0.02	0.02	0.01	0.01
Milk kcal/day	1.70	2.12	3.83	2.99	2.35
Poultry kcal/day available	7.81	6.50	11.94	11.25	7.17
Eggs kcal/day available	2.82	3.48	5.55	5.44	2.83
Alcohol kcal/day	0.21	0.06	0.60	0.13	1.35
Sugary beverages kcal/day	10.33	8.84	4.75	4.17	10.87
Corn kcal/day available	1.32	0.33	1.21	0.81	1.29
potatoes kcal/day available	1.73	2.12	7.81	2.91	0.01
Saturated fatty acids kcal/day	2.90	2.49	2.40	2.90	3.65
Polyunsaturated fatty acids kcal/day	0.58	0.70	0.56	0.44	0.88
Trans fatty acids kcal/day	0.05	0.15	0.03	0.08	0.17
Fruits kcal/day	0.28	0.43	0.34	0.50	0.75
Vegetables kcal/day	0.64	0.68	0.40	0.61	0.84
Nuts and seeds kcal/day	0.01	0.02	0.02	0.03	0.04
Whole grains kcal/day	0.16	0.39	0.22	0.21	0.06
Legumes kcal/day	0.38	0.89	0.77	0.29	0.65
Rice kcal/day available	9.30	10.10	8.34	9.08	7.09
Sweet potatoes kcal/day available	0.12	0.18	0.04	0.25	0.67
Physical activity METs	9.01	10.97	6.78	9.80	10.34
Kcal available/day	13.56	11.05	4.34	7.80	12.42
Child/infant 2SD underweight	8.29	8.08	4.01	4.05	7.08
Discontinued breast feeding <6 mo	0.32	0.20	0.58	0.47	1.52
Total percent weights in BMI formula	77.97	76.44	77.38	76.37	79.09

665
666

667 **Supplementary Table 7. continued**

Risk factors	RCT16 Percent weights	RCT17 Percent weights	RCT18 Percent weights	RCT19 Percent weights	RCT20 Percent weights
Processed meat kcal/day	0.33	0.21	0.29	0.43	0.26
Red meat kcal/day	13.14	7.99	7.48	14.07	11.69
Fish kcal/day	0.02	0.02	0.01	1.97	1.28
Milk kcal/day	3.65	2.02	4.49	4.10	3.12
Poultry kcal/day available	11.06	5.98	9.24	11.22	11.02
Eggs kcal/day available	4.90	3.25	4.17	7.73	5.61
Alcohol kcal/day	4.34	1.52	0.01	4.10	3.80
Sugary beverages kcal/day	1.58	12.85	7.01	3.37	6.49
Corn kcal/day available	2.00	0.42	1.95	0.17	0.77
potatoes kcal/day available	2.98	0.18	3.88	6.20	2.33
Saturated fatty acids kcal/day	1.20	5.43	1.81	0.90	2.87
Polyunsaturated fatty acids kcal/day	0.29	1.50	0.49	0.18	1.07
Trans fatty acids kcal/day	0.05	0.18	0.06	0.03	0.07
Fruits kcal/day	0.23	0.88	0.35	0.12	0.47
Vegetables kcal/day	0.32	1.17	0.54	0.23	0.38
Nuts and seeds kcal/day	0.02	0.05	0.03	0.01	0.04
Whole grains kcal/day	0.03	0.00	0.04	0.12	0.02
Legumes kcal/day	0.17	0.68	0.40	0.09	0.56
Rice kcal/day available	7.18	12.12	9.14	6.79	13.19
Sweet potatoes kcal/day available	0.02	1.13	0.04	0.22	0.21
Physical activity METs	13.29	4.32	12.97	9.16	9.16
Kcal available/day	8.12	9.24	7.45	6.10	4.78
Child/infant 2SD underweight	4.37	5.50	6.14	3.27	2.89
Discontinued breast feeding <6 mo	1.29	2.87	0.49	1.22	0.95
Total percent weights in BMI formula	80.57	79.47	78.50	81.80	83.04

668

669 Appendix 1. Worldwide surveys contributing to the IHME GBD risk
670 factor data.

671 Online only.

672

673 **Appendix 2. Bradford Hill causality criteria based** 674 **assessment methodology detailed**

675 A literature search revealed no published methodological precedents for statistically modeling
676 the correlation between mean BMIs of worldwide countries and subnational
677 regions/provinces/states and their corresponding dietary and other risk factors. Since the
678 Bradford Hill causality criteria¹ (enumerated by the English occupational physician and
679 epidemiologist Sir Austin Bradford Hill) are the gold standard assessment tools to test causality
680 of risk factors related to health outcomes, we explored the data looking for mean BMI to risk
681 factor correlations that could be tested with the Bradford Hill criteria.

682
683 The relevant causality criteria included #1 strength, #2 consistency, #3 dose-response (biological
684 gradient), #4 temporality, #5 analogy, #6 plausibility, #7 experimentation, and #8 coherence. The
685 non applicable criterion is specificity. Since diet, physical activity, and other risk factors may
686 affect many non-communicable disease health outcomes, specificity is not relevant. As Dr.
687 Bradford Hill said, “In short, if specificity exists we may be able to draw conclusions without
688 hesitation; if it is not apparent, we are not thereby necessarily left sitting irresolutely on the
689 fence.”¹

690
691 In considering many candidate methodologies involving univariate and multiple regression
692 analysis, we sought a technique that produced the worldwide BMI versus risk factors multiple
693 regression formula which most accurately predicted the mean BMI of subgroups (e.g.,
694 continents, socio-economic index quartiles, etc.). Such a BMI formula with quantifiable high
695 accuracy in predicting the mean BMI of subgroups would necessarily score high with the

696 Bradford Hill criteria strength, consistency, and dose-response (biological gradient). Of the
697 candidate statistical modeling strategies, the methodology that created the worldwide BMI
698 formula that most accurately estimated subgroup mean BMIs had the following steps:

699

700 1. Transform all the food group risk factors (i.e., alcohol, total sugar, plant foods,
701 animal foods, and fatty acids) from g/day to kilocalories/day (kcal/day).

702 2. Multiply each food group risk factor by its mean kcal/day consumption.

703 3. Multiply each risk factor by the R^2 (coefficient of determination) of its
704 univariate correlation with BMI.

705 4. Adjust for multicollinearities (any animal or plant risk factors in the
706 combination variables that have significant multicollinearity (when an
707 independent variable is highly correlated with one or more of the other
708 independent variables in a multiple regression equation)).

709 5. After performing #2, #3, and #4 above, form combination dietary variables
710 with individual foods each expressed in kcal/day to be added together as the
711 data suggests are appropriate:

712 a. For example: processed meat * processed meat kcal/day * processed
713 meat R^2 * processed meat kcal/day in bottom three SDI quartiles/top
714 quartile + red meat * red meat R^2 * red meat kcal/day * red meat
715 kcal/day in bottom three SDI quartiles/ kcal/day in the top quartile +
716 ...+ eggs * eggs kcal/day, etc.

- 717 b. Explore methods to adjust for multicollinearities based in part on maximizing
718 the closeness of fit of the multiple regression derived BMI formula outputs
719 with the mean BMIs of 37 subgroups of the worldwide data.
- 720 6. Add physical activity before performing multiple regression.
- 721 7. Perform multiple regression analysis with the food combination variables in #5 and
722 physical activity to generate a BMI formula. (e.g., BMI formula= combination
723 variable_a * parameter estimate_a + combination variables_b * parameter estimate_b ...+
724 physical activity * parameter estimate of physical activity.
- 725 8. Explore integrating the variables kcal available, child underweight, and discontinued
726 breast feeding into the BMI formula.
- 727 9. Discard any of these variables that's sign in regression with BMI switches from
728 univariate to the multivariate analysis (e.g., + in univariate and - in multivariate
729 formula).
- 730 10. To simplify the calculations, use the Excel spreadsheet and transfer back and forth to
731 SAS Studio.
- 732 11. To test the BMI formula with subgroups, match the BMI formula mean and standard
733 deviation (SD) with that of mean worldwide BMI:
- 734 a. Adjust the SD of the BMI formula output to equal the SD of the worldwide
735 BMI by multiplying the BMI formula output times the BMI SD / BMI
736 formula SD.
- 737 b. Adjust the BMI formula constant to equal the worldwide population's mean
738 BMI by adding the difference between the mean BMI and the BMI formula
739 constant.

740 12. Explore the BMI formula functionality with different Bradford Hill criteria with
741 standardising the BMI and BMI risk factors or with all variables non standardised.

742
743 We estimated percent weights attributable to each risk factor with the standardised BMI formula
744 from the above methodology with the following additions:

745 1. Equate the sum of the individual food and other risk factor parameter estimates to the
746 total R^2 of the BMI formula:

747 a. Total the absolute values of the parameter estimates for each food and any non
748 food risk factors.

749 b. Divide the worldwide BMI formula R^2 by the sum of the risk factor parameter
750 estimates.

751 c. Multiply the result of b above times each of the individual risk factor parameter
752 estimates.

753 2. Multiply the results in #1 above times 100 to generate percent weights for each risk
754 factor. The sum of the risk factor percent weights will equal the BMI formula R^2 times
755 100.

756
757 To cross validate the generated worldwide BMI formula experimentally, create 20 random
758 subgroups of 100 cohorts with the macro function in SAS. Determine each of the 20 subgroup
759 BMI formulas, and, from those 20 BMI formulas, create a table with: mean, SD, minimum, and
760 maximum for each of the risk factors.

761

762

763 **Bradford Hill causal criteria testing methodology applied to current study**

764 The relevant eight of the nine original Bradford Hill criteria were each scored as:

765 “5” very strongly supporting causality,

766 “4” strongly supporting causality,

767 “3” moderately strongly supporting causality,

768 “2” supporting causality,

769 “1” weakly supporting causality, and

770 “0” not supporting causality

771

772 The scoring for each Bradford Hill causal criterion was as follows:

773 1. Strength: The correlation coefficient, r , of the worldwide multiple regression derived
774 BMI formula with BMI (dependent variable) and BMI risk factors (independent
775 variables) assessed strength.

776 Scoring of strength:

777 5=BMI formula regressed with BMI $r \geq 0.50$ and $P < 0.0001$

778 4=BMI formula regressed with BMI $0.50 > r \geq 0.40$ and $P < 0.0001$

779 3=BMI formula regressed with BMI $0.40 > r \geq 0.30$ and $P < 0.0001$

780 2=BMI formula regressed with BMI $0.30 > r \geq 0.20$ and $P < 0.0001$

781 1=BMI formula regressed with BMI $0.20 > r \geq 0.10$ and $P < 0.0001$

782 0=BMI formula regressed with BMI $r < 0.10$ or $P \geq 0.0001$

783

- 784 2. Consistency: For the purposes of this study, consistency between BMI and BMI formula
785 output was determined by comparing the mean BMI and the mean BMI formula output in
786 each of the following 37 subgroups:
- 787 1) A variable socio-demographic index (SDI)—see Supplementary Table 1 for
788 definition of SDI—divided the world’s population by quartiles of SDI.
 - 789 2) A variable “continents” allowed for analyses of countries from each of the six
790 inhabited continents.
 - 791 3) The four countries (UK, USA, Mexico, and Japan) with subnational data on BMI and
792 the risk factors were grouped and assessed to compare the BMI formula output with
793 the overall mean BMI in those countries.
 - 794 4) Based on the total kcal/day of all foods that increased BMI, a combination variable
795 was constructed and the world’s population divided into quartiles from the highest to
796 lowest total kcal/day.
 - 797 5) Similarly, based on the total kcal/day of all foods that decreased BMI, we divided the
798 world’s population into quartiles from the highest to lowest by total kcal/day.
 - 799 6) Based on physical activity (METs/week), the world’s population was divided into
800 quartiles.
 - 801 7) We evaluated dose response by dividing the BMI formula output into quartiles from
802 the highest to lowest.
 - 803 8) The four countries with subnational data were individually evaluated.
 - 804 9) The first four of the 20 random number generated database subgroups were included
805 in the consistency analysis.
 - 806 10) Male and female were separately assessed.

807 11) Assessing dose-response was done in quartiles by the BMI formula outputs after the
808 BMI formula was harmonized with the mean and SD of worldwide mean BMI. See
809 the dose-response criteria (#4).

810
811 For each of the 37 subgroups, the absolute differences between the means of BMI and the
812 BMI formula output in were totaled (e.g., continent Africa BMI formula output - mean
813 BMI for Africa, etc.).

814 Scoring of consistency for BMI compared with BMI formula output for each of the 37
815 subgroups:

816 5=The mean of the absolute differences between mean BMI and BMI formula output was
817 ≤ 0.40 units.

818 4=The mean of the absolute differences between mean BMI and BMI formula output was
819 ≤ 0.50 units.

820 3=The mean of the absolute differences between mean BMI and BMI formula output was
821 ≤ 0.60 units.

822 2=The mean of the absolute differences between mean BMI and BMI formula output was
823 ≤ 0.70 units.

824 1=The mean of the absolute differences between mean BMI and BMI formula output was
825 ≤ 1.0 units.

826 0=The mean of the absolute differences between mean BMI and BMI formula output was
827 > 1.0 units.

828

829 3. Dose-response (biological gradient): Dr. Bradford Hill thought that a clear dose-response
830 effect on the incidence of disease with exposure to a single risk factor was the clearest
831 evidence of a causal relationship. More recently, it has been realized that cause and effect
832 relationships are often more complex. In this analysis of dose-response, instead of using
833 single risk factor levels related to BMI, levels of a multivariable regression derived BMI
834 formula outputs in quartiles were related to mean BMIs in those quartiles. Dose-response
835 of BMI formula estimates versus mean BMI were included in the testing of consistency
836 above.

837 Scoring of dose-response (biologic gradient) was based on this mean absolute difference
838 when the BMI formula output was divided into quartiles:

839 5= The BMI formula output versus mean BMI absolute differences from each of the four
840 quartiles averages ≤ 0.40 BMI units.

841 4= The BMI formula output versus mean BMI absolute differences from each of the four
842 quartiles averages ≤ 0.50 units.

843 3= The BMI formula output versus mean BMI absolute differences from each of the four
844 quartiles averages ≤ 0.60 BMI units.

845 2= The BMI formula output versus mean BMI absolute differences from each of the four
846 quartiles averages ≤ 0.70 BMI units.

847 1= The BMI formula output versus mean BMI absolute differences from each of the four
848 quartiles averages ≤ 1.0 BMI units.

849 0= The BMI formula output versus mean BMI absolute differences from each of the four
850 quartiles averages > 1.0 BMI units

851

852 4. Temporality: The dictionary defines the noun temporality as: "The state of existing
853 within or having some relationship with time." Bradford Hill said, "Temporality refers to
854 the necessity that the cause to precede the effect in time. This criterion is unarguable,
855 insofar as any claimed observation of causation must involve the putative cause C
856 preceding the putative effect D. It does not, however, follow that a reverse time order is
857 evidence against the hypothesis that C can cause D. Rather, observations in which C
858 followed D merely shows that C could not have caused D in these instances; they provide
859 no evidence for or against the hypothesis that C can cause D in those instances in which it
860 precedes D."

861
862 Dr. Bradford Hill was an occupational physician before the current availability in
863 nutritional epidemiology of data on trends over 28 years of 18 components of worldwide
864 diets along with the global BMI trend. Consequently, now it is fair to test temporality by
865 deriving a standardised multiple regression formula with BMI trend, measured by the
866 slope of the least squared regression line (LSRL) over 1990-2017, as (dependent
867 variable). The independent variables would consist of the LSRL trends over 1990-2017
868 of the same dietary components, physical activity, and other variables as in the original
869 BMI formula. We considered the strength (r) of the BMI trend formula versus the BMI
870 trend an appropriate measure of temporality. Any risk factor was excluded from the trend
871 BMI formula if its sign (+ or -) did not match the sign in the original BMI formula.

872
873 With a standardised multiple regression methodology similar to that of deriving risk
874 factor percent weights above, a standardised BMI trend (dependent variable) versus

875 standardised risk factor trends (independent variables) formula was derived with the risk
876 factor parameter estimates adjusted to equate to trend percent weights.

877 Scoring of temporality: The Pearson correlation coefficient, r , of the resulting multiple
878 regression derived BMI trend versus risk factors trends formula:

879 $5=r \geq 0.50$ and $P < 0.0001$.

880 $4=0.50 > r \geq 0.40$ and $P < 0.0001$.

881 $3=0.40 > r \geq 0.30$ and $P < 0.0001$.

882 $2=0.30 > r \geq 0.20$ and $P < 0.0001$.

883 $1=0.20 > r \geq 0.10$ and $P < 0.0001$.

884 $0=r < 0.10$ or $P \geq 0.0001$

885

886 5. Analogy: High BMI is among the four metabolic risk factors that are strongly associated with
887 cardiovascular diseases, cancers, and other non-communicable diseases. The other major
888 metabolic risk factors for non-communicable diseases are high systolic blood pressure (SBP),
889 high low density lipoprotein cholesterol (LDL-C), and high fasting plasma glucose (FPG).

890

891 We tested analogy by the number of the diet, physical activity, and other variables in the
892 percent weight of risk factor BMI formula that were also in the percent weight of risk factor
893 formulas of the other metabolic factors and including only risk factors concordant in
894 direction (+ or -) with the coefficients of percent weights of risk factors in the BMI formula.

895 Scoring of analogy:

896 5=At least three-quarters of BMI formula risk factors are also in the risk factor formulas
897 of all three of the other metabolic factors (SBP, LDL-C, and FPG) and have concordant
898 signs (+ or -) for each risk factor.

899 4= At least three-quarters of BMI formula risk factors are also in the risk factor formulas
900 of two of the three other metabolic factors (SBP, LDL-C, and FPG) and have concordant
901 signs (+ or -) for each risk factor.

902 3= At least three-quarters of BMI formula risk factors are also in the risk factor formulas
903 of one of the three other metabolic factors (SBP, LDL-C, and FPG) and have concordant
904 signs (+ or -) for each risk factor.

905 2= At least two-thirds of BMI formula risk factors are also in the risk factor formulas of
906 one of the three other metabolic factors (SBP, LDL-C, and FPG) and have concordant
907 signs (+ or -) for each risk factor.

908 1=At least half of BMI formula risk factors are also in the risk factor formulas of one of
909 the three other metabolic factors (SBP, LDL-C, and FPG) and has concordant signs (+ or
910 -) for each risk factor.

911 0= None of the above.

912

913 6. Plausibility: To test plausibility, we looked to find if any of our findings were at variance
914 with the preponderance of studies published. We searched the medical literature
915 particularly for systematic reviews of the relationships of foods and other variables with
916 BMI.

917 Scoring of plausibility:

918 5=None of the current findings were at variance with the preponderance of the medical
919 literature.

920 4= One of the current findings was at variance with the preponderance of the medical
921 literature.

922 3= Two of the current findings were at variance with the preponderance of the medical
923 literature.

924 2=Three of the current findings were at variance with the preponderance of the medical
925 literature.

926 1=Four of the current findings were at variance with the preponderance of the medical
927 literature.

928 0= Five or more of the current findings were at variance with the preponderance of the
929 medical literature.

930

931 7. Experiment: Dr. Bradford Hill thought that evidence drawn from experimentation,
932 including in epidemiologic studies, may lead to the strongest support for causal
933 inference.¹

934

935 We used a cross validation method to assess Bradford Hill's "experiment" criterion.

936 Random number generation of 20 subgroups each with 100 cohorts derived 20

937 standardised BMI formulas to compare with the standardised worldwide BMI formula.

938 Scoring of experiment:

939 5=At least 15 subgroups BMI formulas included all same risk factors with the same signs
940 of coefficients as the worldwide BMI formula.

- 941 4=At least 10 subgroups of BMI formulas included all same risk factors with the same
942 signs of coefficients as the worldwide BMI formula.
- 943 3=All 20 subgroups of BMI formulas included at least three-quarters of same risk factors
944 with the same signs of coefficients as the worldwide BMI formula.
- 945 2=At least 10 subgroups of BMI formulas included at least three-quarters of same risk
946 with the same signs of coefficients factors as the worldwide BMI formula.
- 947 1= At least 10 subgroups of BMI formulas included at least half of same risk factors with
948 the same signs of coefficients as the worldwide BMI formula.
- 949 0=None of the above.
- 950
- 951 8. Coherence: According to Dr. Bradford Hill, "...cause and effect interpretation of our data
952 should not seriously conflict with the generally known facts of the natural history and
953 biology of the disease."¹ In this analysis of BMI associated with BMI formula estimates,
954 coherence was the numerical total score of the above seven relevant causality criteria
955 each on a 0-5 scale. The maximum score was 40.
- 956 Scoring of coherence:
- 957 5=Score on the first seven Bradford Hill causation criteria=35-40.
- 958 4=Score on the first seven Bradford Hill causation criteria=30-34.
- 959 3=Score on the first seven Bradford Hill causation criteria=25-29.
- 960 2=Score on the first seven Bradford Hill causation criteria=20-24.
- 961 1=Score on the first seven Bradford Hill causation criteria=15-24.
- 962 0=Score on the first seven Bradford Hill causation criteria<15.
- 963

964 **Reference**

965 **1. Hill AB. THE ENVIRONMENT AND DISEASE: ASSOCIATION OR CAUSATION?**

966 *Proceedings of the Royal Society of Medicine.* 1965; **58**(5):295-300.

967 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1898525/>

968

969

970

971

972

