medRxiv preprint doi: https://doi.org/10.1101/2021.08.04.21261538; this version posted August 5, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

Title: Association between obesity and hospitalization in mild COVID-19 young adult outpatients in Brazil: a prospective cohort study

Ivaine Tais Sauthier Sartor[1*], Caroline Nespolo de David[1*], Gabriela Heiden Telo[2], Gabriela Oliveira Zavaglia[1], Ingrid Rodrigues Fernandes[1], Luciane Beatriz Kern[1], Márcia Polese-Bonatto[1], Thaís Raupp Azevedo[1], Amanda Paz Santos[1], Walquiria Aparecida Ferreira de Almeida[3], Victor Bertollo Gomes Porto[3], Fernanda Hammes Varela[1,2], Marcelo Comerlato Scotta[1,2], Regis Goulart Rosa[1], Renato T. Stein[1,2].

*These authors contributed equally to this manuscript.

1. Social Responsibility – PROADI-SUS, Hospital Moinhos de Vento, Porto Alegre, Brazil

2. School of Medicine, Pontifícia Universidade Católica do Rio Grande do Sul, Porto Alegre, Brazil

3. General Coordination, Programa Nacional de Imunização, Brazilian Ministry of Health

Corresponding author: Ivaine Tais Sauthier Sartor, ivaine.sartor@hmv.org.br, phone: 55-51-983016858, address: Ramiro Barcelos street, 630/1002, ZIP-Code: 90035-001, Porto Alegre, Brazil

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

medRxiv preprint doi: https://doi.org/10.1101/2021.08.04.21261538; this version posted August 5, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

Abstract

Background/Objectives

The aim of this study was to evaluate the association between obesity and hospitalization in mild COVID-19 adult outpatients in Brazil.

Subjects/Methods

Adults with signs and symptoms suggestive of acute SARS-CoV-2 infection who sought two hospitals (one public and one private) emergency department were prospectively enrolled. Patients with confirmed COVID-19 at inclusion were followed by phone calls at day (D) D7, D14 and D28. Multivariable logistic regression models were employed to explore the association between obesity and other potential predictors for hospitalization.

Results

A total of 1,050 participants were screened, 310 were diagnosed with COVID-19 by RT-PCR. Median age was 37.4 (IQR 29.8-45.0) years, and 186 (60.0%) were female. Duration of symptoms was 3.0 (IQR 2.0-5.0) days, and 10.0 (IQR 8.0-12.0) was the median number of symptoms at inclusion. A total of 98 (31.6%) were obese, and 243 (78.4%) had no previous medical conditions. Twenty three participants (23/310, 7.4%) required hospitalization during the period. After adjusting, obesity (BMI≥30.0 kg/m^2) (OR=2.69, 95%CI 1.63-4.83, P<0.001) and older age (OR=1.05, 95%CI 1.01-1.09, P<0.001), were significantly associated with higher risks of hospitalization.

Conclusions

Obesity, followed by aging, was the main factor associated with hospital admission for COVID-19 in a young population in a low-middle income country. Our findings highlighted the need for actions to promote additional protection for obese population, such as vaccination, and to encourage lifestyle changes.

Keywords: mild COVID-19, noncritical COVID-19, young adults, follow-up, outpatients, hospitalization

medRxiv preprint doi: https://doi.org/10.1101/2021.08.04.21261538; this version posted August 5, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

Introduction

Coronavirus disease 2019 (COVID-19) has brought large numbers of patients to medical attention within a short time span for care of a previously undescribed illness, challenging health services and providers to deliver time-sensitive interventions under difficult circumstances. Since the first cases of COVID-19 were documented in China in late December 2019, the epidemiologic focus has largely been on establishing the severity of disease and outcomes among severe patients[1,2]. Older age and some metabolic disorders such as diabetes were early found to be predictors of hospital admission and mortality in patients with COVID-19[3].

The COVID-19 pandemic has occurred at a time when the prevalence of obesity is increasing all around the world. Obesity is a well-established risk factor for metabolic non-communicable diseases such as type 2 diabetes and hypertension, highly influencing most major cardiovascular diseases. With the worldwide spread of acute respiratory syndrome coronavirus 2 (SARS-CoV-2), obesity has also emerged as an independent risk factor for severe COVID-19, likely related to the adipose tissue inflammation's effects on the immune system, besides all the metabolic dysfunction associated[4]. In a recent United States Department of Health and Human Services/Centers for Disease Control and Prevention publication, overweight and obesity were both identified as risk factors for the need of mechanical ventilation related to COVID-19, while only obesity was associated with hospitalization and death in an adult cohort of inpatients[5]. However, as risk estimated for severe COVID-19-associated outcomes were measured mostly among adults who received care at a hospital, the current knowledge about the impact of obesity on COVID-19 hospital admission may possibly be overestimated due to a selection bias, as most studies evaluated only inpatients.

Few studies were developed in mild COVID-19 adult outpatients, especially in low-middle income countries (LMIC). Given the uncertainty regarding outcomes in this low-risk outpatient population, and considering the exponential rise in the prevalence of obesity and its association with severe forms of the disease, this study aimed to evaluate the association

medRxiv preprint doi: https://doi.org/10.1101/2021.08.04.21261538; this version posted August 5, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

between obesity and hospitalization in mild COVID-19 adult outpatients in Brazil. We also aimed to describe the clinical characteristics of COVID-19 outpatients and identify other predictors of hospitalization.

Materials/Subjects and Methods

1. Study design and population

This is a prospective cohort study with data collected in referral hospitals from Southern Brazil, one public (Hospital Restinga e Extremo Sul) and the other from the private health system (Hospital Moinhos de Vento). From 13th May to 14th September 2020 we assessed adult (≥18 years old) outpatients who had a visit in the emergency department (ED) health system for COVID-19 symptoms. Inclusion criteria were presenting at least one sign or symptom suggestive of COVID-19 (cough, fever, or sore throat), within 14 days of onset of symptoms. We then created a cohort of those with confirmed COVID-19, defined as a positive result on real-time reverse transcriptase (RT-PCR) for SARS-CoV-2.

2. Study procedures

At baseline, clinical, demographic, comorbidities, and signs or symptoms suggestive of COVID-19 data were collected. All participants underwent bilateral nasopharyngeal and oropharyngeal swab collection for SARS-CoV-2 detection via RT-PCR assay. The samples were analyzed in the Molecular Biology Laboratory at Hospital Moinhos de Vento, as described previously[6]. Reported weight and height were used to calculate body mass index (BMI). Participants were followed by phone interviews at 7, 14 and 28 days after seeking care for COVID-19 symptoms (study baseline). Up to ten telephone call attempts were made during a period of three days from the D7, D14 or D28. When necessary, text messages were sent in order to notify that the researchers were trying to contact them. If there was no success in contacting one of the time points, the questions were asked retroactively on the next call. The questions were asked directly to the participant or, in his absence or incapacity, for a person authorized by him at baseline. The participant was considered lost

medRxiv preprint doi: https://doi.org/10.1101/2021.08.04.21261538; this version posted August 5, 2021. The copyright holder for this preprint
(which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
It is made available under a CC-BY-NC-ND 4.0 International license .

when it was not possible to make any contact in different day periods (including failure in 10 call attempts or time exceeded considering 3-day window) within the 28-day follow-up.

During the follow up interviews, participants were asked about the need of hospital admission, use of supplemental oxygen, admission at intensive care units (ICU), and use of invasive mechanical ventilation. All data were collected by trained researchers and using standardized questionnaires developed for the study in the Research Electronic Data Capture software (REDCap)[7,8].

The outcome of interest was unplanned hospital admission in the period of 28 days. In order to predict the outcome, demographic and clinical characteristics as age at time of enrollment (years), sex, days of onset of symptoms, racial or ethnic group, active or second hand smoke at home, hospital at inclusion, number of symptoms and number of medical conditions were considered. In order to evaluate obesity as a predictor for the outcome of interest, we used BMI and participants were classified as non-obese (<30.0 kg/m^2), and obesity (≥30.0 kg/m^2).

3. Statistical analyses

Data normality assumptions were verified using the Shapiro-Wilk test for continuous variables, median values and interquartile ranges (IQR) were calculated, and two-tailed Mann-Whitney-Wilcoxon test was used. Pearson's Chi-square or Fisher's exact tests were used to evaluate proportions. Uni and multivariable logistic regression analyses were performed to assess the predictors of hospitalization considering the follow up period of 28 days after having a COVID-19 health care visit (study baseline). The association between obesity and hospitalization was assessed using multiple logistic regression models with adjustment for relevant covariates (hospital at inclusion; years of education; number of medical conditions (none, one, two or more); days of onset of symptoms to inclusion; number of symptoms (≤8, 9-10, 11-12, ≥13); and cardiovascular diseases, which includes: ischemic heart disease, heart failure and hypertension). We performed additional models including covariates considered risk factors to hospitalization in order to assess consistency

medRxiv preprint doi: https://doi.org/10.1101/2021.08.04.21261538; this version posted August 5, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

and risk of bias. Odds ratios (ORs) with 95% confidence intervals (CIs) were calculated. All data preprocessing and analyses were performed in R 3.5.0 statistical software[9].

4. Ethics

The study was performed in accordance with the Decree 466/12[10] of the National Health Council and Good Clinical Practice Guidelines, after approval by the Institutional Review Board (IRB nº 4.637.933). All participants included in this study provided written informed consent.

Results

In this study 1,050 subjects were screened and 13 were excluded (9 for not meeting inclusion criteria, 2 for not consenting, and 1 withdrew consent, and 1 without SARS-CoV-2 RT-PCR result), and 727 participants with negative RT-PCR results. A total of 310 adults with confirmed SARS-CoV-2 were included. The success in the 28-days follow-up was obtained for 297 (95.8%) participants, as shown in Figure 1.

At baseline, the median participants' age was 37.4 years (IQR, 29.8-45.0). Out of 310 participants, 186 (60.0%) were female, 222 (71.6%) declared themselves caucasian, and 173 (55.8%) were from the public hospital. The median days of symptoms was 3.0 (IQR, 2.0-5.0), and 10.0 (IQR 8.0-12.0) was the median number of symptoms at inclusion. The most commonly reported symptoms were headache (90.6%), cough (87.4%), malaise (77.1%), and myalgia (76.5%). Few participants were active or second hand smoker at home (17.7%), and more than half was overweight (112, 36.1%) or obese (98, 31.6%). Most participants (78.4%) had no previous medical conditions. Hypertension (12.6%), asthma (6.1%) and type 1 or type 2 diabetes (3.5%) were the most frequent baseline comorbidities reported.

At the 28-day follow-up, 23/310 (7.4%) participants required hospitalization. The median BMI of hospitalized participants was 32.4kg/m^2 (IQR, 30.2-34.1), whereas for the outpatients was 27.7kg/m^2 (IQR, 24.3-30.5) (P<0.001). The median of the duration from onset of symptoms at the hospitalizations and at enrollment was 14.0 (IQR, 8.0-17.5) and 9.5 (IQR, 6.0-13.5) days, respectively. During the hospitalization, 15/23 (65.2%) participants required only use of

medRxiv preprint doi: https://doi.org/10.1101/2021.08.04.21261538; this version posted August 5, 2021. The copyright holder for this preprint
(which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
It is made available under a CC-BY-NC-ND 4.0 International license .

158 supplemental oxygen, 6/23 (26.1%) were admitted to ICU, 2/23 (8.7%) used invasive
159 mechanical ventilation, and one participant died (1/23, 4.3%).
160 Participants who required hospitalization were significantly older (P=0.005) and obese
161 (P<0.001) compared with those who did not. No other differences were observed between
162 groups, as shown in Table 1. In multivariable logistic regression analysis age and obesity
163 remained as an important predictor of hospital admission in our cohort (age: OR=1.05,
164 95%CI 1.01-1.09; obesity: OR=2.69, 95%CI 1.63-4.83). When adjusted for hospital of
165 inclusion, years of education, number of medical conditions, cardiovascular diseases, days
166 of onset of symptoms and number of symptoms at enrollment, age and obesity remained
167 associated with hospitalization, as shown in Figure 2.
168 Among the 15 obese participants with COVID-19 admitted at hospital within 28 days of
169 follow up, the median age was 42.0 years (IQR, 37.2-51.7), 9 (60.0%) were male, 8 (53.3%)
170 were admitted at private hospitals. Four (26.6%) participants reported underlying medical
171 conditions (hypertension, diabetes mellitus, asthma, chronic obstructive pulmonary disease
172 or ischemic heart disease). The median days of onset of symptoms to hospitalization was
173 8.0 (IQR, 7.0-9.0), and the median days from the ED sought to hospitalization was 5.5 (IQR,
174 4.0-7.0). Ten (66.7%) participants required supplemental oxygen, 4 (26.7%) were admitted
175 to ICU, one needed invasive mechanical ventilation (6.7%) and one participant died (6.7%).
176
177 **Discussion**
178 Obesity is a recognized risk factor associated with worse outcomes related to COVID-19;
179 however, most of the previous studies were carried out in a population of patients already
180 admitted to the hospital or at a high admission rate due to the disease[11]. In this study, we
181 evaluated the risk of hospital admissions in LMIC, through an outpatient cohort of young
182 patients with symptoms of COVID-19 enrolled early in the course of the disease. Our results
183 identified, for a low-risk and young population, that obesity was the main predictor of
184 hospitalization for COVID-19, with approximately 3-fold increase in the risk of hospital
185 admission in comparison to non-obese patients.

medRxiv preprint doi: https://doi.org/10.1101/2021.08.04.21261538; this version posted August 5, 2021. The copyright holder for this preprint
(which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
It is made available under a CC-BY-NC-ND 4.0 International license .

The world is still struggling to fight the COVID-19 pandemic. According to the World Health Organization (WHO), the mortality rate from COVID-19 is 0.5-1.0%[12], while the hospitalization rate for the disease is 5.0-15.0%. Based on currently available information, the Centers for Disease Control and Prevention has identified severe obesity as an important risk factor for worse prognosis and higher mortality in patients with COVID-19, while any degree of obesity has been associated with COVID-19 poor prognosis. In our study, 7.4% of our sample required hospital admission, similarly to data from literature. Most participants were overweight/obese at baseline, but only obesity predicted the main outcome. In a meta-analysis of 75 studies[13], obesity had a 46.0% higher risk for being COVID-19 positive, a 113.0% higher risk of being hospitalized, and a 48.0% increased mortality. There are many mechanisms that jointly may explain this impact; although the effect of obesity on the immune system remains not completely established, it appears to have an effect independent of coexisting risk factors. Data from literature also identify a dose-response relationship between BMI and COVID-19 worse outcomes[5].

In addition to metabolic factors that seem to impact the severity of COVID-19, advanced age has been identified as one of the major factors associated with worse outcomes in the pandemic, being one of the most important predictors of hospital admission[11]. In our sample, the mean age of participants was younger than most studies in the literature; however, we also identified an increased risk of hospitalization for COVID-19 with each increased year of age. Concerns brought by our data include the fact that the predictors of higher risk for COVID-19 involve populations especially in need of care by health system[5]. The obese and elderly individuals are possibly the ones that will be mostly compromised by the social distancing that the pandemic generates, possibly resulting in worse health habits, such as physical inactivity and unhealthy eating habits, besides increased emotional distress. This could further aggravate the risk related to COVID-19 for this population, unless governmental and health measures, such as vaccination and strategies to increase consumption of healthier foods, stimulate physical activity and promote mental health[13], aimed at the health of this population, starts to be prioritized.

medRxiv preprint doi: https://doi.org/10.1101/2021.08.04.21261538; this version posted August 5, 2021. The copyright holder for this preprint
(which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
It is made available under a CC-BY-NC-ND 4.0 International license .

Our study has limitations. Our analysis was based on a convenience sample. Moreover, our participants represent a single geographic region of a heterogeneous country, such as Brazil. However, worldwide prevalence of obesity is a public health concern and our findings are in line with previous reports[14–17]. As another point to address, the two centers included in this study represent different populations, public and private, which may characterize different clusters and limit the interpretation of our results for the sample as a whole. Also, most of our data represents self-report and may be subject to bias. Despite that, we believe that our results add to the literature as they represent an outpatient prospective cohort of young individuals with low risk for complications related to COVID-19 in a LMIC, identifying important risk predictors for this population. Moreover, a similar impact of obesity was consistently found despite adjusting for several possible confounders.

In conclusion, our data identified that obesity, in addition to age, was the most important risk factor predicting hospital admission for COVID-19 in a young population in Southern Brazil. These results highlight the importance of health support strategies aimed at this population, in order to promote additional protection, such as vaccination, and to encourage lifestyle changes.

Acknowledgments

We thank the Scientific Committee of the Research Support Nucleus (NAP) of Moinhos de Vento Hospital for technical-scientific consultancy. We thank the inclusion personal, laboratory team, and site staff from Hospital Moinhos de Vento and from Hospital Restinga e Extremo Sul.

Funding

This work was supported by the Brazilian Ministry of Health, through the Institutional Development Program of the Brazilian National Health System (PROADI-SUS) in collaboration with Hospital Moinhos de Vento. The Ministry of Health had no role in study

medRxiv preprint doi: https://doi.org/10.1101/2021.08.04.21261538; this version posted August 5, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

241 design, sample and data collection, data analysis and interpretation, as well as in the
242 decision to publish.
243
244 **Competing Interests**
245 The authors declare no competing financial interests.
246
247 **References**
248 1 Grasselli G, Zangrillo A, Zanella A, Antonelli M, Cabrini L, Castelli A *et al.* Baseline
249 Characteristics and Outcomes of 1591 Patients Infected With SARS-CoV-2 Admitted to
250 ICUs of the Lombardy Region, Italy. *JAMA* 2020; **323**: 1574.
251 2 Altonen BL, Arreglado TM, Leroux O, Murray-Ramcharan M, Engdahl R. Characteristics,
252 comorbidities and survival analysis of young adults hospitalized with COVID-19 in New
253 York City. *PloS One* 2020; **15**: e0243343.
254 3 Berlin DA, Gulick RM, Martinez FJ. Severe Covid-19. *N Engl J Med* 2020; **383**: 2451–
255 2460.
256 4 Huang Y, Lu Y, Huang Y-M, Wang M, Ling W, Sui Y *et al.* Obesity in patients with COVID-
257 19: a systematic review and meta-analysis. *Metabolism* 2020; **113**: 154378.
258 5 Kompaniyets L, Goodman AB, Belay B, Freedman DS, Sucosky MS, Lange SJ *et al.*
259 Body Mass Index and Risk for COVID-19–Related Hospitalization, Intensive Care Unit
260 Admission, Invasive Mechanical Ventilation, and Death — United States, March–
261 December 2020. *Morb Mortal Wkly Rep* 2021; **70**: 355–361.
262 6 Polese-Bonatto M, Sartor ITS, Varela FH, Gianinni GLT, Azevedo TR, Kern LB *et al.*
263 Children have similar RT-PCR cycle threshold for SARS-CoV-2 in comparison with adults.
264 *medRxiv* 2021; : 2021.04.20.21255059.
265 7 Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic
266 data capture (REDCap)--a metadata-driven methodology and workflow process for
267 providing translational research informatics support. *J Biomed Inform* 2009; **42**: 377–381.
268 8 Harris PA, Taylor R, Minor BL, Elliott V, Fernandez M, O'Neal L *et al.* The REDCap

medRxiv preprint doi: https://doi.org/10.1101/2021.08.04.21261538; this version posted August 5, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

consortium: Building an international community of software platform partners. *J Biomed Inform* 2019; **95**: 103208.

9 R: The R Project for Statistical Computing. https://www.r-project.org/ (accessed 24 Jun2021).

10 Ministério da Saúde. https://bvsms.saude.gov.br/bvs/saudelegis/cns/2013/res0466_12_12_2012.html (accessed 13 Jul2021).

11 Petrilli CM, Jones SA, Yang J, Rajagopalan H, O'Donnell L, Chernyak Y *et al.* Factors associated with hospital admission and critical illness among 5279 people with coronavirus disease 2019 in New York City: prospective cohort study. *BMJ* 2020; **369**: m1966.

12 Estimating mortality from COVID-19: Scientific brief, 4 August 2020. https://www.who.int/publications-detail-redirect/WHO-2019-nCoV-Sci-Brief-Mortality-2020.1 (accessed 30 Jul2021).

13 Popkin BM, Du S, Green WD, Beck MA, Algaith T, Herbst CH *et al.* Individuals with obesity and COVID-19: A global perspective on the epidemiology and biological relationships. *Obes Rev Off J Int Assoc Study Obes* 2020; **21**: e13128.

14 Sales-Peres SH de C, de Azevedo-Silva LJ, Bonato RCS, Sales-Peres M de C, Pinto AC da S, Santiago Junior JF. Coronavirus (SARS-CoV-2) and the risk of obesity for critically illness and ICU admitted: Meta-analysis of the epidemiological evidence. *Obes Res Clin Pract* 2020; **14**: 389–397.

15 Simonnet A, Chetboun M, Poissy J, Raverdy V, Noulette J, Duhamel A *et al.* High Prevalence of Obesity in Severe Acute Respiratory Syndrome Coronavirus-2 (SARS-CoV-2) Requiring Invasive Mechanical Ventilation. *Obes Silver Spring Md* 2020; **28**: 1195–1199.

16 Nakeshbandi M, Maini R, Daniel P, Rosengarten S, Parmar P, Wilson C *et al.* The impact of obesity on COVID-19 complications: a retrospective cohort study. *Int J Obes* 2020; **44**: 1832–1837.

medRxiv preprint doi: https://doi.org/10.1101/2021.08.04.21261538; this version posted August 5, 2021. The copyright holder for this preprint
(which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
It is made available under a CC-BY-NC-ND 4.0 International license .

297 17 Kang Z, Luo S, Gui Y, Zhou H, Zhang Z, Tian C *et al.* Obesity is a potential risk factor

298 contributing to clinical manifestations of COVID-19. *Int J Obes* 2020; **44**: 2479–2485.

299

medRxiv preprint doi: https://doi.org/10.1101/2021.08.04.21261538; this version posted August 5, 2021. The copyright holder for this preprint
(which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
It is made available under a CC-BY-NC-ND 4.0 International license .

Figure 1. Study flowchart. Thirty three participants out of 297 answered the questions retroactively since there was no success in contact at that time point. Twelve responsible family members were the respondents. (RT-PCR) Reverse transcription polymerase chain reaction. (SARS-CoV-2) Severe acute respiratory syndrome coronavirus 2.

Figure 2. Forestplot of multivariable logistic regression models adjusting age and obesity for the covariates: hospital of inclusion (reference: private hospital at inclusion), years of education (reference: 11 years or less, which corresponds to complete high school education), number of medical conditions (reference: none), cardiovascular diseases (reference: no), days of onset of symptoms (continuous) and number of symptoms (reference: ≤8 symptoms) at enrollment. *Number of medical conditions comprised the sum of cases of: hypertension, asthma, diabetes mellitus type 1 or 2, cancer, chronic obstructive pulmonary disease, ischemic heart disease, heart failure, human immunodeficiency virus and previous transplantation (none, one, two or more). **Cardiovascular diseases comprised the presence of: ischemic heart disease, heart failure or hypertension. (yr) years.

Table 1. Demographic and clinical characteristics of included subjects. (IQR) interquartile range. (yr) years. (BMI) body mass index. (COPD) chronic obstructive pulmonary disease. (ED) emergency department. (ICU) intensive care unit.

Characteristics	Participants at inclusion (n = 310)	Losses (n = 13)	Participants that needed hospitalization (n = 23)	Participants without hospitalization (n = 274)	P	Hospitalized obese participants (n = 15)
Age, median (IQR)	37.4 (29.8-45.0)	44.2 (33.2-47.0)	42.0 (36.5-51.7)	36.8 (29.2-44.0)	**0.005***	42.0 (37.2-51.7)
Female sex, n (%)	186 (60.0)	7 (53.8)	12 (52.2)	167 (60.9)	0.55†	6 (40.0)
Active or second hand smoker at home, n (%)	55 (17.7)	NI	4 (17.4)	51 (18.6)	1.00‡	1 (6.7)
Years of education						
11 yr or less, n (%)	156 (50.3)	NI	13 (56.5)	143 (52.2)	0.85†	8 (53.3)
12 yr or more, n (%)	141 (45.5)	NI	10 (43.5)	131 (47.8)		7 (46.7)
Nutritional status						
Non-obesity (BMI<30.0 kg/m^2), n (%)	201 (64.8)	10 (76.9)	5 (21.7)	**186 (67.9)**	**<0.001†**	-
Obesity (BMI≥30.0 kg/m^2), n (%)	98 (31.6)	3 (23.1)	**15 (65.2)**	80 (29.2)		15 (100.0)
Hospital system at inclusion						
Private hospital, n (%)	137 (44.2)	2 (15.4)	11 (47.8)	124 (45.3)	0.98†	8 (53.3)
Public hospital, n (%)	173 (55.8)	11 (84.6)	12 (52.2)	150 (54.7)		7 (46.7)
Racial or ethnic group						

medRxiv preprint doi: https://doi.org/10.1101/2021.08.04.21261538; this version posted August 5, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

Caucasian, n (%)	222 (71.6)	NI	16 (69.6)	206 (75.2)	0.73†	11 (73.3)
Non-caucasian, n (%)	75 (24.2)	NI	7 (30.4)	68 (24.8)		4 (26.7)
Duration of symptoms at inclusion						
Days, median (IQR)	3.0 (2.0-5.0)	3.0 (2.0-5.0)	3.0 (2.0-3.0)	3.0 (2.0-5.0)	0.30*	2.0 (2.0-3.0)
Number of symptoms						
2-8, n (%)	95 (30.6)	1 (7.7)	8 (34.8)	86 (31.4)	0.69†	5 (33.3)
9-10, n (%)	82 (26.5)	3 (23.1)	7 (30.4)	72 (26.3)		5 (33.3)
11-12, n (%)	74 (23.9)	5 (38.5)	3 (13.0)	66 (24.1)		2 (13.3)
≥13, n (%)	59 (19.0)	4 (30.8)	5 (21.7)	50 (18.2)		3 (20.0)
Underlying medical conditions						
Hypertension, n (%)	39 (12.6)	1 (7.7)	5 (21.7)	33 (12.0)	0.21‡	3 (20.0)
Asthma, n (%)	19 (6.1)	1 (7.7)	1 (4.3)	17 (6.2)	1.00‡	1 (6.7)
Diabetes mellitus type 1 or 2, n (%)	11 (3.5)	2 (15.4)	1 (4.3)	8 (2.9)	0.58‡	1 (6.7)
Cancer, n (%)	5 (1.6)	0 (0.0)	0 (0.0)	5 (1.8)	1.00‡	0 (0.0)
COPD, n (%)	4 (1.3)	0 (0.0)	1 (4.3)	3 (1.1)	0.31‡	1 (6.7)
Ischemic heart disease, n (%)	3 (1.0)	0 (0.0)	1 (4.3)	2 (0.7)	0.24‡	1 (6.7)
Heart failure, n (%)	1 (0.3)	0 (0.0)	0 (0.0)	5 (1.8)	1.00‡	0 (0.0)


medRxiv preprint doi: https://doi.org/10.1101/2021.08.04.21261538; this version posted August 5, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
It is made available under a CC-BY-NC-ND 4.0 International license .


HIV, n (%)	1 (0.3)	0 (0.0)	0 (0.0)	1 (0.4)	1.00‡	0 (0.0)
Previous transplantation, n (%)	1 (0.3)	0 (0.0)	0 (0.0)	1 (0.4)	1.00‡	0 (0.0)
Previous chemotherapy, n (%)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	-	0 (0.0)
Cardiovascular diseases**, n (%)	39 (12.6)	1 (7.7)	5 (21.7)	33 (12.0)	0.21‡	1 (6.7)
Number of medical conditions						
None, n (%)	243 (78.4)	10 (76.9)	17 (73.9)	216 (78.8)	0.63†	11 (73.3)
One, n (%)	52 (16.8)	2 (15.4)	4 (17.4)	46 (16.8)		2 (13.3)
Two or more, n (%)	15 (4.8)	1 (7.7)	2 (8.7)	12 (4.4)		2 (13.3)
Hospitalization						
Yes, n (%)	23 (7.4)	-	-	-	-	-
Days after onset of symptoms to hospitalization, median (IQR)	8.0 (7.0-9.5)	-	8.0 (7.0-9.8)	-	-	8.0 (7.0-9.0)
Days from the ED sought to hospitalization, median (IQR)	5.0 (4.0-7.0)	-	5.0 (4.0-7.0)	-	-	5.5 (4.0-7.0)
Use of supplemental oxygen, n (%)	15 (4.8)	-	15 (65.2)	-	-	10 (66.7)
Admission at ICU, n (%)	6 (1.9)	-	6 (26.1)	-	-	4 (26.7)
Use of invasive mechanical ventilation, n (%)	2 (0.6)	-	2 (8.7)	-	-	1 (6.7)
Deaths, n (%)	1 (0.3)	-	1 (4.3)	-	-	1 (6.7)

medRxiv preprint doi: https://doi.org/10.1101/2021.08.04.21261538; this version posted August 5, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

**Cardiovascular diseases comprised: ischemic heart disease, heart failure and hypertension		NI = not informed		* Mann-Whitney-Wilcoxon test; † Pearson's Chi-squared test; ‡ Fisher's exact test		

medRxiv preprint doi: https://doi.org/10.1101/2021.08.04.21261538; this version posted August 5, 2021. The copyright holder for this preprint **(which was not certified by peer review)** is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
It is made available under a CC-BY-NC-ND 4.0 International license .


1,050 screened participants
13 excluded
9 ineligible
2 did not consent
1 withdrew consent
1 without RT-PCR SARS-CoV-2 result
727 negative RT-PCR results were removed from analysis
310 included participants
13 follow-up losses
297 analysed


Characteristics
OR
95% CI
P
Model 1
Age, yr
1.05
1.01-1.09
0.007
Obesity
2.69
1.63-4.83
<0.001
Model 2
Age, yr
1.05
1.01-1.09
0.016
Obesity
2.93
1.74-5.35
<0.001
Public hospital at inclusion
0.49
0.17-1.34
0.166
Model 3
Age, yr
1.05
1.01-1.09
0.007
Obesity
2.86
1.70-5.22
<0.001
12 yr of education or more
1.65
0.61-4.54
0.326
Model 4
Age, yr
1.07
1.02-1.12
0.005
Obesity
2.81
1.69-5.09
<0.001
*Number of medical conditions
0.69
0.21-1.45
0.287
Model 5
Age, yr
1.06
1.01-1.11
0.008
Obesity
2.74
1.65-4.94
<0.001
**Cardiovascular diseases
0.66
0.13-2.50
0.565
Model 6
Age, yr
1.05
1.02-1.10
0.007
Obesity
2.6
1.57-4.67
<0.001
Days of onset of symptoms to inclusion
0.86
0.65-1.07
0.215
Model 7
Age, yr
1.05
1.01-1.09
0.011
Obesity
2.79
1.68-5.03
<0.001
Number of symptoms
0.8
0.51-1.24
0.333
0 0.5 1 1.5 2 2.5 3 3.5 4 4.5 5 5.5