

Quantifying contact patterns: development and characteristics of the British Columbia COVID-19 population mixing patterns survey (BC-Mix)

Prince A. Adu^{1,2}, Mawuena Binka¹, Bushra Mahmood,³ Dahn Jeong^{1,2}, Terri Buller-Taylor¹, Makuza Jean Damascene^{1,2}, Sarafa Iyaniwura^{1,4}, Notice Ringa^{1,2}, Héctor A. Velásquez García^{1,2}, Stanley Wong¹, Amanda Yu¹, Sofia Bartlett^{1,6}, James Wilton¹, Mike A. Irvine^{1,7}, Michael Otterstatter^{1,2}, Naveed Z. Janjua^{1,2}

¹British Columbia Centre for Disease Control, Vancouver, British Columbia, Canada

²School of Population and Public Health, University of British Columbia, Vancouver, British Columbia, Canada

³Department of Medicine, University of British Columbia, Vancouver, British Columbia, Canada

⁴Department of Mathematics and Institute of Applied Mathematics, University of British Columbia, Vancouver, British Columbia, Canada

⁵Faculty of Health Sciences, Simon Fraser University, Burnaby, British Columbia, Canada

⁶Department of Pathology and Laboratory Medicine, Vancouver, British Columbia, Canada

⁷Faculty of Health Sciences, Simon Fraser University, Burnaby, British Columbia, Canada

Corresponding author:

Naveed Zafar Janjua MBBS, MSc, DrPH
655 W 12th Ave, Vancouver, BC V5Z 4R4
Naveed.janjua@bccdc.ca

ABSTRACT

Introduction

Several non-pharmaceutical interventions such as physical distancing, hand washing, self-isolation, and schools and business closures, were implemented in British Columbia (BC) following the first laboratory-confirmed case of coronavirus disease 2019 (COVID-19) on January 26, 2020, to minimize in-person contacts that could spread infections. The BC COVID-19 Population Mixing Patterns survey (BC-Mix) was established as a surveillance system to measure behaviour and contact patterns in BC over time to inform the timing of the easing/re-imposition of control measures. In this paper, we describe the BC-Mix survey design and the demographic characteristics of respondents.

Methods

The ongoing repeated online survey was launched in September 2020. Participants are mainly recruited through social media platforms (including Instagram, Facebook, YouTube, WhatsApp). A follow up survey is sent to participants two to four weeks after completing the baseline survey. Survey responses are weighted to BC's population by age, sex, geography, and ethnicity to obtain generalizable estimates. Additional indices such as the material and social deprivation index, residential instability, economic dependency, and others are generated using census and location data.

Results

As of July 26, 2021, over 61,000 baseline survey responses were received of which 41,375 were eligible for analysis. Of the eligible participants, about 60% consented to follow up and about 27% provided their personal health numbers for linkage with healthcare databases. Approximately 50% of respondents were female, 39% were 55 years or older, 65% identified as white and 50% had at least a university degree.

Conclusion

The pandemic response is best informed by surveillance systems capable of timely assessment of behaviour patterns. BC-Mix survey respondents represent a large cohort of British Columbians providing near real-time information on behavioural and contact patterns in BC. Data from the BC-Mix survey would inform provincial COVID-19-related control measures.

Introduction

The novel coronavirus disease 2019 (COVID-19), caused by the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), has spread worldwide since December 2019. A global pandemic was declared by the World Health Organization in March 2020 and, as of July 2021, there have been over 200 million cases of COVID-19 infections and over 4.3 million resultant deaths globally (1). As vaccine rollouts continue at varying rates worldwide, physical distancing measures (2) remain among the most effective methods for COVID-19 prevention and control (3). Many governments have put in place physical distancing measures such as travel restrictions, closure of schools and workplaces, and the banning of large group gatherings to interrupt the transmission of SARS-CoV-2. These measures attempt to reduce contact between infected and healthy individuals in order to minimize disease spread and the impact on the healthcare system.

British Columbia (BC) is located on the West Coast of Canada and covers almost a million square kilometres. It has a diverse population of approximately 5.15 million as of July 1, 2020 (4). Public health officials in BC began urging the public to practice physical distancing and avoid any non-essential travel in early March 2020. By March 17, 2020, a public health emergency was declared in the province and various physical distancing measures were implemented (5). These included restriction of indoor and outdoor gatherings, closure of businesses that were unable to meet physical distancing measures, self-isolation requirements after travelling outside the country, and general physical distancing in all public space. While these measures were important for controlling the rapid spread of disease, they also had sweeping economic, social, and mental health impacts.

Assessing the impact of physical distancing measures on person-to-person contact can provide valuable information for refining control measures and help minimize both COVID-19-related disease burden and the related economic, social, and mental health impacts. Early detection of COVID-19 resurgences requires mechanisms for tracking precursors of transmission, including changes in social contacts, mixing patterns and physical distancing behaviours as well as early signals of a COVID-19 spread. Although methods such as mathematical modelling can estimate the potential for resurgences, these methods often lack population-based empirical data on contact patterns, especially on the varying levels of contact patterns exhibited by different demographic groups in the population. These population-specific data could better inform mathematical models by incorporating explicit knowledge of contact patterns that are driving transmission rather than inferring these from reported cases and hospitalizations (6,6). Ultimately, they serve as an evidence-base to guide targeted measures that are amenable to actions by the government to ensure that the COVID-19 cases remain below the resurgence thresholds.

Various studies have assessed the impact of physical distancing measures imposed by governments on local contact patterns and behaviours during the COVID-19 pandemic in

Belgium (7), Greece (8), Kenya (9), Luxembourg (10), the Netherlands (11), and the U.K (12). Such surveys can measure the public's compliance with the physical distancing measures and provide valuable information to inform other public health measures that may be necessary to avoid further waves of COVID-19 infections. In addition, the impact of physical distancing measures on mixing patterns and contact behaviours may vary across different age groups, and by individuals' primary place of activity such as schools or workplaces (8,13–15).

Here, we describe the development the BC COVID-19 Population Mixing Patterns survey (BC-Mix), an ongoing online survey to monitor and assess social contact behaviours and mixing patterns in BC, Canada, during the COVID-19 pandemic. We detail the development of the survey and recruitment of respondents, as well as the characteristics of the participants.

Methods and analysis

Survey design and methodology

The BC-Mix (<http://www.bccdc.ca/our-research/projects/bc-mix-covid-19-survey>) uses a cross-sectional survey design with longitudinal follow-up. Eligible population include residents of BC who are at least 18 years of age. The survey began on September 4, 2021, and as of August 2021 is ongoing. Once a participant has completed the survey for the first time, they are invited for repeated follow-up. The first-time responses are referred to as the 'baseline'. Participants responding to the baseline survey are invited to complete the first follow-up survey after two weeks. Subsequent follow-up surveys are then sent in four-week intervals, following the completion of the previous survey.

Participant recruitment

To capture participants from a broad demographic range, the survey invitation and survey are disseminated through Instagram, Facebook, YouTube, WhatsApp, Twitter, and Google search engine results pages. The Google Ads Audience manager and Facebook Ads manager allow for paid advertisements to be targeted at specific audiences. We use these tools to target the survey advertisement campaigns to only residents of BC who are 18 years and above. We also monitor the demographic profile of survey participants and occasionally use these functions to target recruitment to age groups or sex that may be under-represented (16).

To help capture underrepresented groups, we promote the survey to various ethnic populations. For instance, a South Asian community organization promotes the survey on their social media pages and also sends the survey to individuals on their mailing list. Although the survey is in English, it is also promoted in different languages (specifically, Korean and Farsi) to members of minority community groups in BC on their social media pages. Flyers are also distributed at grocery stores and restaurants particularly including those frequented by minority groups.

Participant and public involvement

The initial version of the BC-Mix survey was first piloted with a randomly selected sample of the BC population and feedback received was incorporated in the final version before the official launch. Methods of recruitment and priority of research questions were also informed by discussions with members of the public and with a community group. We also receive input from survey participants on an ongoing basis through a dedicated e-mail address. We plan to create dashboards and other infographics of the study results on the study website. A newsletter suitable for non-specialist audience will also be sent to participants.

Survey domain and case definitions

The BC-Mix survey instrument was adapted from the POLYMOD study (14) and the Berkeley Interpersonal Contact Study [BICS] (17) and was administered through Qualtrics (18), an online survey tool. The baseline survey comprises 94 questions across six key domains:

1. **Demographic information:** This domain includes age, sex, gender, ethnicity, education, employment, household characteristics, and postal code.
2. **COVID-19 testing and results, symptoms, and health behaviours:** This domain captures COVID-19 testing information, symptoms, and behaviours such as doctor visits following symptoms.
3. **Activities and behaviour in and outside of the home:** This domain captures social contact and mixing behaviours such as number of contacts, location, and duration of contact during the past 24 hours. Other questions in this domain include age and sex of contact, and relationship of respondent to the contact persons, physical distancing behaviour (e.g., handwashing) and personal protective equipment use. Initially, respondents were asked to provide this information for up to three of their reported contacts. We began collecting data for up to 10 contacts from December 11, 2020. Also from December 11, 2020, we began collecting general information about greater than 10 contacts i.e., if a participant reports more than 10 contacts per day, they are asked general questions about these contacts for e.g., age group, duration, and location of the majority of those contacts. If majority of contacts took place at a workplace setting, a follow up question asks respondents to report the type of work setting where the contacts occurred.
4. **Internet and social media use:** This domain captures information on internet and social media use in terms of most frequently used platform and frequency of use.
5. **Perceptions and attitudes around COVID-19:** This domain measures the respondent's perception of the physical distancing measures, and their self-confidence or ability to carry out them.
6. **COVID-19 vaccine acceptance sub-questionnaire:** This sub-questionnaire was added on March 8, 2021. Items from this domain were developed using a vaccine acceptance behavioral framework, which synthesizes constructs from the Theory of Reasoned Action (TRA)(19), Theory of Planned Behavior (TPB)(20,21) and the Health Belief Model (HBM) (22), to understand and predict the uptake of COVID-19 vaccine. According to the TRA, the best single predictor of behaviour is an individual's intention (23). Intentions, in turn, are an

outcome of the individual's attitude toward performing the behavior in question, and/or the individual's perceptions of support from family and friends (subjective norms) for engaging in the behavior (24). Perceived control or self-efficacy, the confidence that one has the ability to perform the intended behavior (25), is another important construct taken from TPB. The TPB assumes that an individual's perception of whether they can successfully engage in a particular behavior often has a direct effect on their intentions, such as getting a vaccine (26). The widely-used HBM, has previously been used to evaluate beliefs and attitudes toward seasonal influenza and pandemic swine flu vaccines as well as the COVID-19 vaccine (27–29). Relevant constructs from HBM were applied to develop questionnaire items to assess perceived threat of contracting the COVID-19, perceived severity of disease if infected and belief in the safety and effectiveness of getting the vaccine. Overall, this sub-questionnaire is meant to provide an understanding of some of the individual level health beliefs, perceptions and attitudes that may influence vaccine uptake. The vaccine acceptance sub-questionnaire has the following the domains: *Attitude (perceived susceptibility, severity, benefits and barriers)*, *Descriptive and Subjective Norms*, *Perceived Control* and *Intention*.

Location data is used to generate other indicators at the area level. For example, the Quebec Material and Social Deprivation combines six indicators related to health and welfare that represent material or social deprivation based on Canadian Census data, including 1) proportion of persons without high school diploma 2) ratio of employment to population 3) average income 4) proportion of persons separated, divorced, widowed 5) proportion of single-parent families and 6) proportion of people living alone (30).

A full list of key variables in the survey and definitions is presented in S1 Table in the Supplementary file.

Analysis, data cleaning and weighting

A survey completion rate of at least 33% of questions, valid non-missing responses for the sex and age questions are required for inclusion for weighting the survey data and further analysis. All duplicates are removed.

To ensure that the BC-Mix sample is representative of the BC population, survey data are weighted to obtain generalizable estimates (Table 1). Using the 2016 Census data (31), the survey is weighted with the following auxiliary variables: age, sex, geography, and ethnicity using the weighting adjustment technique (32) in the following hierarchy: As our first criterion, we consider age, sex, geography and ethnicity as our auxiliary variables. If a record has valid responses for all these variables except the ethnicity variable, then the survey weight is generated using only age, sex, and geography (second criterion). If a record does not meet the first and second criteria, then we apply the third criterion which uses age, sex, and ethnicity as the auxiliary variables. Finally, we use only age and sex as auxiliary variables if a record does not satisfy the first three criteria.

Survey weights are estimated separately for baseline and for each follow-up. To assess participant profile, we computed un-weighted and weighted frequency and percentages of key demographic variables using SAS Software version 9.4. Baseline survey data was used to provide the survey participant profile and in comparison, with BC population profile (Table 1). To assess potential systematic differences between eligible and ineligible responses, a comparison of the baseline eligible participants versus ineligible participants is presented the S2 Table in the Supplementary file. Participant profile of follow up surveys is also presented in S3 Table in the Supplementary file.

Ethics

Informed consent was sought on the survey start page. The study was reviewed and approved by the University of British Columbia Behavioral Research Ethics Board (No: H20-01785).

Results

As of July 26, 2021, there were 61,183 respondents who participated in the baseline survey of which 41,375 were eligible for analysis. There were 15,194 (eligible=10,993) participants in the first follow-up survey, 11,343 (eligible n=8,164) in the second, 8,521 (eligible n=6,375) in the third, 6,487 (eligible n=4,981) in the fourth, 5,014 (eligible=3,891) in the fifth, 4,094 (eligible=3,184) in the sixth, 3,125 (eligible n= 2,417) in the seventh and 2,317 (eligible n=1,760) participants in the eighth follow-up survey (Fig. 1).

Considering the baseline sample (Table 1), there were approximately equal number of male and female (weighted % of female =50.0%). Majority of participants were 55 years or older (weighted %= 39.4%), self identified as White (weighted %= 64.6%), had at least a university degree (weighted %= 50.0%) and lived in the Fraser Health region (weighted %= 36.2%).

Almost 63.8% (unweighted n=20,633) consented to a follow-up after the baseline survey and at least 94.2% (unweighted n=10,357) consented to receiving subsequent follow-up surveys (Table 1 and S3 Table in the Supplementary File). Approximately 27.3% (unweighted n=7,290) of respondents in the baseline provided their personal health numbers for linkage with other healthcare utilization databases.

After weighting, the baseline survey sample is representative of the general BC population in terms of age, sex, health region, and ethnicity (Table 1). The distribution of the eligible participants was also similar to the distribution of ineligible participants in terms of sex, age, race/ethnicity and geography/health region (S2 Table in the Supplementary file).

Discussion

Following the identification of COVID-19 cases in BC, several interventions including physical distancing measures were implemented to limit the spread of COVID-19 in the province. Subsequently, the BC-Mix was developed by the BC Centre for Disease Control (BCCDC)(33) as part of an early warning system for monitoring social and physical interactions between individuals of different age-groups and demography, and to help predict when COVID-19 transmission might further increase. This paper describes BC-Mix survey methods and the profile of survey respondents.

Recent studies similar to the BC-Mix have assessed social contact patterns relevant to the spread and control of COVID-19 in different countries(7–12,34,35) many of which have adapted features of the POLYMOD project (14). The 2020 Belgian CoMix survey (7) is an online longitudinal survey that closely monitors changes in social mixing behaviours among a sample of Belgian adults (aged 18 years and above). The U.K CoMix survey assesses contact patterns of a representative sample of U.K adults. Launched on March 24, 2020, participants are followed up every 2 weeks to monitor changes in their self reported behaviours (12). In Canada, the Quebec-based CONNECT study uses population-based survey to assess social contacts and mixing patterns (34). Brankston and colleagues (35) also used paid panel representative of Canadian adults to construct contact patterns to determine the impact of physical distancing measures on COVID-19 transmission. Most of these studies commissioned market research companies or used survey panels to recruit participants (7,12,17,35). While market companies or survey panels offer a convenient approach to sampling, they have some challenges. Panels are made of membership in loyalty programs or other panels constituting a select group of population and may therefore not represent complete random recruitment from a population of interest.

The use of targeted social media advertisement for participant recruitment has gained prominence in health research (16,36), having been applied in areas such as mental health (37), cannabis use (38), smoking behaviour (39) and in other health related studies (40). For our survey, we use social media advertisement and other recruitment strategies. Although social media-based recruitment does not necessarily generate a random sample of the general population given the characteristics of people who are on social media may differ from those who are not, social media channels like Facebook, Instagram, Twitter, and others have powerful targeting capabilities that allow researchers to target advertisements to users with specific demographic characteristics. They also have the advantage of reaching hard-to-reach populations (37–39).

Quota sampling has been used by other studies to achieve representativeness (7,44). We used two approaches to achieve the same goal: adaptive recruitment through promotion and targeting to specific populations and then post hoc weighting. Our survey tool does not set quotas on recruitment but uses targeted advertisements to improve representativeness.

The following issues should be considered for interpretation of results from BC-Mix. Some population groups are underrepresented in the survey possibly due to lack of access to social media. These are people who are economically marginalised and less likely to have access to a computer/electronic device or to have access to the internet/cellular data, e.g., people living in poverty, people who are unemployed, people who are unhoused, etc. Also, people who are in prison (sentenced or on remand) or people who are under immigration detention may not have access to the internet or cellular devices. Our survey responses may be subject to recall bias since we ask respondents to recall contacts and other behaviours or activities from the previous day. Other studies have used diaries (14) to overcome this weakness but this may be logistically challenging and attrition with this method may be quite high. Another potential bias inherent in our survey is the issue of reporting bias, as respondents may respond in ways consistent with the laws around physical distancing. In addition, the BC-Mix is available only in English, thus excluding individuals who cannot communicate in English. This notwithstanding, according to the 2016 Census, 96.6% of BC's population indicated that they can converse in English (31). Therefore, we do not believe that any bias associated with language would be significant. Another limitation to mention is the large number of recruits that were ineligible and the attrition between successive rounds of survey. This could be related to survey fatigue, or the time required to complete the survey.

Our survey has several strengths. Web-based surveys like the BC-Mix provide timely information for pandemic response (45). Also, during an infectious disease pandemic, web-based surveys offer a more convenient approach to data collection compared to in-person or other modes of data collection. We also found paid advertisements to be more cost effective compared to the cost of panel data from survey companies (36). An additional strength of our study is its large sample size. Our total recruited sample of over 61,000 participants compares to the 1,356 participants in the U.K CoMix study (12), the 9,743 participants in the BICS study (17) study, 1,542 participants in the Belgian CoMix study (7) and the 7,290 participants in the POLYMOD study (14). In addition, because we opted to achieve representativeness post-data collection (at the analysis stage), we were able to consider many important variables besides age and sex in our weighting strategy. It would have been logistically challenging to consider all these variables had we used quota-sampling given that many market research company panels were limited in terms recruitment by age, sex, and geography. Using many auxiliary variables in our weighting strategy increased the representativeness of the BC population.

Conclusion

To our knowledge, the BC-Mix is the first and largest surveillance tool providing real time quantitative data on mixing patterns and contact characteristics in BC and one of the largest in North America. Tools such as the BC-Mix are integral to the COVID-19 pandemic response to provide critical data to inform the timing of loosening or re-imposition of physical distancing measures. Further analyses on contact patterns, relationship of contact patterns with

transmission, disparities in contact patterns, facemask use, are in progress and will be published soon.

Supporting information

S1 Table. BC-Mix variable names and definitions

S2 Table. Comparison of baseline eligible and ineligible participants, frequencies and proportions

S3 Table. Participant profile of BC-Mix follow up surveys: frequencies and proportions (%)

Authors' contributions

Conceptualization: Naveed Z. Janjua, Prince A. Adu

Survey design: Naveed Z. Janjua, Prince A. Adu, Terri Buller-Taylor, Bushra Mahmood

Data curation: Prince A. Adu, Amanda Yu, Stanley Wong

Statistical analysis: Prince A. Adu

Funding acquisition: Naveed Janjua

Methodology: Naveed Z. Janjua, Prince A. Adu, Mawuena Binka, Terri Buller-Taylor, Bushra Mahmood, Sarafa Iyaniwura, Michael Otterstatter

Writing first draft: Prince A. Adu, Dahn Jeong, Mawuena Binka, Terri Buller-Taylor, Sarafa Iyaniwura, Notice Ringa

Writing-review & editing: Naveed Z. Janjua, Héctor A. Velásquez García, Bushra Mahmood, Makuza Jean Damascene, Mawuena Binka, James Wilton, Sofia Bartlett, Michael Otterstatter, Mike Irvine.

Acknowledgement

The authors will like to express their gratitude to Mei Wong and Dr Joan Hu for their methodological guidance.

Funding statement

This work was supported by Michael Smith Foundation for Health Research COVID-19 Research Response Fund (Award #: COV-2020-1183)

Competing interests statement

None declared

Word Count

3,200 words

References

1. Johns Hopkins Coronavirus Resource Center. COVID-19 Map [Internet]. Johns Hopkins Coronavirus Resource Center. [cited 2021 May 23]. Available from: <https://coronavirus.jhu.edu/map.html>
2. Lu H, Stratton CW, Tang Y. Outbreak of pneumonia of unknown etiology in Wuhan, China: The mystery and the miracle. *J Med Virol*. 2020 Apr;92(4):401–2.
3. Islam N, Sharp SJ, Chowell G, Shabnam S, Kawachi I, Lacey B, et al. Physical distancing interventions and incidence of coronavirus disease 2019: natural experiment in 149 countries. *BMJ*. 2020 Jul 15;370:m2743.
4. BC Stats. 2020 Sub-Provincial Population Estimates Highlights. 2020 Jul;4.
5. Joint statement on Province of B.C.'s COVID-19 response, latest updates |BC Gov News [Internet]. 2020 [cited 2021 May 23]. Available from: <https://news.gov.bc.ca/releases/2020HLTH0089-000505>
6. Ferguson N, Laydon DJ, Nedjati Gilani G, Imai N, Ainslie KM, Baguelin M, et al. Report 9: Impact of non-pharmaceutical interventions (NPIs) to reduce COVID19 mortality and healthcare demand. 2020 Jan 1 [cited 2021 Jul 19]; Available from: <https://www.scienceopen.com/document?vid=4e6290c9-0ded-40ed-b858-ba18119863c0>
7. Coletti P, Wambua J, Gimma A, Willem L, Vercruyssen S, Vanhoutte B, et al. CoMix: comparing mixing patterns in the Belgian population during and after lockdown. *Sci Rep*. 2020 Dec 14;10(1):21885.
8. Sypsa V, Roussos S, Paraskevis D, Lytras T, Tsiodras S, Hatzakis A. Effects of Social Distancing Measures during the First Epidemic Wave of Severe Acute Respiratory Syndrome Infection, Greece. *Emerg Infect Dis*. 2021 Feb;27(2):452–62.
9. Quaife M, van Zandvoort K, Gimma A, Shah K, McCreesh N, Prem K, et al. The impact of COVID-19 control measures on social contacts and transmission in Kenyan informal settlements. *BMC Med*. 2020 Oct 5;18(1):316.
10. Latsuzbaia A, Herold M, Bertemes J-P, Mossong J. Evolving social contact patterns during the COVID-19 crisis in Luxembourg. *PLOS ONE*. 2020 Aug 6;15(8):e0237128.
11. Backer JA, Mollema L, Vos RAE, Klinkenberg D, Klis FRM van der, Melker HE de, et al. The impact of physical distancing measures against COVID-19 transmission on contacts and mixing patterns in the Netherlands: repeated cross-sectional surveys in 2016/2017, April 2020 and June 2020. *medRxiv*. 2020 Oct 16;2020.05.18.20101501.
12. Jarvis CI, Van Zandvoort K, Gimma A, Prem K, Auzenberg M, O'Reilly K, et al. Quantifying the impact of physical distance measures on the transmission of COVID-19 in the UK. *BMC Med*. 2020 May 7;18(1):124.

13. Davies NG, Klepac P, Liu Y, Prem K, Jit M, Eggo RM. Age-dependent effects in the transmission and control of COVID-19 epidemics. *Nat Med*. 2020 Aug;26(8):1205–11.
14. Mossong J, Hens N, Jit M, Beutels P, Auranen K, Mikolajczyk R, et al. Social Contacts and Mixing Patterns Relevant to the Spread of Infectious Diseases. *PLOS Med*. 2008 Mar 25;5(3):e74.
15. Wallinga J, Teunis P, Kretzschmar M. Using data on social contacts to estimate age-specific transmission parameters for respiratory-spread infectious agents. *Am J Epidemiol*. 2006 Nov 15;164(10):936–44.
16. Grow A, Perrotta D, Fava ED, Cimentada J, Rampazzo F, Gil-Clavel S, et al. Addressing Public Health Emergencies via Facebook Surveys: Advantages, Challenges, and Practical Considerations. *J Med Internet Res*. 2020 Dec 14;22(12):e20653.
17. Feehan DM, Mahmud AS. Quantifying population contact patterns in the United States during the COVID-19 pandemic. *Nat Commun*. 2021 Feb 9;12(1):893.
18. Qualtrics. Qualtrics [Internet]. Provo, Utah, USA: Qualtrics; 2021. Available from: <https://www.qualtrics.com>
19. Ajzen I, Fishbein M. Understanding attitudes and predicting social behavior Prentice-Hall Inc. Englewood Cliffs, NJ; 1980.
20. Ajzen I. From Intentions to Actions: A Theory of Planned Behavior. In: Kuhl J, Beckmann J, editors. *Action Control: From Cognition to Behavior* [Internet]. Berlin, Heidelberg: Springer; 1985 [cited 2021 Jun 15]. p. 11–39. (SSSP Springer Series in Social Psychology). Available from: https://doi.org/10.1007/978-3-642-69746-3_2
21. Ajzen I. The theory of planned behavior. *Organ Behav Hum Decis Process*. 1991 Dec 1;50(2):179–211.
22. Becker MH. The health belief model and personal health behavior. *Health Educ Monogr*. 1974;2:324–473.
23. Fishbein M, Ajzen I. *Belief, Attitude, Intention, and Behavior: An Introduction to Theory and Research*. Reading, MA: Addison-Wesley; 1975.
24. Sheppard BH, Hartwick J, Warshaw PR. The Theory of Reasoned Action: A Meta-Analysis of Past Research with Recommendations for Modifications and Future Research. *J Consum Res*. 1988 Dec 1;15(3):325–43.
25. Bogoch II, Watts A, Thomas-Bachli A, Huber C, Kraemer MUG, Khan K. Pneumonia of unknown aetiology in Wuhan, China: potential for international spread via commercial air travel. *J Travel Med*. 2020 Mar 13;27(2).
26. Fisher WA, Kohut T, Salisbury CMA, Salvadori MI. Understanding human papillomavirus vaccination intentions: comparative utility of the theory of reasoned action and the theory

- of planned behavior in vaccine target age women and men. *J Sex Med.* 2013 Oct;10(10):2455–64.
27. Santos AJ, Kislaya I, Machado A, Nunes B. Beliefs and attitudes towards the influenza vaccine in high-risk individuals. *Epidemiol Infect.* 2017 Jul;145(9):1786–96.
 28. Teitler-Regev S, Shahrabani S, Benzion U. Factors Affecting Intention among Students to Be Vaccinated against A/H1N1 Influenza: A Health Belief Model Approach. *Adv Prev Med.* 2011 Dec 20;2011:e353207.
 29. Wong MCS, Wong ELY, Huang J, Cheung AWL, Law K, Chong MKC, et al. Acceptance of the COVID-19 vaccine based on the health belief model: A population-based survey in Hong Kong. *Vaccine.* 2021 Feb 12;39(7):1148–56.
 30. Bureau d'information et d'études En Santé Des Populations (BIESP). Deprivation Index [Internet]. INSPQ. 2019 [cited 2021 May 27]. Available from: <https://www.inspq.qc.ca/en/deprivation/material-and-social-deprivation-index>
 31. Government of Canada SC. Census Profile, 2016 Census - British Columbia [Province] and Canada [Country] [Internet]. 2017 [cited 2021 Jul 25]. Available from: <https://www12.statcan.gc.ca/census-recensement/2016/dp-pd/prof/details/page.cfm?Lang=E&Geo1=PR&Code1=59&Geo2=PR&Code2=01&SearchText=Canada&SearchType=Begins&SearchPR=01&B1=All&type=0>
 32. Bethlehem JG. *Applied survey methods: a statistical perspective.* Hoboken, N.J: Wiley; 2009. 375 p. (Wiley series in survey methodology).
 33. BC Centre for Disease Control. What We Do [Internet]. [cited 2021 Jun 10]. Available from: <http://www.bccdc.ca/about/what-we-do>
 34. Institut national de, santé publique du Québec. CONNECT□: étude des contacts sociaux des Québécois - 27 avril 2021 [Internet]. INSPQ. [cited 2021 Jun 17]. Available from: <https://www.inspq.qc.ca/covid-19/donnees/connect/27-avril-2021>
 35. Brankston G, Merkley E, Fisman DN, Tuite AR, Poljak Z, Loewen PJ, et al. Quantifying Contact Patterns in Response to COVID-19 Public Health Measures in Canada. *medRxiv.* 2021 Mar 12;2021.03.11.21253301.
 36. Thornton L, Batterham PJ, Fassnacht DB, Kay-Lambkin F, Calcar AL, Hunt S. Recruiting for health, medical or psychosocial research using Facebook: Systematic review. *Internet Interv.* 2016 May 1;4:72–81.
 37. Choi I, Milne DN, Glozier N, Peters D, Harvey SB, Calvo RA. Using different Facebook advertisements to recruit men for an online mental health study: Engagement and selection bias. *Internet Interv.* 2017 Jun 1;8:27–34.

38. Borodovsky JT, Marsch LA, Budney AJ. Studying Cannabis Use Behaviors With Facebook and Web Surveys: Methods and Insights. *JMIR Public Health Surveill.* 2018 May 2;4(2):e9408.
39. Ramo DE, Prochaska JJ. Broad Reach and Targeted Recruitment Using Facebook for an Online Survey of Young Adult Substance Use. *J Med Internet Res.* 2012 Feb 23;14(1):e1878.
40. Fazzino TL, Rose GL, Pollack SM, Helzer JE. Recruiting U.S. and Canadian College Students via Social Media for Participation in a Web-Based Brief Intervention Study. *J Stud Alcohol Drugs.* 2015 Jan 1;76(1):127–32.
41. Guillory J, Kim A, Murphy J, Bradfield B, Nonnemaker J, Hsieh Y. Comparing Twitter and Online Panels for Survey Recruitment of E-Cigarette Users and Smokers. *J Med Internet Res [Internet].* 2016 Nov 15 [cited 2021 Jun 16];18(11). Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5128722/>
42. Walsh DC, Rudd RE, Moeykens BA, Moloney TW. Social marketing for public health. *Health Aff Proj Hope.* 1993;12(2):104–19.
43. Freimuth VS, Mettger W. Is there a hard-to-reach audience? *Public Health Rep Wash DC* 1974. 1990 Jun;105(3):232–8.
44. Leung K, Jit M, Lau EHY, Wu JT. Social contact patterns relevant to the spread of respiratory infectious diseases in Hong Kong. *Sci Rep.* 2017 Aug 11;7(1):7974.
45. Hlatshwako TG, Shah SJ, Kosana P, Adebayo E, Hendriks J, Larsson EC, et al. Online health survey research during COVID-19. *Lancet Digit Health.* 2021 Feb 1;3(2):e76–7.

Table 1. Participant profile of BC-Mix baseline data (n=41,375), September 04, 2020-July 26, 2021

		Survey				British Columbia population	
		Un-weighted frequency	Un-weighted % (excl. missing)	Weighted frequency	Weighted %	Population frequency	Population %
Sex							
	Male	6,823	16.5	21,293	50.0	1,805,105	48.5
	Female	34,552	83.5	21,261	50.0	1,914,755	51.5
	Missing						
Age							
	18-34	4,978	12.0	11,575	27.2	1,002,745	27
	35-54	12,110	29.3	14,194	33.4	1,251,835	33.7
	55+	24,287	58.7	16,784	39.4	1,465,280	39.4
Race/ethnicity							
	Indigenous	1,757	4.4	2,180	5.3	186,705	5
	Chinese	882	2.2	4,451	10.9	418,035	11.2
	White	35,026	87.5	26,383	64.6	2,448,155	65.8
	South Asian	606	1.5	3,473	8.5	280,470	7.5
	Other	1,766	4.4	4,352	10.7	386,495	10.4
	Missing/Unknown	1,338	n/a	1715	n/a	n/a	n/a
Health region							
	Fraser Health	8,451	26.1	11,793	36.2	1,347,410	36.2
	Interior Health	6,143	19.0	5,336	16.4	595,105	16
	Northern Island	1,825	5.6	1,828	5.6	213,235	5.7
	Vancouver Coastal	7,315	22.6	8,118	24.9	934,055	25.1
	Vancouver Island	8,640	26.7	5,535	17.0	630,055	16.9
	Missing/Unknown	9,001	n/a	9,943	n/a	n/a	n/a
Education							
	Below high school	807	2.5	1,096	3.0	2,301,030	12.5
	Below bachelor	16,928	51.7	15,176	47.0	466,295	61.9
	University degree	15,029	45.9	16,273	50.0	952,535	25.6
	Missing/Unknown	8,611	n/a	10,009	n/a	n/a	n/a
Employment status							
	Employed full-time (30 hours or more/week)	10,654	32.0	13,608	40.8	n/a	n/a
	Employed part-time	2,993	9.0	3,131	9.4	n/a	n/a

	Survey				British Columbia population	
	Un-weighted frequency	Un-weighted % (excl. missing)	Weighted frequency	Weighted %	Population frequency	Populatio %
Self-employed	2,704	8.1	3,013	9.0	n/a	n/a
Unemployed but looking for a job	952	2.9	1,522	4.6	n/a	n/a
Unemployed and not looking for a job	406	1.2	510	1.5	n/a	n/a
Full-time parent, homemaker	879	2.6	740	2.2	n/a	n/a
Retired	12,757	38.3	8,096	24.3	n/a	n/a
Student/Pupil	566	1.7	1,197	3.6	n/a	n/a
Long-term sick or disabled	968	2.9	914	2.7	n/a	n/a
Prefer not to answer	424	1.3	619	1.9	n/a	n/a
Missing/Unknown	8,072	n/a	n/a	n/a	n/a	n/a
Material Deprivation Index						
1 (Privileged)	6,407	22.3	6,100	21.8	n/a	n/a
2	6,475	22.5	5,873	21.1	n/a	n/a
3	6,972	24.2	6,010	21.6	n/a	n/a
4	4,822	16.8	5,187	18.7	n/a	n/a
5 (Deprived)	4,085	14.2	4,656	16.8	n/a	n/a
Missing	1,2614	n/a	n/a	n/a	n/a	n/a
Social Deprivation Index						
1 (Privileged)	4,932	17.2	5,199	17.9	n/a	n/a
2	4,756	16.5	4,752	16.4	n/a	n/a
3	6,311	21.9	6,022	20.8	n/a	n/a
4	5,932	20.6	6,054	20.9	n/a	n/a
5 (Deprived)	6,830	23.8	6,957	24.0	n/a	n/a
Missing	1,2614	n/a	n/a	n/a	n/a	n/a
Follow up consent						
Yes	20,633	63.8	19,051	58.9	n/a	n/a
No	11,689	36.2	13,275	41.1	n/a	n/a
Missing	9,053	n/a	n/a	n/a	n/a	n/a
Data linkage consent						
Yes	7,290	27.3	7,318	26.4	n/a	n/a

	Survey				British Columbia population	
	Un-weighted frequency	Un-weighted % (excl. missing)	Weighted frequency	Weighted %	Population frequency	Population %
No	19,467	72.8	20,362	73.6	n/a	n/a
Missing	14,618	n/a	n/a	n/a	n/a	n/a




Fig. 1. Participant flowchart