

1 **Monitoring socioeconomic inequalities across HIV knowledge, attitudes,** 2 **behaviours and prevention in 18 sub-Saharan African countries**

3
4 *Mohamed Hamidouche^{1,2}, Pearl Anne Ante-Testard^{1,2}, Rachel Baggaley³, Laura Temime^{1,2}, Kévin*
5 *Jean^{1,2,4*}*

6
7 ¹Laboratoire MESuRS, Conservatoire national des Arts et Métiers, Paris, France ;
8 ²Unité PACRI, Institut Pasteur, Conservatoire national des Arts et Métiers, Paris, France ;
9 ³World Health Organization, Global HIV, Hepatitis and STI Programmes, Geneva, Switzerland
10 ⁴MRC Centre for Global Infectious Disease Analysis, Department of Infectious Disease Epidemiology,
11 Imperial College London, London W2 1PG, United Kingdom.

12
13
14 * Correspondance to:
15 Kévin Jean
16 Laboratoire MESuRS, Conservatoire National des Arts et Métiers, 292 rue Saint Martin, 75003, Paris, France
17 kevin.jean@lecnam.net

18
19 Running head: Monitoring inequalities in HIV prevention

20
21 Keywords: HIV; health inequalities; epidemiology; sub-Saharan Africa; HIV prevention;
22 social epidemiology

23
24 Word count:
25 Abstract: 244/250
26 Main text: 3162/3500

27

Objectives: Socioeconomic inequalities in HIV prevention services coverage constitute important barriers to global prevention targets, especially in sub-Saharan Africa (SSA). We aimed at monitoring these inequalities from population-based survey data in 18 SSA countries between 2010 and 2018.

Methods: We defined eight HIV indicators aimed at capturing uptake of HIV prevention services among adult participants. Country-specific wealth-related inequalities were measured using the Relative and Slope Index of Inequalities (RII and SII, respectively) and then pooled using random-effects meta-analyses. We compared inequalities between African regions using the Wilcoxon rank-sum test.

Results: The sample consisted of 358,591 participants (66% women). Despite variability between countries and indicators, the meta-analysis revealed significant levels of relative and absolute inequalities in 6 out of 8 indicators: HIV-related knowledge, positive attitudes toward people living with HIV (PLHIV), condom use at last sexual intercourse, participation to prevention of mother-to-child transmission programs, medical male circumcision and recent HIV testing. The largest inequalities were reported in condom use, with condom use reported 5 times more among the richest versus the poorest (RII=5.02, 95% Confidence interval, CI: 2.79-9.05) and in positive attitudes toward PLHIV, with a 32-percentage point difference between the richest and poorest (SII=0.32, 95% CI: 0.26-0.39). Conversely, no significant inequalities were observed in multi-partnership and HIV seropositivity among youth. Overall, inequalities tended to be larger in West and Central vs. East and Southern African countries.

Conclusions: Despite efforts to scale-up HIV-prevention programs, socioeconomic inequalities remain substantial over the continuum of HIV primary and secondary prevention in several SSA countries.

51 Introduction

52 Despite a sustained decrease over the last decade, sub-Saharan Africa (SSA) continues to bear a
53 disproportionate burden of the global HIV epidemic, with about two thirds of the worldwide
54 number of HIV infections [1]. Many SSA countries are not on track to meet global targets of 90%
55 reduction of new HIV infections by 2030 compared to 2010 [2,3]. There is however a large body
56 of literature supporting the fact that these targets could be reached by effectively combining
57 existing prevention interventions and tools, including behavioural prevention, condoms,
58 voluntary male medical circumcision (VMMC), pre-exposure prophylaxis (PrEP), HIV testing,
59 and prevention of mother-to-child transmission (PMTCT), alongside increased antiretroviral
60 coverage [4].

61 HIV risk, knowledge and attitudes, access to services and HIV outcomes vary largely across
62 gender, age, location or social status. Inequalities, including socio-economic inequalities, have
63 been recognized as a contributor in the failure to meet the UNAIDS 90-90-90 testing and
64 treatment 2020 targets [5], and may also be a factor in the disappointing progress in effective
65 prevention coverage. Indeed, socio-economic inequalities have been reported across HIV
66 knowledge [6], HIV testing [7-9], PMTCT [10], and VMMC [11]. However, there has not been to
67 date a comprehensive large-scale assessment of socioeconomic inequalities across the major
68 prevention pillars to guide the focus of future programming to overcome current gaps. Such an
69 assessment would be valuable, though. Firstly, it would allow to identify, within the main HIV
70 prevention pillars, those that could potentially be at specific risk of generating socioeconomic
71 inequalities in HIV infections. Secondly, a monitoring of national HIV prevention indicators
72 focusing on inequalities would potentially allow to identify and then learn from settings that are
73 specifically performant in terms of equity.

74 In this paper, we used nationally representative surveys conducted across 18 sub-Saharan
75 African countries between 2010 and 2018 to quantify wealth-related inequalities in several
76 indicators aiming at capturing HIV prevention elements.

77

78 **Methods**

79 *Study population and data*

80 We analysed data from the most recent Demographic and Health Surveys (DHS) in 18 Sub-
 81 Saharan African countries (based on data availability as of November 2020). The country
 82 sample was a convenience sample based, and slightly extended, from an earlier research
 83 project [9]. The sample included Burkina Faso, Cameroon, Côte d'Ivoire, the Democratic
 84 Republic of Congo (DRC), Ethiopia, Guinea, Kenya, Lesotho, Liberia, Malawi, Mali, Niger,
 85 Rwanda, Senegal, Sierra Leone, Tanzania, Zambia, and Zimbabwe.

86 DHS are nationally representative surveys based on a multistage design with households as
 87 sampling units. They collect a wide range of demographic and reproduction health indicators
 88 among consenting adults that are interviewed face-to-face by trained interviewers using
 89 standardized questionnaires [12]. The collected data notably include age and gender, as well
 90 as specific data on HIV/AIDS. In particular, some DHS include an HIV serological survey,
 91 for which participants are asked to consent to be tested for HIV (anonymously in most of the
 92 surveys included here). Household wealth was assessed using the DHS wealth index, a
 93 composite measure of living standards that is based on the household's assets (e.g.
 94 refrigerators) and characteristics (e.g., sanitation facilities) [13].

95 All DHS were conducted by national central statistics agencies or research institutes. The
 96 institutions that approved, implemented, or provided funding for the surveys were responsible
 97 for ethical clearance, which guaranteed informed consent and confidentiality of participant's
 98 information.

99

100 *HIV indicators*

101 We defined and analysed a set of eight HIV indicators aimed at capturing HIV knowledge,
 102 attitudes, behaviours and access to and uptake of HIV prevention services. These indicators
 103 were chosen and designed in a pragmatic way based on the information collected in the DHS
 104 across all countries selected for this analysis. These HIV indicators were consistently coded as
 105 binary variables with the level 1 (the level 0, respectively) reflecting a favourable
 106 (unfavourable, respectively) condition regarding HIV. These HIV indicators were:

- 107 i) *HIV-related knowledge*: based on the responses to a set of seven questions related to
 108 HIV transmission and prevention, we defined a categorical indicator reflecting good
 109 HIV-related knowledge (coded as 1 for those answering all questions correctly, 0
 110 otherwise).
- 111 ii) *Positive attitudes towards PLHIV*: based on a set of two questions related to stigma
 112 towards PLHIV, we defined a categorical indicator reflecting positive attitudes
 113 towards PLHIV (coded as 1 for those answering positively to both questions, 0
 114 otherwise).
- 115 iii) *No multipartnership (no or one sexual partner) in the past year*: based on the number
 116 of sexual partners reported in the last 12 months (self-reported, coded as 1 for those
 117 reporting no or one sexual partner, 0 otherwise)
- 118 iv) *Condom use at last sexual intercourse* (self-reported, coded as 1 for those reporting
 119 having used a condom during their last intercourse, 0 otherwise).
- 120 v) *VMMC (among men only)*: based on the self-reported circumcision status and the
 121 qualification of the person who performed the circumcision (health
 122 worker/professional). We only analysed VMMC in 7 countries among of 14 that have

been prioritized for the implementation of VMMC as HIV-prevention intervention in 2007 (mainly in Eastern and Southern Africa) [14].

vi) *Participation in Prevention of Mother to Child Transmission (PMTCT*, among women only): based on self-reported HIV testing during antenatal care among women who gave birth in the 12 months preceding the survey (coded as 1 for those reporting having been tested, 0 otherwise).

vii) *Recent HIV testing*: based on self-reported HIV testing within the past 12 months (coded as 1 for those reporting having been recently tested, 0 otherwise).

viii) *HIV seronegativity among youth*: based on the result of the HIV serological survey among 15-24 years old participants (coded as 1 in seronegative youths, 0 otherwise). trends in HIV prevalence among youth (aged 15-24 years) have been documented to be a surrogate reflecting trends in HIV incidence [15,16]. As the onset of sexual activity in this group is likely to be recent, prevalence reflects recent infection and therefore incidence. We thus used the socioeconomic gradient in HIV seroprevalence among the 15-24 years old as a proxy for the socioeconomic gradient in HIV incidence. To consistently consider a level 1 reflecting a favourable condition and therefore facilitate the interpretation and comparison of inequalities across indicators, the level 1 was chosen to code HIV seronegativity.

Further details on the construction of these HIV-related indicators are provided in Appendix 1

Statistical analysis

For each country survey and each HIV indicator, we calculated percentages of positivity taking into account the survey design and sampling weights. We then assessed within-country socioeconomic inequalities related to each indicator based on participants' relative rank in the

cumulative distribution of the wealth index. Inequalities were measured on both relative and absolute scales using the relative index of inequality (RII) and the slope index of inequality (SII), respectively [17]. The RII expresses the ratio of the predicted outcomes between the richest and the poorest people in the wealth distribution. The SII represents the absolute difference in the predicted proportions of these two extremes. For instance, a RII value of 2 indicates that individuals at the upper extreme of the wealth distribution (the richest) are twice more likely to report the outcome than individuals at the lower extreme (the poorest). A SII value of 0.2 means that the proportion of the outcome is higher by 0.2 (or higher by 20 percentage-points) in the upper extreme of the wealth distribution (the richest) versus the lower extreme (the poorest). Reporting inequalities on both scales is recommended, because conclusions can be skewed when only one or the other is used [18]. Furthermore, the choice of a relative scale over an absolute scale or vice versa carries an implicit normative judgment on what a fair and socially just distribution of health should be [19].

Both RII and SII were obtained by fitting a modified Poisson regression with a robust variance and a log link function to estimate the association between participants' relative wealth rank and each indicator, and by using generalised estimating equations to account for the clustering of observations [20]. We used the Wilcoxon rank-sum test to compare indices of inequalities between West and Central Africa (WCA) versus East and Southern African (ESA) countries (except for *VMMC*, which was studied in ESA only). Lastly, we used random-effects meta-analysis to average inequality estimates across countries [21].

Results

Study population

DHS included in the present work were conducted between 2010 and 2018 (Figure 1), with participation rates ranging from 90% (Ethiopia 2016) to 100% (Rwanda 2014-15) (Table 1). Overall, these studies represented 358,591 participants, (238,055 women and 120,536 men).




Figure 1: Demographic and Health Survey (DHS) year (left) and adult HIV prevalence (right) of the 18 Sub-Saharan African countries (NB: For Kenya, HIV prevalence is unavailable in its last DHS 2014).

Table 1: Distribution of HIV-related indicators for 18 Sub-Saharan African countries (Demographic and Health Surveys). PLHIV: People living with HIV; PMTCT : Prevention of Mother-to-Child Transmission; VMMC: Voluntary Medical Male Circumcision; NA: Not Available. ¹: as reported in the country's DHS final report ; ²: within the past year; ³: at last sexual intercourse; ⁴: among those aged 15-24 y. Countries are ordered west to east.

West and Central Africa										
	Senegal (2017)	Sierra Leone (2013)	Guinea (2018)	Liberia (2013)	Côte d'Ivoire (2011-12)	Mali (2012- 13)	Burkina Faso (2010)	Niger (2012)	Cameroon (2018)	DR Congo (2013-14)
Response rate (%) ¹	92%	96%	98%	96%	92%	94%	97%	91%	97%	98%
Sample size	23 764	23 920	14 991	13 357	15 195	14 823	24 394	15 088	20 505	27 483
HIV-related knowledge	17.7%	19.5%	15.8%	6.2%	14.0%	19.9%	22.7%	11.0%	30.3%	12.5%
Positive Attitude towards PLHIV	32.9%	34.1%	16.0%	34.7%	47.0%	45.1%	34.5%	20.1%	55.1%	35.7%
No multipartnership ²	97.1%	88.1%	95.4%	90.1%	88.1%	95.2%	94.2%	95.6%	90.1%	91.1%
Condom use ³	6.5%	5.0%	8.3%	12.8%	18.5%	3.6%	12.7%	1.1%	29.7%	9.2%
Participation to PMTCT (Females)	67.0%	72.2%	30.7%	76.4%	50.0%	23.2%	40.3%	33.5%	89.4%	25.0%
VMMC (Males)	-	-	-	-	-	-	-	-	-	-
Recent HIV-Testing ²	18.6	14.7%	8.4%	19.1%	13.5%	6.8%	10.3%	6.9%	57.7%	8.6%
HIV-seronegativity among youth ⁴	99.8%	98.9%	99.1%	98.9%	98.7%	99.2%	99.7%	100%	98.7%	99.3%
East and Southern Africa										
	Zambia (2018)	Lesotho (2014)	Zimbabwe (2015)	Rwanda (2014-15)	Malawi (2015- 16)	Tanzania (2011-12)	Kenya (2014)	Ethiopia (2016)		
Response rate (%) ¹	94%	95%	94%	100%	96%	97%	92%	90%		
Sample size	25 815	9 552	18 351	19 714	32 040	19 319	11 909	28 371		
HIV-related knowledge	29.6%	23.8%	40.2%	43.4%	34.7%	28.3%	27.2%	22.2%		
Positive Attitude towards PLHIV	72.8%	80.5%	77.9%	84.3%	81.6%	61.5%	75.2%	39.7%		
No multipartnership ²	92.1%	87.6%	92.9%	98.0%	96.1%	88.9%	93.4%	98.2%		
Condom use ³	18.8%	51.3%	23.4%	12.4%	14.6%	19.3%	21.6%	3.0%		
Participation to PMTCT (Females)	97.3%	98.4%	97.9%	99.4%	97.2%	80.5%	98.0%	57.2%		
VMMC (Males)	40.4%	NA	11.3%	85.4%	9.0%	40.7%	60.0%	15.8%		
Recent HIV-Testing ²	59.8%	52.6%	43.6%	38.8%	43.9%	30.4%	64.9%	20.6%		
HIV-seronegativity among youth ⁴	96.1%	90.3%	95.0%	99.0%	96.9%	98.0%	NA	99.8%		

HIV indicators

Country-specific percentages of positivity for each HIV indicator are presented in Table 1. In ESA countries, good HIV-related knowledge ranged from 22% (Ethiopia) to 43 % (Rwanda) while it was $\leq 20\%$ in 8 out of 10 WCA countries (with the exception of Cameroon, 30% and Burkina Faso, 23%). Similarly, positive attitudes toward PLHIV were $>60\%$ in 7 out of 8 ESA countries (with the exception of Ethiopia, 40%) while the highest level in WCA was 55% (Cameroon). Reporting no multipartnership in the past 12 months was consistently high across countries, ranging from 88% (Sierra Leone, Côte d'Ivoire and Lesotho) to 98% (Rwanda and Ethiopia). Reported condom use at last sexual intercourse was highest in Lesotho (51%). In other countries reported condom use was lower, ranging from 1% (Niger) to 23% (Zimbabwe). Overall, higher uptake of HIV testing in the past 12 months was observed in ESA, with uptake $>20\%$ in all countries, as compared to in WCA, where uptake was $<20\%$ in 9 out of 10 countries. Similarly, ESA countries reported levels of participation to PMTCT $>95\%$ in 6 out of 8 countries (with the exception of Ethiopia, 57% and Tanzania, 81%), while it was $<90\%$ in all WCA countries, with large heterogeneity (from 23% in Mali to 89% in Cameroon). As mentioned earlier, VMMC was reported in ESA countries only, with levels ranging from 9% (Malawi) to 85% (Rwanda).

Socioeconomic inequalities in the HIV indicators

Measures of relative and absolute inequalities pooled across countries are presented in Table 2. Significant levels of relative and absolute inequalities in favour of the richest were observed for HIV-related knowledge, positive attitudes toward PLHIV, condom use at last sex, participation to PMTCT, VMMC and recent HIV testing. In contrast, we observed inequalities in the other direction in reporting no multipartnership and in HIV-seronegativity among youth (that is, the poorest reported lower levels of multipartnership and exhibited lower levels of HIV seropositivity), although these inequalities were borderline significant and of low magnitude as

compared to the other indicators. The ranking of the indicators by level of inequalities varied between the relative and the absolute scales. On the relative scale, the highest level was observed for condom use, with the richest being approximatively 5 times more likely (RII = 5.02; 95 Confidence interval, CI, 2.79-9.05) to use condom than the poorest people. In the absolute scales, the largest gap between the richest and the poorest people was observed in positive attitudes toward PLHIV, with a difference of 32 percentage-points (SII=0.32; 95% CI 0.26-0.39). On the relative scale, levels of inequalities were significantly higher in WCA as compared to ESA countries for the following indicators: HIV-related knowledge, positive attitude towards PLHIV, condom use, participation to PMTCT and recent HIV testing (all rank-sum test p -values <0.05). Absolute levels of inequalities were also higher in WCA versus ESA countries for participation to PMTCT (0.43 versus 0.12, respectively, $p=0.009$) and for recent HIV testing, although this was borderline significant (0.16 versus 0.09, respectively, $p=0.06$).

A more detailed comparison of the levels of inequalities across indicators and across countries is displayed in Figure 2. In WCA countries, the countries exhibiting the higher levels of inequalities were generally not the same when considering the relative or the absolute scale. In contrast, in ESA countries, Ethiopia tended to exhibit the highest levels of inequalities on both the relative and absolute scales. We also observed significant inequalities in favour of the poor in HIV-seronegativity among youth in some countries, especially on the absolute scale (for example in Sierra Leone, Zambia, Rwanda), although their magnitudes were lower than for other indicators.

Table 3 presents the correlations between RII (log-transformed) and SII values across HIV-related indicators. We observed significant positive correlations between relative- and absolute inequalities metrics for most of the indicators (Pearson correlation coefficients >0.6 in 6 out of 8 indicators), apart from condom use and VMMC presenting no evidence of correlation.

Table 2: Pooled relative and absolute wealth-related inequalities in various HIV-related indicators in sub-Saharan Africa, per region and overall. RII: relative index of inequality; SII : slope index of inequality; PLHIV: People living with HIV; PMTCT : Prevention of Mother-to-Child Transmission; VMMC: Voluntary Medical Male Circumcision. ¹: rank-sum test p-value; ²: within the past year; ³: at last sexual intercourse; ⁴: among those aged 15-24 y.

	West and Central Africa		East and Southern Africa		P-value ¹	Overall (all countries)	
	Pooled RII [95% CI]	I2	Pooled RII [95% CI]	I2		Pooled RII [95% CI]	I2
HIV-related knowledge	3.13 [2.26;4.34]	52%	1.89 [1.33;2.69]	70%	p = 0.0145	2.45 [1.91;3.14]	72%
Positive Attitude towards PLHIV	2.49 [1.74;3.56]	78%	1.60 [1.19;2.15]	76%	p = 0.016	1.99 [1.57;2.53]	81%
No multipartnership ²	0.99 [0.97;1.00]	0%	0.99 [0.98;1.01]	0%	p = 0.32	0.99 [0.98;1.00]	0%
Condom use ³	8.08 [3.87;16.87]	77%	2.80 [1.13;6.95]	88%	p = 0.007	5.02 [2.79;9.05]	88%
Participation to PMTCT (Females)	2.87 [1.60;5.15]	82%	1.09 [0.90;1.33]	51%	p = 0.002	1.70 [1.20;2.43]	85%
VMMC (Males)	-	-	2.08 [1.38;3.15]	59%	-	-	-
Recent HIV-Testing ²	4.47 [2.13;9.40]	90%	1.24 [0.85;1.81]	76%	p = 0.003	2.43 [1.47;4.04]	92%
HIV-seronegativity among youth ⁴	0.99 [0.98;0.99]	0%	0.98 [0.95;1.02]	0%	p = 0.45	0.99 [0.98;1.00]	0%
	West and Central Africa		East and Southern Africa		P-value ¹	Overall (all countries)	
	Pooled SII [95% CI]	I2	Pooled SII [95% CI]	I2		Pooled SII [95% CI]	I2
HIV-related knowledge	0.18 [0.13;0.23]	98%	0.18 [0.11;0.25]	98%	p = 0.89	0.18 [0.14;0.22]	98%
Positive Attitude towards PLHIV	0.33 [0.23;0.43]	99%	0.31 [0.19;0.44]	100%	p = 0.76	0.32 [0.26;0.39]	99%
No multipartnership ²	-0.02 [-0.04;-0.00]	98%	-0.01 [-0.02;0.01]	96%	p = 0.32	-0.01 [-0.03;-0.00]	97%
Condom use ³	0.17 [0.11;0.23]	98%	0.15 [0.10;0.20]	96%	p = 0.69	0.16 [0.12;0.20]	98%
Participation to PMTCT (Females)	0.43 [0.31;0.55]	97%	0.12 [-0.07;0.32]	99%	p = 0.009	0.29 [0.17;0.42]	99%
VMMC (Males)	-	-	0.20 [0.11;0.29]	97%	-	-	-
Recent HIV-Testing ²	0.16 [0.11;0.21]	97%	0.09 [-0.01;0.19]	99%	p = 0.06	0.13 [0.08;0.18]	99%
HIV-seronegativity among youth ⁴	-0.01 [-0.02;-0.01]	92%	-0.01 [-0.05;0.03]	98%	p = 1.00	-0.01 [-0.03;0.00]	97%



246

247 **Figure 2 : Relative (top) and absolute (bottom) wealth-related inequalities in various HIV-related**
 248 **indicators across 18 sub-Saharan African countries.** Countries are ordered wet to east. RII: relative
 249 index of inequality; SII : slope index of inequality; PLHIV: People living with HIV; PMTCT : Prevention
 250 of Mother-to-Child Transmission; VMMC: Voluntary Medical Male Circumcision.

Table 3 : Correlation between country-level relative and absolute wealth-related inequalities, per HIV indicator. RII: relative index of inequality; SII: slope index of inequality; PLHIV: People living with HIV; PMTCT : Prevention of Mother-to-Child Transmission; VMMC: Voluntary Medical Male Circumcision. 1: log-transformed; 2: within the past year; 3: at last sexual intercourse; 4: among those aged 15-24 y.

	Correlation between RII ¹ and SII values,	
	Pearson r	p-value
HIV-related knowledge	0.607	0.0075
Positive Attitude towards PLHIV	0.716	<0.001
No multipartnership ²	0.992	<0.001
Condom use ³	0.064	0.7996
Participation to PMTCT (Females)	0.809	<0.001
VMMC (Males)	-0.098	0.8347
Recent HIV-Testing ²	0.641	0.0041
HIV-seronegativity among youth ⁴	1.000	<0.001

DISCUSSION

In this study, relying on cross-sectional population-based surveys, we provide an extensive assessment of socio-economic inequalities in various HIV indicators capturing knowledge, attitudes, behaviours and access to and uptake of prevention services in a large set of sub-Saharan African countries. We document important levels of wealth-related inequalities, both on the relative and absolute scales, in HIV-related knowledge, positive attitudes toward PLHIV, condom use, participation to PMTCT, uptake of VMMC and recent HIV testing. The magnitude of these inequalities varies across countries and indicators. However, inequalities tend to be more marked in WCA than in ESA countries. In contrast, we document low overall levels of inequalities in multi-partnership and HIV seropositivity among youth, a surrogate of HIV incidence.

The arithmetic behind inequality measures

The results we provide here allow two types of comparisons: cross-country comparisons of inequalities given a specific HIV indicator, and within-country comparisons of inequalities

across HIV indicators. However, first bearing in mind some arithmetic considerations is important when comparing RII and SII across indicators and countries [22,23]. Indeed, for any indicator, the overall level in the population should be considered when interpreting relative and absolute inequalities measures. Relative inequalities tend to be larger at low overall levels of the considered indicator, while absolute inequalities tend to be larger at intermediate levels. This arithmetic explains, for instance, why condom use in Niger (overall self-reported level: 1.1%) acknowledges dramatic relative inequalities (RII >100) but relatively limited absolute inequalities (SII = 0.07). This illustrates the importance of reporting both relative and absolute measures when reporting inequalities, as conclusions would be quite different if based on only one or the other indicators in this single example. However, our analysis of correlation showed that the levels of inequalities tended to correlate across both scales for most of the indicators we used.

Cross-country comparisons of inequalities

Heterogeneity across countries regarding the inequalities reported here may be driven by a potentially large spectrum of socioeconomic, epidemiological or healthcare-related factors. Nevertheless, we observed that geography explained some of this heterogeneity: higher levels of wealth-related inequalities were observed in countries in WCA compared to those in ESA, especially on the relative scale. Several factors may explain higher levels of inequalities in WCA. This may be related to the historical prominence of the private healthcare sector in this region, which generally implies user fees that may act as a barrier to access to care for the poorest [24]. Higher levels of inequalities are also possibly linked to the smaller overall HIV epidemics in WCA compared to ESA. Indeed, ESA countries had higher HIV prevalence than WCA countries in our sample, except for Ethiopia, which precisely had the highest levels of wealth-related inequalities within ESA countries. Previous results suggested that HIV-related inequalities correlated with epidemiological rather than macro-economic factors [25]. In addition, the country-level HIV prevalence has been reported to correlate with levels of HIV donor and national funding for HIV

prevention services [26]. Moreover, donor funding tended to prioritise prevention programs targeting key populations in WCA, as new HIV infections are often predominantly in these populations in the region, whilst in ESA HIV prevention efforts are broader encompassing both key populations and the general populations [27]. While key populations deserve a priority focus for HIV prevention in WCA, additional efforts should be considered to overcome the socioeconomic disparities in knowledge and access to HIV testing and prevention services more broadly.

Comparisons of inequalities across HIV indicators

Overall, we did not observe large socioeconomic disparities in reporting having multiple sexual partners, consistently with recent results [28]. Large absolute and relative inequalities, overall, remain regarding lack of knowledge and stigmatizing attitudes toward PLHIV, and this may undermine HIV prevention, care and treatment [29]. Concerning inequalities are observed in condom use, PMTCT, VMMC and HIV testing, especially in WCA countries. This is of particular concern because these are interventions that can prevent new infections, directly for condom use and VMMC, or when linked to care and treatment for PMTCT and HIV testing. Large inequalities in condom use may reflect difficulties to access condoms free of charge. However, PMTCT, VMMC and HIV testing are also interventions that are usually provided at no direct cost for the individual, which underlines that providing these services without cost to clients is not the only factor to ensure equitable access to prevention interventions. Lessons should be drawn from the experiences of PMTCT or HIV testing programs in ESA countries that are currently offered to all at no cost, without producing measurable health inequalities. For instance, Rwanda has successfully integrated HIV services within the existing health system and assured delivery of services in remote areas. These strategies have strengthened the country response to HIV and have most probably contributed to limiting health inequalities [30].

Contrary to what one could expect considering the inequalities disfavoring the poorest in terms of access to HIV prevention service that we report here, we did not find any evidence of

socioeconomic inequalities regarding HIV prevalence among the youth, an indicator we used as a proxy for HIV incidence. This apparently inconsistent result may however be linked to the complex and changing social epidemiology of HIV. Many early studies reported that higher SEP tended to be associated with a higher prevalence of HIV infection, making this distribution unusual as compared to many health outcomes [31,32]. The absence of overall socioeconomic gradient we report here may represent a transient state where inequalities are inverting, from a higher risk for the wealthiest to a higher risk for the poorest [33,34]. Indeed, a recent study relying on a >20-years follow-up among a population-based open cohort in rural Uganda documented a widening socioeconomic gradient over time, with a higher risk of incident HIV infection among the poorest [35]. An effort to more systematically monitor socio-economic inequalities in HIV incidence is thus of high importance to detect possibly changing SEP gradients. It is anticipated that the newer-generation surveys, such as the population-based HIV impact assessment surveys (<https://phia.icap.columbia.edu/>) which integrate HIV incidence assays, may provide useful data for this monitoring.

Main strengths and limitations of the study

This study has several limitations and strengths. The between-country comparison may be affected by the period of data collection for each national survey (from 2010 to 2018). Large HIV programs have been implemented over this study period and may have reduced inequalities over time [9]. Our analysis was limited to indicators pragmatically built on data collected as part of the DHS. As such, it matches only partially the recently proposed unifying framework of the HIV prevention cascade [36]. For instance, an important aspect absent from the DHS data, and therefore from our analysis, relates to knowledge and use of antiretroviral treatment for individual and collective health benefits. For the same reason, our analysis ignores important prevention interventions, such as pre-exposure prophylaxis (PrEP), or interventions targeted to key populations. However, to our knowledge, this study constitutes the first effort to quantify

both relative and absolute socioeconomic inequalities on a large set of HIV-related indicators collected from large, representative surveys conducted in numerous SSA countries.

Conclusions

In conclusion, this study provides a comprehensive monitoring of socioeconomic inequalities in HIV knowledge, attitudes, behaviours and prevention in sub-Saharan Africa. In other fields such as child health, such monitoring has provided important insight in the way policies may be tailored to the patterns of inequalities in order to best address them [37]. We hope that this study will help in the strategical articulation of HIV prevention approaches that is required for fulfilling the focus on reducing inequalities that the renewed 2021-2026 Global AIDS Strategy adopted [38].

Authors' contributions

MH, PAA-T and KJ conceived and planned the study with input from LT. MH and PAA-T collated and processed DHS data. MH and KJ conducted the analysis and produced output figures and tables with input from PAA-T and LT. All authors contributed to the interpretation of the results. MH and KJ wrote the first draft of the report and all authors contributed to subsequent revisions.

REFERENCES

- 1 GBD 2017 HIV collaborators. Global, regional, and national incidence, prevalence, and mortality of HIV, 1980-2017, and forecasts to 2030, for 195 countries and territories: a systematic analysis for the Global Burden of Diseases, Injuries, and Risk Factors Study 2017. *Lancet HIV* 2019; 6:e831–e859.
- 2 2020 Global AIDS Update — Seizing the moment — Tackling entrenched inequalities to end epidemics. <https://www.unaids.org/en/resources/documents/2020/global-aids-report> (accessed 8 Mar2021).
- 3 Sartorius B, VanderHeide JD, Yang M, Goosmann EA, Hon J, Haeuser E, *et al.* Subnational mapping of HIV incidence and mortality among individuals aged 15–49 years in sub-Saharan Africa, 2000–18: a modelling study. *The Lancet HIV* 2021; 8:e363–e375.

- 377 4 Krishnaratne S, Hensen B, Cordes J, Enstone J, Hargreaves JR. Interventions to strengthen the HIV
378 prevention cascade: a systematic review of reviews. *The Lancet HIV* 2016; 3:e307–e317.
- 379 5 Evidence review: Implementation of the 2016–2021 UNAIDS strategy: on the fast-track to end
380 aids. Geneva, Switzerland: UNAIDS; 2020. [https://www.unaids.org/sites/default/files/UNAIDS-](https://www.unaids.org/sites/default/files/UNAIDS-2016-2021-Strategy-Evidence-review_en.pdf)
381 [2016-2021-Strategy-Evidence-review_en.pdf](https://www.unaids.org/sites/default/files/UNAIDS-2016-2021-Strategy-Evidence-review_en.pdf) (accessed 28 May2021).
- 382 6 Burgoyne AD, Drummond PD. Knowledge of HIV and AIDS in women in sub-Saharan Africa. *Afr J*
383 *Reprod Health* 2008; 12:14–31.
- 384 7 Jean K, Anglaret X, Moh R, Lert F, Dray-Spira R. Barriers to HIV Testing in Côte d’Ivoire: The Role
385 of Individual Characteristics and Testing Modalities. *PLoS ONE* 2012; 7:e41353.
- 386 8 Kirakoya-Samadoulougou F, Jean K, Maheu. Uptake of HIV Testing in Burkina Faso: An
387 Assessment of Individual and Community-level Determinants. *BMC Public Health* 2017; (In
388 revision).
- 389 9 Ante-Testard PA, Benmarhnia T, Bekelync A, Baggaley R, Ouattara E, Temime L, *et al.* Temporal
390 trends in socioeconomic inequalities in HIV testing: an analysis of cross-sectional surveys from 16
391 sub-Saharan African countries. *The Lancet Global Health* 2020; 8:e808–e818.
- 392 10 Hlarlath MO, Grede N, de Pee S, Bloem M. Economic and Social Factors are Some of the Most
393 Common Barriers Preventing Women from Accessing Maternal and Newborn Child Health
394 (MNCH) and Prevention of Mother-to-Child Transmission (PMTCT) Services: A Literature Review.
395 *AIDS Behav* 2014; 18:516–530.
- 396 11 Gummerson E, Maughan-Brown B, Venkataramani A. Who is taking up voluntary medical male
397 circumcision? Early evidence from Tanzania. *AIDS* 2013; 27:2657–2659.
- 398 12 Corsi DJ, Neuman M, Finlay JE, Subramanian SV. Demographic and health surveys: a profile. *Int J*
399 *Epidemiol* 2012; 41:1602–1613.
- 400 13 Rutstein SO, Johnson K. DHS Comparative Reports. The DHS Wealth Index. Calverton, Maryland,
401 U.S.A: ORC Macro; 2004.
- 402 14 Njeuhmeli E, Schnure M, Vazzano A, Gold E, Stegman P, Kripke K, *et al.* Using mathematical
403 modeling to inform health policy: A case study from voluntary medical male circumcision scale-
404 up in eastern and southern Africa and proposed framework for success. *PLOS ONE* 2019;
405 14:e0213605.
- 406 15 Kim AA, Hallett T, Stover J, Gouws E, Musinguzi J, Mureithi PK, *et al.* Estimating HIV Incidence
407 among Adults in Kenya and Uganda: A Systematic Comparison of Multiple Methods. *PLoS One*
408 2011; 6. doi:10.1371/journal.pone.0017535
- 409 16 Mahy M, Garcia-Calleja JM, Marsh KA. Trends in HIV prevalence among young people in
410 generalised epidemics: implications for monitoring the HIV epidemic. *Sex Transm Infect* 2012;
411 88:i65–i75.
- 412 17 Moreno-Betancur M, Latouche A, Menvielle G, Kunst AE, Rey G. Relative index of inequality and
413 slope index of inequality: a structured regression framework for estimation. *Epidemiology* 2015;
414 26:518–527.

- 415 18 King NB, Harper S, Young ME. Use of relative and absolute effect measures in reporting health
416 inequalities: structured review. *BMJ* 2012; 345:e5774.
- 417 19 Harper S, King NB, Meersman SC, Reichman ME, Breen N, Lynch J. Implicit Value Judgments in
418 the Measurement of Health Inequalities. *Milbank Q* 2010; 88:4–29.
- 419 20 Zou G. A Modified Poisson Regression Approach to Prospective Studies with Binary Data. *Am J*
420 *Epidemiol* 2004; 159:702–706.
- 421 21 Viechtbauer W. Conducting Meta-Analyses in R with the metafor Package. *Journal of Statistical*
422 *Software* 2010; 36:1–48.
- 423 22 Houweling TA, Kunst AE, Huisman M, Mackenbach JP. Using relative and absolute measures for
424 monitoring health inequalities: experiences from cross-national analyses on maternal and child
425 health. *Int J Equity Health* 2007; 6:15.
- 426 23 Mackenbach JP, Martikainen P, Menvielle G, Gelder R de. The arithmetic of reducing relative and
427 absolute inequalities in health: a theoretical analysis illustrated with European mortality data. *J*
428 *Epidemiol Community Health* 2016; 70:730–736.
- 429 24 Ridde V. From institutionalization of user fees to their abolition in West Africa: a story of pilot
430 projects and public policies. *BMC Health Services Research* 2015; 15:S6.
- 431 25 Ante-Testard PA, Temime L, Jean K. Epidemiological rather than macro-economic factors
432 correlate with socioeconomic inequalities in HIV testing in 16 sub-Saharan African countries.
433 *medRxiv* 2021; :2021.09.22.21263940.
- 434 26 Amico P, Aran C, Avila C. HIV Spending as a Share of Total Health Expenditure: An Analysis of
435 Regional Variation in a Multi-Country Study. *PLoS One* 2010; 5.
436 doi:10.1371/journal.pone.0012997
- 437 27 Funding for HIV and AIDS. Avert. 2021.[https://www.avert.org/professionals/hiv-around-](https://www.avert.org/professionals/hiv-around-world/global-response/funding)
438 [world/global-response/funding](https://www.avert.org/professionals/hiv-around-world/global-response/funding) (accessed 23 Jul2021).
- 439 28 Ali MM, Merdad L, Bellizzi S. Socioeconomic variations in risky sexual behavior among
440 adolescents in 14 sub-Saharan Africa countries who report ever having had sex. *Int J Equity*
441 *Health* 2021; 20:11.
- 442 29 Nyblade L, Mingkwan P, Stockton MA. Stigma reduction: an essential ingredient to ending AIDS
443 by 2030. *Lancet HIV* 2021; 8:e106–e113.
- 444 30 Binagwaho A, Kankindi I, Kayirangwa E, Nyemazi JP, Nsanzimana S, Morales F, *et al.* Transitioning
445 to Country Ownership of HIV Programs in Rwanda. *PLOS Medicine* 2016; 13:e1002075.
- 446 31 Hargreaves JR, Glynn JR. Educational attainment and HIV-1 infection in developing countries: a
447 systematic review. *Trop Med Int Health* 2002; 7:489–498.
- 448 32 Fox AM. The social determinants of HIV serostatus in sub-Saharan Africa: an inverse relationship
449 between poverty and HIV? *Public Health Rep* 2010; 125 Suppl 4:16–24.
- 450 33 Hargreaves JR, Davey C, White RG. Does the “inverse equity hypothesis” explain how both
451 poverty and wealth can be associated with HIV prevalence in sub-Saharan Africa? *J Epidemiol*
452 *Community Health* 2013; 67:526–529.

- 453 34 Hargreaves JR, Davey C, Fearon E, Hensen B, Krishnaratne S. Trends in Socioeconomic
454 Inequalities in HIV Prevalence among Young People in Seven Countries in Eastern and Southern
455 Africa. *PLoS ONE* 2015; 10:e0121775.
- 456 35 Santelli JS, Chen I, Makumbi F, Wei Y, Nalugoda F, Lutalo T, *et al.* Household wealth and HIV
457 incidence over time, rural Uganda, 1994–2018. *AIDS* 2021; 35:1835–1843.
- 458 36 Hargreaves JR, Delany-Moretlwe S, Hallett TB, Johnson S, Kapiga S, Bhattacharjee P, *et al.* The
459 HIV prevention cascade: integrating theories of epidemiological, behavioural, and social science
460 into programme design and monitoring. *The Lancet HIV* 2016; 3:e318–e322.
- 461 37 Victora CG, Joseph G, Silva ICM, Maia FS, Vaughan JP, Barros FC, *et al.* The Inverse Equity
462 Hypothesis: Analyses of Institutional Deliveries in 286 National Surveys. *Am J Public Health* 2018;
463 108:464–471.
- 464 38 End Inequalities. End AIDS. Global AIDS Strategy 2021-2026. Geneva, Switzerland: UNAIDS; 2021.
465 <https://www.unaids.org/en/resources/documents/2021/2021-2026-global-AIDS-strategy>
466 (accessed 28 May2021).
- 467