| 1        |   |
|----------|---|
| 2        |   |
| 3        |   |
| 4        | Deleterious drugs in COVID-19: a rapid systematic review and  |
| 5        | meta-analysis   |
| 6        |   |
| 7        |   |
| 8        | Michael Holder <sup>1*</sup> , Catherine Heeney <sup>1¶</sup> , Stephen Malden <sup>1¶</sup> , Uditha Perera <sup>1¶</sup> , Aziz Sheikh <sup>1</sup> |
| 9        |   |
| 10       |   |
| 11       |   |
|          |   |
| 12       | <sup>1</sup> Usher Institute, University of Edinburgh, Edinburgh, United Kingdom  |
| 13       |   |
| 14       |   |
| 15       | * Corresponding author  |
| 16       | E-mail: michael holder?@nhs scot (MH)   |
| 10       |   |
| 17       |   |
| 18       |   |
| 19<br>20 | <sup>¶</sup> These authors contributed equally to this work.  |

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

## 1 Abstract

2 Background: Concerns have been expressed about a number of drugs that

- 3 potentially worsen outcomes in patients with COVID-19. We sought to identify all
- 4 potentially deleterious drug groups in COVID-19 and critically assess the
- 5 underpinning strength of evidence pertaining to the harmful effects of these drugs.

6 Methods and findings: We performed a rapid systematic review, searching

- 7 Medline, Embase and two COVID-19 portfolios (WHO COVID-19 database and
- 8 NIH iSearch COVID-19 portfolio) for papers and preprints related to primary
- 9 studies investigating drugs identified as potentially deleterious. Primary outcomes
- 10 were direct measures of susceptibility to infection, disease severity and mortality.
- 11 Study quality was assessed using the National Heart, Lung, and Blood Institute
- 12 quality assessment tools. Random-effects meta-analyses were used for data
- 13 synthesis with further subgroup analyses where possible for specific outcome, study
- 14 design, statistical adjustment and drug groups when two were combined.
- 15 Sensitivity analyses were performed by removing any studies at high risk of bias
- 16 and by publication status.
- 17 49 observational studies (15 peer-reviewed papers and 34 preprints) reported
- 18 primary outcomes for eight drug groups hypothesised to be deleterious. Meta-
- 19 analysis showed that acute inpatient corticosteroid use was associated with
- 20 increased mortality (OR 2.22, 95% CI 1.26-3.90), however this result appeared to
- 21 have been biased by confounding via indication. One subgroup analysis indicated
- 22 an association between immunosuppressant use and susceptibility to COVID-19
- among case control and cross-sectional studies (OR 1.29, 95% CI 1.19-1.40) but this
- 24 was not found with cohort studies (OR 1.11, 95% CI 0.86-1.43). Studies which
- 25 adjusted for multiple confounders showed that people taking angiotensin-
- 26 converting-enzyme inhibitors (ACEIs) or angiotensin-II-receptor blockers (ARBs)
- 27 required a lower level of care (OR 0.85, 95% CI 0.74-0.98). Furthermore, studies

- 1 which combined these two drug groups in their analysis demonstrated an
- 2 association with a lower mortality (OR 0.68, 95% CI 0.55-0.85).
- 3 **Conclusions:** We found minimal high quality or consistent evidence that any drug
- 4 groups increase susceptibility, severity or mortality in COVID-19. Converse to initial
- 5 hypotheses, we found some evidence that regular use of ACEIs and ARBs prior to
- 6 infection may be effective in reducing the level of care required, such as requiring
- 7 intensive care, in patients with COVID-19.
- 8

# 1 Introduction

| 2  | The ongoing Coronavirus 2019 (COVID-19) pandemic has already had a profound   |  |  |  |  |  |  |
|--|---|--|--|--|--|--|--|
| 3  | global impact. As of 3 <sup>rd</sup> March 2021, there had been over 114 million confirmed  |  |  |  |  |  |  |
| 4  | cases and over 2.5 million deaths worldwide (1). SARS-CoV-2 is one of seven   |  |  |  |  |  |  |
| 5  | coronaviruses known to infect humans (2). It is, together with severe acute   |  |  |  |  |  |  |
| 6  | respiratory syndrome coronavirus (SARS-CoV) and the Middle East respiratory   |  |  |  |  |  |  |
| 7  | syndrome coronavirus (MERS-CoV), also one of three zoonotic human   |  |  |  |  |  |  |
| 8  | betacoronaviruses to have emerged in the last 20 years (3). The genome sequence of  |  |  |  |  |  |  |
| 9  | SARS-CoV-2 was found to be 79% similar to SARS-CoV (4); both viruses enter  |  |  |  |  |  |  |
| 10   | human cells via angiotensin-converting enzyme 2 (ACE2), a factor linked to human-   |  |  |  |  |  |  |
| 11   | to-human transmission (5). ACE2 received considerable attention following its   |  |  |  |  |  |  |
| 12   | discovery as the entry for SARS-CoV, and has previously been purported to play a  |  |  |  |  |  |  |
| 13   | protective role in disease, as opposed to the deleterious role of angiotensin-  |  |  |  |  |  |  |
| 14   | converting enzyme (ACE) (6). Conversely, some hypothesized that drugs which   |  |  |  |  |  |  |
| 15   | upregulate ACE2 may increase the risk of severe SARS-CoV-2 infection (7).   |  |  |  |  |  |  |
|  |   |  |  |  |  |  |  |
| 16   | American to the second of lengum COVID 10 cases require admission to an   |  |  |  |  |  |  |
| 16   | Approximately five percent of known COVID-19 cases require admission to an  |  |  |  |  |  |  |
| 16<br>17   | Approximately five percent of known COVID-19 cases require admission to an intensive care unit (ICU) (8). Many of these patients develop acute respiratory  |  |  |  |  |  |  |
| 16<br>17<br>18   | Approximately five percent of known COVID-19 cases require admission to an intensive care unit (ICU) (8). Many of these patients develop acute respiratory distress syndrome (ARDS) and require mechanical ventilation (MV) or other  |  |  |  |  |  |  |
| 16<br>17<br>18<br>19   | Approximately five percent of known COVID-19 cases require admission to an<br>intensive care unit (ICU) (8). Many of these patients develop acute respiratory<br>distress syndrome (ARDS) and require mechanical ventilation (MV) or other<br>respiratory support (9). Cytokine storms (a large release of pro-inflammatory   |  |  |  |  |  |  |
| 16<br>17<br>18<br>19<br>20   | Approximately five percent of known COVID-19 cases require admission to an<br>intensive care unit (ICU) (8). Many of these patients develop acute respiratory<br>distress syndrome (ARDS) and require mechanical ventilation (MV) or other<br>respiratory support (9). Cytokine storms (a large release of pro-inflammatory<br>proteins) have been implicated in the development of ARDS, as well as multi-organ  |  |  |  |  |  |  |
| 16<br>17<br>18<br>19<br>20<br>21   | Approximately five percent of known COVID-19 cases require admission to an<br>intensive care unit (ICU) (8). Many of these patients develop acute respiratory<br>distress syndrome (ARDS) and require mechanical ventilation (MV) or other<br>respiratory support (9). Cytokine storms (a large release of pro-inflammatory<br>proteins) have been implicated in the development of ARDS, as well as multi-organ<br>failure, with a correlation between cytokine levels and mortality (10).   |  |  |  |  |  |  |
| <ol> <li>16</li> <li>17</li> <li>18</li> <li>19</li> <li>20</li> <li>21</li> <li>22</li> </ol>   | Approximately five percent of known COVID-19 cases require admission to an<br>intensive care unit (ICU) (8). Many of these patients develop acute respiratory<br>distress syndrome (ARDS) and require mechanical ventilation (MV) or other<br>respiratory support (9). Cytokine storms (a large release of pro-inflammatory<br>proteins) have been implicated in the development of ARDS, as well as multi-organ<br>failure, with a correlation between cytokine levels and mortality (10).<br>Since the beginning of the pandemic, both expert opinion and small-scale   |  |  |  |  |  |  |
| <ol> <li>16</li> <li>17</li> <li>18</li> <li>19</li> <li>20</li> <li>21</li> <li>22</li> <li>23</li> </ol>   | Approximately five percent of known COVID-19 cases require admission to an<br>intensive care unit (ICU) (8). Many of these patients develop acute respiratory<br>distress syndrome (ARDS) and require mechanical ventilation (MV) or other<br>respiratory support (9). Cytokine storms (a large release of pro-inflammatory<br>proteins) have been implicated in the development of ARDS, as well as multi-organ<br>failure, with a correlation between cytokine levels and mortality (10).<br>Since the beginning of the pandemic, both expert opinion and small-scale<br>observational studies have contributed to a number of drug groups being  |  |  |  |  |  |  |
| <ol> <li>16</li> <li>17</li> <li>18</li> <li>19</li> <li>20</li> <li>21</li> <li>22</li> <li>23</li> <li>24</li> </ol>                                     | Approximately five percent of known COVID-19 cases require admission to an<br>intensive care unit (ICU) (8). Many of these patients develop acute respiratory<br>distress syndrome (ARDS) and require mechanical ventilation (MV) or other<br>respiratory support (9). Cytokine storms (a large release of pro-inflammatory<br>proteins) have been implicated in the development of ARDS, as well as multi-organ<br>failure, with a correlation between cytokine levels and mortality (10).<br>Since the beginning of the pandemic, both expert opinion and small-scale<br>observational studies have contributed to a number of drug groups being  |  |  |  |  |  |  |
| <ol> <li>16</li> <li>17</li> <li>18</li> <li>19</li> <li>20</li> <li>21</li> <li>22</li> <li>23</li> <li>24</li> <li>25</li> </ol>                         | Approximately five percent of known COVID-19 cases require admission to an<br>intensive care unit (ICU) (8). Many of these patients develop acute respiratory<br>distress syndrome (ARDS) and require mechanical ventilation (MV) or other<br>respiratory support (9). Cytokine storms (a large release of pro-inflammatory<br>proteins) have been implicated in the development of ARDS, as well as multi-organ<br>failure, with a correlation between cytokine levels and mortality (10).<br>Since the beginning of the pandemic, both expert opinion and small-scale<br>observational studies have contributed to a number of drug groups being<br>postulated to alter the progression of the disease, whether positively or negatively.   |  |  |  |  |  |  |
| <ol> <li>16</li> <li>17</li> <li>18</li> <li>19</li> <li>20</li> <li>21</li> <li>22</li> <li>23</li> <li>24</li> <li>25</li> </ol>                         | Approximately five percent of known COVID-19 cases require admission to an<br>intensive care unit (ICU) (8). Many of these patients develop acute respiratory<br>distress syndrome (ARDS) and require mechanical ventilation (MV) or other<br>respiratory support (9). Cytokine storms (a large release of pro-inflammatory<br>proteins) have been implicated in the development of ARDS, as well as multi-organ<br>failure, with a correlation between cytokine levels and mortality (10).<br>Since the beginning of the pandemic, both expert opinion and small-scale<br>observational studies have contributed to a number of drug groups being<br>postulated to alter the progression of the disease, whether positively or negatively.<br>At various times, conflicting advice regarding non-steroidal anti-inflammatory   |  |  |  |  |  |  |
| <ol> <li>16</li> <li>17</li> <li>18</li> <li>19</li> <li>20</li> <li>21</li> <li>21</li> <li>22</li> <li>23</li> <li>24</li> <li>25</li> <li>26</li> </ol> | Approximately five percent of known COVID-19 cases require admission to an<br>intensive care unit (ICU) (8). Many of these patients develop acute respiratory<br>distress syndrome (ARDS) and require mechanical ventilation (MV) or other<br>respiratory support (9). Cytokine storms (a large release of pro-inflammatory<br>proteins) have been implicated in the development of ARDS, as well as multi-organ<br>failure, with a correlation between cytokine levels and mortality (10).<br>Since the beginning of the pandemic, both expert opinion and small-scale<br>observational studies have contributed to a number of drug groups being<br>postulated to alter the progression of the disease, whether positively or negatively.<br>At various times, conflicting advice regarding non-steroidal anti-inflammatory<br>drugs (NSAIDs) (11-13), corticosteroids (14, 15) and other immunosuppressants (16-   |  |  |  |  |  |  |
| <ol> <li>16</li> <li>17</li> <li>18</li> <li>19</li> <li>20</li> <li>21</li> <li>22</li> <li>23</li> <li>24</li> <li>25</li> <li>26</li> <li>27</li> </ol> | Approximately five percent of known COVID-19 cases require admission to an<br>intensive care unit (ICU) (8). Many of these patients develop acute respiratory<br>distress syndrome (ARDS) and require mechanical ventilation (MV) or other<br>respiratory support (9). Cytokine storms (a large release of pro-inflammatory<br>proteins) have been implicated in the development of ARDS, as well as multi-organ<br>failure, with a correlation between cytokine levels and mortality (10).<br>Since the beginning of the pandemic, both expert opinion and small-scale<br>observational studies have contributed to a number of drug groups being<br>postulated to alter the progression of the disease, whether positively or negatively.<br>At various times, conflicting advice regarding non-steroidal anti-inflammatory<br>drugs (NSAIDs) (11-13), corticosteroids (14, 15) and other immunosuppressants (16-<br>18), ACE Inhibitors (ACEIs) and Angiotensin Receptor Blockers (ARBs) (19) have |  |  |  |  |  |  |

- 1 been reported in the scientific literature and media, leading to considerable
- 2 confusion for both patients and healthcare professionals.
- 3 Especially during the early stages of the pandemic, most of the advice stemmed
- 4 from outcomes in other respiratory infections including severe acute respiratory
- 5 syndrome (SARS) (20, 21) and Middle East respiratory syndrome (MERS) (14) or
- 6 hypotheses based on the mechanism of viral entry into cells (19, 22, 23) and general
- 7 effects on immunity (17, 18). In addition, many scientific papers have been
- 8 published prior to peer review to speed up information sharing, which has led to
- 9 questions surrounding the scientific validity of some of these findings (24).
- 10 We aimed to identify, critically appraise and synthesise the evidence on drugs
- 11 which may be deleterious in COVID-19.

## 12 Methods

- 13 This study is reported as per PRISMA guidelines on the reporting of systematic
- 14 reviews (25). In response to the exponential rate at which new research on this topic
- 15 is being published, a rapid review methodology was adopted following the
- 16 Cochrane rapid review protocol (26) with the aim of producing reliable results in a
- 17 timely fashion.

### **18** Search strategy

19 As the availability of studies increased, our search strategy evolved iteratively to 20 capture as much as possible of the available evidence base. This study consisted of a 21 three-stage search strategy involving literature from four databases using the 22 PECOS format (Population, Exposure, Comparison, Outcome, Study design): Embase, Medline, World Health Organization (WHO) COVID-19 Database (27) and 23 24 National Institutes for Health (NIH) iSearch COVID-19 Portfolio (28). The searches 25 were performed in April 2020 and included all papers between December 2019 and 26 April 2020. We first searched the Medline and Embase databases simultaneously 27 through Ovid, using a combination of Medical Subject Headings (MESH) and

- 1 keywords, to capture papers related to the drugs already identified as potentially
- 2 harmful in exploratory searches, as well as a string to identify new drugs.
- 3 (Appendix 1: search terms). The primary goal of this search was to amass a list of
- 4 drugs hypothesized as potentially harmful in COVID-19 in order to inform the
- 5 subsequent searches.
- 6 The second search was performed on the WHO COVID-19 Database (27), a database
- 7 aimed at compiling all the available publications on COVID-19. This search used
- 8 terms to capture evidence related to all drugs identified prior to the second search.
- 9 Due to the rapidity of emerging evidence, we felt we could not rely solely on peer-
- 10 reviewed publications as this would severely limit our findings and we decided to
- 11 include preprints a priori, which at the time were the primary source of information
- 12 available on COVID-19 epidemiology and pharmacovigilance (29). The third and
- 13 final search was, therefore, undertaken for preprints on the NIH iSearch COVID-19
- 14 Portfolio database, which included Research Square, medRxiv, chemRxiv, arXiv,
- bioRxiv and Social Science Research Network. This search was screened for primarydata only.

### 17 Screening

As per the Cochrane rapid review guidelines (26), all abstracts were screened by the first reviewer (MH). A second reviewer screened any excluded papers (SM and CH), while a third reviewer helped to settle any disagreements (UP). All studies included were then screened as a full text in the same manner, with all included full texts being moved on to extraction. Finally, this process was repeated for reference lists from all relevant systematic reviews identified from the searches.

24 The first two searches were initially screened with the aim of capturing a

- 25 comprehensive list of drugs hypothesized to be harmful and therefore included
- 26 opinion pieces. These searches were then screened a second time for papers with
- 27 primary data using a second set of inclusion and exclusion criteria (Appendix 2:

- 1 inclusion & exclusion criteria). The third search was screened only for primary data
- 2 using the second set of criteria.

### 3 Data extraction

- 4 Data extraction was performed using a tailored data extraction template designed to
- 5 capture relevant data pertaining to study design, setting, demographics and
- 6 findings (Appendix 5). Data from each study were extracted by the first reviewer
- 7 (MH) and checked independently by a second reviewer (UP).

### 8 **Quality assessment**

- 9 Quality assessment was performed using the National Heart, Lung, and Blood
- 10 Institute (NHLBI) checklists (30). We chose this tool for three reasons: it is a
- 11 validated and reliable tool; it contains checklists to assess a large range of
- 12 observational study designs;(13) and it is relatively simple and quick to administer
- 13 in comparison to other quality assessment tools, which aligned with the rationale
- 14 for adopting a rapid review methodology. Two of the four reviewers (MH and
- 15 SM/CH/UP) checked the quality of each study against the relevant checklist and
- 16 reached consensus through discussion when in disagreement.
- 17 No formal assessment of strength of evidence by outcome was undertaken;
- 18 however, we considered our findings in line with the key aspects of the grading of
- 19 recommendations assessment, development and evaluation (GRADE) (31).

### 20 Outcome definitions

- 21 Drug group analysis was split into three categories: susceptibility to infection;
- 22 severity of disease; and mortality. Susceptibility was defined as testing positive,
- 23 clinical diagnosis or hospitalisation (if compared with untested individuals) as these
- 24 all compared case numbers against those without the disease. Severity of disease
- 25 was split further where possible into either a diagnostic index or level of care.
- 26 Diagnostic index of severity was defined as any combination of factors used to

1 differentiate severe or critical disease from mild or moderate, or diagnostic criteria

- 2 for ARDS. Level of care was defined as hospitalisation (if compared with
- 3 community care of COVID-19 cases), ventilation (mechanical or non-invasive
- 4 ventilation), ICU care or mortality in combination with other outcomes including
- 5 remaining as an inpatient. Mortality was defined as any measure of mortality
- 6 outside of or within hospital, either within a set time frame or at any point during
- 7 data collection.
- 8 Although hospitalisation rates would normally imply more severe disease than
- 9 simply testing positive, in many countries testing was largely only available in
- 10 hospitals (32-34). If a study compared those admitted to hospital due to COVID-19
- 11 with untested individuals, this was considered analogous with testing positive for
- 12 the purpose of analysis. However, if the control group contained positive cases
- 13 tested in the community, then this was considered as requiring a higher level of
- 14 care.
- 15 Due to the common pathway affected by both ACEIs and ARBs, and the
- 16 recommendation against co-prescription (35), these drug groups were combined in
- 17 analyses. Henceforth, when combined, ACEIs and ARBs will be referred to as renin-
- 18 angiotensin-aldosterone blockers (RAASBs).
- For the purposes of this review, we considered immunosuppressants as a class of drugs used primarily to suppress the human immune system. Most included studies investigated immunosuppressants in this manner, although some studies analysed individual immunosuppressants. Where corticosteroids were grouped
- 23 together with other immunosuppressants, these studies were included here.

### 24 Narrative synthesis

- 25 For groups of outcomes for a particular drug group that had fewer than four effect
- 26 estimates with calculable and relevant odds ratios (ORs), these were synthesised

- 1 narratively. Study description tables were used to summarise study characteristics
- 2 and compare findings.

### 3 Statistical analysis

4 Any drug groups with a sufficient number of comparable studies were synthesised 5 using meta-analyses. Considering the implications of random-effects models on 6 statistical power (36), outcomes were deemed eligible for meta-analysis if more than 7 four studies were identified that investigated the same drug group. When sufficient 8 numbers of results were available, these were then run with sub-group analyses to 9 attempt to address any heterogeneity in the results and investigate which 10 confounding factors may have affected findings. The subgroup analyses included 11 study design, level of adjustment for confounders, acute doses or regular users, the 12 two subsets of disease severity defined above and whether ACEIs and ARBs were 13 analysed together or separately. Additionally, preprint studies that had not yet been 14 formally peer-reviewed were also removed during sensitivity analyses.

15 We included studies that either reported an OR, relative risk (RR) or the raw data

16 that allowed the calculation of OR in the meta-analyses. If available, we used an

17 adjusted OR, otherwise we used an unadjusted OR, either reported in the paper or

18 manually calculated. If the calculated OR exhibited a different result from the

19 conclusion reached by the paper, this was not included in the results.

If more than one outcome was reported, these were all included if they were in different categories (i.e. severe disease and mortality). If multiple outcome subsets were reported within the same categories, the outcome with the highest number of patients in the exposed group, and therefore greatest weight, was used. The only deviation from this was within the severity of disease category, where the diagnostic index would take precedence to reduce any potential selection bias for increased level of care.

- 1 We used MetaXL (Version 5.3; EpiGear International Pty Ltd) to run random-effects
- 2 meta-analyses and create forest plots and funnel plots. Results were reported with a
- 3 95% confidence interval (CI). We used Cochrane's Q and I<sup>2</sup> test to assess for
- 4 heterogeneity. An I<sup>2</sup> <30% was considered low heterogeneity, ≥30% but <50% was
- 5 considered moderate and ≥50% was considered high heterogeneity (37). Sensitivity
- 6 analyses were performed on all models by removing each study to assess their risk
- 7 of bias on the pooled effect estimates and heterogeneity. Publication bias was
- 8 assessed for any meta-analyses that contained 10 or more studies by visually
- 9 inspecting funnel plots for asymmetry.

10

# 1 Results

## 2 Eligible studies identified:

- 3 From the first two searches, we identified eight drug groups that were hypothesized
- 4 to be deleterious in COVID-19 from 178 peer-reviewed publications comprising
- 5 ACEIs, ARBs, corticosteroids, Immunosuppressants, mineralocorticoid receptor
- 6 antagonists (MCRAs), NSAIDs, Statins and Thiazolidinediones (TZDs).
- 7 From the combined total of papers from all three searches, we identified 63 papers
- 8 with primary data related to these drugs, with 49 measuring at least one primary
- 9 outcome included in the study (Fig 1).



1

2 Fig 1. PRISMA Flow Chart. PRISMA - Preferred Reporting Items for Systematic Reviews 3 and Meta-Analyses

#### **Study characteristics:** 4

5 51% of the papers identified were from China (n=25). The remainder were either

- 6 from Europe (Denmark, France, Italy, Spain and United Kingdom (UK)), North
- 7 America (United States of America (USA)) or other parts of Asia (South Korea)

- 1 (Table 1). The majority of papers were cohort studies (n=29) and the remainder were
- 2 either cross-sectional studies (n=9), case-control studies (n=8), case series (n=2) or a
- 3 hybrid design (n=1). There were no experimental or quasi-experimental studies
- 4 included in this review. Thirteen studies investigated more than one relevant drug,
- 5 with one study (38) presenting data for all eight drug groups.
- 6 Table 1. Geographical Demographics of Included Studies and Peer-Review Status

|             | Number of      |  |  |  |
|-------------|----------------|--|--|--|
| Study       | studies (peer- |  |  |  |
| Country     | reviewed)      |  |  |  |
| China       | 25 (10)        |  |  |  |
| USA         | 9 (2)          |  |  |  |
| UK          | 4 (0)          |  |  |  |
| Italy       | 3 (2)          |  |  |  |
| Spain       | 3 (1)          |  |  |  |
| France      | 2 (0)          |  |  |  |
| South Korea | 2 (0)          |  |  |  |
| Denmark     | 1 (0)          |  |  |  |
| Total       | 49 (15)        |  |  |  |

7 USA - United States of America, UK - United Kingdom.

### 8 Exposure status and outcome measures:

9 Most studies analysed outcomes for patients taking drugs regularly prior to

10 contracting COVID-19. Studies that analysed ACEI, ARB, MCRA, statin and TZD

11 usage measured outcomes only from those taking the drugs regularly, although one

12 study compared outcomes of continued use of ACEIs and ARBs after admission

- 13 with withdrawal of treatment (39). One study analysed acute treatment with the
- 14 immunosuppressant tocilizumab (40), and one analysed acute treatment with the
- 15 NSAID celecoxib (41). All the remaining studies, which looked at the use of drugs
- 16 for treating COVID-19 focussed on corticosteroids. There was a mix of studies

- 1 assessing acute treatment doses of corticosteroids and those taking long-term
- 2 corticosteroids on severity of disease. All studies analysing susceptibility to
- 3 infection with corticosteroids assessed those taking long-term steroids and all
- 4 studies analysing mortality assessed those receiving acute doses.

### 5 **Quality assessment:**

- 6 The majority of the studies were assessed as being of fair quality (45%), followed by
- 7 good quality (35%); the remaining studies were assessed to be of high risk of bias
- 8 and rated as poor quality (20%). Generally, studies were downgraded due to limited
- 9 justifications of sample sizes used and insufficient controlling for potential
- 10 confounders.
- 11 Sensitivity analysis by publication status:

As a considerable portion of the included studies were published as preprints that had not formally been peer-reviewed, we undertook sensitivity analysis on any of these studies eligible for meta-analysis to determine their influence on the overall effect estimates when included in random-effects models. None of the included preprints were found to significantly influence any of these results, described below, and they were therefore included in the analyses. Results of the sensitivity

18 analyses can be viewed in Appendix 7.

### 19 MCRAs, NSAIDs, Statins and TZDs

For four of the drug groups (MCRAs, NSAIDs, statins and TZDs), meta-analysis
revealed no evidence of protection or harm for any of the groups of outcomes. None
of these drug groups had any studies analysing mortality, and only NSAIDs had
enough studies analysing severity of disease to perform a meta-analysis (Appendix
6 – Fig 14). No pooled effect estimates showed any statistically significant evidence
of harm in patients with COVID-19 (Table 2) (Appendix 6 – Figs 15-17).

26

#### 1 Table 2. Summary of results for MCRAs, NSAIDs, statins and TZDs

| Drug    | Studies | Susceptibility | Pooled     | Severity | Pooled     | Mortality | Pooled |
|---------|---------|----------------|------------|----------|------------|-----------|--------|
|         | (n)     | (n)            | effect     | (n)      | effect     | (n)       | effect |
| NSAIDs  | 9       | 5              | OR 1.04    | 6        | OR 0.90    | -         | -      |
|         |         |                | (95% CI    |          | (95% CI    |           |        |
|         |         |                | 0.91-1.18) |          | 0.66-1.22) |           |        |
| Statins | 5       | 4              | OR 1.04    | 2        | -          | -         | -      |
|         |         |                | (95% CI    |          |            |           |        |
|         |         |                | 0.87-1.25) |          |            |           |        |
| TZDs    | 4       | 4              | OR 1.04    | 1        | -          | -         | -      |
|         |         |                | (95% CI    |          |            |           |        |
|         |         |                | 0.61-1.79) |          |            |           |        |
| MCRAs   | 2       | 2              | -          | -        | -          | -         | -      |
|         |         |                |            |          |            |           |        |

2 NSAIDs - non-steroidal anti-inflammatory drugs, TZDs - thiazolidinediones, MCRAs -3 mineralocorticoid receptor antagonists. Effects of drug groups on susceptibility

4 Results of studies looking into the effect of drug groups on the susceptibility to

5 COVID-19 infection are summarised in Appendix 3 - Table 3.

#### RAASB 6

7 Ten studies were used in the meta-analysis of susceptibility to COVID-19 with

8 RAASBs (33, 38, 42-49). Seven stratified their results into those taking ACEIs and

9 those taking ARBs, and three combined the two drug groups (Fig 2).

10 The total pooled effect estimate from all the studies showed no evidence that

RAASBs affect the risk of contracting COVID-19 (OR 0.94; 95% CI 0.82-1.07). Visual 11

12 inspection of the funnel plot showed no evidence of publication bias (Appendix 4 -

13 Fig 12). When split into subgroups for drug group, study design and statistical

14 adjustment, this result was unchanged (Appendix 6 – Figs 18-20).





2 Fig 2. Forest plot for susceptibility to COVID-19 for those taking regular RAASBs.

3 ACEI – angiotensin-converting enzyme inhibitor, ARB – angiotensin-II-receptor blocker, OR

- 4 - odds ratio, CI - confidence interval.
- 5 Corticosteroids

6 Four studies investigated the effects of corticosteroids on susceptibility to COVID-19

- 7 (38, 44, 48, 49). One study, rated as good quality, measured only those taking
- 8 inhaled corticosteroids (38), therefore meta-analysis was not conducted. This study
- 9 was also the only one to report an increased risk of contracting COVID-19 amongst
- 10 patients taking corticosteroids, with the remaining three studies reporting null
- 11 effects (Appendix 3 – Table 3).
- 12 *Immunosuppressants*
- 13 Five studies (32, 38, 44, 46, 48) analysing the effect of immunosuppressants on
- 14 susceptibility to COVID-19 were included in the meta-analysis (Fig 3).
- 15 The initial analysis showed no statistically significant evidence of harm or benefit
- (OR 1.11; 95% CI 0.86 1.43). Sensitivity analysis showed removal of Rentsch, CT et 16
- 17 al resulted in a statistically significant movement of the pooled effect in the direction
- 18 of harm (OR 1.29; 95% CI 1.19 - 1.40; I<sup>2</sup> 0%), but was 96.9% weighted towards the
- 19 result from Mancia, G et al. When analysed as subgroups based on study design, we

- 1 found the same result for case-control and cross-sectional studies (OR 1.29; 95% CI
- 2 1.19 1.40) (**Fig 3**).
- 3 The only study looking at susceptibility that was not included in the meta-analysis
- 4 (50) looked at individual immunosuppressive drugs compared against each other,
- 5 finding janus kinase inhibitor usage was more prevalent among rheumatology
- 6 patients with COVID-19, whereas no evidence of a protective or harmful effect of
- 7 methotrexate was observed.



- 9 Fig 3. Forest plot for effect of immunosuppressants on susceptibility to COVID-19,
- 10 stratified by study design. *OR* odds ratio, *CI* confidence interval.

### 1 Effects of drug groups on severity

- 2 Results of studies looking into the effect of drug groups on the severity of COVID-
- 3 19 infection are summarised in Appendix 3 **Table 4**.

### 4 RAASB

- 5 Among the total pool of studies, there was no evidence of benefit or harm with
- 6 RAASBs (OR 0.93; 95% CI 0.77 1.12) (Fig 4). No individual studies affected either
- 7 the pooled effect estimate or the heterogeneity. Visual inspection of the funnel plot
- 8 showed no evidence of publication bias (Appendix 4 Fig 13).



9

10 Fig 4. Forest plot of effect of RAASBs on severity of COVID-19. ACEI – angiotensin-

11 converting enzyme inhibitor, ARB – angiotensin-II-receptor blocker, OR – odds ratio, CI –

- 12 confidence interval.
- 13 Subgroup analyses consisting of three subgroups: either no or insufficient
- 14 adjustment and compared against a general population; compared against
- 15 hypertensive controls; and what we considered sufficiently adjusted results

- 1 (adjustment for at least age, sex and one or more co-morbidities) was performed.
- 2 This showed statistically significant evidence of a protective effect from ACEIs and
- 3 ARBs for adjusted results (OR 0.86; 95% CI 0.77 0.96) (Fig 5). Sensitivity analysis
- 4 showed that the removal of the result from Bean, DM et al (51) removed the
- 5 statistically significant pooled effect estimate (OR 0.90; 95% CI 0.80 1.02).



6

- 7 Fig 5. Forest plot of effect of RAASBs on severity of COVID-19 stratified by
- 8 adjustment. ACEI angiotensin-converting enzyme inhibitor, ARB angiotensin-II-receptor
- 9 blocker, OR odds ratio, CI confidence interval.

- 1 Creating diagnostic severity index or higher level of care subgroups showed
- 2 statistically non-significant and heterogeneous results (OR 0.84; 95% CI 0.56 1.27; I<sup>2</sup>
- 3 45% and OR 0.94; 95% CI 0.77 1.15; I<sup>2</sup> 62%, respectively). When these two groups
- 4 were analysed separately and subgroup analysis was run for level of adjustment,
- 5 this showed a statistically non-significant result for adjusted studies using the
- 6 severity index (OR 0.77; 95% CI 0.39 1.49). However, further statistically significant
- 7 evidence of benefit for RAASBs among adjusted studies in regard to the highest
- 8 level of care needed was observed (OR 0.85; 95% CI 0.74 0.98) (Fig 6). Again,
- 9 removal of the result from Bean, DM et al (51) made the result statistically non-
- 10 significant. This stratification also showed statistically significant evidence of harm
- 11 from unadjusted studies (OR 1.63; 95% CI 1.16 2.31).



12

- 13 Fig 6 Forest plot of effect of RAASBs on the highest level of care needed, stratified by
- 14 level of adjustment. ACEI angiotensin-converting enzyme inhibitor, ARB angiotensin-II-
- 15 receptor blocker, OR odds ratio, CI confidence interval.

- 1 Corticosteroids
- 2 Eight studies with verifiable data assessed the risk of corticosteroids increasing the
- 3 risk of developing severe disease (49, 52-58). The majority were cohort studies,
- 4 which varied by study quality and other methodological factors (Appendix 3 –
- 5 **Table 4**).
- 6 Four cohort studies measured outcomes from acute treatment doses of
- 7 corticosteroids, reaching different conclusions; however, one study did not have a
- 8 calculable OR and so a meta-analysis was not possible.
- 9 The four studies which used long-term steroids as their exposure were analysed
- 10 together in a meta-analysis. One study ran different analyses for oral and inhaled
- 11 steroids. As these can be co-prescribed, this analysis was run twice with each model
- 12 including either the inhaled or oral corticosteroid measurement. Neither model
- 13 showed statistically significant evidence of harm or benefit for severity of COVID-19
- 14 for those taking long-term steroids (inhaled corticosteroids = OR 1.40; 95% CI 0.81 –
- 15 2.39; I<sup>2</sup> 36%; oral corticosteroids = OR 1.36; 95% CI 0.76 2.45; I<sup>2</sup> 44% (**Fig** 7)).
- 16

![](_page_20_Figure_17.jpeg)

18 Fig 7. Forest plot of effect of regular corticosteroids on severity of COVID-19. OR –

- 19 odds ratio, CI confidence interval.
- 20 Immunosuppressants
- 21 Meta-analysis was conducted on four studies (46, 54, 56, 59) of severity of COVID-19
- 22 in those taking immunosuppressants. All four were cohort studies and all scored

- 1 poorly on aspects of controlling for confounding during quality assessment
- 2 (Appendix 3 **Table 4**).
- 3 The pooled effect estimate showed no statistically significant evidence of benefit or
- 4 harm (OR 0.68; 0.27 1.71) (**Fig** 8).

![](_page_21_Figure_5.jpeg)

6 Fig 8. Forest plot for effect of immunosuppressants on severity of COVID-19. OR –

7 odds ratio, CI – confidence interval, ARDS – acute respiratory distress syndrome.

- 8 The studies not included in this meta-analysis reached opposing conclusions, but in
- 9 very different scenarios. The only study to look at acute usage of tocilizumab (40)
- 10 was a case control study, finding evidence of benefit in terms of requiring
- 11 ventilation (OR 0.42: 95% CI 0.2 0.89) or ICU care (OR 0.17; 95% CI 0.06-0.48). An
- 12 additional case control study (60), found that those on immunosuppressants had
- 13 more severe COVID-19 when compared with their family members who tested
- 14 positive for SARS-CoV-2, but no OR was available nor calculable.

### 15 Effects of drug groups on mortality

- 16 Results of studies looking into the effect of drug groups on mortality in COVID-19
- 17 infection are summarised in Appendix 3 **Table 5**.
- 18 RAASB
- 19 Eight studies investigated death rates among those taking RAASBs (61-68) with six
- 20 grouping them together and two analysing ACEIs and ARBs separately (Appendix
- 21 3 **Table 5**).
- 22 None of these studies adjusted their results, although three used a hypertensive
- 23 control group. When subgroup analysis was undertaken to reflect these, it showed

- 1 no evidence of harm or benefit against hypertensive controls (OR 0.91; 95% CI 0.62 –
- 2 1.32), the general population (OR 1.11; 95% CI 0.67 1.85) or for the pooled effect
- 3 overall (OR 0.97; 95% CI 0.74 1.29) (Fig 9).

![](_page_22_Figure_4.jpeg)

4

Fig 9. Forest plot of effect of RAASBs on mortality, stratified by control group. ACEI –
angiotensin-converting enzyme inhibitor, ARB – angiotensin-II-receptor blocker, OR – odds
ratio, CI – confidence interval.

- 8 Subgroup analysis of studies which combined RAASBs showed a pooled effect
- 9 estimate exhibiting statistically significant evidence of benefit (OR 0.68; 95% CI 0.55
- 10 0.85) with low heterogeneity (I<sup>2</sup> 5%) (Fig 10). During sensitivity analysis, the result
- 11 became statistically non-significant if the results from either Ip, A et al or Sánchez-
- 12 Álvarez, JE et al were removed. These two studies were also the only two in this
- 13 analysis that were rated as poor during the quality assessment.

perpetuity. It is made available under a CC-BY-NC 4.0 International license .

![](_page_23_Figure_2.jpeg)

1

### 2 Fig 10. Forest plot for effect of RAASBs on mortality in COVID-19, stratified by drug

3 group. ACEI – angiotensin-converting enzyme inhibitor, ARB – angiotensin-II-receptor

4 blocker, OR – odds ratio, CI – confidence interval.

5 Two studies were not included in the meta-analysis, each reporting conflicting

6 results with regards to mortality (69, 70) (Appendix 3 – **Table 5**).

- 7 Corticosteroids
- 8 Seven (62, 67, 71-75) of the 10 studies assessing mortality in those given
- 9 corticosteroids were included in the meta-analysis. All of these analysed acute doses
- 10 used to treat COVID-19 admissions in hospital (Appendix 3 **Table 5**). Two studies
- 11 (55, 76) reported no evidence of harm or benefit but it was not possible to extract
- 12 summary data for pooled analysis. The remaining study (54) compared hazard
- 13 ratios (HRs) for high and low dose corticosteroids, finding a low-dose had no
- 14 evidence of either harm or benefit (HR 1.26; 95% CI 0.61 2.58), but a high-dose
- 15 showed evidence of increased mortality (HR 3.5; 95% CI 1.79 6.86).

- 1 Meta-analysis of the remaining studies showed evidence of harm (OR 2.22; 95% 1.26
- 2 3.90) (Fig 11). The authors of three of the studies highlighted confounding by
- 3 indication as a likely factor considerably affecting their results, and when these
- 4 studies were removed, the pooled effect estimate showed there was no clear
- 5 evidence of an effect (OR 1.48; 95% CI 0.73 3.00; I<sup>2</sup> 67%).

![](_page_24_Figure_6.jpeg)

7 Fig 11. Forest plot of effect of corticosteroids on mortality. OR – odds ratio, CI –

- 8 confidence interval.
- 9 Immunosuppressants
- 10 Two studies analysed mortality rates in those taking immunosuppressants. (40, 59).
- 11 Neither found statistically significant evidence of increased or decreased mortality
- 12 (OR 0.88; 95% CI 0.17 4.46 and OR 0.90; 95% CI 0.24 3.42, respectively)
- 13 (Appendix 3 **Table 5**).
- 14
- 15
- 16

## 1 **Discussion**

## 2 Statement of principal findings

3 This rapid systematic review and meta-analysis attempted to identify and assess 4 any potentially deleterious drug groups in covid-19 susceptibility and prognosis. In 5 terms of susceptibility to infection, we found no pooled effect estimates and one 6 subgroup analysis with statistically significant evidence of increased susceptibility 7 (Fig 3), which would be in keeping with many other infections and a known risk of 8 immunosuppressant use (77). Hence, we found no evidence to suggest that those 9 without COVID-19 should stop taking their medication to reduce their risk of 10 contracting the disease. Stopping immunosuppressants may also result in a flare-up 11 of a person's underlying condition, which can result in increased risk of infection 12 (78). Furthermore, the withdrawal of any necessary medications could cause harm 13 from the disease the drug was being used to treat, as well as from a potentially 14 severe form of COVID-19 for people with hypertension and diabetes who are 15 already at increased risk (79, 80).

16 We found no evidence that long-term use of any of these drugs increases the 17 severity of disease, nor the mortality rates. One subgroup analysis of RAASBs 18 showed a statistically significant relationship between RAASB usage and requiring 19 a higher level of hospital care (Fig 6). However, this was found in an unadjusted 20 subgroup where the adjusted subgroup showed evidence of protection, highlighting 21 the effect of known confounders, such as age and cardiovascular disease (81), on the 22 results from these observational studies. This evidence of protection against more 23 severe disease was also shown by the pooled effect of all adjusted studies analysing 24 severity (Fig 5).

### 25 Strengths and limitations

26 This rapid systematic review and meta-analysis has synthesised the best available 27 evidence from the first few months of the pandemic for each of these drugs when

1 taken by those with or at risk of COVID-19. This study has, within the limitations

2 imposed by time and the unparalleled research publication rate, aimed to compile a

3 comprehensive list of drugs that were hypothesized to be deleterious in peer-

4 reviewed papers at the time of the searches.

5 This review however has several limitations. The rapidity with which research is 6 being produced and published affected our study in a number of ways. Firstly, we 7 acknowledge that while a rapid review is justifiable on the grounds of swifter 8 completion, it is less robust than a systematic review. Secondly, the studies included 9 in this review were also completed within a short space of time in order to attempt 10 to learn from clinical practice as quickly as possible. As a result of this, many papers 11 that would be relevant to this review, including papers with stronger evidence and 12 more robust methodology than some of those included may now be available. 13 Finally, the research landscape is changing so quickly that the studies included in 14 this review were all published prior to conducting the last search. This will bias our 15 results towards the studies that were produced more quickly and in countries that experienced larger COVID-19 case numbers earlier on in this outbreak. These 16 countries may have large differences in their demographics as well as clinical 17 18 protocols for both pre-existing co-morbidities and COVID-19, which could impact 19 the generalisability of our findings.

20 Another factor heightening the risk of bias, errors and methodological inadequacies, 21 is the lack of peer-reviewed articles and the use of preprints in this review. We 22 conducted a sensitivity analysis in January 2021 to analyse whether the removal of 23 papers that had not made it through the peer-review process at this point affected 24 the results, but there were no significant changes to the pooled outcomes (Appendix 25 7). Nevertheless, we must emphasise that because we have included preprints, the 26 results from this review are only to highlight potential outcomes to evaluate with 27 further studies and should not be used to alter clinical practice.

1 As all of the evidence in this review is from observational studies, this introduces a

- 2 risk of bias from confounding factors, despite many studies attempting to
- 3 statistically adjust for these. Any confounders not adjusted for or as yet unknown
- 4 will have affected our results. One example of this is the discovery that people with
- 5 a Black, Asian or minority ethnic background are at an increased risk of severe
- 6 COVID-19 (82), something that was not apparent at the beginning of the outbreak
- 7 and was therefore only adjusted for in five of the studies included in the review.
- 8 Other biases evident in the studies included confounding by indication, most
- 9 evident among the studies looking at acute treatment of COVID-19 with
- 10 corticosteroids, and misclassification bias in studies where it was possible the
- 11 control group had been buying the drug of interest over-the-counter or had stopped
- 12 taking it immediately prior to admission to hospital. This happened with NSAIDS
- 13 for example.
- 14 Using aspects of the GRADE assessment, the quality of evidence presented in this
- 15 review could be classified as low, primarily due to inconsistency, indirectness,
- 16 imprecision and risk of bias across each outcome (31).

### 17 Interpretation in light of the wider literature

18 One of the reasons RAASBs, MCRAs, NSAIDs, Statins and TZDs were hypothesized 19 to be harmful is their potentially upregulating effect on ACE2, this being the entry 20 point into cells for SARS-CoV-2. Although there are studies which have shown that 21 all of these drugs may upregulate ACE2, particularly RAASBs, the evidence is 22 largely from animal studies, and some of it is contradictory (83). We found no 23 evidence to suggest any of these drugs worsen outcomes or increase the 24 susceptibility to contracting COVID-19 and some evidence to suggest RAASBs 25 could be protective. Whilst it is unclear if this is related to their effect on ACE2, it 26 appears that the protective nature of ACE2 may in fact provide a target for treating 27 COVID-19 (84). Studies have also shown similar protective effects of ACEIs (85), 28 ARBs and statins (86) in non-COVID pneumonias.

1 Another prominent reason for drugs to be hypothesized to worsen outcomes was 2 immunosuppression. Advice emerged early on advising against using steroids in 3 COVID-19 (14), but this was at odds with the experiences of front-line clinical 4 workers in China who were using it in a large number of cases (87). Guidance 5 surrounding long-term immunosuppressants in those with COVID-19 was also 6 issued (88), however at the same time it appeared that dampening the immune 7 response may have a role in treating acute infections (89). With the discovery that 8 interleukin 6 (IL-6) was elevated in patients with severe COVID-19 (90), IL-6 9 antagonists such as tocilizumab started to be used therapeutically (91). We only 10 found one study which analysed the outcomes of those treated with tocilizumab 11 and this found a reduction of ICU admissions and need for ventilation when 12 patients were given tocilizumab (40). This was an observational study where the 13 patients were "highly selected" so, although the results were adjusted for 14 confounders, there remains a risk of bias. 15 The majority of studies included in this review measured outcomes in patients

16 taking drugs regularly. We only found one study (39) comparing patients who had 17 the drug withheld during the acute infection with those that continued to take them. 18 This study found no effect on disease severity or mortality but found statistically 19 significant reduction in viral clearance and length of stay in those in whom RAASBs 20 were withheld during their hospital stay. This area requires more research, as any 21 conclusions we draw from this review can only be applied to those who take these 22 medications prior to the onset of COVID-19. In order to guide clinical practice, 23 studies comparing initiated, continued and withheld medications, ideally 24 randomised controlled trials (RCTs), are paramount (92). 25 With most drug groups included in this review, dosage and timing was not usually

26 measured or analysed. One study of corticosteroids compared the effect of the 27 cumulative dose of steroids and the authors concluded that mortality with high-28 dose steroids was higher than with low-dose steroids (54). Another study which 29 found favourable outcomes for corticosteroids as a treatment for COVID-19

- 1 hypothesized that the timing was also important, opting for low dose steroids early
- 2 on in the disease progression (93). These hypotheses were similarly at risk of
- 3 confounding by indication, with those going on to develop ARDS more likely to get
- 4 higher dose steroids at a later point in time.

### 5 Implications for policy, practice and research

- 6 This review contains exclusively observational studies, many of which have not
- 7 been peer-reviewed, and so we do not recommend any changes to normal
- 8 prescribing practice. However, we have found no evidence to suggest medications
- 9 should be stopped solely due to contracting COVID-19. This goes along with
- 10 guidelines published for many of these drug groups (94-96) and the findings of
- 11 other rapid reviews (28, 97).
- 12 For those taking the drugs highlighted in this review, the binary decision between
- 13 stopping and continuing their medications, coupled with the contrasting advice
- 14 from experts, the media and governments, helped fuel the false dichotomy that
- 15 drug groups were either harmful or not. Due to the large number of factors that can
- 16 affect outcomes in COVID-19, the harm or benefit derived from starting, stopping or
- 17 continuing drugs will affect individuals differently, with co-morbidities,
- 18 demographics, other medications and individual response to the virus playing a
- 19 role in the progression of the disease. However, this review raises the prospect that
- 20 drug groups that have been hypothesized to be deleterious in COVID-19, may have
- 21 the potential to be beneficial in certain circumstances. This is evidenced further by a
- 22 number of clinical trials, either planned, in progress or completed, involving drugs
- 23 or drug groups identified during this review (98-103).
- 24 Furthermore, where trials are unfeasible, as in the case where the aim is to examine
- 25 the effects of long-term drug use on COVID-19, good quality, large-scale
- 26 observational studies will be necessary to form the best possible evidence. RCTs will
- 27 understandably be focussed on finding drugs used to treat COVID-19, the results of

- 1 which are beginning to be published (98). However, trials looking at treatments
- 2 with drugs included in this review may help inform their use in other contexts.

### 3 Conclusions

- 4 We have highlighted the gaps in the research, such as the lack of RCTs and
- 5 mortality data for NSAIDs, immunosuppressants and long-term steroid use. We
- 6 have proposed that there should be some focus in future research on these gaps as
- 7 well as the use of ACE2, or drugs which upregulate it, as a potential target for
- 8 treating COVID-19.
- 9 This study found a total of eight drug groups hypothesized to be deleterious in
- 10 COVID-19. The available data from the first few months of the current pandemic
- 11 suggest that there is little to no evidence these drug groups increase susceptibility,
- 12 severity or mortality in COVID-19 and we found some evidence that ACEIs and
- 13 ARBs may be protective in preventing a more severe disease.

## 14 **References**

15 1. Johns Hopkins University. Coronavirus Resource Centre 2020 [cited 2020 3 16 April]. Available from: <u>https://coronavirus.jhu.edu/</u>. 17 2. Centers for Disease Control and Prevention. Human Coronavirus Types 2020 18 [cited 2020 Jul 11]. Available from: https://www.cdc.gov/coronavirus/types.html. 19 European Centre For Disease Prevention and Control. Coronaviruses 2020 3. 20 [cited 2020 Jul 11]. Available from: https://www.ecdc.europa.eu/en/covid-19/latest-21 evidence/coronaviruses. 22 4. Lu R, Zhao X, Li J, Niu P, Yang B, Wu H, et al. Genomic characterisation and 23 epidemiology of 2019 novel coronavirus: implications for virus origins and receptor 24 binding. Lancet. 2020;395(10224):565-74. 25 5. Wu F, Zhao S, Yu B, Chen YM, Wang W, Song ZG, et al. A new coronavirus 26 associated with human respiratory disease in China. Nature. 2020;579(7798):265-9. 27 6. Li X, Molina-Molina M, Abdul-Hafez A, Uhal V, Xaubet A, Uhal BD. 28 Angiotensin converting enzyme-2 is protective but downregulated in human and 29 experimental lung fibrosis. Am J Physiol Lung Cell Mol Physiol. 2008;295(1):L178-30 85. 31 7. Fang L, Karakiulakis G, Roth M. Are patients with hypertension and diabetes 32 mellitus at increased risk for COVID-19 infection? The Lancet Respiratory Medicine.

33 2020;8(4):e21.

- perpetuity. It is made available under a CC-BY-NC 4.0 International license .
- 1 8. Wu Z, McGoogan JM. Characteristics of and Important Lessons From the
- 2 Coronavirus Disease 2019 (COVID-19) Outbreak in China: Summary of a Report of
- 3 72314 Cases From the Chinese Center for Disease Control and Prevention. JAMA.
- 4 2020;323(13):1239-42.
- 5 9. Murthy S, Gomersall CD, Fowler RA. Care for Critically Ill Patients With
- 6 COVID-19. JAMA. 2020;323(15):1499-500.
- 7 10. Ye Q, Wang B, Mao J. The pathogenesis and treatment of the 'Cytokine Storm'
- 8 in COVID-19. J Infect. 2020;80(6):607-13.
- 9 11. Medicines and Healthcare Regulatory Agency. Ibuprofen use and
- 10 Coronavirus (COVID-19) 2020 [cited 2020 2 April]. Available from:
- 11 https://www.gov.uk/government/news/ibuprofen-use-and-covid19coronavirus.
- 12 12. Day M. Covid-19: European drugs agency to review safety of ibuprofen. BMJ.
- 13 2020;368:m1168.
- 14 13. Reality Check Team, BBC Monitoring. Coronavirus and ibuprofen: Separating
- 15 fact from fiction: BBC News,; 2020 [cited 2020 Jul 14]. Available from:
- 16 https://www.bbc.co.uk/news/51929628.
- 17 14. Russell CD, Millar JE, Baillie JK. Clinical evidence does not support
- corticosteroid treatment for 2019-nCoV lung injury. The Lancet. 2020;395(10223):473-5.
- 20 15. Hartmann-Boyce J, Hobbs R. Inhaled Steroids in Asthma during the COVID-
- 21 19 Outbreak: CEBM; 2020 [cited 2020 Jul 14]. Available from:
- https://www.cebm.net/covid-19/inhaled-steroids-in-asthma-during-the-covid-19 outbreak/.
- 24 16. Kovarik JM, Burtin P. Immunosuppressants in advanced clinical development
- 25 for organ transplantation and selected autoimmune diseases. Expert Opin Emerg
- 26 Drugs. 2003;8(1):47-62.
- 27 17. Manez R, Kusne S, Linden P, Gonzalez-Pinto I, Bonet H, Kramer D, et al.
- 28 Temporary withdrawal of immunosuppression for life-threatening infections after
- 29 liver transplantation. Transplantation. 1994;57(1):149-51.
- 30 18. Conforti C, Giuffrida R, Dianzani C, Di Meo N, Zalaudek I. COVID-19 and
- psoriasis: Is it time to limit treatment with immunosuppressants? A call for action.Dermatologic Therapy. 2020.
- 33 19. Ferrario CM, Jessup J, Chappell MC, Averill DB, Brosnihan KB, Tallant EA, et
- 34 al. Effect of angiotensin-converting enzyme inhibition and angiotensin II receptor
- 35 blockers on cardiac angiotensin-converting enzyme 2. Circulation.
- 36 2005;111(20):2605-10.
- 20. Day M. Covid-19: ibuprofen should not be used for managing symptoms, say
  doctors and scientists. BMJ. 2020;368:m1086.
- 39 21. Little P. Non-steroidal anti-inflammatory drugs and covid-19. BMJ.
- 40 2020;368:m1185.
- 41 22. Aronson J, Ferner RE. Angiotensin converting enzyme (ACE) inhibitors and
- 42 angiotensin receptor blockers in COVID-19: CEBM; 2020 [cited 2020 Jul 14].
- 43 Available from: <u>https://www.cebm.net/covid-19/angiotensin-converting-enzyme-</u>
- 44 <u>ace-inhibitors-and-angiotensin-receptor-blockers-in-covid-19/</u>.

perpetuity. It is made available under a CC-BY-NC 4.0 International license .

- 1 23. Cure E, Cumhur Cure M. Angiotensin-converting enzyme inhibitors and
- 2 angiotensin receptor blockers may be harmful in patients with diabetes during
- 3 COVID-19 pandemic. Diabetes Metab Syndr. 2020;14(4):349-50.
- 4 24. Sharm MS, S; Kelland, K. Coronavirus and the risks of 'speed science': World
- 5 Economic Forum; 2020 [cited 2020 2 April]. Available from:
- 6 <u>https://www.weforum.org/agenda/2020/03/speed-science-coronavirus-covid19-</u>
- 7 <u>research-academic</u>.
- 8 25. Moher D, Liberati A, Tetzlaff J, Altman DG, Group P. Preferred reporting
- 9 items for systematic reviews and meta-analyses: the PRISMA statement. PLoS Med.
- 10 2009;6(7):e1000097.
- 11 26. Garritty C, Gartlehner G, Kamel C, King V, Nussbaumer-Streit B, Stevens A, et
- al. Cochrane Rapid Reviews. Interim Guidance from the Cochrane Rapid ReviewsMethods Group. 2020.
- 14 27. World Health Organization. Global research on coronavirus disease (COVID-
- 15 19) 2020 [cited 2020 2 April]. Available from:
- 16 https://www.who.int/emergencies/diseases/novel-coronavirus-2019/global-research-
- 17 <u>on-novel-coronavirus-2019-ncov</u>.
- 18 28. World Health Organization. COVID-19 and the use of angiotensin-converting
- 19 enzyme inhibitors and receptor blockers 2020 [cited 2020 Jul 29]. Available from:
- https://www.who.int/publications/i/item/covid-19-and-the-use-of-angiotensin converting-enzyme-inhibitors-and-receptor-blockers.
- 22 29. Gianola S, Jesus TS, Bargeri S, Castellini G. Characteristics of academic
- 23 publications, preprints, and registered clinical trials on the COVID-19 pandemic.
- 24 PLOS ONE. 2020;15(10):e0240123.
- 25 30. National Heart Lung and Blood Institute. Study Quality Assessment Tools
- 26 2014 [cited 2020 June]. Available from: <u>https://www.nhlbi.nih.gov/health-</u>
- 27 <u>topics/study-quality-assessment-tools</u>.
- 28 31. Guyatt GH, Oxman AD, Vist GE, Kunz R, Falck-Ytter Y, Alonso-Coello P, et
- al. GRADE: an emerging consensus on rating quality of evidence and strength of
   recommendations. BMJ. 2008;336(7650):924-6.
- 31 32. Gisondi P, Zaza G, Del Giglio M, Rossi M, Iacono V, Girolomoni G. Risk of
- 32 hospitalization and death from COVID-19 infection in patients with chronic plaque
- 33 psoriasis receiving a biologic treatment and renal transplant recipients in
- 34 maintenance immunosuppressive treatment. J Am Acad Dermatol. 2020;83(1):285-7.
- 35 33. Khawaja AP, Warwick AN, Hysi PG, Kastner A, Dick A, Khaw PT, et al.
- Associations with covid-19 hospitalisation amongst 406,793 adults: the UK Biobank
   prospective cohort study. medRxiv. 2020. [preprint]
- 38 34. Yan H, Valdes AM, Vijay A, Wang S, Liang L, Yang S, et al. Role of Drugs
- 39 Affecting the Renin-Angiotensin-Aldosterone System on Susceptibility and Severity
- 40 of COVID-19: A Large Case-Control Study from Zheijang Province, China.
- 41 medRxiv. 2020. [preprint]
- 42 35. National Institute for Health and Care Excellence. Renin-angiotensin system
- 43 drugs: dual therapy 2015 [cited 2020 Jul 20]. Available from:
- 44 <u>https://www.nice.org.uk/advice/ktt2/chapter/evidence-context</u>.

#### perpetuity. It is made available under a CC-BY-NC 4.0 International license .

1 Jackson D, Turner R. Power analysis for random-effects meta-analysis. Res 36. 2 Synth Methods. 2017;8(3):290-302. 3 Higgins JP, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, et al. Chapter 37. 4 10: Analysing data and undertaking meta-analyses. Cochrane handbook for 5 systematic reviews of interventions. Version 6 ed: John Wiley & Sons; 2019. 6 38. Mancia G, Rea F, Ludergnani M, Apolone G, Corrao G. Renin-Angiotensin-7 Aldosterone System Blockers and the Risk of Covid-19. N Engl J Med. 8 2020;382(25):2431-40. 9 39. Huang L, Chen Z, Ni L, Chen L, Zhou C, Gao C, et al. Impact of Angiotensin-10 converting Enzyme Inhibitors and Angiotensin Receptor Blockers on Inflammatory Responses and Viral Clearance in COVID-19 Patients: A Multicenter Retrospective 11 12 Cohort Study. Research Square. 2020. [preprint] 13 Roumier M, Paule R, Groh M, Vallee A, Ackermann F. Interleukin-6 blockade 40. 14 for severe COVID-19. medRxiv. 2020. [preprint] 15 Hong W, Chen Y, You K, Tan S, Wu F, Tao J, et al. Celebrex adjuvant therapy 41. 16 on COVID-19: An experimental study. medRxiv. 2020. [preprint] 17 Caraballo C, McCullough M, Fuery M, Chouairi F, Keating C, Ravindra N, et 42. al. COVID-19 Infections and Outcomes in a Live Registry of Heart Failure Patients 18 19 Across an Integrated Health Care System. medRxiv. 2020. [preprint] 20 Dooley H, Lee K, Freidin M, Hemani G, Roberts A, Lavigne du Cadet J, et al. 43. 21 ACE Inhibitors, ARBs and Other Anti-Hypertensive Drugs and Novel COVID-19: 22 An Association Study from the COVID Symptom Tracker App in 2,215,386 23 Individuals. SSRN Electronic Journal. 2020. [preprint] 24 Huh K, Ji W, Kang M, Hong J, Bae GH, Lee R, et al. Association of previous 44. 25 medications with the risk of COVID-19: a nationwide claims-based study from 26 South Korea. medRxiv. 2020. [preprint] 27 Kolin DA, Kulm S, Elemento O. Clinical and Genetic Characteristics of Covid-45. 28 19 Patients from UK Biobank. medRxiv. 2020. [preprint] 29 Rentsch CT, Kidwai-Khan F, Tate JP, Park LS, King JT, Skanderson M, et al. 46. 30 Covid-19 Testing, Hospital Admission, and Intensive Care Among 2,026,227 United 31 States Veterans Aged 54-75 Years. medRxiv. 2020. [preprint] 32 Reynolds HR, Adhikari S, Pulgarin C, Troxel AB, Iturrate E, Johnson SB, et al. 47. 33 Renin-Angiotensin-Aldosterone System Inhibitors and Risk of Covid-19. N Engl J 34 Med. 2020;382(25):2441-8. 35 48. Shah SJ, Barish PN, Prasad PA, Kistler AL, Neff N, Kamm J, et al. Clinical features, diagnostics, and outcomes of patients presenting with acute respiratory 36 37 illness: a comparison of patients with and without COVID-19. medRxiv. 2020. [preprint] 38 39 49. Yan H, Valdes AM, Vijay A, Wang S, Liang L, Yang S, et al. Role of Drugs Affecting the Renin-Angiotensin-Aldosterone System on Susceptibility and Severity 40 41 of COVID-19: A Large Case-Control Study from Zheijang Province, China. medRxiv. 2020. [preprint] 42 43 Michelena X, Borrell H, Lopez-Corbeto M, Lopez-Lasanta M, Moreno E, 50. 44 Pascual-Pastor M, et al. Incidence of COVID-19 in a cohort of adult and paediatric

perpetuity. It is made available under a CC-BY-NC 4.0 International license .

- 1 patients with rheumatic diseases treated with targeted biologic and synthetic
- 2 disease-modifying anti-rheumatic drugs. Semin Arthritis Rheum. 2020;50(4):564-70.
- 3 51. Bean D, Kraljevic Z, Searle T, Bendayan R, Pickles A, Folarin A, et al. ACE-
- 4 inhibitors and Angiotensin-2 Receptor Blockers are not associated with severe
- 5 SARS- COVID19 infection in a multi-site UK acute Hospital Trust. medRxiv. 2020.
- 6 [preprint]
- 7 52. Argenziano MG, Bruce SL, Slater CL, Tiao JR, Baldwin MR, Barr RG, et al.
- 8 Characterization and clinical course of 1000 Patients with COVID-19 in New York:
- 9 retrospective case series. medRxiv. 2020. [preprint]
- 10 53. Cao C, Chen M, Li Y, Yu L, Huang W, Qian G, et al. Clinical Features and
- 11 Predictors for Patients with Severe SARS-CoV-2 Pneumonia: A Retrospective
- 12 Multicenter Cohort Study. SSRN Electronic Journal. 2020. [preprint]
- 13 54. Li X, Xu S, Yu M, Wang K, Tao Y, Zhou Y, et al. Risk factors for severity and
- 14 mortality in adult COVID-19 inpatients in Wuhan. J Allergy Clin Immunol.
- 15 2020;146(1):110-8.
- 16 55. Liu Y, Sun W, Li J, Chen L, Wang Y, Zhang L, et al. Clinical features and
- 17 progression of acute respiratory distress syndrome in coronavirus disease 2019.
- 18 medRxiv. 2020. [preprint]
- 19 56. Robilotti EV, Babady NE, Mead PA, Rolling T, Perez-Johnston R, Bernardes M,
- 20 et al. Determinants of Severity in Cancer Patients with COVID-19 Illness. medRxiv.
- 21 2020. [preprint]
- 22 57. Wang D, Wang J, Jiang Q, Yang J, Li J, Gao C, et al. No Clear Benefit to the Use
- of Corticosteroid as Treatment in Adult Patients with Coronavirus Disease 2019 : A
   Retrospective Cohort Study. medRxiv. 2020. [preprint]
- 25 58. Lui GC-Y, Yip TC-F, Wong VW-S, Chow VC-Y, Ho TH-Y, Li TC-M, et al.
- 26 Adverse Clinical Outcomes in COVID-19 Versus SARS: A Territory-Wide Cohort
- 27 Study in Hong Kong. SSRN Electronic Journal. 2020. [preprint]
- 28 59. Monreal E, Maza SSdl, Gullón P, Natera-Villalba E, Chico-García JL, Beltrán-
- 29 Corbellini Á, et al. Immunosuppression is associated with a lower risk of moderate
- to severe acute respiratory distress syndrome in COVID-19. Research Square. 2020.[preprint]
- 32 60. Zhu L, Gong N, Liu B, Lu X, Chen D, Chen S, et al. Coronavirus Disease 2019
- 33 Pneumonia in Immunosuppressed Renal Transplant Recipients: A Summary of 10
- 34 Confirmed Cases in Wuhan, China. Eur Urol. 2020;77(6):748-54.
- 35 61. Benelli G, Buscarini E, Canetta C, La Piana G, Merli G, Scartabellati A, et al.
- 36 SARS-COV-2 comorbidity network and outcome in hospitalized patients in Crema,
- 37 Italy. medRxiv. 2020. [preprint]
- 38 62. Chen M, Fan Y, Wu X, Zhang L, Guo T, Deng K, et al. Clinical Characteristics
- 39 And Risk Factors For Fatal Outcome in Patients With 2019-Coronavirus Infected
- 40 Disease (COVID-19) in Wuhan, China. SSRN Electronic Journal. 2020. [preprint]
- 41 63. Guo T, Fan Y, Chen M, Wu X, Zhang L, He T, et al. Cardiovascular
- 42 Implications of Fatal Outcomes of Patients With Coronavirus Disease 2019 (COVID-
- 43 19). JAMA Cardiol. 2020.

perpetuity. It is made available under a CC-BY-NC 4.0 International license .

- 1 Ip A, Parikh K, Parrillo JE, Mathura S, Hansen E, Sawczuk IS, et al. 64.
- 2 Hypertension and Renin-Angiotensin-Aldosterone System Inhibitors in Patients
- 3 with Covid-19. medRxiv. 2020. [preprint]
- 4 Li J, Wang X, Chen J, Zhang H, Deng A. Association of Renin-Angiotensin 65.
- 5 System Inhibitors With Severity or Risk of Death in Patients With Hypertension
- 6 Hospitalized for Coronavirus Disease 2019 (COVID-19) Infection in Wuhan, China.
- 7 JAMA Cardiol. 2020.
- 8 66. Richardson S, Hirsch JS, Narasimhan M, Crawford JM, McGinn T, Davidson
- 9 KW, et al. Presenting Characteristics, Comorbidities, and Outcomes Among 5700
- 10 Patients Hospitalized With COVID-19 in the New York City Area. JAMA.
- 2020;323(20):2052-9. 11
- 12 Sánchez-Álvarez JE, Fontán MP, Martín CJ, Pelícano MB, Reina CJC, Prieto 67.
- 13 ÁMS, et al. Status of SARS-CoV-2 infection in patients on renal replacement
- 14 therapy. Report of the COVID-19 Registry of the Spanish Society of Nephrology 15
- (SEN). Nefrología (English Edition). 2020;40(3):272-8.
- 16 Yang G, Tan Z, Zhou L, Yang M, Peng L, Liu J, et al. Angiotensin II Receptor 68.
- 17 Blockers and Angiotensin-Converting Enzyme Inhibitors Usage is Associated with
- Improved Inflammatory Status and Clinical Outcomes in COVID-19 Patients With 18
- 19 Hypertension. medRxiv. 2020. [preprint]
- Lee H-Y, Ahn J, Kang CK, Won S-H, Park J-H, Kang CH, et al. Association of 20 69.
- 21 Angiotensin II Receptor Blockers and Angiotensin-Converting Enzyme Inhibitors on
- 22 COVID-19-Related Outcome. SSRN Electronic Journal. 2020. [preprint]
- 23 Zhang P, Zhu L, Cai J, Lei F, Qin J-J, Xie J, et al. Association of Inpatient Use of 70.
- 24 Angiotensin Converting Enzyme Inhibitors and Angiotensin II Receptor Blockers
- 25 with Mortality Among Patients With Hypertension Hospitalized With COVID-19.
- Circ res. 2020. 26
- 27 Cao J, Tu WJ, Cheng W, Yu L, Liu YK, Hu X, et al. Clinical Features and Short-71.
- 28 term Outcomes of 102 Patients with Coronavirus Disease 2019 in Wuhan, China.
- 29 Clin Infect Dis. 2020;71(15):748-55.
- 30 Liang M, Chen P, He M, Tang J, He X, Zhou Y, et al. Corticosteroid Treatment 72.
- 31 in Critically Ill Patients with COVID-19: A Retrospective Cohort Study. Research 32 Square. 2020. [preprint]
- 33 73. Wu C, Chen X, Cai Y, Xia J, Zhou X, Xu S, et al. Risk Factors Associated With
- 34 Acute Respiratory Distress Syndrome and Death in Patients With Coronavirus
- 35 Disease 2019 Pneumonia in Wuhan, China. JAMA Intern Med. 2020.
- 36 Zhang S, Zhao J, Wu Z, Shang Y, Zheng J, Meng M, et al. Potential Predictors 74.
- 37 for Disease Progression and Medication Evaluation of 2019 Novel Coronavirus-
- 38 Infected Pneumonia in Wuhan, China. SSRN Electronic Journal. 2020. [preprint]
- Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical course and risk factors 39 75.
- 40 for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective
- 41 cohort study. Lancet. 2020;395(10229):1054-62.
- 42 Lu X, Chen T, Wang Y, Wang J, Yan F. Adjuvant corticosteroid therapy for 76.
- 43 critically ill patients with COVID-19. Crit Care. 2020;24(1):241.
medRxiv preprint doi: https://doi.org/10.1101/2021.09.17.21262724; this version posted September 21, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

#### perpetuity. It is made available under a CC-BY-NC 4.0 International license .

- 1 77. George MP, Masur H, Norris KA, Palmer SM, Clancy CJ, McDyer JF.
- 2 Infections in the immunosuppressed host. Ann Am Thorac Soc. 2014;11 Suppl
- 3 4(Suppl 4):S211-20.
- 4 78. Ceribelli A, Motta F, De Santis M, Ansari AA, Ridgway WM, Gershwin ME, et
- 5 al. Recommendations for coronavirus infection in rheumatic diseases treated with
- 6 biologic therapy. J Autoimmun. 2020;109:102442.
- 7 79. Zhu L, She ZG, Cheng X, Qin JJ, Zhang XJ, Cai J, et al. Association of Blood
- 8 Glucose Control and Outcomes in Patients with COVID-19 and Pre-existing Type 2
- 9 Diabetes. Cell Metab. 2020;31(6):1068-77 e3.
- 10 80. Gao C, Cai Y, Zhang K, Zhou L, Zhang Y, Zhang X, et al. Association of
- 11 hypertension and antihypertensive treatment with COVID-19 mortality: a
- 12 retrospective observational study. Eur Heart J. 2020;41(22):2058-66.
- 13 81. Guan WJ, Liang WH, Zhao Y, Liang HR, Chen ZS, Li YM, et al. Comorbidity
- and its impact on 1590 patients with COVID-19 in China: a nationwide analysis. Eur
- 15 Respir J. 2020;55(5):2000547.
- 16 82. Raisi-Estabragh Z, McCracken C, Bethell MS, Cooper J, Cooper C, Caulfield
- 17 MJ, et al. Greater risk of severe COVID-19 in Black, Asian and Minority Ethnic
- 18 populations is not explained by cardiometabolic, socioeconomic or behavioural
- 19 factors, or by 25(OH)-vitamin D status: study of 1326 cases from the UK Biobank.
- 20 Journal of Public Health. 2020;42(3):451-60.
- 21 83. Dambha-Miller H, Albasri A, Hodgson S, Wilcox CR, Khan S, Islam N, et al.
- 22 Currently prescribed drugs in the UK that could upregulate or downregulate ACE2
- 23 in COVID-19 disease: a systematic review. BMJ Open. 2020;10(9):e040644.
- 24 84. Lumpuy-Castillo J, Lorenzo-Almorós A, Pello-Lázaro AM, Sánchez-Ferrer C,
- 25 Egido J, Tuñón J, et al. Cardiovascular Damage in COVID-19: Therapeutic
- Approaches Targeting the Renin-Angiotensin-Aldosterone System. Int J Mol Sci.2020;21(18).
- 28 85. Chu C, Zeng S, Hasan AA, Hocher C-F, Krämer BK, Hocher B. Comparison of
- 29 infection risks and clinical outcomes in patients with and without SARS-CoV-2 lung
- 30 infection under renin–angiotensin–aldosterone system blockade: Systematic review
- 31 and meta-analysis. British Journal of Clinical Pharmacology.n/a(n/a).
- 32 86. Mortensen E, Anzueto A. Prior use of both a statin and ARB is associated with
- 33 lower mortality for patients hospitalized with pneumonia. European Respiratory
- 34 Journal. 2016;48(suppl 60):OA3329.
- 35 87. Wang D, Hu B, Hu C, Zhu F, Liu X, Zhang J, et al. Clinical Characteristics of
- 36 138 Hospitalized Patients With 2019 Novel Coronavirus-Infected Pneumonia in
- 37 Wuhan, China. JAMA. 2020;323(11):1061-9.
- 38 88. Wang C, Rademaker M, Baker C, Foley P. COVID-19 and the use of
- 39 immunomodulatory and biologic agents for severe cutaneous disease: An
- 40 Australia/New Zealand consensus statement. The Australasian journal of
- 41 dermatology. 2020;07.
- 42 89. Mehta P, McAuley DF, Brown M, Sanchez E, Tattersall RS, Manson JJ, et al.
- 43 COVID-19: consider cytokine storm syndromes and immunosuppression. Lancet.
- 44 2020;395(10229):1033-4.

medRxiv preprint doi: https://doi.org/10.1101/2021.09.17.21262724; this version posted September 21, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

perpetuity. It is made available under a CC-BY-NC 4.0 International license .

- 1 90. Chen L, Liu HG, Liu W, Liu J, Liu K, Shang J, et al. [Analysis of clinical
- 2 features of 29 patients with 2019 novel coronavirus pneumonia]. Zhonghua Jie He
- 3 He Hu Xi Za Zhi. 2020;43(0):E005.
- 4 91. Michot JM, Albiges L, Chaput N, Saada V, Pommeret F, Griscelli F, et al.
- 5 Tocilizumab, an anti-IL-6 receptor antibody, to treat COVID-19-related respiratory
- 6 failure: a case report. Ann Oncol. 2020;31(7):961-4.
- 7 92. Gnanenthiran SR, Borghi C, Burger D, Charchar F, Poulter NR, Schlaich MP, et
- 8 al. Prospective meta-analysis protocol on randomised trials of renin-angiotensin
- 9 system inhibitors in patients with COVID-19: an initiative of the International
- 10 Society of Hypertension. BMJ Open. 2021;11(2):e043625.
- 11 93. Wang Y, Jiang W, He Q, Wang C, Wang B, Zhou P, et al. Early, low-dose and
- 12 short-term application of corticosteroid treatment in patients with severe COVID-19
- 13 pneumonia: single-center experience from Wuhan, China. medRxiv. 2020. [preprint]
- 14 94. Bozkurt B, Kovacs R, Harrington B. Joint HFSA/ACC/AHA Statement
- 15 Addresses Concerns Re: Using RAAS Antagonists in COVID-19. J Card Fail.
- 16 2020;26(5):370.

17 95. British Society for Rheumatology. COVID-19 guidance 2020 [cited 2020 Jul 29].

- 18 Available from: https://www.rheumatology.org.uk/practice-quality/covid-19-
- 19 guidance.
- 20 96. Guys & St Thomas' NHS Foundation Trust. Dermatology and coronavirus:
- 21 frequently asked questions 2020 [cited 2020 Jul 29]. Available from:
- 22 <u>https://www.guysandstthomas.nhs.uk/our-services/dermatology/dermatology-and-</u>
- 23 <u>coronavirus-frequently-asked-questions.aspx#na</u>.
- 24 97. National Institute for Health and Care Excellence. COVID-19 rapid evidence
- 25 summary: Long-term use of non-steroidal anti-inflammatory drugs (NSAIDs) for
- people with or at risk of COVID-19 2020 [cited 2020 Jul 29]. Available from:
   https://www.nice.org.uk/advice/es25/chapter/Key-messages.
- 28 98. Horby P, Lim WS, Emberson J, Mafham M, Bell J, Linsell L, et al. Effect of
- 29 Dexamethasone in Hospitalized Patients with COVID-19: Preliminary Report.
- 30 medRxiv. 2020:2020.06.22.20137273. [preprint]
- 31 99. CTU Bern. CORON-ACT started enrollment 2020 [cited 2020 Jul 28]. Available32 from:
- 33 <u>https://www.ctu.unibe.ch/about\_us/news/coron\_act\_started\_enrollment/index\_eng.</u>
   34 html.
- 35 100. Beale R, Carter B, Douthwaite S, WIlliams S, Mehta M, Lynch E, et al. Lipid
- 36 Ibuprofen versus standard of care for acute hypoxaemic respiratory failure due to
- 37 COVID-19: a multicentre, randomised, controlled trial (The Liberate Trial) BTD
- 38 Health2020 [cited 2020 Jul 28]. Available from: <u>https://btd-health.com/</u>.
- 39 101. Chen L. Atorvastatin as Adjunctive Therapy in COVID-19 (STATCO19) NIH
- 40 U.S. National Library of Medicine2020 [cited 2020 Jul 28]. Available from:
- 41 https://clinicaltrials.gov/ct2/show/NCT04380402.
- 42 102. Jardine M, Wilcox A. Controlled evaLuation of Angiotensin Receptor Blockers
- 43 for COVID-19 respIraTorY Disease (CLARITY): NIH U.S. National Library of
- 44 Medicine; 2020 [cited 2020 Jul 28]. Available from:
- 45 <u>https://clinicaltrials.gov/ct2/show/NCT04394117</u>.

- 103. United in Research. The PACE Study: United Health Group; 2020 [cited 2020
- Jul 28]. Available from: <u>https://www.unitedinresearch.com/studies/2</u>.

### Appendix 1: search terms 1

# 2 First search (Ovid)

| 3  | 1.  | Coronavirus Infections/ or covid-19.mp. or Betacoronavirus/                     |
|----|-----|---|
| 4  | 2.  | Pneumonia, Viral/ or Severe Acute Respiratory Syndrome/ or sars-cov-2.mp.       |
| 5  |     | or SARS Virus/  |
| 6  | 3.  | Pandemics/ or ncov-2019.mp.   |
| 7  | 4.  | 2019-nCov.mp. or Spike Glycoprotein, Coronavirus/                               |
| 8  | 5.  | 1 or 2 or 3 or 4  |
| 9  | 6.  | Analgesics/ or Anti-Inflammatory Agents, Non-Steroidal/ or non-                 |
| 10 |     | steroidal.mp or Cyclooxygenase 2 Inhibitors/                                    |
| 11 | 7.  | Ibuprofen.mp. or ibuprofen/   |
| 12 | 8.  | Angiotensin-Converting Enzyme Inhibitor?.mp. or Angiotensin-Converting          |
| 13 |     | Enzyme Inhibitors/  |
| 14 | 9.  | ACE Inhibitor?.mp.  |
| 15 | 10. | Antihypertensive Agents/ or Angiotensin Receptor Antagonists/ or                |
| 16 |     | ARB?.mp. or Angiotensin II Type 1 Receptor Blockers/                            |
| 17 | 11. | Angiotensin Receptor Blocker?.mp.   |
| 18 | 12. | Angiotensin II Receptor Blocker?.mp.  |
| 19 | 13. | Steroid?.mp. or Steroids/   |
| 20 | 14. | Corticosteroid?.mp. or Adrenal Cortex Hormones/                                 |
| 21 | 15. | Immunosuppressant?.mp. or Immunosuppressive Agents/                             |
| 22 | 16. | ((drug? or medication? or prescri*) adj10 (harm or safe* or risk or worse* or   |
| 23 |     | sever* or discontinue? or unsafe or avoid)).mp. [mp=ti, ab, ot, nm, hw, fx, kf, |
| 24 |     | ox, px, rx, ui, an, sy, tn, dm, mf, dv, kw, dq]                                 |
| 25 | 17. | 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15 or 16                      |
| 26 | 18. | 5 and 17  |
| 27 | 19. | Limit 18 to yr="2019-Current"   |
| 28 |     |   |

### Second search (WHO COVID-19 database) 1

- 2 1. NSAID\*
- 3 2. Non-steroid\*
- 3. Analgesi\* 4
- 4. Ibuprofen 5
- 5. ACE Inhibitor\* 6
- 7 6. ACE-I\*
- 8 7. ACEI\*
- 9 8. Angiotensin Converting Enzyme Inhibitor\*
- 10 9. Renin-Angiotensin-Aldosterone System Inhibitor\*
- 11 10. Renin-Angiotensin-Aldosterone System Blocker\*
- 12 11. RAAS Inhibitor\*
- 12. RAAS Blocker\* 13
- 14 13. \*pril
- 14. Antihypertensive\* 15
- 15. Angiotensin Receptor Blocker\* 16
- 16. ARB\* 17
- 18 17. \*sartan
- 19 18. \*steroid\*
- 20 19. \*cortico\*
- 20. Immunosuppressant\* 21
- 22 21. Immunosuppressive\*
- 23 22. Immunomodulat\*
- 24 23. Immune modulat\*
- 25 24. Antimineralocorticoid\*
- 26 25. Aldosterone antagonist\*
- 26. Spironolactone 27
- 28 27. MCRA\*
- 29 28. MRA\*
- 29. Mineralocorticoid Receptor Antagonist\* 30

- 1 30. \*statin\*
- 2 31. HMG CoA Reductase Inhibitor\*
- 3 32. Thiazolidinedione\*
- 4 33. \*glitazone\*
- Third search (NIH iSearch COVID-19 portfolio) 5
- 1. NSAID 6
- 7 2. Non-steroid
- 8 3. Analgesi\*
- 9 4. Ibuprofen
- 10 5. ACE Inhibitor
- 11 6. ACE-I
- 12 7. ACEI
- 13 8. Angiotensin Converting Enzyme Inhibitor
- 14 9. "Renin-Angiotensin-Aldosterone System Inhibitor"
- 15 10. "Renin-Angiotensin-Aldosterone System Blocker"
- 11. "RAAS Inhibitor" 16
- 17 12. "RAAS Blocker"
- 13. \*pril 18
- 19 14. Antihypertensive
- 20 15. Angiotensin Receptor Blocker
- 21 16. ARB
- 22 17. \*sartan
- 23 18. \*steroid\*
- 24 19. \*cortico\*
- 25 20. Immunosuppressant
- 26 21. Immunosuppressive
- 27 22. Immunomodulat\*
- 28 23. "Immune modulator"
- 29 24. Antimineralocorticoid

- 25. "Aldosterone antagonist" 1
- 2 26. Spironolactone
- 3 27. MCRA
- 4 28. MRA
- 29. "Mineralocorticoid Receptor Antagonist" 5
- 30. \*statin 6
- 31. "HMG CoA Reductase Inhibitor" 7
- 8 32. Thiazolidinedione
- 9 33. \*glitazone
- 10

### Appendix 2: inclusion & exclusion criteria 1

### **First set** 2

#### Inclusion criteria: 3

| 4  | 1.   | Papers looking at whether a medication or medications are potentially  |
|--|--|--|
| 5  |  | unsafe if prescribed for patients with confirmed or suspected COVID-19 or  |
| 6  |  | SARS-Cov-2 infection   |
| 7  | 2.   | Papers looking at whether a medication or medications should be  |
| 8  |  | discontinued in patients who have confirmed or suspected COVID-19 or   |
| 9  |  | SARS-Cov-2 infection and would otherwise take them regularly   |
| 10   | 3.   | Papers looking at the overall safety or benefits of prescribing or   |
| 11   |  | discontinuing a medication or medications that have been identified as   |
| 12   |  | potentially unsafe for patients under criteria 1 or 2  |
| 13   | 4.   | Paper addressing or responding to any paper already included in reference  |
| 14   |  | to the safety of a medication or medications   |
| 15   |  |  |
| 10   |  |  |
| 16   | Exclus   | ion criteria:  |
|  |  |  |
| 17   | 1.   | Paper does not mention any specific medications or groups of medications   |
| 17<br>18   | 1.<br>2.   | Paper does not mention any specific medications or groups of medications<br>Paper not related to current COVID-19 outbreak   |
| 17<br>18<br>19   | 1.<br>2.<br>3.   | Paper does not mention any specific medications or groups of medications<br>Paper not related to current COVID-19 outbreak<br>Published before November 2019   |
| 17<br>18<br>19<br>20   | 1.<br>2.<br>3.<br>4.   | Paper does not mention any specific medications or groups of medications<br>Paper not related to current COVID-19 outbreak<br>Published before November 2019<br>Paper not written in English   |
| <ol> <li>17</li> <li>18</li> <li>19</li> <li>20</li> <li>21</li> </ol>   | 1.<br>2.<br>3.<br>4.<br>5.   | Paper does not mention any specific medications or groups of medications<br>Paper not related to current COVID-19 outbreak<br>Published before November 2019<br>Paper not written in English<br>Only looking at unlicensed uses of medications, where no medication  |
| 17<br>18<br>19<br>20<br>21<br>22   | 1.<br>2.<br>3.<br>4.<br>5.   | Paper does not mention any specific medications or groups of medications<br>Paper not related to current COVID-19 outbreak<br>Published before November 2019<br>Paper not written in English<br>Only looking at unlicensed uses of medications, where no medication<br>involved has already been identified as posing a potential risk when  |
| <ol> <li>17</li> <li>18</li> <li>19</li> <li>20</li> <li>21</li> <li>22</li> <li>23</li> </ol>                         | 1.<br>2.<br>3.<br>4.<br>5.   | Paper does not mention any specific medications or groups of medications<br>Paper not related to current COVID-19 outbreak<br>Published before November 2019<br>Paper not written in English<br>Only looking at unlicensed uses of medications, where no medication<br>involved has already been identified as posing a potential risk when<br>prescribed for its licensed use(s)  |
| <ol> <li>17</li> <li>18</li> <li>19</li> <li>20</li> <li>21</li> <li>22</li> <li>23</li> <li>24</li> </ol>             | <ol> <li>1.</li> <li>2.</li> <li>3.</li> <li>4.</li> <li>5.</li> <li>6.</li> </ol> | Paper does not mention any specific medications or groups of medications<br>Paper not related to current COVID-19 outbreak<br>Published before November 2019<br>Paper not written in English<br>Only looking at unlicensed uses of medications, where no medication<br>involved has already been identified as posing a potential risk when<br>prescribed for its licensed use(s)<br>Only looking at medications being used for the treatment of COVID-19,   |
| <ol> <li>17</li> <li>18</li> <li>19</li> <li>20</li> <li>21</li> <li>22</li> <li>23</li> <li>24</li> <li>25</li> </ol> | <ol> <li>1.</li> <li>2.</li> <li>3.</li> <li>4.</li> <li>5.</li> <li>6.</li> </ol> | Paper does not mention any specific medications or groups of medications<br>Paper not related to current COVID-19 outbreak<br>Published before November 2019<br>Paper not written in English<br>Only looking at unlicensed uses of medications, where no medication<br>involved has already been identified as posing a potential risk when<br>prescribed for its licensed use(s)<br>Only looking at medications being used for the treatment of COVID-19,<br>where no medication involved has already been identified as posing a |

- 1 7. No new data, hypothesis, guidance or recommendations stated related to
- 2 COVID-19 for any individual relevant medications

# 3 Second set

4 Inclusion criteria:

| 5  | 1.     | Studies looking at the outcomes of patients with COVID-19 or SARS-Cov-2         |
|----|--------|---|
| 6  |        | infection who were, or would normally be, taking medications that have          |
| 7  |        | been identified in the preliminary search as potentially harmful if prescribed  |
| 8  |        | for patients with confirmed or suspected COVID-19 or SARS-Cov-2 infection       |
| 9  |        | (see list)  |
| 10 | 2.     | Studies looking at the outcomes of patients with COVID-19 or SARS-Cov-2         |
| 11 |        | infection when given medications that have been identified in the               |
| 12 |        | preliminary search as potentially harmful if prescribed for patients with       |
| 13 |        | confirmed or suspected COVID-19 or SARS-Cov-2 infection                         |
| 14 | Exclus | sion criteria:  |
| 15 | 1.     | Paper does not mention any medications or groups of medications already         |
| 16 |        | identified as potentially harmful   |
| 17 | 2.     | Paper not related to current COVID-19 outbreak                                  |
| 18 | 3.     | Published before November 2019  |
| 19 | 4.     | Paper not written in English  |
| 20 | 5.     | Paper has no primary data   |
| 21 | 6.     | Study not performed in vivo   |
| 22 | 7.     | It is unclear from the paper if the clinical outcomes follow the administration |
| 23 |        | of a medication or group of medications already identified as potentially       |
| 24 |        | harmful   |
| 25 | 8.     | There is no measure against non-COVID-19 or non-drug exposed control            |

# Appendix 3: study tables

Table 3. Summary of studies looking into the effect of drug groups on the susceptibility to COVID-19 infection.

| First Author    | Design | Country        | Drug Group | Setting       | Population                               | Sample Size | Specific drugs | Outcome<br>definitions  | Result                          | Statistic                           | 95% CI                                   | Quality | Confounders<br>adjusted/matc<br>hed for   | Peer-review<br>or Preprint | Drugs in<br>meta-analyses |
|-----------------|--------|----------------|------------|---------------|--|-------------|----------------|---|---------------------------------|-------------------------------------|--|---------|---|----------------------------|---------------------------|
| Caraballo,<br>C | XSS    | USA            | RAASB      | Commu<br>nity | Heart failure<br>registry                | 26703       | Combined       | Symptomatic patients receiving positive test result                       | Protective                      | OR 0.68                             | 0.49-0.94                                | Poor    | None  | PP                         | Yes                       |
| Dooley,<br>H    | XSS    | UK             | RAASB      | Commu<br>nity | Self-<br>reporting<br>COVID app<br>users | 2215386     | Combined       | Self-reported positive PCR<br>test  | NSE                             | OR 0.87                             | 0.66-1.13                                | Good    | None  | PP                         | Yes                       |
| Huh, K          | CCS    | South<br>Korea | RAASB      | All           | Patients<br>tested for<br>COVID          | 65149       | ACEI &<br>ARB  | Positive test for SARS-CoV-<br>2 NOS                                      | ACEI:<br>NSE<br>ARB:<br>Harmful | ACEI: OR<br>1.25<br>ARB: OR<br>1.13 | ACEI:<br>0.91-1.71<br>ARB: 1.01-<br>1.26 | Good    | Age, sex, co-<br>morbidities,<br>location,<br>healthcare usage<br>and other drugs | PP                         | Yes                       |
| Khawaja,<br>AP  | CoS    | UK             | RAASB      | All           | UK Biobank                               | 406793      | ACEI &<br>ARB  | Hospitalization with positive<br>test compared with general<br>population | NSE                             | ACEI: OR<br>1.17<br>ARB: OR<br>1.00 | ACEI:<br>0.90-1.52<br>ARB: 0.70-<br>1.42 | Fair    | Age, sex,<br>ethnicity and<br>hypertension<br>status                              | PP                         | Yes                       |
| Kolin, D        | CoS    | UK             | RAASB      | All           | UK Biobank                               | 502536      | ACEI &<br>ARB  | Positive test compared with general population                            | NSE                             | ACEI: RR<br>1.32<br>ARB: RR<br>1.37 | ACEI:<br>0.95-1.84<br>ARB: 0.94-<br>1.98 | Fair    | Age, sex, BMI,<br>systolic BP, race<br>and deprivation                            | PP                         | Yes                       |

| First Author    | Design | Country | Drug Group | Setting<br>Population   | Sample Size | Specific drugs | Outcome<br>definitions                     | Result | Statistic                           | 95% CI                                   | Quality | Confounders<br>adjusted/matc<br>hed for   | Peer-review<br>or Preprint | Drugs in<br>meta-analyses |
|-----------------|--------|---------|------------|---|-------------|----------------|--|--------|-------------------------------------|--|---------|---|----------------------------|---------------------------|
| Mancia,<br>G    | CCS    | Italy   | RAASB      | All Patients with<br>confirmed or<br>suspected<br>COVID       | 6272        | ACEI &<br>ARB  | Diagnosed as per interim<br>WHO guidelines | NSE    | ACEI: OR<br>0.96<br>ARB: OR<br>0.95 | ACEI:<br>0.87-1.07<br>ARB: 0.86-<br>1.05 | Good    | Age, sex,<br>location and co-<br>morbidities  | PR                         | Yes                       |
| Rentsch,<br>CT  | CoS    | USA     | RAASB      | Hospital VA Birth<br>Cohort<br>registry (54-<br>75 years old) | 3789        | Combined       | Positive test for SARS-CoV-<br>2 NOS       | NSE    | OR: 0.98                            | 0.78-1.23                                | Good    | Age, sex, race,<br>co-morbidities,<br>vital signs,<br>medications and<br>health<br>behaviours | PP                         | Yes                       |
| Reynolds,<br>HR | CoS    | USA     | RAASB      | All Patients<br>tested for<br>COVID                           | 12594       | ACEI &<br>ARB  | At least one positive test                 | NSE    | ACEI: OR<br>0.92<br>ARB: OR<br>1.00 | ACEI:<br>0.79-1.08<br>ARB: 0.86-<br>1.15 | Good    | Age, sex, race,<br>BMI, smoking<br>history, co-<br>morbidities and<br>treatment               | PR                         | Yes                       |
| Shah, SJ        | CoS    | USA     | RAASB      | Hospital Respiratory<br>admissions<br>tested for<br>COVID     | 316         | Combined       | Positive test compared with negative test  | NSE    | OR: 1.24                            | 0.49-3.13                                | Fair    | None  | РР                         | Yes                       |

| First Author | Design | Country        | Drug Group<br>Setting | Population  | Sample Size | Specific drugs      | Outcome<br>definitions  | Result                             | Statistic                   | 95% CI                                   | Quality | Confounders<br>adjusted/matc<br>hed for   | Peer-review | Drugs in<br>meta-analyses |
|--------------|--------|----------------|-----------------------|---|-------------|---------------------|---|------------------------------------|-----------------------------|--|---------|---|-------------|---------------------------|
| Yan, H       | CCS    | China          | RAASB Hospital        | Admissions<br>with COVID                            | 610         | ACEI &<br>ARB       | Hospitalized compared with<br>general population; Severe or<br>critical compared with non-<br>severe. Severe defined as one<br>of: RR ≥30/min, SpO2<br>≤93%, FiO2 ≤300mmHg,<br>critical defined as one of:<br>requiring MV, shock,<br>requiring ICU care for organ<br>support | ACEI:<br>NSE<br>ARB:<br>Protective | ACEI: OR<br>0.65<br>OR 0.24 | ACEI:<br>0.26-1.57<br>ARB: 0.17-<br>0.34 | Fair    | Age, sex and<br>BMI   | PP          | Yes                       |
| Huh, K       | CCS    | South<br>Korea | CS All                | Patients<br>tested for<br>COVID                     | 65149       |                     | Positive test for SARS-CoV-<br>2 NOS  | Protective                         | OR 0.86                     | 0.79-0.94                                | Good    | Age, sex, co-<br>morbidities,<br>location,<br>healthcare usage<br>and other drugs | PP          | Yes                       |
| Mancia,<br>G | CCS    | Italy          | CS All                | Patients with<br>confirmed or<br>suspected<br>COVID | 6272        | Inhaled<br>steroids | Diagnosed as per interim<br>WHO guidelines  | Harmful                            | OR 1.52                     | 1.37-1.68                                | Good    | Age, sex,<br>location and co-<br>morbidities                                      | PR          | Yes                       |
| Shah, SJ     | CoS    | USA            | CS Hospital           | Respiratory<br>admissions<br>tested for<br>COVID    | 316         |                     | Positive test compared with negative test   | NSE                                | OR 1.77                     | 0.63-4.91                                | Fair    | None  | PP          | Yes                       |

| First Author  | Design | Country        | Drug Group | Setting  | Population   | Sample Size | Specific drugs                     | Outcome<br>definitions  | Result  | Statistic                            | 95% CI                                    | Quality | Confounders<br>adjusted/matc<br>hed for   | Peer-review<br>or Dranrint | Drugs in<br>meta-analyses |
|---------------|--------|----------------|------------|----------|--|-------------|------------------------------------|---|---------|--------------------------------------|---|---------|---|----------------------------|---------------------------|
| Yan, H        | CCS    | China          | CS         | Hospital | Admissions<br>with COVID   | 610         |                                    | Hospitalized compared with<br>general population; Severe or<br>critical compared with non-<br>severe. Severe defined as one<br>of: RR ≥30/min, SpO2<br>≤93%, FiO2 ≤300mmHg,<br>critical defined as one of:<br>requiring MV, shock,<br>requiring ICU care for organ<br>support | NSE     | OR 1.20                              | 0.49-2.94                                 | Fair    | Age, sex and<br>BMI   | PP                         | Yes                       |
| Gisondi,<br>P | XSS    | Italy          | IS         | All      | Patients with<br>psoriasis on<br>biologic<br>treatment or<br>renal<br>transplant<br>patients | 1223        |                                    | Admission to hospital with<br>COVID-19 compared with<br>general population  | NSE     | OR 0.35                              | 0.05-2.52                                 | Poor    | None  | PR                         | Yes                       |
| Huh, K        | CCS    | South<br>Korea | IS         | All      | Patients<br>tested for<br>COVID  | 65149       | Mycopheno<br>late and<br>sirolimus | Positive test for SARS-CoV-<br>2 NOS  | NSE     | Myco: OR<br>1.00<br>Siro: OR<br>1.16 | Myco:<br>0.56-1.79<br>Siro: 0.15-<br>8.87 | Good    | Age, sex, co-<br>morbidities,<br>location,<br>healthcare usage<br>and other drugs | PP                         | Yes                       |
| Mancia,<br>G  | CCS    | Italy          | IS         | All      | Patients with<br>confirmed or<br>suspected<br>COVID  | 6272        |                                    | Diagnosed as per interim<br>WHO guidelines  | Harmful | OR 1.30                              | 1.20-1.42                                 | Good    | Age, sex,<br>location and co-<br>morbidities                                      | PR                         | Yes                       |

| First Author   | Design | Country        | Drug Group<br>Setting | gumg    | Population   | Sample Size | Specific drugs | Outcome<br>definitions                         | Result     | Statistic | 95% CI    | Quality | Confounders<br>adjusted/matc<br>hed for   | Peer-review<br>or Preprint | Drugs in<br>meta-analyses |
|----------------|--------|----------------|-----------------------|---------|--|-------------|----------------|--|------------|-----------|-----------|---------|---|----------------------------|---------------------------|
| Rentsch,<br>CT | CoS    | USA            | IS Ho                 | ospital | VA Birth<br>Cohort<br>registry (54-<br>75 years old) | 3789        |                | Positive test                                  | NSE        | OR 0.82   | 0.55-1.22 | Good    | Age, sex, race,<br>co-morbidities,<br>vital signs,<br>medications and<br>health<br>behaviours | PP                         | Yes                       |
| Shah, SJ       | CoS    | USA            | IS Ho                 | ospital | Respiratory<br>admissions<br>tested for<br>COVID     | 316         |                | Positive test compared with negative test      | NSE        | OR 1.57   | 0.61-4.08 | Fair    | None  | PP                         | Yes                       |
| Huh, K         | CCS    | South<br>Korea | NSAID AI              | 11      | Patients<br>tested for<br>COVID                      | 65149       |                | Positive test for SARS-CoV-<br>2 NOS           | Protective | OR 0.90   | 0.84-0.96 | Good    | Age, sex, co-<br>morbidities,<br>location,<br>healthcare usage<br>and other drugs             | PP                         | Yes                       |
| Kolin, D       | CoS    | UK             | NSAID AI              | 11      | UK Biobank   | 502536      |                | Positive test compared with general population | NSE        | RR 1.02   | 0.79-1.22 | Fair    | Age, sex, BMI,<br>systolic BP, race<br>and deprivation  | PP                         | Yes                       |
| Mancia,<br>G   | CCS    | Italy          | NSAID AI              | 11      | Patients with<br>confirmed or<br>suspected<br>COVID  | 6272        |                | Diagnosed as per interim<br>WHO guidelines     | NSE        | OR 1.06   | 0.98-1.15 | Good    | Age, sex,<br>location and co-<br>morbidities  | PR                         | Yes                       |

| First Author    | Design | Country        | Drug Group | Setting       | Population   | Sample Size | Specific drugs | Outcome<br>definitions                              | Result  | Statistic | 95% CI    | Quality | Confounders<br>adjusted/matc<br>hed for   | Peer-review | Drugs in | meta-analyses |
|-----------------|--------|----------------|------------|---------------|--|-------------|----------------|---|---------|-----------|-----------|---------|---|-------------|----------|---------------|
| Rentsch,<br>CT  | CoS    | USA            | NSAID      | Hospital      | l VA Birth<br>Cohort<br>registry (54-<br>75 years old) | 3789        |                | Positive test                                       | Harmful | OR 1.27   | 1.02-1.58 | Good    | Age, sex, race,<br>co-morbidities,<br>vital signs,<br>medications and<br>health<br>behaviours | PP          | Yes      |               |
| Yan, H          | CCS    | China          | NSAID      | Hospital      | Admissions<br>with COVID                               | 610         | Aspirin        | Hospitalized compared with general population       | NSE     | OR 1.25   | 0.66-2.39 | Fair    | Age, sex and<br>BMI   | PP          | Yes      |               |
| Caraballo,<br>C | XSS    | USA            | Statin     | Commu<br>nity | Heart failure<br>registry                              | 26703       |                | Symptomatic patients receiving positive test result | NSE     | OR 0.76   | 0.56-1.05 | Poor    | None  | PP          | Yes      |               |
| Huh, K          | CCS    | South<br>Korea | Statin     | All           | Patients<br>tested for<br>COVID                        | 65149       |                | Positive test for SARS-CoV-<br>2 NOS                | NSE     | OR 0.99   | 0.89-1.10 | Good    | Age, sex, co-<br>morbidities,<br>location,<br>healthcare usage<br>and other drugs             | PP          | Yes      |               |
| Mancia,<br>G    | CCS    | Italy          | Statin     | All           | Patients with<br>confirmed or<br>suspected<br>COVID    | 6272        |                | Diagnosed as per interim<br>WHO guidelines          | NSE     | OR 1.02   | 0.94-1.10 | Good    | Age, sex,<br>location and co-<br>morbidities  | PR          | Yes      |               |
| Yan, H          | CCS    | China          | Statin     | Hospital      | Admissions<br>with COVID                               | 610         |                | Hospitalized compared with general population       | Harmful | OR 2.50   | 1.48-4.21 | Fair    | Age, sex and<br>BMI   | PP          | Yes      |               |

| First Author   | Design | Country        | Drug Group | Setting  | Population  | Sample Size | Specific drugs | Outcome<br>definitions  | Result  | Statistic | 95% CI    | Quality | Confounders<br>adjusted/matc<br>hed for   | Peer-review | or rreprint<br>Drugs in | meta-analyses |
|----------------|--------|----------------|------------|----------|---|-------------|----------------|---|---------|-----------|-----------|---------|---|-------------|-------------------------|---------------|
| Huh, K         | CCS    | South<br>Korea | TZD        | All      | Patients<br>tested for<br>COVID                     | 65149       |                | Positive test for SARS-CoV-<br>2 NOS  | NSE     | OR 1.00   | 0.70-1.41 | Good    | Age, sex, co-<br>morbidities,<br>location,<br>healthcare usage<br>and other drugs | PP          | Ye                      | S             |
| Kolin, D       | CoS    | UK             | TZD        | All      | UK Biobank  | 502536      |                | Positive test compared with<br>general population; Being<br>admitted to hospital<br>compared with those testing<br>positive but remaining in the<br>community | NSE     | OR 0.94   | 0.35-2.57 | Fair    | Age, sex, BMI,<br>systolic BP, race<br>and deprivation                            | PP          | Ye                      | S             |
| Mancia,<br>G   | CCS    | Italy          | TZD        | All      | Patients with<br>confirmed or<br>suspected<br>COVID | 6272        |                | Diagnosed as per interim<br>WHO guidelines; Critical or<br>fatal disease, critical defined<br>as receiving MV   | Harmful | OR 1.81   | 1.23-2.67 | Good    | Age, sex,<br>location and co-<br>morbidities                                      | PR          | Ye                      | S             |
| Yan, H         | CCS    | China          | TZD        | Hospital | Admissions<br>with COVID                            | 610         |                | Hospitalized compared with general population   | NSE     | OR 0.35   | 0.11-1.13 | Fair    | Age, sex and<br>BMI   | PP          | Ye                      | s             |
| Khawaja,<br>AP | CoS    | UK             | MCRA       | All      | UK Biobank  | 406793      |                | Hospitalization with positive<br>test compared with general<br>population   | NSE     | OR 1.54   | 0.62-3.85 | Fair    | Age, sex,<br>ethnicity and<br>hypertension<br>status                              | PP          | Na<br>ive               | rrat          |
| Mancia,<br>G   | CCS    | Italy          | MCRA       | All      | Patients with<br>confirmed or<br>suspected<br>COVID | 6272        |                | Diagnosed as per interim<br>WHO guidelines  | NSE     | OR 0.90   | 0.75-1.07 | Good    | Age, sex,<br>location and co-<br>morbidities                                      | PR          | Na<br>ive               | rrat          |

CoS: Cohort Study, CCS: Case Control Study, XSS: Cross Sectional Study, CaseS: Case Series, NSE: No significant evidence, NVD: No verifiable data, HR: Hazard Ratio, OR: Odds Ratio, HD: high dose, LD: low dose, PR: Peer reviewed, PP: Preprint

| First Author       | Design | Country | Drug Group | Setting      | Population  | Sample Size | Acute or<br>Regular | Specific drugs     | Outcome<br>definitions  | Result                                      | Statistic                             | 95% CI                                  | Quality | Confounders<br>adjusted/matc<br>hed for | Peer-review or<br>PP | Drugs in meta-<br>analyses |
|--------------------|--------|---------|------------|--------------|---|-------------|---------------------|--------------------|---|---|---------------------------------------|---|---------|---|----------------------|----------------------------|
| Argenzia<br>no, MG | CoS    | USA     | RAA<br>SB  | Hospi<br>tal | ED and hospital<br>presentations<br>with COVID      | 1000        | Regul<br>ar         | Combin<br>ed       | Transfer to ICU   | NSE   | OR 1.11                               | 0.81-1.53                               | Fair    | None                                    | PP                   | Yes                        |
| Bean,<br>DM        | CoS    | UK      | RAA<br>SB  | Hospi<br>tal | Hospital<br>admissions with<br>COVID                | 1200        | Regul<br>ar         | Combin<br>ed       | Death or admission to critical<br>care unit for organ support<br>within 21 days of symptom<br>onset   | l Prot<br>ectiv<br>e                        | OR 0.63                               | 0.47-0.84                               | Good    | Co-morbidities, age and sex             | PP                   | Yes                        |
| Benelli,<br>G      | CoS    | Italy   | RAA<br>SB  | Hospi<br>tal | Admissions with<br>COVID                            | 411         | Regul<br>ar         | ACEIs<br>&<br>ARBs | Requiring NIV or CPAP   | AC<br>EI:<br>Har<br>mful<br>AR<br>B:<br>NSE | ACEI:<br>OR 1.95<br>ARB: OR<br>1.68   | ACEI:<br>1.06-3.60<br>ARB:<br>0.94-2.99 | Fair    | None                                    | PP                   | Yes                        |
| Dauchet,<br>L      | XSS    | France  | RAA<br>SB  | All          | Patients with<br>confirmed or<br>suspected<br>COVID | 288         | Regul<br>ar         | ACEIs<br>&<br>ARBs | ICU admission; Being put<br>through to emergency line<br>due to symptoms  | AC<br>EI:<br>NSE<br>AR<br>B:<br>Har<br>mful | ACEI:<br>SPR 0.95<br>ARB:<br>SPR 1.56 | ACEI:<br>0.55-1.64<br>ARB:<br>1.02-2.39 | Good    | None                                    | РР                   | Yes                        |
| Feng, Y            | XSS    | China   | RAA<br>SB  | Hospi<br>tal | Admissions with<br>COVID                            | 476         | Regul<br>ar         | Combin<br>ed       | Critical compared with<br>moderate: critical defined as<br>respiratory failure requiring<br>MV, shock or requiring ICU<br>care. Moderate defined as<br>fever, cough and pneumonia<br>on chest CT but RR<30/min,<br>SpO2>93% and FiO2<br>>300mmHg (i.e not severe) | Prot<br>ectiv<br>e                          | OR 0.17                               | 0.05-0.52                               | Fair    | None                                    | PR                   | Yes                        |

### Table 4. Summary of studies looking into the effect of drug groups on the severity of COVID-19 infection.

| First Author    | Design    | Country | Drug Group | Setting      | Population  | Sample Size | Acute or<br>Regular | Specific drugs     | Outcome<br>definitions  | Result<br>Statistic                     | 95% CI                                    | Quality | Confounders<br>adjusted/matc<br>hed for  | Peer-review or<br>PP | Drugs in meta-<br>analyses |
|-----------------|-----------|---------|------------|--------------|---|-------------|---------------------|--------------------|---|---|---|---------|--|----------------------|----------------------------|
| Feng, Z         | CoS       | China   | RAA<br>SB  | Hospi<br>tal | Admissions with<br>COVID                            | 564         | Regul<br>ar         | Combin<br>ed       | Severe or critical compared<br>with non-severe. Severe<br>defined as one of: RR<br>≥30/min, SpO2 ≤93%, FiO2<br>≤300mmHg, critical defined<br>as one of: requiring MV,<br>shock, requiring ICU care for<br>organ support                       | NSE OR 0.41                             | 0.05-3.19                                 | Good    | Propensity score<br>matched and adjusted<br>for age, gender,<br>smoking history, co-<br>morbidities and lung<br>CT score | РР                   | Yes                        |
| Kolin, D        | CoS       | UK      | RAA<br>SB  | All          | UK Biobank  | 502536      | Regul<br>ar         | ACEIs<br>&<br>ARBs | Being admitted to hospital<br>compared with those testing<br>positive but remaining in the<br>community   | NSE ACEI:<br>0.56<br>ARB:<br>0.60       | ACEI:<br>0.26-1.22<br>ARB:<br>0.25-1.42   | Fair    | Age, sex, BMI, systolic<br>BP, race and<br>deprivation   | PP                   | Yes                        |
| Li, J           | Case<br>S | China   | RAA<br>SB  | Hospi<br>tal | Hypertensive<br>COVID<br>admissions                 | 1178        | Regul<br>ar         | Combin<br>ed       | Severe compared with non-<br>severe. Severe defined as one<br>of: RR $\geq$ 30/min, SpO2<br>$\leq$ 93%, FiO2 $\leq$ 300mmHg,<br>lung infiltrates more than<br>50% within 24 to 48 hours,<br>respiratory failure and/or<br>multi-organ failure | NSE OR 1.11                             | 0.71-1.73                                 | Good    | None   | PR                   | Yes                        |
| Li, X           | CoS       | China   | RAA<br>SB  | Hospi<br>tal | Admissions with<br>COVID                            | 548         | Regul<br>ar         | Combin<br>ed       | Severe or critical compared<br>with non-severe. Severe<br>defined as per IDSA & ATS<br>guidelines for CAP   | NSE OR 0.86                             | 0.45-1.61                                 | Good    | Age, sex, blood tests,<br>antiviral usage  | PR                   | Yes                        |
| Liu,<br>Yingxia | CoS       | China   | RAA<br>SB  | Hospi<br>tal | Hypertensive<br>COVID<br>admissions                 | 78          | Regul<br>ar         | ACEIs<br>&<br>ARBs | Severe compared with non-<br>severe as per NHCPRC<br>2019-nCoV guidelines   | NSE ACEI:<br>OR 0.57<br>ARB: OF<br>0.54 | ACEI:<br>0.11-2.95<br>R ARB:<br>0.25-1.16 | Fair    | Sex  | PP                   | Yes                        |
| Mancia,<br>G    | CCS       | Italy   | RAA<br>SB  | All          | Patients with<br>confirmed or<br>suspected<br>COVID | 6272        | Regul<br>ar         | ACEIs<br>&<br>ARBs | Critical or fatal disease,<br>critical defined as receiving<br>MV   | NSE ACEI:<br>OR 0.91<br>ARB: OF<br>0.83 | ACEI:<br>0.69-1.21<br>ARB:<br>0.63-1.10   | Good    | Age, sex, location and co-morbidities  | PR                   | Yes                        |

| First Author    | Design | Country | Drug Group | Setting      | Population                                       | Sample Size | Acute or<br>Regular | Specific drugs     | Outcome<br>definitions  | Result<br>Statistic                     | 95% CI                                  | Quality | Confounders<br>adjusted/matc<br>hed for  | Peer-review or<br>PP | Drugs in meta-<br>analyses |
|-----------------|--------|---------|------------|--------------|--|-------------|---------------------|--------------------|---|---|---|---------|--|----------------------|----------------------------|
| Meng, J         | CoS    | China   | RAA<br>SB  | Hospi<br>tal | Hypertensive<br>COVID<br>admissions              | 42          | Regul<br>ar         | Combin<br>ed       | Severe disease or death,<br>severe defined as per<br>NHCPRC 2019-nCoV<br>guidelines   | NSE OR 0.28                             | 0.07-1.12                               | Fair    | None   | PR                   | Yes                        |
| Rentsch,<br>CT  | CoS    | USA     | RAA<br>SB  | Hospi<br>tal | VA Birth Cohort<br>registry (54-75<br>years old) | 3789        | Regul<br>ar         | Combin<br>ed       | Hospitalization compared<br>with community care after<br>testing positive   | NSE OR 1.15                             | 0.71-1.87                               | Good    | Age, sex, race, co-<br>morbidities, vital signs,<br>medications and health<br>behaviours | PP                   | Yes                        |
| Reynolds,<br>HR | CoS    | USA     | RAA<br>SB  | All          | Patients tested<br>for COVID                     | 12594       | Regul<br>ar         | ACEIs<br>&<br>ARBs | ICU admission, MV or death  | NSE ACEI:<br>OR 0.90<br>ARB: OR<br>0.99 | ACEI:<br>0.70-1.14<br>ARB:<br>0.79-1.25 | Good    | Age, sex, race, BMI,<br>smoking history, co-<br>morbidities and<br>treatment             | PR                   | Yes                        |
| Yan, H          | CCS    | China   | RAA<br>SB  | Hospi<br>tal | Admissions with<br>COVID                         | 610         | Regul<br>ar         | ACEIs<br>&<br>ARBs | Severe or critical compared<br>with non-severe. Severe<br>defined as one of: RR<br>≥30/min, SpO2 ≤93%, FiO2<br>≤300mmHg, critical defined<br>as one of: requiring MV,<br>shock, requiring ICU care for<br>organ support | NSE ACEI:<br>OR 1.23<br>ARB: OR<br>0.77 | ACEI:<br>0.19-7.93<br>ARB:<br>0.36-1.63 | Fair    | Age, sex and BMI   | PP                   | Yes                        |
| Yang, G         | CoS    | China   | RAA<br>SB  | Hospi<br>tal | Hypertensive<br>COVID<br>admissions              | 126         | Regul<br>ar         | Combin<br>ed       | Severe or critical compared<br>with non-severe. Severe<br>defined as one of: RR<br>≥30/min, SpO2 ≤93%, FiO2<br>≤300mmHg, critical defined<br>as one of: requiring MV,<br>shock, requiring ICU care for<br>organ support | NSE OR 0.73                             | 0.34-1.58                               | Fair    | None   | PP                   | Yes                        |
| Zeng, Z         | CoS    | China   | RAA<br>SB  | Hospi<br>tal | Hypertensive<br>COVID<br>Admissions              | 75          | Regul<br>ar         | Combin<br>ed       | Severe compared with non-<br>severe. Severe defined as per<br>IDSA & ATS guidelines for<br>CAP  | NSE OR 2.46                             | 0.94-6.45                               | Poor    | None   | PP                   | Yes                        |

| First Author       | Design | Country | Drug Group | Setting      | Population                                      | Sample Size | Acute or<br>Regular | Specific drugs                         | Outcome<br>definitions  | Result             | Statistic                           | 95% CI                                   | Quality | Confounders<br>adjusted/matc<br>hed for           | Peer-review or<br>PP | Drugs in meta-<br>analyses                     |
|--------------------|--------|---------|------------|--------------|---|-------------|---------------------|--|---|--------------------|-------------------------------------|--|---------|---|----------------------|--|
| Argenzia<br>no, MG | CoS    | USA     | CS         | Hospi<br>tal | ED and hospital<br>presentations<br>with COVID  | 1000        | Regul<br>ar         | Inhaled,<br>nasal,<br>oral<br>steroids | Transfer to ICU   | NSE                | OR 1.06                             | 0.59-1.90                                | Fair    | None  | PP                   | Yes  |
| Cao, C             | CoS    | China   | CS         | Hospi<br>tal | Admissions with<br>mild-moderate<br>COVID       | 58          | Acute               | Methyl-<br>prednis<br>olone            | Severe compared with non-<br>severe. Severe defined as one<br>of: $RR \ge 33/min$ , $SpO2 \le$<br>94% at rest, $PO2/FiO2\le$<br>300mmHg, requiring MV,<br>developing shock or multiple<br>organ failure requiring ICU | Prot<br>ectiv<br>e | OR 0.14                             | 0.02-0.80                                | Fair    | None  | PP                   | Narrat<br>ive                                  |
| Li, X              | CoS    | China   | CS         | Hospi<br>tal | Admissions with<br>COVID                        | 548         | Regul<br>ar         | Oral                                   | Severe or critical compared<br>with non-severe. Severe<br>defined as per IDSA & ATS<br>guidelines for CAP   | NSE                | Oral: OR<br>0.51<br>Inh: OR<br>0.69 | Oral:<br>0.09-2.84<br>Inh: 0.11-<br>4.16 | Good    | Age, sex, blood tests,<br>antiviral usage         | PR                   | Yes  |
| Liu,<br>Yanli      | CoS    | China   | CS         | Hospi<br>tal | Admissions with COVID                           | 109         | Acute               |  | Berlin definition of ARDS   | NSE                | OR 1.38                             | 0.64-2.98                                | Fair    | None  | PP                   | Narrat<br>ive                                  |
| Lui, GCY           | CoS    | China   | CS         | Hospi<br>tal | Admissions with<br>COVID                        | 114         | Acute               | Methyl-<br>prednis<br>olone            | ICU admission, MV or death  | Har<br>mful        | HR 4.17                             | 1.23-<br>14.11                           | Poor    | Age, sex, co-<br>morbidities and blood<br>markers | PP                   | OR<br>incalc<br>ulable<br>or<br>inaccu<br>rate |
| Robilotti,<br>EV   | CoS    | USA     | CS         | Hospi<br>tal | Cancer patients<br>with<br>symptomatic<br>COVID | 423         | Regul<br>ar         | Steroids<br>or<br>lympho<br>paenia     | Requiring high-flow oxygen<br>or MV   | NSE                | OR 1.52                             | 0.92-2.50                                | Fair    | Age, race, treatment<br>and co-morbidities        | PP                   | Yes  |
| Wang, D            | CoS    | China   | CS         | Hospi<br>tal | Admissions with<br>COVID                        | 115         | Acute               | Methyl-<br>prednis<br>olone            | ICU admission or death  | NSE                | OR 2.16                             | 0.49-9.43                                | Poor    | Age, sex, co-<br>morbidities and blood<br>markers | PP                   | Narrat<br>ive                                  |

| First Author   | Design | Country | Drug Group | Setting      | Population                                       | Sample Size | Acute or<br>Regular | Specific drugs | Outcome<br>definitions  | Result             | Statistic | 95% CI         | Quality | Confounders<br>adjusted/matc<br>hed for  | Peer-review or<br>PP | Drugs in meta-<br>analyses                     |
|----------------|--------|---------|------------|--------------|--|-------------|---------------------|----------------|---|--------------------|-----------|----------------|---------|--|----------------------|--|
| Yan, H         | CCS    | China   | CS         | Hospi<br>tal | Admissions with<br>COVID                         | 610         | Regul<br>ar         |                | Hospitalized compared with<br>general population; Severe or<br>critical compared with non-<br>severe. Severe defined as one<br>of: RR ≥30/min, SpO2<br>≤93%, FiO2 ≤300mmHg,<br>critical defined as one of:<br>requiring MV, shock,<br>requiring ICU care for organ<br>support | Har<br>mful        | OR 7.56   | 1.17-<br>48.93 | Fair    | Age, sex and BMI   | PP                   | Yes  |
| Yi, P          | CoS    | China   | CS         | Hospi<br>tal | Admissions with<br>COVID                         | 100         | Acute               |                | Severe or critical compared<br>with non-severe. Severe<br>defined as one of: RR<br>≥30/min, SpO2 ≤93%, FiO2<br>≤300mmHg, critical defined<br>as one of: requiring MV,<br>shock, requiring ICU care for<br>organ support   | NSE                | -         | -              | Fair    | Age and BMI  | PP                   | OR<br>incalc<br>ulable<br>or<br>inaccu<br>rate |
| Li, X          | CoS    | China   | IS         | Hospi<br>tal | Admissions with<br>COVID                         | 548         | Regul<br>ar         |                | Severe or critical compared<br>with non-severe. Severe<br>defined as per IDSA & ATS<br>guidelines for CAP   | NSE                | OR 0.35   | 0.11-4.16      | Good    | Age, sex, blood tests,<br>antiviral usage  | PR                   | Yes  |
| Monreal,<br>E  | CoS    | Spain   | IS         | Hospi<br>tal | Admissions with<br>COVID                         | 138         | Regul<br>ar         |                | Berlin definition for ARDS  | Prot<br>ectiv<br>e | OR 0.16   | 0.05-0.52      | Fair    | Age, sex and time of onset   | PP                   | Yes  |
| Rentsch,<br>CT | CoS    | USA     | IS         | Hospi<br>tal | VA Birth Cohort<br>registry (54-75<br>years old) | 3789        | Regul<br>ar         |                | Hospitalization compared<br>with community care after<br>testing positive   | NSE                | OR 1.62   | 0.75-3.50      | Good    | Age, sex, race, co-<br>morbidities, vital signs,<br>medications and health<br>behaviours | PP                   | Yes  |

| First Author       | Design | Country | Drug Group | Setting           | Population                                      | Sample Size | Acute or<br>Regular | Specific drugs                                   | Outcome<br>definitions  | Result                                   | Statistic   | 95% CI  | Quality | Confounders<br>adjusted/matc<br>hed for         | Peer-review or<br>PP | Drugs in meta-<br>analyses                     |
|--------------------|--------|---------|------------|-------------------|---|-------------|---------------------|--|---|--|---|---|---------|---|----------------------|--|
| Robilotti,<br>EV   | CoS    | USA     | IS         | Hospi<br>tal      | Cancer patients<br>with<br>symptomatic<br>COVID | 423         | Regul<br>ar         | Chemot<br>herapy                                 | Requiring high-flow oxygen<br>or MV   | NSE                                      | OR 1.16   | 0.76-1.78   | Fair    | Age, race, treatment<br>and co-morbidities      | PP                   | Yes  |
| Roumier,<br>M      | CCS    | France  | IS         | Hospi<br>tal      | Admissions with<br>COVID                        | 60          | Acute               | Tocilizu<br>mab                                  | Requiring MV  | Prot<br>ectiv<br>e                       | OR 0.42   | 0.20-0.89   | Fair    | Matched for age, sex and severity               | PP                   | No   |
| Zhu, L             | CCS    | China   | IS         | Hospi<br>tal      | Renal transplant<br>admissions with<br>COVID    | 10          | Regul<br>ar         |  | Severe compared with non-<br>severe. Severe defined as per<br>7th edition of NHCPRC 7th<br>version; Time from onset to<br>shedding (unclear); Recovery<br>compared with death   | Har<br>mful                              | -   | -   | Fair    | None  | PR                   | OR<br>incalc<br>ulable<br>or<br>inaccu<br>rate |
| Argenzia<br>no, MG | CoS    | USA     | NSAI<br>D  | Hospi<br>tal      | ED and hospital presentations with COVID        | 1000        | Regul<br>ar         |  | Transfer to ICU   | NSE                                      | OR 0.78   | 0.55-1.11   | Fair    | None  | PP                   | Yes  |
| Castro,<br>VM      | XSS    | USA     | NSAI<br>D  | Com<br>munit<br>y | Patients tested<br>for COVID                    | 2271        | Regul<br>ar         | Ibuprof<br>en,<br>naproxe<br>n,<br>ketorola<br>c | Being admitted to hospital<br>compared with those testing<br>positive but remaining in the<br>community   | Prot<br>ectiv<br>e<br>and<br>har<br>mful | Ibu: OR<br>0.65<br>Nap: OR<br>0.39<br>Ket: OR<br>1.70 | Ibu: 0.42-<br>0.99<br>Nap:<br>0.14-0.93<br>Ket: 1.03-<br>2.78 | Good    | Age, sex, race,<br>Charlston index,<br>location | PP                   | Yes  |
| Hong, W            | CoS    | China   | NSAI<br>D  | Hospi<br>tal      | Admissions with<br>COVID                        | 44          | Acute               | Celecox  | Progression to severe or<br>critical disease. Severe<br>defined as one of: RR<br>≥30/min, SpO2 ≤93%, FiO2<br>≤300mmHg. Critical defined<br>as one of: requiring MV,<br>shock, requiring ICU care for<br>organ support | NSE                                      | , _   | -   | Poor    | None  | PP                   | No   |

| First Author       | Design | Country     | Drug Group | Setting      | Population   | Sample Size | Acute or<br>Regular | Specific drugs | Outcome<br>definitions  | Result                              | Statistic | 95% CI    | Quality | Confounders<br>adjusted/matc<br>hed for  | Peer-review or<br>PP | Drugs in meta-<br>analyses |
|--------------------|--------|-------------|------------|--------------|--|-------------|---------------------|----------------|---|-------------------------------------|-----------|-----------|---------|--|----------------------|----------------------------|
| Kolin, D           | CoS    | UK          | NSAI<br>D  | All          | UK Biobank   | 502536      | Regul<br>ar         |                | Being admitted to hospital<br>compared with those testing<br>positive but remaining in the<br>community   | NSE                                 | RR 0.75   | 0.47-1.18 | Fair    | Age, sex, BMI, systolic<br>BP, race and<br>deprivation                                   | PP                   | Yes                        |
| Kragholm<br>, K    | CCS    | Denmar<br>k | NSAI<br>D  | All          | National registry<br>(>30 years old<br>without heart<br>failure) | 1872        | Regul<br>ar         | Ibuprof<br>en  | Severe disease, ICU<br>admission or death. Severe<br>disease defined as ICD-10<br>code for severe adult<br>respiratory syndrome   | NSE                                 | OR 1.57   | 0.72-3.38 | Fair    | Age, sex and co-<br>morbidities  | PP                   | Yes                        |
| Rentsch,<br>CT     | CoS    | USA         | NSAI<br>D  | Hospi<br>tal | VA Birth Cohort<br>registry (54-75<br>years old)                 | 3789        | Regul<br>ar         |                | Hospitalization compared<br>with community care after<br>testing positive   | Har<br>mful                         | OR 1.18   | 0.74-1.89 | Good    | Age, sex, race, co-<br>morbidities, vital signs,<br>medications and health<br>behaviours | PP                   | Yes                        |
| Yan, H             | CCS    | China       | NSAI<br>D  | Hospi<br>tal | Admissions with<br>COVID   | 610         | Regul<br>ar         | Aspirin        | Severe or critical compared<br>with non-severe. Severe<br>defined as one of: RR<br>≥30/min, SpO2 ≤93%, FiO2<br>≤300mmHg, critical defined<br>as one of: requiring MV,<br>shock, requiring ICU care for<br>organ support | Prot<br>ectiv<br>and<br>har<br>mful | OR 0.51   | 0.12-2.13 | Fair    | Age, sex and BMI   | PP                   | Yes                        |
| Argenzia<br>no, MG | CoS    | USA         | Statin     | Hospi<br>tal | ED and hospital<br>presentations<br>with COVID                   | 1000        | Regul<br>ar         |                | Transfer to ICU   | NSE                                 | OR 1.20   | 0.89-1.63 | Fair    | None   | PP                   | Narrat<br>ive              |
| Yan, H             | CCS    | China       | Statin     | Hospi<br>tal | Admissions with<br>COVID   | 610         | Regul<br>ar         |                | Severe or critical compared<br>with non-severe. Severe<br>defined as one of: RR<br>≥30/min, SpO2 ≤93%, FiO2<br>≤300mmHg, critical defined<br>as one of: requiring MV,<br>shock, requiring ICU care for<br>organ support | Har<br>mful                         | OR 0.98   | 0.32-2.99 | Fair    | Age, sex and BMI   | PP                   | Narrat<br>ive              |

| First Author | Design | Country | Drug Group | Setting      | Population               | Sample Size | Acute or<br>Regular | Specific drugs | Outcome<br>definitions   | Result | Statistic | 95% CI         | Quality | Confounders<br>adjusted/matc<br>hed for | Peer-review or<br>PP | Drugs in meta-<br>analyses |
|--------------|--------|---------|------------|--------------|--------------------------|-------------|---------------------|----------------|--|--------|-----------|----------------|---------|---|----------------------|----------------------------|
| Yan, H       | CCS    | China   | TZD        | Hospi<br>tal | Admissions with<br>COVID | 610         | Regul<br>ar         |                | Severe or critical compared<br>with non-severe. Severe<br>defined as one of: RR<br>≥30/min, SpO2 ≤93%, FiO2<br>≤300mmHg, critical defined<br>as one of: requiring MV | NSF    | E OR 4.52 | 0.34-<br>59.54 | Fair    | Age, sex and BMI                        | PP                   | Yes                        |
|              |        |         |            |              |                          |             |                     |                | shock, requiring ICU care fo<br>organ support  | r      |           |                |         |   |                      |                            |

CoS: Cohort Study, CCS: Case Control Study, XSS: Cross Sectional Study, CaseS: Case Series, NSE: No significant evidence, NVD: No verifiable data, HR: Hazard Ratio, OR: Odds Ratio, HD: high dose, LD: low dose, PR: Peer reviewed, PP: Preprint

| First Author                   | Design | Country        | Drug Group | Setting<br>Population  | Sample Size | Acute or<br>Regular | <b>Outcome</b><br>definitions                     | Result     | Statistic                           | 95% CI                                   | Quality | Confounders<br>adjusted/<br>matched for | Peer-review<br>or Preprint | Drugs in<br>meta-<br>analyses |
|--------------------------------|--------|----------------|------------|--|-------------|---------------------|---|------------|-------------------------------------|--|---------|---|----------------------------|-------------------------------|
| Benelli,<br>G                  | CoS    | Italy          | RAA<br>SB  | Hospital Admissions<br>with COVID                                | 411         | Regular             | Deaths compared with discharges and inpatients    | NSE        | ACEI: OR<br>1.39<br>ARB: OR<br>1.54 | ACEI:<br>0.67-2.86<br>ARB: 0.79-<br>2.98 | Fair    | None                                    | PP                         | Yes                           |
| Chen,<br>Ming                  | CCS    | China          | RAA<br>SB  | Hospital Deaths and<br>discharges<br>from hospital<br>with COVID | 123         | Regular             | Deaths compared with discharges                   | NSE        | OR 1.13                             | 0.28-4.54                                | Fair    | None                                    | PP                         | Yes                           |
| Guo, T                         | XSS    | China          | RAA<br>SB  | Hospital Admissions<br>with COVID                                | 187         | Regular             | Deaths compared with discharges                   | NSE        | OR 1.70                             | 0.63-4.58                                | Fair    | None                                    | PR                         | Yes                           |
| Ip, A                          | CCS    | USA            | RAA<br>SB  | Hospital Hypertensive<br>COVID<br>discharges and<br>deaths       | 1216        | Regular             | Deaths compared with discharges                   | Protective | OR 0.66                             | 0.51-0.85                                | Poor    | None                                    | PP                         | Yes                           |
| Lee,<br>HY                     | CoS    | South<br>Korea | RAA<br>SB  | All Patients with<br>lab confirmed<br>COVID                      | 8266        | Regular             | Mortality within 60 days                          | NSE        | HR 1.07                             | 0.68-1.65                                | Good    | Age, sex and co-<br>morbidities         | PP                         | Narrative                     |
| Li, J                          | CaseS  | China          | RAA<br>SB  | Hospital Hypertensive<br>COVID<br>admissions                     | 1178        | Regular             | In hospital death<br>compared with<br>discharges  | NSE        | OR 0.76                             | 0.44-1.33                                | Good    | None                                    | PR                         | Yes                           |
| Richard<br>son, S              | CaseS  | USA            | RAA<br>SB  | Hospital Hypertensive<br>COVID<br>discharges and<br>deaths       | 5700        | Regular             | In hospital death<br>compared with<br>discharges  | NSE        | ACEI: OR<br>1.34<br>ARB: OR<br>1.21 | ACEI:<br>0.94-1.91<br>ARB: 0.89-<br>1.65 | Fair    | None                                    | PR                         | Yes                           |
| Sánchez<br>-<br>Álvarez,<br>JE | XSS    | Spain          | RAA<br>SB  | All Renal<br>replacement<br>patients with<br>COVID               | 375         | Regular             | In hospital death<br>compared with cured<br>cases | Protective | OR 0.59                             | 0.39-0.90                                | Poor    | None                                    | PR                         | Yes                           |

### Table 5. Summary of studies looking into the effect of drug groups on the mortality from COVID-19 infection

| First Author  | Design      | Country | Drug Group | Setting  | Population  | Sample Size | Acute or<br>Regular | Outcome<br>definitions   | Result                    | Statistic                 | 95% CI                                 | Quality | Confounders<br>adjusted/<br>matched for  | Peer-review<br>or Preprint | Drugs in<br>meta-<br>analyses          |
|---------------|-------------|---------|------------|----------|---|-------------|---------------------|--|---------------------------|---------------------------|--|---------|--|----------------------------|--|
| Yang, G       | CoS         | China   | RAA<br>SB  | Hospital | Hypertensive<br>COVID<br>admissions                     | 126         | Regular             | Deaths compared with discharges and inpatients   | NSE                       | OR 0.32                   | 0.07-1.51                              | Fair    | None   | PP                         | Yes                                    |
| Zhang,<br>P   | CoS         | China   | RAA<br>SB  | Hospital | Hypertensive<br>COVID<br>admissions                     | 1128        | Regular             | 28 day mortality   | Protective                | HR 0.37                   | 0.15-0.89                              | Good    | Matched for age, sex,<br>symptoms, co-<br>morbidities, CT<br>findings, CRP and<br>creatinine. Adjusted<br>for location,<br>treatment, D-Dimer<br>and unilateral lesion | PR                         | Narrative                              |
| Cao, J        | CoS         | China   | CS         | Hospital | Admissions<br>with COVID                                | 102         | Acute               | All deaths occuring in<br>data collection period<br>compared with those<br>discharged                    | NSE                       | OR 2.06                   | 0.70-6.09                              | Fair    | None   | PR                         | Yes                                    |
| Chen,<br>Ming | CCS         | China   | CS         | Hospital | Deaths and<br>discharges<br>from hospital<br>with COVID | 123         | Acute               | Deaths compared with discharges  | Harmful                   | OR 4.75                   | 1.78-12.66                             | Fair    | None   | PP                         | Yes                                    |
| Li, X         | CoS         | China   | CS         | Hospital | Admissions<br>with COVID                                | 548         | Acute               | Deaths at any point from<br>admission until final date<br>of follow up (4 weeks<br>after last admission) | HD:<br>Harmful<br>LD: NSE | HD: HR 3.5<br>LD: HR 1.26 | HD: 1.79-<br>6.86<br>LD: 0.61-<br>2.58 | Good    | Age, sex, blood tests,<br>antiviral usage  | PR                         | Narrative                              |
| Liang,<br>M   | CoS/<br>CCS | China   | CS         | Hospital | Admissions<br>with COVID                                | 120         | Acute               | Death during hospital stay   | NSE                       | OR 0.65                   | 0.10-4.14                              | Good    | None   | PP                         | Yes                                    |
| Liu,<br>Yanli | CoS         | China   | CS         | Hospital | Admissions<br>with COVID                                | 109         | Acute               | Deaths compared with discharges  | NSE                       | -                         | -                                      | Fair    | None   | PP                         | OR<br>incalculable<br>or<br>inaccurate |
| Lu, X         | CoS         | China   | CS         | ICU      | Critical<br>COVID<br>discharges and<br>deaths           | 244         | Acute               | 28 day mortality   | NSE                       | -                         | -                                      | Good    | Propensity score<br>matched and adjusted<br>for age, SpO2 and<br>lymphocyte count  | PP                         | OR<br>incalculable<br>or<br>inaccurate |

| First Author                   | Design | Country | Drug Group | Setting Population                                 | Sample Size | Acute or<br>Regular | Outcome<br>definitions  | Result     | Statistic | 95% CI    | Quality | Confounders<br>adjusted/<br>matched for | Peer-review<br>or Preprint | Drugs in<br>meta-<br>analyses |
|--------------------------------|--------|---------|------------|--|-------------|---------------------|---|------------|-----------|-----------|---------|---|----------------------------|-------------------------------|
| Sánchez<br>-<br>Álvarez,<br>JE | XSS    | Spain   | CS         | All Renal<br>replacement<br>patients with<br>COVID | 375         | Acute               | In hospital death<br>compared with cured<br>cases                   | Harmful    | OR 1.81   | 1.12-2.91 | Poor    | None                                    | PR                         | Yes                           |
| Wu, C                          | CoS    | China   | CS         | Hospital COVID<br>patients with<br>ARDS            | 201         | Acute               | Death at any point up to<br>18 days after recruitment               | Protective | HR 0.38   | 0.20-0.72 | Fair    | None                                    | PR                         | Yes                           |
| Zhang,<br>S                    | CoS    | China   | CS         | Hospital Admissions<br>with COVID                  | 262         | Acute               | 28 day mortality  | Harmful    | OR 4.97   | 2.85-8.45 | Poor    | None                                    | PP                         | Yes                           |
| Zhou, F                        | CoS    | China   | CS         | Hospital Admissions<br>with COVID                  | 191         | Acute               | Deaths compared with discharges                                     | Harmful    | OR 3.18   | 1.63-6.19 | Poor    | None                                    | PR                         | Yes                           |
| Monreal<br>, E                 | CoS    | Spain   | IS         | Hospital Admissions<br>with COVID                  | 138         | Regular             | Deaths compared with<br>those with a final<br>outcome               | NSE        |           |           | Fair    | Age, sex and time of onset              | PP                         | Yes                           |
| Roumie<br>r, M                 | CCS    | France  | IS         | Hospital Admissions<br>with COVID                  | 60          | Acute               | In-hospital deaths within<br>2 days of end of<br>recruitment period | NSE        |           |           | Fair    | Matched for age, sex<br>and severity    | PP                         | No                            |

CoS: Cohort Study, CCS: Case Control Study, XSS: Cross Sectional Study, CaseS: Case Series, NSE: No significant evidence, NVD: No verifiable data, HR: Hazard Ratio, OR: Odds Ratio, HD: high dose, LD: low dose, PR: Peer reviewed, PP: Preprint

# **Appendix 4: funnel plots**



Fig 12. Funnel plot to assess for publication bias from studies measuring susceptibility to COVID-19 in those taking RAASBs. ACEI – angiotensin-converting enzyme inhibitor, ARB - angiotensin-II-receptor blocker, OR - odds ratio.



Fig 13. Funnel plot to assess for publication bias from studies measuring severity of COVID-19 in those taking RAASBs. ACEI – angiotensin-converting enzyme inhibitor, ARB - angiotensin-II-receptor blocker, OR - odds ratio.

## Appendix 5: data collection table headings

| Result (#)  |
|---|
| Study (#)   |
| First Author  |
| Journal   |
| Type (Peer reviewed or preprint)                            |
| Date  |
| Country   |
| Setting (Hospital, community or both)                       |
| Sample Size   |
| Drug Group  |
| Specific Drug(s)  |
| Regular or Acute  |
| Study Design  |
| Population Demographic                                      |
| Cited by (#)  |
| Outcome (1-6)   |
| Significance (harm, benefit or no statistically significant |
| evidence)   |
| Odds Ratio/Relative Risk                                    |
| p-value   |
| Confidence Intervals  |
| Control Group   |
| Subgroup(s) Affected  |
| Identified Confounding Factors                              |
| Comments  |

## **Appendix 6: forest plots**



### Fig 14. Forest plot for effect of NSAIDs on severity of COVID-19. NSAID - non-steroidal

anti-inflammatory drug, OR - odds ratio, CI - confidence interval.



Fig 15. Forest plot for effect of NSAIDs on susceptibility to COVID-19. NSAID - nonsteroidal anti-inflammatory drug, OR – odds ratio, CI – confidence interval.



Fig 16. Forest plot for effect of statins on susceptibility to COVID-19. OR – odds ratio, CI - confidence interval.

medRxiv preprint doi: https://doi.org/10.1101/2021.09.17.21262724; this version posted September 21, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.





### Fig 17 . Forest plot for effect of TZDs on susceptibility to COVID-19. TZD -

thiazolidinediones, OR – odds ratio, CI – confidence interval.



**Fig 18. Forest plot of susceptibility to COVID-19 for those taking regular RAAS blockers, split by drug group.** *ACEI* – *angiotensin-converting enzyme inhibitor, ARB* – *angiotensin-II-receptor blocker, OR* – *odds ratio, CI* – *confidence interval.* 



**Fig 19. Forest plot of susceptibility to COVID-19 for those taking regular RAAS blockers, split by study design.** *ACEI* – *angiotensin-converting enzyme inhibitor, ARB* – *angiotensin-II-receptor blocker, OR* – *odds ratio, CI* – *confidence interval.* 





Fig 20. Forest plot of susceptibility to COVID-19 for those taking regular RAAS blockers, split by level of adjustment. ACEI – angiotensin-converting enzyme inhibitor, ARB – angiotensin-II-receptor blocker, OR – odds ratio, CI – confidence interval.

# Appendix 7: sensitivity analysis for removal of preprints

| Peer-Reviewed Before | Subsequently Peer- | Not Peer-Reviewed |
|----------------------|--------------------|-------------------|
| Study                | Reviewed           |                   |
| Cao, J               | Argenziano, MG     | Benelli, G        |
| Feng, Y              | Bean, DM           | Cao, C            |
| Gisondi, P           | Caraballo, C       | Castro, VM        |
| Guo, T               | Hong, W            | Chen, M           |
| Li, J                | Huh, K             | Dauchet, L        |
| Li, X                | Kolin, DA          | Dooley, H         |
| Mancia, G            | Kragholm, K        | Feng, Z           |
| Meng, J              | Lu, X              | Ip, A             |
| Reynolds, HR         | Michelena, X       | Khawaja, AP       |
| Richardson, S        | Monreal, E         | Lee, HY           |
| Sánchez-Álvarez, JE  | Robilotti, EV      | Liang, M          |
| Wu, C                | Shah, SJ           | Liu, Yingxia      |
| Zhang, P             | Wang, D            | Liu, Yanli        |
| Zhou, F              | Yang, G            | Lui, GCY          |
| Zhu, L               | Yi, P              | Rentsch, CT       |
|                      |                    | Roumier, M        |
|                      |                    | Yan, H            |
|                      |                    | Zeng, Z           |
|                      |                    | Zhang, S          |

# **RAAS** blockers






medRxiv preprint doi: https://doi.org/10.1101/2021.09.17.21262724; this version posted September 21, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC 4.0 International license .









medRxiv preprint doi: https://doi.org/10.1101/2021.09.17.21262724; this version posted September 21, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC 4.0 International license .

## **Corticosteroids**

No meta-analysis was performed for susceptibility with corticosteroids.

Severity with acute corticosteroids leaves Wang, D only: OR 2.16 (95% CI 0.49 -9.43).



## **Immunosuppressants**

Susceptibility with immunosuppressants had one study not peer reviewed (Rentsch CT et al.) and sensitivity analysis was run for this already within the study.

medRxiv preprint doi: https://doi.org/10.1101/2021.09.17.21262724; this version posted September 21, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC 4.0 International license .



No meta-analysis was performed for mortality related to immunosuppressants.

## **NSAIDs**, statins, TZDs



No meta-analyses were performed for mortality related to either NSAIDs, statins or TZDs.

No meta-analyses were performed for severity related to either statins or TZDs.

medRxiv preprint doi: https://doi.org/10.1101/2021.09.17.21262724; this version posted September 21, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC 4.0 International license.



No meta-analyses were performed for MCRAs.