

Identifying molecular mediators of the relationship between body mass index and endometrial cancer risk: a Mendelian randomization analysis

Emma Hazelwood^{1,2}, Eleanor Sanderson^{1,2}, Vanessa Y Tan^{1,2}, Katherine S Ruth³, Timothy M Frayling⁴, Niki Dimou⁵, Marc J Gunter⁵, Laure Dossus⁵, Claire Newton⁶, Neil Ryan^{6,7}, Dimitri J Pournaras⁸, Tracy A O'Mara⁹, George Davey Smith^{1,2}, Richard M Martin^{1,2,10}, James Yarmolinsky^{1,2}

¹MRC Integrative Epidemiology Unit, University of Bristol, Bristol, UK

²Bristol Medical School, University of Bristol, Bristol, UK

³University of Exeter Medical School, University of Exeter, Exeter, UK

⁴Genetics of Complex Traits, College of Medicine and Health, University of Exeter, Exeter, UK

⁵Nutrition and Metabolism Branch, International Agency for Research on Cancer, Lyon, France

⁶Department of Gynecology, St Michaels Hospital University Hospitals Bristol NHS Foundation Trust, Bristol, UK

⁷The Academic Women's Health Unit, Translational Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK

⁸Department of Upper GI and Bariatric/Metabolic Surgery, North Bristol NHS Trust, Southmead Hospital, Bristol, UK

⁹Department of Genetics and Computational Biology, QIMR Berghofer Medical Research Institute, Brisbane, Queensland, Australia

¹⁰National Institute for Health Research Bristol Biomedical Research Centre, University of Bristol, University Hospitals Bristol and Weston NHS Foundation Trust, Bristol, UK

Corresponding author:

25 James Yarmolinsky, PhD
26 MRC Integrative Epidemiology Unit
27 Population Health Sciences
28 Bristol Medical School
29 University of Bristol
30 Bristol, UK
31 james.yarmolinsky@bristol.ac.uk

Abstract

Background: Endometrial cancer is the most common gynaecological cancer in high-income countries. Elevated body mass index (BMI) is an established modifiable risk factor for this condition and is estimated to confer a larger effect on endometrial cancer risk than any other cancer site. However, the molecular mechanisms underpinning this association remain unclear. We used Mendelian randomization (MR) to evaluate the causal role of 14 molecular risk factors (hormonal, metabolic, and inflammatory markers) in endometrial cancer risk. We then evaluated and quantified the potential mediating role of these molecular traits in the relationship between BMI and endometrial cancer.

Methods and Findings: Genetic instruments to proxy 14 molecular risk factors and BMI were constructed by identifying single-nucleotide polymorphisms (SNPs) reliably associated ($P < 5.0 \times 10^{-8}$) with each respective risk factor in previous genome-wide association studies (GWAS). Summary statistics for the association of these SNPs with overall and subtype-specific endometrial cancer risk (12,906 cases and 108,979 controls) were obtained from a GWAS meta-analysis of the Endometrial Cancer Association Consortium (ECAC), Epidemiology of Endometrial Cancer Consortium (E2C2), and UK Biobank. SNPs were combined into multi-allelic models and odds ratios (ORs) and 95% confidence intervals (95% CIs) were generated using inverse-variance weighted random-effects models. The mediating roles of the molecular risk factors in the relationship between BMI and endometrial cancer were then estimated using multivariable MR. In MR analyses, there was strong evidence that BMI (OR per SD increase: 1.88, 95% CI: 1.69 to 2.09, $P = 3.87 \times 10^{-31}$), total testosterone (OR per inverse normal transformed nmol/L increase: 1.64, 95% CI: 1.43 to 1.88, $P = 1.71 \times 10^{-12}$), bioavailable testosterone (OR per inverse normal transformed nmol/L increase: 1.46, 95% CI: 1.29 to 1.65, $P = 3.48 \times 10^{-9}$), fasting insulin (OR per natural log transformed pmol/L increase: 3.93, 95% CI: 2.29 to 6.74, $P = 7.18 \times 10^{-7}$) and sex hormone-binding globulin (SHBG, OR per inverse normal transformed nmol/L increase: 0.71, 95% CI:

0.59 to 0.85, $P = 2.07 \times 10^{-4}$) had a causal effect on endometrial cancer risk. Additionally, there was suggestive evidence that total serum cholesterol (OR per mg/dL increase: 0.90, 95% CI: 0.81 to 1.00, $P = 4.01 \times 10^{-2}$) had an effect on endometrial cancer risk. In mediation analysis using multivariable MR, we found evidence for a mediating role of fasting insulin (19% total effect mediated, 95% CI: 5 to 34%, $P = 9.17 \times 10^{-3}$), bioavailable testosterone (15% mediated, 95% CI: 10 to 20%, $P = 1.43 \times 10^{-8}$), and SHBG (7% mediated, 95% CI: 1 to 12%, $P = 1.81 \times 10^{-2}$) in the relationship between BMI and endometrial cancer risk. The primary limitations of this analysis include the assumption of linear relationships across univariable and multivariable analyses and the restriction of analyses to individuals of European ancestry.

Conclusions: Our comprehensive Mendelian randomization analysis provides insight into potential causal mechanisms linking BMI with endometrial cancer risk and suggests pharmacological targeting of insulinemic and hormonal traits as a potential strategy for the prevention of endometrial cancer.

Introduction

Endometrial cancer is the most common gynaecological cancer in high-income countries and the second most common globally [1, 2]. In 2020, there were 417,367 new cases diagnosed and 97,370 endometrial cancer-related deaths worldwide [3]. In contrast to several other cancer types where incidence rates have been declining over the past two decades, the global incidence of endometrial cancer continues to increase [4-8].

Elevated body mass index (BMI) is an established risk factor for endometrial cancer and is estimated to confer a larger effect on risk of this malignancy than any other cancer type [9-11]. A recent meta-analysis of 30 prospective studies reported that each 5 kg/m² increase in BMI was associated with a 54% (95% CI: 47 to 61%) higher risk of endometrial cancer [12-14]. It is estimated that excess adiposity accounts for 34% of global endometrial cancer cases, with the increasing incidence of endometrial cancer mirroring rising levels of obesity worldwide [15-17]. Lifestyle and dietary interventions encouraging maintenance of a healthy weight therefore remain cornerstones for the primary prevention of endometrial cancer [9]. Alongside weight management strategies, greater characterisation of the molecular mechanisms underpinning an effect of excess adiposity on endometrial cancer could provide a complementary approach to cancer prevention through the development of pharmacological interventions targeting these traits in high-risk groups.

Observational epidemiological studies have reported associations between several hormonal, metabolic, and inflammatory factors linked to obesity and endometrial cancer, including bioavailable testosterone, sex hormone-binding globulin (SHBG), oestradiol and fasting insulin [18-22]. However, conventional observational studies are susceptible to residual confounding (due to unmeasured or imprecisely measured confounders), reverse causation, and other forms of bias which undermine robust

causal inference. Therefore, the causal nature of these risk factors, and thus their suitability as effective intervention targets for endometrial cancer prevention, remains unclear.

Mendelian randomization (MR) is an analytical approach that uses germline genetic variants as instruments (“proxies”) for risk factors to evaluate the causal effects of these factors on disease outcomes in observational settings [23, 24]. Since germline genetic variants are randomly assorted at meiosis, MR analyses should be less prone to confounding by lifestyle and environmental factors than conventional observational studies. Furthermore, since germline genetic variants are fixed at conception, MR analyses are not subject to reverse causation bias. The statistical power and precision of MR analysis can be increased by employing a “two-sample MR” framework in which summary genetic association data from two independent samples – one representing genetic variant-exposure associations and one representing genetic variant-outcome associations – are synthesised to estimate causal effects [25].

Recent MR studies have suggested potential causal relationships between circulating levels of several molecular traits, including low-density lipoprotein (LDL) cholesterol, insulin, total and bioavailable testosterone, and sex hormone-binding globulin (SHBG) and endometrial cancer risk, and have confirmed a causal role of BMI in endometrial cancer risk [17, 26-32]. However, many previously reported molecular risk factors for endometrial cancer from conventional observational studies remain untested in an MR framework, meaning the causal relevance of these factors in disease onset is unclear. Additionally, no MR studies to date have attempted to quantify the potential mediating role of these factors in the relationship between BMI and endometrial cancer risk.

Given the unclear causal relevance of previously reported molecular traits in endometrial cancer aetiology, we used a two-sample MR approach to evaluate the causal role of 14 endogenous sex hormones, metabolic traits, and inflammatory markers in endometrial cancer risk (overall and in

112

113 endometrioid and non-endometrioid subtypes). We then used multivariable MR to evaluate and
114 quantify the mediating role of these molecular traits in the relationship between BMI and endometrial
115 cancer risk.

Methods

Our analytical strategy was as follows: first, we attempted to corroborate previous MR findings that there was evidence of a causal relationship between BMI and endometrial cancer risk (overall and by histological subtype); second, we examined evidence for a causal relationship between previously reported molecular factors and endometrial cancer risk (overall and by histological subtype); third, we evaluated the causal relationship between BMI and those molecular risk factors that were confirmed to influence endometrial cancer risk (overall and by histological subtype); finally, we performed a mediation analysis to quantify the proportion of the total effect of BMI on endometrial cancer risk that was mediated by each identified trait.

Endometrial cancer study population

Summary genetic association data on overall and subtype-specific endometrial cancer risk were obtained from a genome-wide association study (GWAS) of 12,906 cases (including 8,758 endometrioid and 1,230 non-endometrioid endometrial cancer cases) and up to 108,979 controls of European ancestry [31]. This meta-GWAS combined 17 previously reported studies from the Endometrial Cancer Association Consortium (ECAC), the Epidemiology of Endometrial Cancer Consortium (E2C2), and UK Biobank, with four studies contributing samples to more than one genotyping project. Participants were recruited from Australia, Belgium, Germany, Poland, Sweden, the UK, and the USA and associations were adjusted for principal components of ancestry. Genotyping was performed using one of several Illumina arrays and imputation was performed using the 1000 Genomes Phase 3 reference panel [33]. Further information on this meta-GWAS including study-specific genotyping, imputation, and quality control procedures is provided in **S2 Appendix**.

138

139 *Identification of previously reported molecular risk factors for endometrial cancer*

140 We performed two pragmatic searches of the literature using PubMed. The first search
141 identified previously published MR analyses of molecular risk factors for endometrial cancer. The second
142 search identified narrative or systematic reviews of potential molecular mechanisms underpinning the
143 relationship between obesity and endometrial cancer (additional information on search strategies used
144 in literature reviews is presented in **S3 Appendix**). Combined, these literature reviews identified 20
145 unique molecular traits which could mediate the effect of BMI on endometrial cancer risk, of which 14
146 had suitable genetic instruments available. These traits include nine metabolic factors (low-density
147 lipoprotein (LDL) cholesterol, high-density lipoprotein (HDL) cholesterol, total serum cholesterol,
148 triglycerides, blood glucose, fasting insulin, insulin-like growth factor 1 (IGF-1), adiponectin, and leptin);
149 three endogenous sex hormones or traits that regulate their bioactivity (total and bioavailable
150 testosterone, and SHBG); and two inflammatory markers (interleukin-6 (IL-6) and C-reactive protein
151 (CRP), measured as high-sensitivity CRP) (**Fig 1**) [29, 34-41]. Summary genetic association data on BMI
152 were obtained from a GWAS of 681,275 individuals of European ancestry [42]. Additional information on
153 participant demographics and covariates included in adjustment strategies across each GWAS are
154 presented in **S4 Table**. All studies contributing data to these analyses had the relevant institutional
155 review board approval from each country, in accordance with the Declaration of Helsinki, and all
156 participants provided informed consent.

157

158 *Statistical analyses*

MR analysis can generate unbiased estimates of causal effects of risk factors on disease outcomes if the following assumptions are met: (i) the instrument strongly associates with the exposure (“relevance”), (ii) there is no confounding of the instrument-outcome relationship (“exchangeability”), and (iii) the instrument only affects the outcome through the exposure (“exclusion restriction”) (**Fig. 2**) [43].

To construct genetic instruments for BMI and previously reported molecular risk factors, we obtained single-nucleotide polymorphisms (SNPs) reliably ($P < 5 \times 10^{-8}$) and independently ($r^2 < 0.001$) associated with each trait. To construct a genetic instrument for leptin, we restricted genetic variants to *cis*-acting SNPs (i.e. in or within $\pm 100\text{kb}$ from the gene encoding the protein). For leptin, IL-6 and CRP analyses, SNPs were permitted to be in weak linkage disequilibrium (LD) ($r^2 < 0.10$) to maximise instrument strength. For all traits where instruments consisted of SNPs in weak LD (i.e. leptin, IL-6 and CRP), standard errors for causal estimates were inflated to account for correlation between SNPs with reference to the 1000 Genomes Phase 3 reference panel [33, 44].

For traits instrumented by a single SNP, the Wald ratio was used to generate effect estimates and the delta method was used to approximate standard errors [45]. For traits instrumented by two or more SNPs, inverse-variance weighted (IVW) random-effects models were used to estimate causal effects [45]. A Bonferroni correction was applied as a heuristic to account for multiple testing in MR analyses for the 15 risk factors (14 molecular traits and BMI) investigated. Results below this threshold were classified as “strong evidence” ($P < 3.33 \times 10^{-3}$ ($0.05/15$ traits)), whereas results between this threshold and $P < 0.05$ were classified as “suggestive evidence”.

When using genetic instruments, there is potential for horizontal pleiotropy - when a genetic variant influences an outcome through a biological pathway independent to the exposure, a violation of the “exclusion restriction” criterion [46]. We evaluated the presence of horizontal pleiotropy by

performing various sensitivity analyses. First, for instruments consisting of ≥ 10 SNPs, we re-calculated causal estimates obtained from IVW models using MR-Egger regression, weighted median estimation, and weighted mode estimation (additional information on these sensitivity analyses is provided in **S2 Appendix**) [46-48]. Each of these models makes different assumptions regarding the nature of horizontal pleiotropy in the genetic instrument and therefore performing all three can provide complementary support to IVW models in evaluating the presence of horizontal pleiotropy. These models were not employed when instruments consisted of < 10 SNPs because of their reduced statistical power to detect horizontal pleiotropy in these settings (additional information on these sensitivity analyses is provided in **S2 Appendix**). Second, we performed “leave-one-out” analyses for all findings showing strong or suggestive evidence of effects in IVW models ($P < 0.05$) for traits where instruments consisted of ≥ 10 SNPs and findings were consistent across MR-Egger, weighted median, and weighted mode sensitivity analyses or where instruments consisted of < 10 SNPs. This approach sequentially removes each SNP from an instrument and then re-calculates the overall effect estimate to examine robustness of findings to individual influential SNPs in IVW models.

Instruments were derived from sex-combined GWAS for all traits other than those related to endogenous sex hormones to maximise statistical power where there was limited evidence of sex-specificity of SNP associations. As a sensitivity analysis we also re-performed MR analyses using sex-specific instruments where possible. For BMI, all analyses with strong or suggestive evidence for an effect ($P < 0.05$) were repeated using genome-wide significant ($P < 5.0 \times 10^{-8}$) variants identified in female-specific analyses. Likewise, for fasting insulin and CRP analyses, the effect estimates and standard errors of SNPs used to instrument these traits were replaced with female-specific values where there was previous evidence of sex-specificity of associations (trait-specific criteria for identifying sex-specific effects are presented in **S2 Appendix**). Findings from sex-specific sensitivity analyses are presented in **Tables S8, S17, S28-29**. Finally, Steiger filtering was performed across all analyses to

identify and subsequently remove any SNPs which explained more variance in the outcome than the exposure (i.e. suggesting misspecification of the causal direction between traits) [49].

Mediation analysis

For all molecular traits that were identified as being on the causal pathway between BMI and endometrial cancer risk, we used multivariable MR to generate estimates of the direct effect (i.e. the remaining effect of the exposure on the outcome when the effect of the candidate mediator on the outcome has been adjusted for) and indirect effect (i.e. the effect of the exposure on the outcome through the candidate mediator) using the product of coefficients method [50]. The proportion of the total effect of BMI on endometrial cancer risk (“proportion mediated”) that was mediated by each molecular trait was calculated using these estimates. In the case of fasting insulin, due to weak instrument bias, several different approaches were employed to attempt to maximise conditional instrument strength (for further information on these analyses see **S2 Appendix**). Standard errors for the proportion mediated were calculated using the delta method [51]. In addition, we aimed to perform additional mediation analyses combining all mediators into a single model to examine the extent to which these mediators influenced endometrial cancer independently or via shared biological pathways (presumed relationships between BMI, fasting insulin, SHBG, bioavailable testosterone, and endometrial cancer risk are presented in **Fig 3**). When all putative mediators were combined into a single model with BMI, however, there was persistent weak instrument bias. Of various alternate approaches examined to minimise this bias, the restriction of models to pairs of mediators (without inclusion of BMI) was found to generate the largest conditional F-statistics for each mediator included in the model (for further information on these analyses see **S2 Appendix**).

228

229 *Sample overlap sensitivity analyses*

230 There was moderate sample overlap (52.2-62.4%) across some analyses which can bias MR
231 estimates toward the confounded observational estimate in the presence of weak instrument bias [52]
232 (**S5 Table**). This bias can be inflated by “Winner’s curse”, in which weights for genetic instruments are
233 derived from discovery samples that overlap with outcome samples. Though instruments in this analysis
234 were constructed from genome-wide significant variants ($P < 5.0 \times 10^{-8}$) which should minimise the
235 possibility of weak instrument bias, we performed the following sensitivity analyses to evaluate whether
236 our findings could be influenced by sample overlap: i) for analyses examining the effect of blood glucose
237 on endometrial cancer risk, we re-performed MR analyses using alternate GWAS data for this trait
238 where there was no sample overlap [53, 54]; ii) for analyses examining the effect of BMI on total
239 testosterone, bioavailable testosterone, SHBG and endometrial cancer, we re-performed MR analyses
240 using alternate GWAS data for BMI where there was no sample overlap [55]; and iii) for analyses
241 examining the effect of total testosterone, bioavailable testosterone, SHBG and IGF-1 on endometrial
242 cancer (where suitable alternate GWAS data were not available), we re-constructed instruments for sex
243 hormones using more conservative P -value thresholds ($P < 5.0 \times 10^{-9}$, $P < 5.0 \times 10^{-10}$) to minimise the
244 potential for inclusion of weak instruments into analyses and then re-performed MR analyses. Similarly,
245 in mediation analysis, due to the presence of sample overlap and possible influence of Winner’s curse,
246 for any trait with sample overlap in the same multivariable MR model the analysis was repeated with a
247 more stringent P value ($P < 5.0 \times 10^{-9}$) used for instrument construction.

248 All statistical analyses were performed using R (Vienna, Austria) version 4.0.2. Additional
249 information on statistical packages used across various analyses is presented in **S2 Appendix**.

Results

Evaluating the effect of BMI on endometrial cancer risk

In MR analyses, there was strong evidence for an effect of BMI on risk of overall endometrial cancer (odds ratio (OR) per SD (4.7 kg/m²) increase in BMI: 1.88, 95% confidence interval (CI): 1.69 to 2.09, $P = 3.87 \times 10^{-31}$) (**Fig 4, Table 2**). This finding was consistent across sensitivity analyses examining evidence of horizontal pleiotropy, including MR-Egger, weighted median, and weighted mode models, in analyses using a female-specific BMI instrument, in analyses exploring potential Winner's curse bias in instrument construction, and in the leave-one-out analysis (**S7 Figure, S8 Table, S9 Table**).

In subtype-stratified analyses, there was evidence to support an effect of BMI on risk of both endometrioid (OR per SD (4.7 kg/m²) increase in BMI: 1.89, 95% CI: 1.65 to 2.16, $P = 1.67 \times 10^{-20}$) and non-endometrioid endometrial cancer (OR per SD (4.7 kg/m²) increase in BMI: 1.67, 95% CI: 1.19 to 2.35, $P = 3.03 \times 10^{-3}$) (**Fig 4, Table 2**). These findings were robust to sensitivity analyses for endometrioid endometrial cancer; however, findings were less consistent for non-endometrioid endometrial cancer in sensitivity analyses using female-specific BMI instruments (**S10-11 Figure, S8-9 Table**). Therefore, only overall and endometrioid endometrial cancer were included in follow-up analyses.

Evaluating the effect of previously reported molecular risk factors on endometrial cancer risk

When examining the effect of previously reported molecular risk factors on overall endometrial cancer risk, there was strong evidence for an effect of total testosterone (OR per increase in inverse-normal transformed (INT) nmol/L total testosterone: 1.64, 95% CI: 1.43 to 1.88, $P = 1.71 \times 10^{-12}$), bioavailable testosterone (OR per increase in INT nmol/L bioavailable testosterone: 1.46, 95% CI: 1.29 to 1.65, $P = 3.48 \times 10^{-9}$), fasting insulin (OR per increase in natural log transformed pmol/L fasting insulin

3.93, 95% CI: 2.29 to 6.74, $P = 7.18 \times 10^{-7}$), and SHBG (OR per increase in INT nmol/L SHBG 0.71, 95% CI: 0.59 to 0.85, $P = 2.07 \times 10^{-4}$) on endometrial cancer risk (**Fig 5; Table 3**). In addition, there was suggestive evidence for an effect of total serum cholesterol (OR per increase in SD (41.7 mg/dL) total serum cholesterol 0.90, 95% CI: 0.81 to 1.00, $P = 4.01 \times 10^{-2}$) on overall endometrial cancer risk. These findings were consistent across sensitivity analyses (**S12-S16 Figure, S17 Table, S8-S9 Table**).

In subtype-stratified analyses, there was strong evidence to support an effect of total testosterone (OR per increase in INT nmol/L total testosterone: 1.60, 95% CI: 1.36 to 1.87, $P = 8.70 \times 10^{-9}$), bioavailable testosterone (OR per increase in INT nmol/L bioavailable testosterone: 1.46, 95% CI: 1.29 to 1.65, $P = 3.48 \times 10^{-9}$), fasting insulin (OR per increase in natural log transformed pmol/L fasting insulin: 4.64, 95% CI: 2.30 to 9.36, $P = 1.84 \times 10^{-5}$), and SHBG (OR per increase in INT nmol/L SHBG: 0.65, 95% CI: 0.54 to 0.80, $P = 3.31 \times 10^{-5}$) on endometrioid endometrial cancer risk (**Fig 5; Table 3**). Findings were consistent across all sensitivity analyses (**S19-S22 Figure, S9 Table, S17 Table**).

Evaluating the effect of BMI on previously reported molecular risk factors

When examining the effect of BMI on molecular traits confirmed to have a causal effect on endometrial cancer (either overall or endometrioid subtype), there was evidence for an effect of BMI on fasting insulin (change in natural log transformed fasting insulin: 0.17, 95% CI: 0.15 to 0.19, $P = 1.51 \times 10^{-74}$), SHBG (change in INT SHBG: -0.17, 95% CI: -0.19 to -0.16, $P = 4.86 \times 10^{-125}$), bioavailable testosterone (change in INT bioavailable testosterone: 0.26, 95% CI: 0.23 to 0.29, $P = 9.97 \times 10^{-68}$), total testosterone (change in INT total testosterone: 0.08, 95% CI: 0.05 to 0.11, $P = 9.04 \times 10^{-10}$), and CRP (change in ln-transformed CRP: 0.35, 95% CI: 0.32 to 0.38, $P = 2.67 \times 10^{-127}$) (**Fig 6; Table 4**). The direction of effect in MR analyses examining the effect of BMI on total testosterone was inconsistent when employing a weighted mode model, suggesting the potential presence of horizontal pleiotropy. Although there was

little evidence for a causal effect of BMI on total serum cholesterol in the IVW model, there was some evidence for an effect across all three MR sensitivity analysis models, suggesting that horizontal pleiotropy may be biasing the IVW estimate towards the null. All other findings were consistent across the various sensitivity analyses (**S23-S27 Figure, S8 Table, S17-S18 Table**).

Mendelian randomization mediation analysis

In mediation analyses evaluating the potential mediating role of molecular traits previously shown to be on the causal pathway between BMI and endometrial cancer, there was evidence for a mediating role of bioavailable testosterone (15% mediated, 95% CI: 10 to 20%, $P = 1.43 \times 10^{-8}$), fasting insulin (11% of total effect mediated, 95% CI: 1 to 21%, $P = 2.89 \times 10^{-2}$), and SHBG (7% mediated, 95% CI: 1 to 12%, $P = 1.81 \times 10^{-2}$) in the relationship between BMI and overall endometrial cancer risk (**Table 5**). There was also evidence for a mediating role of bioavailable testosterone (15% mediated, 95% CI: 9 to 22%, $P = 2.15 \times 10^{-6}$) and fasting insulin (16% mediated, 95% CI: 1 to 21%, $P = 2.89 \times 10^{-2}$) in the relationship between BMI and endometrioid endometrial cancer risk (**Table 5**). Although there was little evidence for a mediating role of SHBG in the relationship between sex-combined BMI and endometrioid endometrial cancer (2% mediated, 95% CI: -9 to 14%, $P = 6.87 \times 10^{-1}$), in the female-specific BMI sensitivity analysis there was strong evidence for a mediating role of female-specific SHBG in the relationship between BMI and endometrioid endometrial cancer (8% mediated, 95% CI: 3 to 13%, $P = 3.38 \times 10^{-3}$). Other than this, findings were consistent across sex-specific BMI, fasting insulin and CRP sensitivity analyses (**S28-S29 Table**).

The conditional F-statistic for both fasting insulin ($F=2$) and BMI ($F=6$) in the multivariable MR performed to evaluate the “proportion mediated” by fasting insulin was < 10 , indicating that there may

be weak instrument bias in these analyses (i.e. over- or underestimation of the “proportion mediated” by fasting insulin) [56]. When re-performing the “proportion mediated” analysis for fasting insulin using an alternative approach (i.e. using an alternative fasting insulin instrument with a larger sample size and limiting the number of SNPs included in the BMI instrument to the 100 with the strongest evidence of association using an LD threshold $r^2 < 0.001$), we found that fasting insulin mediated 19% (95% CI: 5 to 34%, $P = 9.17 \times 10^{-3}$) of the relationship between BMI and overall endometrial cancer risk and 21% (95% CI: 5 to 38%, $P = 1.17 \times 10^{-2}$) of the relationship between BMI and endometrioid endometrial cancer risk (**S30 Table**).

In mediation analyses combining pairs of mediators into a single model, the effect of fasting insulin on overall endometrial cancer risk attenuated (~40% log OR reduction) when SHBG (a presumed downstream mediator of fasting insulin) was included in the model (OR per increase in natural log transformed pmol/L fasting insulin: 2.28, 95% CI: 1.34 to 3.86, $P = 2.85 \times 10^{-3}$). The effect of SHBG on overall endometrial cancer attenuated fully when bioavailable testosterone (a presumed downstream mediator of SHBG) was included in the model (OR per increase in INT nmol/L SHBG: 1.08, 95% CI: 0.86 to 1.36, $P = 5.00 \times 10^{-1}$) (**S34 Table**). The effect of fasting insulin on overall endometrial cancer was strongly attenuated when bioavailable testosterone was included in the model (OR per increase in natural log transformed pmol/L fasting insulin: 1.22, 95% CI: 0.48 to 3.11, $P = 6.78 \times 10^{-1}$), which could reflect mediation of the effect of fasting insulin on endometrial cancer via bioavailable testosterone or the presence of conditionally weak instruments in this model. This could result in over- or underestimation of the proportion of the effect of fasting insulin mediated by bioavailable testosterone.

For the endometrioid histological subtype, using the same approach, the effect of fasting insulin on endometrioid endometrial cancer did not markedly change (~14% logOR reduction) when SHBG was included in the model (OR per increase in natural log transformed pmol/L fasting insulin: 3.74, 95% CI:

0.74 to 19.01, $P = 1.56 \times 10^{-1}$). However, the effect of SHBG on endometrioid endometrial cancer attenuated fully when bioavailable testosterone was included in the model (OR per increase in INT nmol/L SHBG: 1.16, 95% CI: 0.81 to 1.65, $P = 4.12 \times 10^{-1}$). As with analyses of overall endometrial cancer, when fasting insulin and bioavailable testosterone were combined into a single model, the effect of fasting insulin on endometrioid endometrial cancer attenuated toward the null (OR per increase in natural log transformed pmol/L fasting insulin: 1.05, 95% CI: 0.36 to 3.03, $P = 9.33 \times 10^{-1}$), potentially reflecting mediation via bioavailable testosterone or persistent weak instrument bias in this model.

Discussion

Our systematic Mendelian randomization (MR) analysis of 14 previously reported molecular risk factors and body mass index (BMI) in 12,906 endometrial cancer cases and 108,979 controls provided evidence for roles of elevated BMI, fasting insulin, total and bioavailable testosterone, and sex hormone-binding globulin (SHBG) in risk of overall and endometrioid endometrial cancer. In mediation analyses, we found evidence that fasting insulin, bioavailable testosterone concentrations, and SHBG partially mediated the effect of BMI on overall endometrial cancer risk. When combining pairs of mediators together into a single model, we found evidence that an effect of fasting insulin on endometrial cancer was partially mediated by SHBG levels and that an effect of SHBG on endometrial cancer was largely mediated by bioavailable testosterone levels. An effect of fasting insulin on endometrial cancer risk was also strongly attenuated upon adjustment for bioavailable testosterone levels which could reflect mediation of this effect by bioavailable testosterone or conditionally weak instrument bias for fasting insulin concentrations in this analysis. Our analyses found little evidence that several previously reported molecular risk factors, including several metabolic factors (e.g. LDL-C, HDL-C, IGF-1, adiponectin, leptin) and inflammatory markers (CRP, IL-6), were causally implicated in overall or endometrioid endometrial cancer risk.

Several of the findings in this analysis are consistent with evidence from prior conventional observational and MR analyses. For example, the effect of BMI on endometrial cancer risk and the stronger evidence of an effect on endometrioid, as compared to non-endometrioid, endometrial cancer is well-established in the literature and has been shown previously in an MR analysis that used an alternative strategy for instrument construction. Our findings supporting a causal effect of BMI on endometrial cancer risk (OR 1.88, 95% CI: 1.69 to 2.09 per SD (4.7 kg/m²) increase) are larger in magnitude than those from pooled analyses of conventional observational analyses (e.g. the World

Cancer Research Fund (WCRF) pooled analysis of 26 prospective studies: RR 1.50, 95% CI 1.42 to 1.59 per 5.0 kg/m² increase), consistent with previous comparisons of observational and MR estimates across other cancer sites [57, 58]. Smaller magnitudes of effect in observational analyses may reflect regression dilution bias from single time-point measurements of BMI and/or reverse causation from cancer-induced weight loss, whereas MR estimates reflect accumulated exposure across the life-course and are unlikely to be influenced by reverse causation [59].

In agreement with previous MR analyses, our results suggest a causal role of fasting insulin, total and bioavailable testosterone, and SHBG in endometrial cancer risk, although these previous reports either employed smaller sample sizes than this analysis (e.g. fasting insulin analyses were performed in 1,287 endometrial cancer cases vs 12,906 cases in our analysis) or used somewhat differing methods to examine instrumental variable assumptions [28-30]. The restriction of an effect of BMI to bioavailable (and not total) testosterone is in agreement with previous observational studies which have suggested that BMI influences testosterone levels through decreased production of SHBG rather than a direct effect on testosterone production [60-64]. Additionally, important mediating roles of fasting insulin, bioavailable testosterone, and SHBG in the relationship between BMI and endometrial cancer are consistent with studies of bariatric surgery which have suggested protective effects of this procedure against endometrial cancer risk, along with reductions in insulin and bioavailable testosterone levels, and increases in SHBG levels [65-73]. Our findings supporting a role of BMI on these traits are also consistent with the important endocrine function of adipose tissue, which is involved in sex steroid metabolism [62, 74-79].

Potential aetiological roles of the molecular mediators identified in this analysis are consistent with the “unopposed oestrogen” hypothesis which postulates that endometrial carcinogenesis is driven by excess endogenous or exogenous oestrogen levels that are unopposed by progesterone [80-82]. We

were unable to incorporate oestrogen into this analysis as we were unable to identify reliable genetic instruments for this trait. All three of the molecular mediators highlighted in this analysis, however, are known to influence oestrogen: bioavailable testosterone is aromatized to oestradiol; SHBG binds with high-affinity to both oestradiol and bioavailable testosterone [82-87]; and insulin increases androgen and decreases SHBG production [88-91]. We found that an inverse effect of SHBG on endometrial cancer risk was largely attenuated upon adjustment for bioavailable testosterone, suggesting a protective effect of SHBG may be driven via binding of biologically active fractions of circulating testosterone. The attenuation of an effect of fasting insulin on endometrial cancer upon adjustment for bioavailable testosterone could reflect mediation of this effect or the presence of conditionally weak instrument bias in this model. In support of the latter explanation, there is biological evidence that hyperinsulinemia and insulin resistance influence endometrial cancer via oestrogen-independent pathways. For example, insulin has been shown to bind directly to endometrial cells and promote proliferation, and can activate two pathways known to have an important role in carcinogenesis – the phosphatidylinositol-3-kinase-protein kinase B/Akt (PI3K-PKB/Akt) and Ras/Raf/mitogen-activated protein kinase (Ras/Raf/MAPK) pathways [91-96].

Some findings from this MR analysis are not in agreement with evidence from previous conventional observational studies. For example, our analyses found little evidence to support causal roles of several metabolic traits (e.g. circulating HDL-cholesterol, triglycerides, adiponectin, leptin) and inflammatory markers (CRP, IL-6) in endometrial cancer risk, despite these traits being linked to endometrial cancer risk in conventional observational analyses [18-22]. Several of these traits (e.g. HDL-cholesterol, LDL-cholesterol, triglycerides) represent highly correlated metabolic perturbations associated with the obese phenotype which may be too clustered to disentangle using conventional multivariable regression methods [97]. Consequently, some of the divergence in findings across previous conventional observational studies and this MR analysis could reflect residual confounding in the

former. Another potential explanation for divergence in findings is the susceptibility of conventional observational studies to reverse causation (i.e., latent, undiagnosed endometrial cancer influencing levels of a presumed exposure). For example, a previously reported association of circulating IL-6 concentrations with endometrial cancer risk could reflect IL-6 secretion by endometrial cancer-associated fibroblasts rather than a role of IL-6 in endometrial cancer development [98, 99].

We were unable to replicate a previously reported MR-based inverse association of LDL cholesterol levels and endometrial cancer risk in the Endometrial Cancer Association Consortium (IVW OR per SD increase in LDL cholesterol: 0.90, 95% CI: 0.85 to 0.95, $P = 8.39 \times 10^{-5}$). In the previous analysis, SNPs were permitted to be in weak linkage disequilibrium (LD) (pairwise correlation $r^2 < 0.05$ vs $r^2 < 0.001$ in our analysis) and a Heterogeneity in Dependent Instruments (HEIDI) test was performed to identify potentially pleiotropic SNPs, resulting in the removal of 6 such SNPs from the 146 SNPs initially used as an instrument. We attempted to replicate these previously reported findings using a more stringent r^2 threshold (i.e. $r^2 < 0.001$) followed by use of the HEIDI test (resulting in the removal of 2 potentially pleiotropic SNPs) which resulted in a causal estimate that was closer in magnitude to that previously reported (IVW OR 0.93, 95% CI: 0.86 to 1.00, $P = 4.10 \times 10^{-2}$) (**S45 Table**). However, there was greater imprecision in our estimate compared to this previous analysis which could reflect the more liberal LD threshold employed in this earlier analysis.

Our MR analysis provides key insights into potential molecular pathways linking excess adiposity to endometrial cancer risk. This analysis has several strengths including the use of a systematic approach to collate previously reported molecular risk factors for endometrial cancer; the appraisal of their causal relevance in overall and endometrioid endometrial cancer aetiology using an MR framework which should be less prone to conventional issues of confounding and cannot be influenced by reverse causation; the employment of several complementary sensitivity analyses to rigorously assess for

violations of MR assumptions; and the use of a summary data-based MR approach which permitted us to leverage large-scale GWAS data from several studies, enhancing statistical power and precision of causal estimates.

There are several limitations to our analysis. First, we were unable to evaluate the role of six previously reported molecular risk factors for endometrial cancer due to the absence of reliable genetic instruments for these traits. These six risk factors included oestradiol which is believed to be an important molecular mediator of the effect of BMI on endometrial cancer risk[9]. Second, some of the effect estimates for SNPs included in genetic instruments were obtained from discovery GWAS and have not been replicated in an independent sample which can result in “Winner’s curse” bias. There was sample overlap in this analysis across certain traits; however, the use of conventionally strong ($P < 5.0 \times 10^{-8}$) instruments for these traits and the consistency of most findings in sensitivity analyses examining their robustness to potential Winner’s curse bias suggests that this phenomenon was unlikely to markedly influence the results presented in this analysis. Third, although sex-specific sensitivity analyses were performed where data were available, some prior GWAS used in this analysis did not examine for heterogeneity of SNP effects by sex which prevented evaluation of the effect of certain traits on endometrial cancer risk using sex-specific instruments. Fourth, univariable and multivariable MR analyses presented here assume that relationships between exposures and outcomes are linear, although it has been previously suggested that the relationship between BMI and endometrial cancer may best be explained by a non-linear model [12, 100]. Fifth, our analysis was almost exclusively restricted to individuals of European ancestry to minimise bias from population stratification, which may limit the generalisability of our findings to non-European populations. Finally, while various sensitivity analyses were performed to examine violations of exchangeability and exclusion restriction criteria, these assumptions are unverifiable.

With the global incidence of overweight and obesity projected to increase and challenges in implementing successful weight loss strategies, pharmacological approaches targeting molecular mediators of the effect of obesity on endometrial cancer development may offer a viable approach for cancer prevention in high-risk groups [101-105]. Metformin, a safe and inexpensive first-line treatment for type 2 diabetes, could serve as a promising chemoprevention agent for endometrial cancer as it increases insulin sensitivity, thus reversing insulin resistance and lowering fasting insulin levels, alongside inhibiting endometrial proliferation [9, 106]. In addition, unlike some other oral hypoglycaemic medications, metformin users show a tendency toward sustained weight loss [107]. Bioavailable testosterone and SHBG also present potential pharmacological targets, though the multifaceted function of these hormones means that targeting these traits may result in adverse effects [108-113]. Phase II clinical trials examining the efficacy of a combination of contraceptive intrauterine devices, metformin, and weight loss interventions as a non-invasive treatment option for individuals with obesity with early-stage endometrial cancer have had encouraging results, and weight loss has been shown to improve oncological outcomes in women with endometrial cancer undergoing progestin treatment [114, 115].

Our systematic evaluation of 14 previously reported candidate mediators of the effect of BMI on endometrial cancer risk identifies fasting insulin, bioavailable testosterone, and SHBG as plausible mediators of this relationship. While we were unable to entirely disentangle the independent effects of these three traits, identification of a potential mediating role of these traits (and, in particular, fasting insulin) in endometrial carcinogenesis is nonetheless informative for the development of pharmacological interventions targeting these traits for cancer prevention. In this respect, future assessment of the effect of drugs which target molecular mediators identified in this analysis using a “drug-target Mendelian randomization” approach could inform on the potential efficacy of the repurposing of medications for endometrial cancer prevention.

Conclusion

Our comprehensive Mendelian randomization analysis provides insight into potential causal mechanisms linking excess adiposity to endometrial cancer risk. We show that lifelong cumulative elevated BMI causes a larger increased risk than that reported in previous conventional observational studies. We found strong evidence for a mediating role of fasting insulin, bioavailable testosterone and SHBG in the effect of BMI on endometrial cancer risk. These results suggest pharmacological targeting of insulin-related and hormonal traits as a potential strategy for the prevention of endometrial cancer.

Funding

EH is supported by a Cancer Research UK Population Research Committee Studentship (C18281/A30905). EH, VT, GDS, RMM, KR and JY are supported by Cancer Research UK (C18281/A29019) programme grant (the Integrative Cancer Epidemiology Programme). EH, ES, VT, GDS, RMM, and JY are part of the Medical Research Council Integrative Epidemiology Unit at the University of Bristol which is supported by the Medical Research Council (MC_UU_00011/1, MC_UU_00011/3, MC_UU_00011/6, and MC_UU_00011/4) and the University of Bristol. JY is supported by a Cancer Research UK Population Research Postdoctoral Fellowship (C68933/A28534). RMM is also supported by the NIHR Bristol Biomedical Research Centre which is funded by the NIHR and is a partnership between University Hospitals Bristol and Weston NHS Foundation Trust and the University of Bristol. TMF is funded by the MRC (MR/T002239/1). TAO'M is funded by a National Health and Medical Research Council Investigator Fellowship (APP1173170).

Disclaimers: The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health and Social Care; where authors are identified as personnel of the International Agency for Research on Cancer/World Health Organization, the authors alone are responsible for the views expressed in this article and they do not necessarily represent the decisions, policy, or views of the International Agency for Research on Cancer/World Health Organization.

Acknowledgements

The endometrial cancer genome-wide association analyses were supported by the National Health and Medical Research Council of Australia (APP552402, APP1031333, APP1109286, APP1111246 and APP1061779), the U.S. National Institutes of Health (R01-CA134958), European Research Council (EU FP7 Grant), Wellcome Trust Centre for Human Genetics (090532/Z/09Z) and Cancer Research UK. OncoArray genotyping of ECAC cases was performed with the generous assistance of the Ovarian Cancer

Association Consortium (OCAC), which was funded through grants from the U.S. National Institutes of Health (CA1X01HG007491-01 (C.I. Amos), U19-CA148112 (T.A. Sellers), R01-CA149429 (C.M. Phelan) and R01-CA058598 (M.T. Goodman); Canadian Institutes of Health Research (MOP-86727 (L.E. Kelemen)) and the Ovarian Cancer Research Fund (A. Berchuck). We particularly thank the efforts of Cathy Phelan. OncoArray genotyping of the BCAC controls was funded by Genome Canada Grant GPH-129344, NIH Grant U19 CA148065, and Cancer UK Grant C1287/A16563. All studies and funders are listed in O'Mara et al (2018). The authors would like to thank the participants of the 1958 British birth cohort (BS58), Academic Medical Centre Amsterdam Premature Atherosclerosis Study (AMC-PAS), Age, Gene/Environment Susceptibility (AGES) study, Anglo-Scandinavian Cardiac Outcome Trial (ASCOT), Aro-Express Biobank Study (AE), Atherosclerosis Risk in Communities (ARIC) study, Atherosclerotic Disease, Vascular Function, and Genetic Epidemiology (ADVANCE) study, Asian Indian Diabetic Heart Study (AIDHS) /Sikh Diabetes Study (SDS), Avon Longitudinal Study of Parents and Children (ALSPAC), Baltimore Longitudinal Study of Aging (BLSA), British Genetics of Hypertension (BRIGHT) Study, Busselton Health Study (BSN), C Reactive Protein Coronary Heart Disease Genetics Collaboration (CCGC), Cancer Prostate in Sweden 1 (CAPS1) study, Cancer Prostate in Sweden 2 (CAPS2) study, CardioGenics study, Cardiovascular Health Study (CHS), Cardiovascular Health Study (CHS), Cardiovascular Risk in Young Finns Study (YFS), Carotid Intima Media Thickness (IMT) and IMT-Progression as Predictors of Vascular Events in a High Risk European Population (IMPROVE) study, Cebu Longitudinal Health and Nutrition Survey (CLHNS), Charles Bronfman Institute for Personalized Medicine (IPM) BioMe Biobank Program, Cohorte Lausannoise (CoLaus) study, Cohorts for Heart and Aging Research in Genomic Epidemiology (CHARGE) Inflammation Working Group (CIWG), CROATIA-Vis study, Croatian Study (CROAS), Danish National Birth Cohort (DNBC), Data from an Epidemiological Study on Insulin Resistance syndrome (DESIR), Diabetes Epidemiology: Collaborative analysis of Diagnostic criteria in Europe (DECODE) study, Diabetes Genetics Initiative (DGI), DiaGen consortium, Dose-Responses to Exercise Training (DR's EXTRA) study, Dundee study, Dutch and Belgian

542 Lung Cancer Screening Trial (NELSON), Echinococcus Multilocularis and Internal Diseases in Leutkirch
 543 (EMIL) study, Edinburgh Artery Study (EAS), Efficiency and safety of varying frequency of whole blood
 544 donation (INTERVAL) study, Ely Study, Endometrial Cancer Association Consortium (ECAC), Epidemiology
 545 of Endometrial Cancer Consortium (E2C2), EpiHealth cohort, Erasmus Rucphen Family (ERF) study,
 546 Estonian Biobank, Estonian Genome Center of University of Tartu (EGCUT) study, European Prospective
 547 Investigation into Cancer and Nutrition (EPIC) study, Family Blood Pressure Program (FBPP), Family
 548 Blood Pressure Program (FBPP), Family Heart Study (FamHS), Fast Revascularisation during Instability in
 549 Coronary artery disease (FRISCII) study, Fendland study, Finland-United States Investigation of NIDDM
 550 Genetics (FUSION) study, FinMetSeq, Finnish Cardiovascular Study (FINCAVAS), Finnish Diabetes
 551 Prevention Study (DPS), Finnish Genetic Study of Arrhythmic Events (FinGesture), Finnish National
 552 Diabetes Prevention Program (FIN-D2D), Finnish Twin Cohort (FTC), Finrisk study, Framingham Heart
 553 Study (FRAM), Genetic Investigation of ANthropometric Traits (GIANT), Gene-Lifestyle interactions And
 554 Complex traits Involved in Elevated disease Risk Study (GLACIER), Genetic determinants of Obesity and
 555 Metabolic syndrome (DILGOM) study, Genetic Epidemiology Network of Arteriopathy (GENOA) study,
 556 Genetic Predisposition of Coronary Heart Disease in Patients Verified with Coronary Angiogram
 557 (COROGENE), Genetics of Diabetes Audit and Research Study (GoDartS), GenomEUtwin, German
 558 Myocard Infarct Family Study I (GerMiFSI), German Myocard Infarct Family Study II (GerMiFSII), Global
 559 Lipids Consortium (GLGC), Gonburg Osteoporosis and Obesity Determinants Study (GOOD), Health 2000
 560 study, Health and Retirement Study (HRS), Health, Aging, and Body Composition Study (Health ABC),
 561 Health, Risk Factors, Training and Genetics (HERITAGE) Family Study, Heinz Nixdorf Recall (HNR),
 562 Hellenic study of Interactions between Single nucleotide polymorphisms and Eating in Atherosclerosis
 563 Susceptibility (THISEAS), Helsinki Birth Cohort Study (HBCS), Heredity and Phenotype Intervention (HAPI)
 564 Heart Study, Invecchiare in Chianti (InCHIANTI) study, Kingston Gene-by-environment; subset of
 565 International Collaborative Study of Hypertension in Blacks (GXE), Kooperative Gesundheitsforschung in

566 der Region Augsburg (KORA) study, Korcula study, Leiden Longevity Study (LLS), Leipzig adults study,
 567 Leipzig kids study, Lifelines cohort, London Life Sciences Prospective Population Study (LOLIPOP),
 568 LUdwigshafen Risk and Cardiovascular Health (LURIC) study, MAGIC (the Meta-Analyses of Glucose and
 569 Insulin-related traits Consortium), Malmo Preventive Project-Re-examination (MPP-RES) study, Malmo
 570 Diet and Cancer Study (MDC), Medical research Council (MRC) National Survey of Health & Development
 571 (MRC NSHD), Medical Research Council (MRC) National Survey of Health and Development (NSHD),
 572 Medical Research Council/Uganda Virus Research Institute General Population Cohort (MRC/UVRIGPC),
 573 Metabolic Syndrome in Men (METSIM) study, Microisolates in South Tyrol Study (MICROS), Molecular
 574 Genetics of Schizophrenia (MGS), MONica Risk, Genetics, Archiving and Monograph (MORGAM),
 575 Multiethnic Cohort Study (MEC), Myocardial Infarction Genetics Consortium (Migen), Nerlands Study of
 576 Depression and Anxiety (NESDA), Nerlands Twin Register (NTR), Nijmegen Bladder Cancer Study and
 577 Nijmegen Biomedical Study (RUNMC), Nord- Trøndelag Health Study (HUNT), Northern Finland Birth
 578 Cohort 1966 (NFBC66), Northern Finland Birth Cohort 1986 (NFBC86), Northern Sweden Population
 579 Health Study (NSPHS), Nurse's Health Study (NHS), Orkney Complex Disease Study (ORCADES), Pakistan
 580 Risk of Myocardial Infarction Study (PROMIS), Precocious Coronary Artery Disease (PROCARDIS) study,
 581 Prevention of RENal and Vascular ENd-stage Disease (PREVEND) study, Prospective Investigation of
 582 Vasculature in Uppsala Seniors (PIVUS) study, PROspective study of Pravastatin in Elderly at Risk for
 583 vascular disease (PROSPER/PHASE), Prostate, Lung, Colorectal and Ovarian Cancer Screening Trial
 584 (PLCO), Quebec Family Study (QFS), Relationship between Insulin Sensitivity and Cardiovascular disease
 585 Study (RISC), Rotterdam study, SardiNIA Project, Seychelles Family Study (TANDEM), Sorbs study,
 586 Spanish Town study (SPT), STabilization of Arosclerotic plaque By Initiation of darapLadlb rapY
 587 (STABILITY), Stanley cohort, Stockholm Coronary Arosclerosis Risk Factor (SCARF) study, Stockholm Heart
 588 Epidemiology Program (SHEEP), Studies of Epidemiology and Risk factors in Cancer Heredity / UK
 589 Ovarian Cancer Population Study (SEARCH/UKOPS), Study of Health in Pomerania (SHIP), Suivi

590 Temporaire Annuel Non Invasif de la Santé des Lorrains Assurés Sociaux (STANISLAS) cohort, Swedish
591 And Singapore Breast Association Consortium (SASBAC), Swedish Twin Registry (STR), Taichi Consortium,
592 Tracking Adolescents' Individual Lives Survey (TRAILS), Tromsø Study, Twin study at Queensland Institute
593 of Medical Research (QIMR), Twins UK, UK Biobank, Uppsala Longitudinal Study of Adult Men (ULSAM),
594 Vitamin D and Type 2 Diabetes (D2d) study, Wellcome Trust Case Control Consortium (WTCCC),
595 Whitehall study, Women's Genome Health Study (WGHS), and Women's Health Initiative (WHI) for their
596 participation in these studies along with the principal investigators for generating the data utilised for
597 this analysis and for making these data available in the public domain. Data on glycaemic traits have
598 been contributed by MAGIC investigators and have been downloaded from
599 www.magicinvestigators.org, and data on oestradiol and blood glucose have been contributed by the
600 Neale lab and have been downloaded from www.nealelab.is.

References

1. Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA: A Cancer Journal for Clinicians*. 2018;68(6):394-424. doi: <https://doi.org/10.3322/caac.21492>.
2. Howlader N NA, Krapcho M, Miller D, Bishop K, Kosary CL, Yu M, Ruhl J, Tatalovich Z, Mariotto A, Lewis DR, Chen HS, Feuer EJ, Cronin KA (eds). *SEER Cancer Statistics Review, 1975-2014*, National Cancer Institute. Bethesda, MD, https://seercancer.gov/csr/1975_2014/. Based on November 2016 SEER data submission, posted to the SEER web site, April 2017.
3. Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A, et al. Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries. *CA Cancer J Clin*. 2021;71(3):209-49. Epub 2021/02/05. doi: 10.3322/caac.21660. PubMed PMID: 33538338.
4. Sheikh MA, Althouse AD, Freese KE, Soisson S, Edwards RP, Welburn S, et al. USA Endometrial Cancer Projections to 2030: should we be concerned? *Future Oncology*. 2014;10(16):2561-8. doi: 10.2217/fon.14.192. PubMed PMID: 25531045.
5. Lortet-Tieulent J, Ferlay J, Bray F, Jemal A. International Patterns and Trends in Endometrial Cancer Incidence, 1978–2013. *JNCI: Journal of the National Cancer Institute*. 2017;110(4):354-61. doi: 10.1093/jnci/djx214.
6. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2016. *CA: A Cancer Journal for Clinicians*. 2016;66(1):7-30. doi: <https://doi.org/10.3322/caac.21332>.

- 622 7. Zhang S, Gong T-T, Liu F-H, Jiang Y-T, Sun H, Ma X-X, et al. Global, Regional, and National
623 Burden of Endometrial Cancer, 1990-2017: Results From the Global Burden of Disease Study,
624 2017. *Front Oncol.* 2019;9:1440-. doi: 10.3389/fonc.2019.01440. PubMed PMID: 31921687.
- 625 8. Mullins MA, Cote ML. Beyond Obesity: The Rising Incidence and Mortality Rates of
626 Uterine Corpus Cancer. *J Clin Oncol.* 2019;37(22):1851-3. Epub 2019/06/20. doi:
627 10.1200/JCO.19.01240. PubMed PMID: 31232669.
- 628 9. Onstad MA, Schmandt RE, Lu KH. Addressing the Role of Obesity in Endometrial Cancer
629 Risk, Prevention, and Treatment. *J Clin Oncol.* 2016;34(35):4225-30. Epub 2016/11/07. doi:
630 10.1200/JCO.2016.69.4638. PubMed PMID: 27903150.
- 631 10. Lega IC, Lipscombe LL. Review: Diabetes, Obesity, and Cancer—Pathophysiology and
632 Clinical Implications. *Endocrine Reviews.* 2019;41(1):33-52. doi: 10.1210/endrev/bnz014.
- 633 11. Reeves GK, Pirie K, Beral V, Green J, Spencer E, Bull D, et al. Cancer incidence and
634 mortality in relation to body mass index in the Million Women Study: cohort study. *BMJ.*
635 2007;335(7630):1134-. Epub 2007/11/06. doi: 10.1136/bmj.39367.495995.AE. PubMed PMID:
636 17986716.
- 637 12. Aune D, Navarro Rosenblatt DA, Chan DS, Vingeliene S, Abar L, Vieira AR, et al.
638 Anthropometric factors and endometrial cancer risk: a systematic review and dose-response
639 meta-analysis of prospective studies. *Ann Oncol.* 2015;26(8):1635-48. Epub 2015/03/21. doi:
640 10.1093/annonc/mdv142. PubMed PMID: 25791635.
- 641 13. World Cancer Research Fund/American Institute for Cancer Research. Continuous
642 Update Project Expert Report 2018. Diet, nutrition, physical activity and ovarian cancer.
643 Available at dietandcancerreport.org.

14. World Cancer Research Fund/American Institute for Cancer Research. Continuous Update Project Expert Report 2018. Diet, nutrition, physical activity and oesophageal cancer. Available at dietandcancerreport.org.
15. Nevadunsky NS, Van Arsdale A, Strickler HD, Moadel A, Kaur G, Levitt J, et al. Obesity and age at diagnosis of endometrial cancer. *Obstet Gynecol*. 2014;124(2 Pt 1):300-6. Epub 2014/07/09. doi: 10.1097/aog.0000000000000381. PubMed PMID: 25004350.
16. McCullough ML, Patel AV, Patel R, Rodriguez C, Feigelson HS, Bandera EV, et al. Body mass and endometrial cancer risk by hormone replacement therapy and cancer subtype. *Cancer Epidemiol Biomarkers Prev*. 2008;17(1):73-9. Epub 2008/01/12. doi: 10.1158/1055-9965.Epi-07-2567. PubMed PMID: 18187388.
17. Painter JN, O'Mara TA, Marquart L, Webb PM, Attia J, Medland SE, et al. Genetic Risk Score Mendelian Randomization Shows that Obesity Measured as Body Mass Index, but not Waist:Hip Ratio, Is Causal for Endometrial Cancer. *Cancer epidemiology, biomarkers & prevention : a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology*. 2016;25(11):1503-10. Epub 2016/08/22. doi: 10.1158/1055-9965.EPI-16-0147. PubMed PMID: 27550749.
18. Dossus L, Rinaldi S, Becker S, Lukanova A, Tjonneland A, Olsen A, et al. Obesity, inflammatory markers, and endometrial cancer risk: a prospective case-control study. *Endocrine-Related Cancer*. 2010;17(4):1007. doi: 10.1677/erc-10-0053.
19. Friedenreich CM, Langley AR, Speidel TP, Lau DCW, Courneya KS, Csizmadia I, et al. Case-control study of inflammatory markers and the risk of endometrial cancer. *European Journal of*

665 Cancer Prevention. 2013;22(4):374-9. doi: 10.1097/CEJ.0b013e32835b3813. PubMed PMID:
 666 00008469-201307000-00013.

667 20. Wang T, Rohan TE, Gunter MJ, Xue X, Wactawski-Wende J, Rajpathak SN, et al. A
 668 Prospective Study of Inflammation Markers and Endometrial Cancer Risk in Postmenopausal
 669 Hormone Nonusers. Cancer Epidemiology Biomarkers & Prevention. 2011;20(5):971-7. doi:
 670 10.1158/1055-9965.Epi-10-1222.

671 21. Lindemann K, Vatten LJ, Ellstrøm-Eng M, Eskild A. Serum lipids and endometrial cancer
 672 risk: Results from the HUNT-II study. International Journal of Cancer. 2009;124(12):2938-41.
 673 doi: <https://doi.org/10.1002/ijc.24285>.

674 22. Swanson CA, Potischman N, Barrett RJ, Berman ML, Mortel R, Twiggs LB, et al.
 675 Endometrial cancer risk in relation to serum lipids and lipoprotein levels. Cancer Epidemiol
 676 Biomarkers Prev. 1994;3(7):575-81. Epub 1994/10/01. PubMed PMID: 7827588.

677 23. Yarmolinsky J, Wade KH, Richmond RC, Langdon RJ, Bull CJ, Tilling KM, et al. Causal
 678 Inference in Cancer Epidemiology: What Is the Role of Mendelian Randomization? Cancer
 679 Epidemiology Biomarkers & Prevention. 2018;27(9):995-1010. doi: 10.1158/1055-
 680 9965.Epi-17-1177.

681 24. Davey Smith G, Ebrahim S. 'Mendelian randomization': can genetic epidemiology
 682 contribute to understanding environmental determinants of disease?*. International Journal of
 683 Epidemiology. 2003;32(1):1-22. doi: 10.1093/ije/dyg070.

684 25. Pierce BL, Burgess S. Efficient design for Mendelian randomization studies: subsample
 685 and 2-sample instrumental variable estimators. Am J Epidemiol. 2013;178(7):1177-84. Epub

2013/07/19. doi: 10.1093/aje/kwt084. PubMed PMID: 23863760; PubMed Central PMCID: PMCPMC3783091.

26. Thompson DJ, O'Mara TA, Glubb DM, Painter JN, Cheng T, Folkard E, et al. CYP19A1 fine-mapping and Mendelian randomization: estradiol is causal for endometrial cancer. *Endocr Relat Cancer*. 2016;23(2):77-91. Epub 2015/11/18. doi: 10.1530/erc-15-0386. PubMed PMID: 26574572; PubMed Central PMCID: PMCPMC4697192.

27. Kho P-F, Amant F, Annibali D, Ashton K, Attia J, Auer PL, et al. Mendelian randomization analyses suggest a role for cholesterol in the development of endometrial cancer. *International Journal of Cancer*. 2021;148(2):307-19. doi: <https://doi.org/10.1002/ijc.33206>.

28. Nead KT, Sharp SJ, Thompson DJ, Painter JN, Savage DB, Semple RK, et al. Evidence of a Causal Association Between Insulinemia and Endometrial Cancer: A Mendelian Randomization Analysis. *J Natl Cancer Inst*. 2015;107(9). Epub 2015/07/03. doi: 10.1093/jnci/djv178. PubMed PMID: 26134033; PubMed Central PMCID: PMCPMC4572886.

29. Ruth KS, Day FR, Tyrrell J, Thompson DJ, Wood AR, Mahajan A, et al. Using human genetics to understand the disease impacts of testosterone in men and women. *Nature Medicine*. 2020;26(2):252-+. doi: 10.1038/s41591-020-0751-5. PubMed PMID: WOS:000512529400005.

30. Mullee A, Dimou N, Allen N, O'Mara T, Gunter MJ, Murphy N. Testosterone, sex hormone-binding globulin, insulin-like growth factor-1 and endometrial cancer risk: observational and Mendelian randomization analyses. *Br J Cancer*. 2021;125(9):1308-17. Epub 2021/08/08. doi: 10.1038/s41416-021-01518-3. PubMed PMID: 34363033; PubMed Central PMCID: PMCPMC8548546.

- 708 31. O'Mara TA, Glubb DM, Amant F, Annibaldi D, Ashton K, Attia J, et al. Identification of nine
709 new susceptibility loci for endometrial cancer. *Nat Commun.* 2018;9(1):3166. Epub 2018/08/11.
710 doi: 10.1038/s41467-018-05427-7. PubMed PMID: 30093612; PubMed Central PMCID:
711 PMCPMC6085317.
- 712 32. Dimou NL, Papadimitriou N, Mariosa D, Johansson M, Brennan P, Peters U, et al.
713 Circulating adipokine concentrations and risk of five obesity-related cancers: A Mendelian
714 randomization study. *Int J Cancer.* 2021;148(7):1625-36. Epub 2020/10/11. doi:
715 10.1002/ijc.33338. PubMed PMID: 33038280; PubMed Central PMCID: PMCPMC7894468.
- 716 33. Auton A, Abecasis GR, Altshuler DM, Durbin RM, Abecasis GR, Bentley DR, et al. A global
717 reference for human genetic variation. *Nature.* 2015;526(7571):68-74. doi:
718 10.1038/nature15393.
- 719 34. Willer CJ, Schmidt EM, Sengupta S, Peloso GM, Gustafsson S, Kanoni S, et al. Discovery
720 and refinement of loci associated with lipid levels. *Nature Genetics.* 2013;45(11):1274-+. doi:
721 10.1038/ng.2797. PubMed PMID: WOS:000326384100007.
- 722 35. Scott RA, Lagou V, Welch RP, Wheeler E, Montasser ME, Luan JA, et al. Large-scale
723 association analyses identify new loci influencing glycemic traits and provide insight into the
724 underlying biological pathways. *Nature Genetics.* 2012;44(9):991-+. doi: 10.1038/ng.2385.
725 PubMed PMID: WOS:000308491200010.
- 726 36. Neale B. GWAS results. UK Biobank website. [updated 1st August 2018; cited 2020
727 9/12/20].

- 728 37. Sinnott-Armstrong N, Tanigawa Y, Amar D, Mars NJ, Aguirre M, Venkataraman GR, et al.
729 Genetics of 38 blood and urine biomarkers in the UK Biobank. *bioRxiv*. 2019:660506. doi:
730 10.1101/660506.
- 731 38. Georgakis MK, Malik R, Gill D, Franceschini N, Sudlow CLM, Dichgans M. Interleukin-6
732 signaling effects on ischemic stroke and other cardiovascular outcomes: a Mendelian
733 Randomization study. *medRxiv*. 2019:19007682. doi: 10.1101/19007682.
- 734 39. Dastani Z, Hivert M-F, Timpson N, Perry JRB, Yuan X, Scott RA, et al. Novel Loci for
735 Adiponectin Levels and Their Influence on Type 2 Diabetes and Metabolic Traits: A Multi-Ethnic
736 Meta-Analysis of 45,891 Individuals. *PLOS Genetics*. 2012;8(3):e1002607. doi:
737 10.1371/journal.pgen.1002607.
- 738 40. Sun BB, Maranville JC, Peters JE, Stacey D, Staley JR, Blackshaw J, et al. Genomic atlas of
739 the human plasma proteome. *Nature*. 2018;558(7708):73-9. doi: 10.1038/s41586-018-0175-2.
- 740 41. Ligthart S, Vaez A, Vösa U, Stathopoulou MG, de Vries PS, Prins BP, et al. Genome
741 Analyses of >200,000 Individuals Identify 58 Loci for Chronic Inflammation and Highlight
742 Pathways that Link Inflammation and Complex Disorders. *Am J Hum Genet*. 2018;103(5):691-
743 706. Epub 2018/11/06. doi: 10.1016/j.ajhg.2018.09.009. PubMed PMID: 30388399; PubMed
744 Central PMCID: PMC6218410.
- 745 42. Yengo L, Sidorenko J, Kemper KE, Zheng Z, Wood AR, Weedon MN, et al. Meta-analysis
746 of genome-wide association studies for height and body mass index in ~700000 individuals of
747 European ancestry. *Hum Mol Genet*. 2018;27(20):3641-9. Epub 2018/08/21. doi:
748 10.1093/hmg/ddy271. PubMed PMID: 30124842; PubMed Central PMCID: PMCPMC6488973.

43. Davies NM, Holmes MV, Davey Smith G. Reading Mendelian randomisation studies: a guide, glossary, and checklist for clinicians. *BMJ*. 2018;362:k601. Epub 2018/07/14. doi: 10.1136/bmj.k601. PubMed PMID: 30002074; PubMed Central PMCID: PMCPMC6041728
- interests and declare that we have no competing interests.
44. Burgess S, Zuber V, Valdes-Marquez E, Sun BB, Hopewell JC. Mendelian randomization with fine-mapped genetic data: Choosing from large numbers of correlated instrumental variables. *Genetic Epidemiology*. 2017;41(8):714-25. doi: <https://doi.org/10.1002/gepi.22077>.
45. Hemani G, Bowden J, Davey Smith G. Evaluating the potential role of pleiotropy in Mendelian randomization studies. *Hum Mol Genet*. 2018;27(R2):R195-r208. Epub 2018/05/18. doi: 10.1093/hmg/ddy163. PubMed PMID: 29771313; PubMed Central PMCID: PMCPMC6061876.
46. Bowden J, Davey Smith G, Haycock PC, Burgess S. Consistent Estimation in Mendelian Randomization with Some Invalid Instruments Using a Weighted Median Estimator. *Genet Epidemiol*. 2016;40(4):304-14. Epub 2016/04/12. doi: 10.1002/gepi.21965. PubMed PMID: 27061298; PubMed Central PMCID: PMCPMC4849733.
47. Hartwig FP, Davey Smith G, Bowden J. Robust inference in summary data Mendelian randomization via the zero modal pleiotropy assumption. *International journal of epidemiology*. 2017;46(6):1985-98. doi: 10.1093/ije/dyx102. PubMed PMID: 29040600.
48. Bowden J, Davey Smith G, Burgess S. Mendelian randomization with invalid instruments: effect estimation and bias detection through Egger regression. *Int J Epidemiol*. 2015;44(2):512-25. Epub 2015/06/08. doi: 10.1093/ije/dyv080. PubMed PMID: 26050253; PubMed Central PMCID: PMCPMC4469799.

- 771 49. Hemani G, Tilling K, Davey Smith G. Orienting the causal relationship between
772 imprecisely measured traits using GWAS summary data. *PLoS Genet.* 2017;13(11):e1007081.
773 Epub 2017/11/18. doi: 10.1371/journal.pgen.1007081. PubMed PMID: 29149188; PubMed
774 Central PMCID: PMCPMC5711033.
- 775 50. Burgess S, Thompson SG. Multivariable Mendelian Randomization: The Use of
776 Pleiotropic Genetic Variants to Estimate Causal Effects. *American Journal of Epidemiology.*
777 2015;181(4):251-60. doi: 10.1093/aje/kwu283.
- 778 51. Cheung MWL. Comparison of methods for constructing confidence intervals of
779 standardized indirect effects. *Behavior Research Methods.* 2009;41(2):425-38. doi:
780 10.3758/BRM.41.2.425.
- 781 52. Haycock PC, Burgess S, Wade KH, Bowden J, Relton C, Davey Smith G. Best (but oft-
782 forgotten) practices: the design, analysis, and interpretation of Mendelian randomization
783 studies. *Am J Clin Nutr.* 2016;103(4):965-78. doi: 10.3945/ajcn.115.118216. PubMed PMID:
784 26961927.
- 785 53. Kettunen J, Demirkan A, Würtz P, Draisma HH, Haller T, Rawal R, et al. Genome-wide
786 study for circulating metabolites identifies 62 loci and reveals novel systemic effects of LPA. *Nat*
787 *Commun.* 2016;7:11122. Epub 2016/03/24. doi: 10.1038/ncomms11122. PubMed PMID:
788 27005778.
- 789 54. Prins BP, Kuchenbaecker KB, Bao Y, Smart M, Zabaneh D, Fatemifar G, et al. Genome-
790 wide analysis of health-related biomarkers in the UK Household Longitudinal Study reveals
791 novel associations. *Sci Rep.* 2017;7(1):11008. Epub 2017/09/10. doi: 10.1038/s41598-017-
792 10812-1. PubMed PMID: 28887542; PubMed Central PMCID: PMCPMC5591265.

- 793 55. Locke AE, Kahali B, Berndt SI, Justice AE, Pers TH, Felix R, et al. Genetic studies of body
794 mass index yield new insights for obesity biology. *Nature*. 2015;518(7538):197-U401. doi:
795 10.1038/nature14177. PubMed PMID: WOS:000349190300031.
- 796 56. Sanderson E, Spiller W, Bowden J. Testing and Correcting for Weak and Pleiotropic
797 Instruments in Two-Sample Multivariable Mendelian Randomisation. *bioRxiv*.
798 2020:2020.04.02.021980. doi: 10.1101/2020.04.02.021980.
- 799 57. World Cancer Research Fund/American Institute for Cancer Research. Continuous
800 Update Project Expert Report 2018. Diet, nutrition, physical activity and endometrial cancer.
801 Available at dietandcancerreport.org.
- 802 58. Mariosa D, Carreras-Torres R, Martin RM, Johansson M, Brennan P. Commentary: What
803 can Mendelian randomization tell us about causes of cancer? *International Journal of*
804 *Epidemiology*. 2019;48(3):816-21. doi: 10.1093/ije/dyz151.
- 805 59. Morris TT, Heron J, Sanderson E, Smith GD, Tilling K. Interpretation of mendelian
806 randomization using one measure of an exposure that varies over time. *medRxiv*.
807 2021:2021.11.18.21266515. doi: 10.1101/2021.11.18.21266515.
- 808 60. Poddar M, Chetty Y, Chetty VT. How does obesity affect the endocrine system? A
809 narrative review. *Clinical Obesity*. 2017;7(3):136-44. doi: <https://doi.org/10.1111/cob.12184>.
- 810 61. Danforth KN, Eliassen AH, Tworoger SS, Missmer SA, Barbieri RL, Rosner BA, et al. The
811 association of plasma androgen levels with breast, ovarian and endometrial cancer risk factors
812 among postmenopausal women. *International journal of cancer*. 2010;126(1):199-207. doi:
813 10.1002/ijc.24709. PubMed PMID: 19569181.

- 814 62. Penttilä T-L, Koskinen P, Penttilä T-A, Anttila L, Irjala K. Obesity regulates bioavailable
815 testosterone levels in women with or without polycystic ovary syndrome. *Fertility and Sterility*.
816 1999;71(3):457-61. doi: [https://doi.org/10.1016/S0015-0282\(98\)00473-7](https://doi.org/10.1016/S0015-0282(98)00473-7).
- 817 63. Lim SS, Norman RJ, Davies MJ, Moran LJ. The effect of obesity on polycystic ovary
818 syndrome: a systematic review and meta-analysis. *Obesity Reviews*. 2013;14(2):95-109. doi:
819 <https://doi.org/10.1111/j.1467-789X.2012.01053.x>.
- 820 64. Burger HG, Dudley EC, Cui J, Dennerstein L, Hopper JL. A prospective longitudinal study
821 of serum testosterone, dehydroepiandrosterone sulfate, and sex hormone-binding globulin
822 levels through the menopause transition. *J Clin Endocrinol Metab*. 2000;85(8):2832-8. Epub
823 2000/08/18. doi: 10.1210/jcem.85.8.6740. PubMed PMID: 10946891.
- 824 65. Upala S, Anawin S. Bariatric surgery and risk of postoperative endometrial cancer: a
825 systematic review and meta-analysis. *Surgery for Obesity and Related Diseases*. 2015;11(4):949-
826 55. doi: <https://doi.org/10.1016/j.soard.2014.09.024>.
- 827 66. Sjöström L, Gummesson A, Sjöström CD, Narbro K, Peltonen M, Wedel H, et al. Effects of
828 bariatric surgery on cancer incidence in obese patients in Sweden (Swedish Obese Subjects
829 Study): a prospective, controlled intervention trial. *Lancet Oncol*. 2009;10(7):653-62. Epub
830 2009/06/27. doi: 10.1016/s1470-2045(09)70159-7. PubMed PMID: 19556163.
- 831 67. Anderwald CH, Tura A, Promintzer-Schifferl M, Prager G, Stadler M, Ludvik B, et al.
832 Alterations in gastrointestinal, endocrine, and metabolic processes after bariatric Roux-en-Y
833 gastric bypass surgery. *Diabetes Care*. 2012;35(12):2580-7. Epub 2012/08/28. doi:
834 10.2337/dc12-0197. PubMed PMID: 22923664; PubMed Central PMCID: PMC3507557.

- 835 68. Sdralis E, Argentou M, Mead N, Kehagias I, Alexandridis T, Kalfarentzos F. A Prospective
836 Randomized Study Comparing Patients with Morbid Obesity Submitted to Sleeve Gastrectomy
837 With or Without Omentectomy. *Obesity Surgery*. 2013;23(7):965-71. doi: 10.1007/s11695-013-
838 0925-z.
- 839 69. Beiglböck H, Fellingner P, Ranzenberger-Haider T, Itariu B, Prager G, Kautzky-Willer A, et
840 al. Pre-operative Obesity-Associated Hyperandrogenemia in Women and Hypogonadism in Men
841 Have No Impact on Weight Loss Following Bariatric Surgery. *Obesity Surgery*. 2020;30(10):3947-
842 54. doi: 10.1007/s11695-020-04761-4.
- 843 70. Ernst B, Wilms B, Thurnheer M, Schultes B. Reduced circulating androgen levels after
844 gastric bypass surgery in severely obese women. *Obes Surg*. 2013;23(5):602-7. Epub
845 2012/11/29. doi: 10.1007/s11695-012-0823-9. PubMed PMID: 23188476.
- 846 71. Bastounis EA, Karayiannakis AJ, Syrigos K, Zbar A, Makri GG, Alexiou D. Sex hormone
847 changes in morbidly obese patients after vertical banded gastroplasty. *Eur Surg Res*.
848 1998;30(1):43-7. Epub 1998/03/11. doi: 10.1159/000008556. PubMed PMID: 9493693.
- 849 72. Dashti SG, English DR, Simpson JA, Karahalios A, Moreno-Betancur M, Biessy C, et al.
850 Adiposity and Endometrial Cancer Risk in Postmenopausal Women: A Sequential Causal
851 Mediation Analysis. *Cancer Epidemiol Biomarkers Prev*. 2021;30(1):104-13. Epub 2020/10/04.
852 doi: 10.1158/1055-9965.Epi-20-0965. PubMed PMID: 33008875.
- 853 73. Pournaras DJ, Osborne A, Hawkins SC, Vincent RP, Mahon D, Ewings P, et al. Remission
854 of type 2 diabetes after gastric bypass and banding: mechanisms and 2 year outcomes. *Ann*
855 *Surg*. 2010;252(6):966-71. Epub 2010/11/26. doi: 10.1097/SLA.0b013e3181efc49a. PubMed
856 PMID: 21107106.

- 857 74. ter Horst KW, Gilijamse PW, Koopman KE, de Weijer BA, Brands M, Kootte RS, et al.
858 Insulin resistance in obesity can be reliably identified from fasting plasma insulin. *International*
859 *Journal of Obesity*. 2015;39(12):1703-9. doi: 10.1038/ijo.2015.125.
- 860 75. Brochu M, Mathieu M-E, Karelis AD, Doucet É, Lavoie M-E, Garrel D, et al. Contribution
861 of the Lean Body Mass to Insulin Resistance in Postmenopausal Women With Visceral Obesity:
862 A Monet Study. *Obesity*. 2008;16(5):1085-93. doi: <https://doi.org/10.1038/oby.2008.23>.
- 863 76. Trapp EG, Chisholm DJ, Freund J, Boutcher SH. The effects of high-intensity intermittent
864 exercise training on fat loss and fasting insulin levels of young women. *International Journal of*
865 *Obesity*. 2008;32(4):684-91. doi: 10.1038/sj.ijo.0803781.
- 866 77. Ross R, Janssen I, Dawson J, Kungl A-M, Kuk JL, Wong SL, et al. Exercise-Induced
867 Reduction in Obesity and Insulin Resistance in Women: a Randomized Controlled Trial. *Obesity*
868 *Research*. 2004;12(5):789-98. doi: <https://doi.org/10.1038/oby.2004.95>.
- 869 78. Wei S, Schmidt MD, Dwyer T, Norman RJ, Venn AJ. Obesity and Menstrual Irregularity:
870 Associations With SHBG, Testosterone, and Insulin. *Obesity*. 2009;17(5):1070-6. doi:
871 <https://doi.org/10.1038/oby.2008.641>.
- 872 79. Tchernof A, Després J-P. Sex steroid hormones, sex hormone-binding globulin, and
873 obesity in men and women. *Hormone and Metabolic Research*. 2000;32(11/12):526-36.
- 874 80. Henderson BE, Ross RK, Pike MC, Casagrande JT. Endogenous hormones as a major
875 factor in human cancer. *Cancer Res*. 1982;42(8):3232-9. Epub 1982/08/01. PubMed PMID:
876 7046921.
- 877 81. Key TJ, Pike MC. The dose-effect relationship between 'unopposed' oestrogens and
878 endometrial mitotic rate: its central role in explaining and predicting endometrial cancer risk. *Br*

879 J Cancer. 1988;57(2):205-12. Epub 1988/02/01. doi: 10.1038/bjc.1988.44. PubMed PMID:
880 3358913; PubMed Central PMCID: PMCPMC2246441.

881 82. Kaaks R, Lukanova A, Kurzer MS. Obesity, endogenous hormones, and endometrial
882 cancer risk: a synthetic review. Cancer Epidemiol Biomarkers Prev. 2002;11(12):1531-43. Epub
883 2002/12/24. PubMed PMID: 12496040.

884 83. Bélanger C, Luu-The V, Dupont P, Tchernof A. Adipose tissue intracrinology: potential
885 importance of local androgen/estrogen metabolism in the regulation of adiposity. Horm Metab
886 Res. 2002;34(11-12):737-45. Epub 2003/03/28. doi: 10.1055/s-2002-38265. PubMed PMID:
887 12660892.

888 84. Somboonporn W, Davis SR. Testosterone Effects on the Breast: Implications for
889 Testosterone Therapy for Women. Endocrine Reviews. 2004;25(3):374-88. doi:
890 10.1210/er.2003-0016.

891 85. Bulun SE, Lin Z, Imir G, Amin S, Demura M, Yilmaz B, et al. Regulation of Aromatase
892 Expression in Estrogen-Responsive Breast and Uterine Disease: From Bench to Treatment.
893 Pharmacological Reviews. 2005;57(3):359-83. doi: 10.1124/pr.57.3.6.

894 86. Simó R, Sáez-López C, Barbosa-Desongles A, Hernández C, Selva DM. Novel insights in
895 SHBG regulation and clinical implications. Trends Endocrinol Metab. 2015;26(7):376-83. Epub
896 2015/06/06. doi: 10.1016/j.tem.2015.05.001. PubMed PMID: 26044465.

897 87. Grasa MDM, Gulfo J, Camps N, Alcalá R, Monserrat L, Moreno-Navarrete JM, et al.
898 Modulation of SHBG binding to testosterone and estradiol by sex and morbid obesity. Eur J
899 Endocrinol. 2017;176(4):393-404. Epub 2017/01/13. doi: 10.1530/eje-16-0834. PubMed PMID:
900 28077498.

88. Suba Z. Interplay between insulin resistance and estrogen deficiency as co- activators in carcinogenesis. *Pathol Oncol Res.* 2012;18(2):123-33. Epub 2011/10/11. doi: 10.1007/s12253-011-9466-8. PubMed PMID: 21984197.
89. Maturana MA, Spritzer PM. Association between hyperinsulinemia and endogenous androgen levels in peri- and postmenopausal women. *Metabolism.* 2002;51(2):238-43. doi: <https://doi.org/10.1053/meta.2002.29997>.
90. Weinberg ME, Manson JE, Buring JE, Cook NR, Seely EW, Ridker PM, et al. Low sex hormone-binding globulin is associated with the metabolic syndrome in postmenopausal women. *Metabolism.* 2006;55(11):1473-80. doi: <https://doi.org/10.1016/j.metabol.2006.06.017>.
91. Mu N, Zhu Y, Wang Y, Zhang H, Xue F. Insulin resistance: a significant risk factor of endometrial cancer. *Gynecol Oncol.* 2012;125(3):751-7. Epub 2012/03/28. doi: 10.1016/j.ygyno.2012.03.032. PubMed PMID: 22449736.
92. Haeusler RA, McGraw TE, Accili D. Biochemical and cellular properties of insulin receptor signalling. *Nature Reviews Molecular Cell Biology.* 2018;19(1):31-44. doi: 10.1038/nrm.2017.89.
93. Gunter MJ, Hoover DR, Yu H, Wassertheil-Smoller S, Manson JE, Li J, et al. A prospective evaluation of insulin and insulin-like growth factor-I as risk factors for endometrial cancer. *Cancer epidemiology, biomarkers & prevention : a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology.* 2008;17(4):921-9. doi: 10.1158/1055-9965.EPI-07-2686. PubMed PMID: 18398032.

94. Ogawa K, Sun C, Horii A. Exploration of genetic alterations in human endometrial cancer and melanoma: distinct tumorigenic pathways that share a frequent abnormal PI3K/AKT cascade. *Oncol Rep.* 2005;14(6):1481-5. Epub 2005/11/08. PubMed PMID: 16273242.
95. Slomovitz BM, Coleman RL. The PI3K/AKT/mTOR pathway as a therapeutic target in endometrial cancer. *Clin Cancer Res.* 2012;18(21):5856-64. Epub 2012/10/20. doi: 10.1158/1078-0432.Ccr-12-0662. PubMed PMID: 23082003.
96. Weigelt B, Warne PH, Lambros MB, Reis-Filho JS, Downward J. PI3K Pathway Dependencies in Endometrioid Endometrial Cancer Cell Lines. *Clinical Cancer Research.* 2013;19(13):3533-44. doi: 10.1158/1078-0432.Ccr-12-3815.
97. Phillips AN, Smith GD. How independent are “independent” effects? relative risk estimation when correlated exposures are measured imprecisely. *Journal of Clinical Epidemiology.* 1991;44(11):1223-31. doi: 10.1016/0895-4356(91)90155-3.
98. Subramaniam KS, Tham ST, Mohamed Z, Woo YL, Mat Adenan NA, Chung I. Cancer-associated fibroblasts promote proliferation of endometrial cancer cells. *PLoS One.* 2013;8(7):e68923. Epub 2013/08/08. doi: 10.1371/journal.pone.0068923. PubMed PMID: 23922669; PubMed Central PMCID: PMC3724864.
99. Subramaniam KS, Omar IS, Kwong SC, Mohamed Z, Woo YL, Mat Adenan NA, et al. Cancer-associated fibroblasts promote endometrial cancer growth via activation of interleukin-6/STAT-3/c-Myc pathway. *Am J Cancer Res.* 2016;6(2):200-13. Epub 2016/05/18. PubMed PMID: 27186396; PubMed Central PMCID: PMC4859653.

- 941 100. Renehan AG, Soerjomataram I, Tyson M, Egger M, Zwahlen M, Coebergh JW, et al.
942 Incident cancer burden attributable to excess body mass index in 30 European countries.
943 International Journal of Cancer. 2010;126(3):692-702. doi: <https://doi.org/10.1002/ijc.24803>.
- 944 101. Ochner CN, Tsai AG, Kushner RF, Wadden TA. Treating obesity seriously: when
945 recommendations for lifestyle change confront biological adaptations. The Lancet Diabetes &
946 Endocrinology. 2015;3(4):232-4. doi: 10.1016/S2213-8587(15)00009-1.
- 947 102. Evert AB, Franz MJ. Why Weight Loss Maintenance Is Difficult. Diabetes Spectrum.
948 2017;30(3):153-6. doi: 10.2337/ds017-0025.
- 949 103. Abarca-Gómez L, Abdeen ZA, Hamid ZA, Abu-Rmeileh NM, Acosta-Cazares B, Acuin C, et
950 al. Worldwide trends in body-mass index, underweight, overweight, and obesity from 1975 to
951 2016: a pooled analysis of 2416 population-based measurement studies in 128·9 million
952 children, adolescents, and adults. The Lancet. 2017;390(10113):2627-42. doi: 10.1016/S0140-
953 6736(17)32129-3.
- 954 104. Wang Y, Beydoun MA, Min J, Xue H, Kaminsky LA, Cheskin LJ. Has the prevalence of
955 overweight, obesity and central obesity levelled off in the United States? Trends, patterns,
956 disparities, and future projections for the obesity epidemic. International Journal of
957 Epidemiology. 2020;49(3):810-23. doi: 10.1093/ije/dyz273.
- 958 105. Pineda E, Sanchez-Romero LM, Brown M, Jaccard A, Jewell J, Galea G, et al. Forecasting
959 Future Trends in Obesity across Europe: The Value of Improving Surveillance. Obesity Facts.
960 2018;11(5):360-71. doi: 10.1159/000492115.
- 961 106. Zhou G, Myers R, Li Y, Chen Y, Shen X, Fenyk-Melody J, et al. Role of AMP-activated
962 protein kinase in mechanism of metformin action. J Clin Invest. 2001;108(8):1167-74. Epub

963 2001/10/17. doi: 10.1172/jci13505. PubMed PMID: 11602624; PubMed Central PMCID:
 964 PMCPMC209533.

965 107. Knowler WC, Fowler SE, Hamman RF, Christophi CA, Hoffman HJ, Brenneman AT, et al.
 966 10-year follow-up of diabetes incidence and weight loss in the Diabetes Prevention Program
 967 Outcomes Study. *Lancet*. 2009;374(9702):1677-86. Epub 2009/11/03. doi: 10.1016/s0140-
 968 6736(09)61457-4. PubMed PMID: 19878986; PubMed Central PMCID: PMCPMC3135022.

969 108. Boccardo F, Guarneri D, Rubagotti A, Casertelli GL, Bentivoglio G, Conte N, et al.
 970 Endocrine effects of tamoxifen in postmenopausal breast cancer patients. *Tumori*.
 971 1984;70(1):61-8. Epub 1984/02/29. PubMed PMID: 6538707.

972 109. Hu R, Hilakivi-Clarke L, Clarke R. Molecular mechanisms of tamoxifen-associated
 973 endometrial cancer (Review). *Oncol Lett*. 2015;9(4):1495-501. Epub 2015/03/20. doi:
 974 10.3892/ol.2015.2962. PubMed PMID: 25788989; PubMed Central PMCID: PMCPMC4356269.

975 110. Fleming CA, Heneghan HM, O'Brien D, McCartan DP, McDermott EW, Prichard RS. Meta-
 976 analysis of the cumulative risk of endometrial malignancy and systematic review of endometrial
 977 surveillance in extended tamoxifen therapy. *British Journal of Surgery*. 2018;105(9):1098-106.
 978 doi: 10.1002/bjs.10899.

979 111. Berruti A, Terzolo M, Pia A, Angeli A, Dogliotti L. Mitotane associated with etoposide,
 980 doxorubicin, and cisplatin in the treatment of advanced adrenocortical carcinoma. *Cancer*.
 981 1998;83(10):2194-200. doi: [https://doi.org/10.1002/\(SICI\)1097-
 982 0142\(19981115\)83:10<2194::AID-CNCR19>3.0.CO;2-3](https://doi.org/10.1002/(SICI)1097-0142(19981115)83:10<2194::AID-CNCR19>3.0.CO;2-3).

983 112. Theile D, Haefeli WE, Weiss J. Effects of adrenolytic mitotane on drug elimination
 984 pathways assessed in vitro. *Endocrine*. 2015;49(3):842-53. doi: 10.1007/s12020-014-0517-2.

- 985 113. Farland LV, Mu F, Eliassen AH, Hankinson SE, Tworoger SS, Barbieri RL, et al. Menstrual
986 cycle characteristics and steroid hormone, prolactin, and growth factor levels in premenopausal
987 women. *Cancer Causes Control*. 2017;28(12):1441-52. Epub 2017/11/01. doi: 10.1007/s10552-
988 017-0971-2. PubMed PMID: 29086892; PubMed Central PMCID: PMC5718047.
- 989 114. Janda M, Robledo KP, Gebiski V, Armes JE, Alizart M, Cummings M, et al. Complete
990 pathological response following levonorgestrel intrauterine device in clinically stage 1
991 endometrial adenocarcinoma: Results of a randomized clinical trial. *Gynecologic Oncology*.
992 2021;161(1):143-51. doi: <https://doi.org/10.1016/j.ygyno.2021.01.029>.
- 993 115. Barr CE, Ryan NAJ, Derbyshire AE, Wan YL, MacKintosh ML, McVey RJ, et al. Weight Loss
994 During Intrauterine Progestin Treatment for Obesity-associated Atypical Hyperplasia and Early-
995 Stage Cancer of The Endometrium. *Cancer Prev Res (Phila)*. 2021;14(11):1041-50. Epub
996 2021/08/06. doi: 10.1158/1940-6207.Capr-21-0229. PubMed PMID: 34348914.
- 997 116. Lagou V, Mägi R, Hottenga J-J, Grallert H, Perry JRB, Bouatia-Naji N, et al. Sex-dimorphic
998 genetic effects and novel loci for fasting glucose and insulin variability. *Nature Communications*.
999 2021;12(1):24. doi: 10.1038/s41467-020-19366-9.
- 1000 117. Chen J, Spracklen CN, Marenne G, Varshney A, Corbin LJ, Luan Ja, et al. The trans-
1001 ancestral genomic architecture of glycemic traits. *Nature Genetics*. 2021;53(6):840-60. doi:
1002 10.1038/s41588-021-00852-9.
- 1003 118. Larsson SC, Carter P, Vithayathil M, Kar S, Mason AM, Burgess S. Insulin-like growth
1004 factor-1 and site-specific cancers: A Mendelian randomization study. *Cancer Medicine*.
1005 2020;9(18):6836-42. doi: <https://doi.org/10.1002/cam4.3345>.

1006 119. Locke AE, Steinberg KM, Chiang CWK, Service SK, Havulinna AS, Stell L, et al. Exome
1007 sequencing of Finnish isolates enhances rare-variant association power. *Nature*.
1008 2019;572(7769):323-8. Epub 2019/07/31. doi: 10.1038/s41586-019-1457-z. PubMed PMID:
1009 31367044.

1010 120. Folkersen L, Gustafsson S, Wang Q, Hansen DH, Hedman ÅK, Schork A, et al. Genomic
1011 evaluation of circulating proteins for drug target characterisation and precision medicine.
1012 bioRxiv. 2020:2020.04.03.023804. doi: 10.1101/2020.04.03.023804.

1013 121. C Reactive Protein Coronary Heart Disease Genetics Collaboration (CCGC). Association
1014 between C reactive protein and coronary heart disease: mendelian randomisation analysis
1015 based on individual participant data. *BMJ*. 2011;342:d548. doi: 10.1136/bmj.d548.

1016

Figure Legends:

Fig 1. Flowchart detailing the process of identifying previously reported risk factors with suitable genetic instruments.

TNF- α = tumour necrosis factor- α , IGFBP-1 = insulin-like growth factor-binding protein-1, LDL = low-density lipoprotein, HDL = high-density lipoprotein, IGF-1 = insulin-like growth factor-1, IL-6 = interleukin-6, CRP = C-reactive protein, SHBG = sex hormone-binding globulin.

Fig 2. DAG demonstrating the core assumptions of Mendelian randomization.

DAG = directed acyclic diagram, G = genetic instrument, E = exposure, O = outcome, C = confounding factors. Arrows labelled 1, 2, and 3 represent the three core assumptions of MR: (1) the instrument strongly associates with the exposure (“relevance”); (2) there is no confounding of the instrument-outcome relationship (“exchangeability”); and (3) the instrument only affects the outcome through the exposure (“exclusion restriction”). MR uses genetic instruments to proxy exposures in order to strengthen causal inference in observational epidemiological settings. As these genetic instruments are randomly inherited at meiosis, they should not be affected by conventional confounding factors like environmental, lifestyle, and behavioural traits. In addition, since germline genetic variants are fixed at conception and cannot be altered by subsequent exposures, they are not susceptible to reverse causation. Finally, germline genotype can be measured relatively precisely using modern genotyping technologies which minimises measurement error. Collectively, these properties of germline genetic variants (along with technologies that measure them) permit MR analyses to minimise many of the sources of bias which can undermine robust causal inference in conventional observational epidemiological analyses.

Fig 3. DAG demonstrating the proposed causal interactions of BMI, SHBG, fasting insulin, and bioavailable testosterone on endometrial cancer (overall and the endometrioid histological subtype).

DAG = directed acyclic diagram, BMI = body mass index, SHBG = sex hormone-binding globulin.

Fig 4. Mendelian randomization analysis of BMI on overall and subtype-specific endometrial cancer risk.

Results of MR analyses examining the effect of adult BMI on risk of overall and subtype-specific endometrial cancer risk.

Fig 5. Mendelian randomization analysis of total serum cholesterol, fasting insulin, total testosterone, bioavailable testosterone, and sex hormone-binding globulin (SHBG) on overall and endometrioid endometrial cancer risk.

LDL = low-density lipoprotein, HDL = high-density lipoprotein, IGF-1 = insulin-like growth factor-1, IL-6 = interleukin-6, CRP = C-reactive protein, SHBG = sex hormone-binding globulin. (A) Results of MR analyses examining the effects of previously reported molecular risk factors on risk of overall endometrial cancer risk. (B) Results of MR analyses examining the effects of previously reported molecular risk factors on risk of endometrioid subtype endometrial cancer risk.

Fig 6. Mendelian randomization analysis of adult BMI on previously reported endometrial cancer risk factors.

SHBG = sex hormone-binding globulin, LDL = low-density lipoprotein, CRP = C-reactive protein.

1062 **Table 1. Details of the instruments used for exposures.**

Exposure	GWAS	Sample size	Number of SNPs	R ²	F-statistic	Sex-specificity
Adult BMI	Yengo et al. [42]	681,275	507	0.078	57,847	Combined
LDL-cholesterol	Willer et al. [34]	188,577	81	0.182	15,002	Combined
HDL-cholesterol	Willer et al. [34]	188,577	89	0.055	10,978	Combined
Triglyceride	Willer et al. [34]	188,577	55	0.052	9,811	Combined
Total serum cholesterol	Willer et al. [34]	188,577	88	0.063	12,696	Combined
Glucose	Neale et al. [36]	361,194	109	0.036	11,776	Female
Fasting insulin (unadjusted for BMI)	Lagou et al. [116]	98,210	14	0.005	523	Combined
Fasting insulin (adjusted for BMI)	Chen et al. [117]	150,571	14	0.006	865	Combined
IGF-1 (cis and trans variants)	Sinnott-Armstrong et al. [37]	358,072	413	0.036	13,367	Combined
IGF-1 (cis variants)	Larsson et al. [118]	358,072	1	0.002	814	Combined

IL-6	Georgakis et al. [38]	204,402	7	0.004	911	Combined
Adiponectin (cis and trans variants)	Locke et al. [119]	14,172	3	0.023	328	Combined
Adiponectin (cis variants)	Locke et al. [119]	14,172	3	0.023	334	Combined
Leptin	Folkersen et al. [120]	30,931	1	0.001	34	Combined
CRP (cis and trans variants)	Ligthart et al. [41]	204,402	45	0.035	7,414	Combined
CRP (cis variants)	C Reactive Protein Coronary Heart Disease Genetics Collaboration (CCGC) [121]	105,476	4	0.010	1,030	Combined
Total testosterone	Ruth et al. [29]	230,454	131	0.052	10,103	Female
Bioavailable testosterone	Ruth et al. [29]	188,507	147	0.054	10,599	Female

SHBG	Ruth et al. [29]	189,473	215	0.122	26,286	Female
------	------------------	---------	-----	-------	--------	--------

1063 BMI is scaled to an SD increase (4.7 kg/m²). For the analysis involving the plasma proteome, due to the
 1064 requirement of increased statistical power in order to overcome the multiple testing burden, alternative
 1065 summary genetic association data for BMI were obtained from a genome-wide association study of
 1066 681,275 individuals of European ancestry (note that this summary genetic data could not be used for
 1067 other analyses due to substantial overlap of participants with summary genetic data of other traits) [42].
 1068 The CRP GWAS included some individuals of non-European ancestry and adjusted for ancestry where
 1069 applicable. BMI = body mass index, LDL = low-density lipoprotein, HDL = high density lipoprotein, IGF-1 =
 1070 Insulin-like growth factor-1, IL-6 = interleukin-6, CRP = C-reactive protein, SHBG = sex hormone-binding
 1071 globulin, LD = linkage disequilibrium. For instrument construction of IGF-1 (cis and trans variants), a *P*
 1072 value of 5 x 10⁻⁶ was used.

Table 2. Results of MR analyses examining the effect of adult BMI on risk of overall and subtype-specific endometrial cancer risk

Outcome	Method	OR (95% CI)	P value
Overall	IVW	1.88 (1.69 to 2.09)	3.87×10^{-31}
endometrial	Weighted median	1.89 (1.58 to 2.26)	5.18×10^{-12}
cancer	Weighted mode	1.82 (1.35 to 2.44)	9.88×10^{-5}
	MR Egger	2.03 (1.54 to 2.67)	8.41×10^{-7}
Endometrioid	IVW	1.89 (1.65 to 2.16)	1.67×10^{-20}
endometrial	Weighted median	1.99 (1.60 to 2.46)	3.24×10^{-10}
cancer	Weighted mode	1.96 (1.42 to 2.69)	4.67×10^{-5}
	MR Egger	2.15 (1.52 to 3.03)	1.74×10^{-5}
Non-endometrioid	IVW	1.67 (1.19 to 2.35)	3.03×10^{-3}
endometrial	Weighted median	2.29 (1.24 to 4.22)	8.24×10^{-3}
cancer	Weighted mode	1.89 (0.77 to 4.63)	1.63×10^{-1}
	MR Egger	2.25 (0.99 to 5.07)	5.28×10^{-2}

ORs are shown per increase in SD (4.7 kg/m²) BMI. BMI = body mass index, IVW = inverse-variance weighted.

1078 **Table 3. Results of MR analyses examining the effect of previously reported and novel potential**
 1079 **molecular risk factors and overall and endometrioid subtype endometrial cancer risk.**

Exposure	Outcome	Method	OR (95% CI)	P value
LDL- cholesterol	Overall endometrial cancer	IVW	0.95 (0.87 to 1.04)	3.05×10^{-1}
		Weighted median	0.92 (0.82 to 1.03)	1.31×10^{-1}
		Weighted mode	0.91 (0.82 to 1.02)	9.54×10^{-2}
	Endometrioid endometrial cancer	MR Egger	0.90 (0.79 to 1.03)	1.35×10^{-1}
		IVW	0.98 (0.89 to 1.08)	6.70×10^{-1}
		Weighted median	0.96 (0.84 to 1.10)	5.61×10^{-1}
		Weighted mode	0.93 (0.83 to 1.06)	2.79×10^{-1}
		MR Egger	0.93 (0.81 to 1.07)	3.31×10^{-1}
		IVW	1.09 (0.97 to 1.23)	1.48×10^{-1}
		Weighted median	1.10 (0.96 to 1.26)	1.89×10^{-1}
HDL- cholesterol	Overall endometrial cancer	Weighted mode	1.07 (0.94 to 1.23)	3.11×10^{-1}
		MR Egger	1.08 (0.86 to 1.35)	4.99×10^{-1}
	Endometrioid endometrial cancer	IVW	1.04 (0.90 to 1.19)	6.05×10^{-1}
		Weighted median	0.99 (0.84 to 1.16)	8.76×10^{-1}
		Weighted mode	1.03 (0.88 to 1.21)	7.23×10^{-1}
		MR Egger	0.92 (0.71 to 1.20)	5.61×10^{-1}
	Overall endometrial cancer	IVW	0.95 (0.84 to 1.06)	3.55×10^{-1}
		Weighted median	0.87 (0.75 to 1.01)	7.75×10^{-2}
		Weighted mode	0.91 (0.79 to 1.04)	1.61×10^{-1}
		MR Egger	0.83 (0.69 to 1.00)	6.03×10^{-2}
Triglyceride	Overall endometrial cancer	IVW	0.95 (0.84 to 1.06)	3.55×10^{-1}
		Weighted median	0.87 (0.75 to 1.01)	7.75×10^{-2}
		Weighted mode	0.91 (0.79 to 1.04)	1.61×10^{-1}
		MR Egger	0.83 (0.69 to 1.00)	6.03×10^{-2}
	Endometrioid endometrial cancer	IVW	0.95 (0.83 to 1.09)	4.65×10^{-1}

Total serum cholesterol	Overall endometrial cancer	Weighted median	0.92 (0.78 to 1.08)	3.21×10^{-1}
		Weighted mode	0.91 (0.77 to 1.08)	2.87×10^{-1}
		MR Egger	0.87 (0.70 to 1.07)	2.02×10^{-1}
		IVW	0.90 (0.81 to 1.00)	4.01×10^{-2}
		Weighted median	0.80 (0.71 to 0.90)	2.08×10^{-4}
		Weighted mode	0.82 (0.73 to 0.91)	5.97×10^{-4}
	Endometrioid endometrial cancer	MR Egger	0.84 (0.71 to 0.98)	3.09×10^{-2}
		IVW	0.91 (0.82 to 1.02)	9.31×10^{-2}
		Weighted median	0.81 (0.71 to 0.93)	3.40×10^{-3}
		Weighted mode	0.84 (0.74 to 0.97)	1.64×10^{-2}
		MR Egger	0.86 (0.72 to 1.01)	7.60×10^{-2}
		IVW	0.95 (0.79 to 1.14)	5.64×10^{-1}
Glucose	Overall endometrial cancer	Weighted median	0.97 (0.80 to 1.17)	7.36×10^{-1}
		Weighted mode	0.93 (0.78 to 1.11)	4.47×10^{-1}
		MR Egger	0.99 (0.73 to 1.34)	9.29×10^{-1}
		IVW	0.90 (0.73 to 1.11)	3.26×10^{-1}
		Weighted median	0.98 (0.77 to 1.23)	8.31×10^{-1}
		Weighted mode	0.98 (0.80 to 1.20)	8.29×10^{-1}
	Endometrioid endometrial cancer	MR Egger	0.94 (0.67 to 1.33)	7.39×10^{-1}
		IVW	3.93 (2.29 to 6.74)	7.18×10^{-7}
		Weighted median	3.49 (1.60 to 7.62)	1.67×10^{-3}
		Weighted mode	3.55 (0.85 to 14.78)	1.06×10^{-1}
		MR Egger	8.28 (0.67 to 102.10)	1.25×10^{-1}
		IVW	4.64 (2.30 to 9.36)	1.84×10^{-5}
Fasting insulin	Overall endometrial cancer	IVW		
		Weighted median		
		Weighted mode		
		MR Egger		
		IVW		
		Weighted median		
Endometrioid endometrial cancer	Endometrioid endometrial cancer	Weighted mode		
		MR Egger		
		IVW		
		Weighted median		
		Weighted mode		
		MR Egger		

IGF-1 (cis and trans variants)	Overall endometrial cancer	Weighted median	3.93 (1.56 to 9.93)	3.80×10^{-3}
		Weighted mode	3.28 (0.69 to 15.62)	1.61×10^{-1}
		MR Egger	21.59 (0.78 to 593.93)	9.66×10^{-2}
		IVW	0.93 (0.85 to 1.06)	2.60×10^{-1}
		Weighted median	1.01 (0.85 to 1.20)	8.96×10^{-1}
	Endometrioid endometrial cancer	Weighted mode	1.22 (0.89 to 1.67)	2.28×10^{-1}
		MR Egger	1.17 (0.85 to 1.60)	3.49×10^{-1}
		IVW	0.89 (0.77 to 1.03)	1.12×10^{-1}
		Weighted median	1.03 (0.84 to 1.25)	8.06×10^{-1}
		Weighted mode	1.32 (0.92 to 1.90)	1.30×10^{-1}
IGF-1 (cis variants)	Overall endometrial cancer	Wald ratio	1.20 (0.79 to 1.82)	3.92×10^{-1}
	Endometrioid endometrial cancer	Wald ratio	1.40 (0.85 to 2.28)	1.84×10^{-1}
IL-6 (scaled to natural log transformed mg/L change in CRP)	Overall endometrial cancer	IVW	0.90 (0.66 to 1.21)	4.80×10^{-1}
	Endometrioid endometrial cancer	IVW	0.86 (0.60 to 1.23)	4.01×10^{-1}
Adiponectin (cis and trans variants)	Overall endometrial cancer	IVW	0.92 (0.79 to 1.08)	3.17×10^{-1}
	Endometrioid endometrial cancer	IVW	0.94 (0.78 to 1.13)	4.99×10^{-1}
Adiponectin (cis variants)	Overall endometrial cancer	IVW	0.95 (0.83 to 1.08)	3.94×10^{-1}
	Endometrioid endometrial cancer	IVW	1.00 (0.86 to 1.16)	9.92×10^{-1}
Leptin	Overall endometrial cancer	Wald Ratio	1.03 (0.68 to 1.54)	8.96×10^{-1}

	Endometrioid endometrial cancer	Wald Ratio	0.88 (0.54 to 1.42)	5.99×10^{-1}
CRP (cis and trans variants)	Overall endometrial cancer	IVW	1.07 (0.94 to 1.22)	3.03×10^{-1}
		Weighted median	0.97 (0.84 to 1.12)	6.76×10^{-1}
		Weighted mode	1.02 (0.91 to 1.14)	7.80×10^{-1}
		MR Egger	0.96 (0.80 to 1.16)	6.99×10^{-1}
	Endometrioid endometrial cancer	IVW	1.12 (0.96 to 1.30)	1.39×10^{-1}
		Weighted median	1.03 (0.87 to 1.23)	6.92×10^{-1}
		Weighted mode	1.04 (0.90 to 1.20)	6.34×10^{-1}
		MR Egger	0.97 (0.78 to 1.20)	7.69×10^{-1}
CRP (cis variants)	Overall endometrial cancer	IVW	0.98 (0.85 to 1.13)	7.52×10^{-1}
	Endometrioid endometrial cancer	IVW	0.98 (0.83 to 1.16)	8.02×10^{-1}
Total	Overall endometrial cancer	IVW	1.64 (1.43 to 1.88)	1.71×10^{-12}
testosterone		Weighted median	1.67 (1.39 to 2.01)	3.95×10^{-8}
		Weighted mode	1.74 (1.38 to 2.20)	8.33×10^{-6}
		MR Egger	1.81 (1.38 to 2.38)	4.17×10^{-5}
	Endometrioid endometrial cancer	IVW	1.60 (1.36 to 1.87)	8.70×10^{-9}
		Weighted median	1.81 (1.45 to 2.26)	2.05×10^{-7}
		Weighted mode	1.88 (1.42 to 2.48)	2.34×10^{-5}
		MR Egger	1.74 (1.26 to 2.41)	1.02×10^{-3}
Bioavailable testosterone	Overall endometrial cancer	IVW	1.46 (1.29 to 1.65)	3.48×10^{-9}
		Weighted median	1.47 (1.20 to 1.82)	2.46×10^{-4}
		Weighted mode	1.51 (1.19 to 1.93)	1.16×10^{-3}
		MR Egger	1.90 (1.46 to 2.47)	5.63×10^{-6}
	Endometrioid endometrial cancer	IVW	1.46 (1.26 to 1.69)	3.08×10^{-7}

		Weighted median	1.42 (1.14 to 1.76)	1.97×10^{-3}
		Weighted mode	1.60 (1.20 to 2.13)	1.59×10^{-3}
		MR Egger	1.79 (1.31 to 2.43)	3.08×10^{-7}
SHBG	Overall endometrial cancer	IVW	0.71 (0.59 to 0.85)	2.07×10^{-4}
		Weighted median	0.64 (0.48 to 0.86)	2.54×10^{-3}
		Weighted mode	0.69 (0.53 to 0.89)	4.97×10^{-3}
		MR Egger	0.62 (0.46 to 0.84)	2.52×10^{-3}
	Endometrioid endometrial cancer	IVW	0.65 (0.54 to 0.80)	3.31×10^{-5}
		Weighted median	0.60 (0.43 to 0.86)	4.90×10^{-3}
		Weighted mode	0.58 (0.40 to 0.82)	2.50×10^{-3}
		MR Egger	0.61 (0.44 to 0.84)	3.33×10^{-3}

1080 ORs are shown per increase in inverse normal transformed nmol/L SHBG, natural log transformed
 1081 pmol/L fasting insulin, inverse normal transformed nmol/L total testosterone, inverse normal
 1082 transformed nmol/L bioavailable testosterone, SD (38.7 mg/dL) LDL-cholesterol, nmol/L IGF-1, mmol/L
 1083 blood glucose, natural log transformed CRP mg/L IL-6, natural log transformed ug/ml adiponectin for
 1084 combined instrument, natural log transformed ug/ml cis-only adiponectin, natural log transformed mg/L
 1085 CRP, mg/dL triglyceride, SD (41.7 mg/dL) total serum cholesterol, mg/dL HDL-cholesterol, pg/mL leptin.
 1086 LDL = low-density lipoprotein, HDL = high density lipoprotein, IGF-1 = Insulin-like growth factor-1, IL-6 =
 1087 interleukin-6, CRP = C-reactive protein, SHBG = sex hormone-binding globulin, IVW = inverse-variance
 1088 weighted. For instrument construction of IGF-1 (cis and trans variants), a P value of 5×10^{-6} was used.

1089

1090 **Table 4. Results of MR analyses examining the effect of BMI on previously reported molecular risk**
 1091 **factors.**

Outcome	Method	Effect estimate (95% CI)	P value
Total serum cholesterol	IVW	-0.03 (-0.06 to 0.01)	1.22×10^{-1}
	Weighted median	-0.06 (-0.11 to -0.01)	2.98×10^{-2}
	Weighted mode	-0.08 (-0.15 to 0.00)	4.46×10^{-2}
	MR Egger	-0.08 (-0.16 to 0.00)	3.89×10^{-2}
Fasting insulin	IVW	0.17 (0.15 to 0.19)	1.51×10^{-74}
	Weighted median	0.18 (0.15 to 0.21)	8.51×10^{-31}
	Weighted mode	0.18 (0.12 to 0.24)	1.88×10^{-9}
	MR Egger	0.20 (0.16 to 0.25)	1.30×10^{-16}
Total testosterone	IVW	0.08 (0.05 to 0.11)	9.04×10^{-10}
	Weighted median	0.06 (0.03 to 0.09)	4.95×10^{-4}
	Weighted mode	-0.01 (-0.08 to 0.07)	8.60×10^{-1}
	MR Egger	0.02 (-0.05 to 0.09)	5.23×10^{-1}
Bioavailable testosterone	IVW	0.26 (0.23 to 0.29)	9.97×10^{-68}
	Weighted median	0.24 (0.20 to 0.28)	1.72×10^{-38}
	Weighted mode	0.09 (0.01 to 0.17)	2.41×10^{-2}
	MR Egger	0.16 (0.08 to 0.24)	4.36×10^{-5}
SHBG	IVW	-0.17 (-0.19 to -0.16)	4.86×10^{-125}
	Weighted median	-0.16 (-0.18 to -0.15)	8.85×10^{-77}
	Weighted mode	-0.15 (-0.18 to -0.12)	8.43×10^{-20}
	MR Egger	-0.13 (-0.17 to -0.09)	3.11×10^{-11}

1092 BMI is scaled to an SD increase (4.7 kg/m^2). Effect estimate represents change in SD (41.7 mg/dL) total
1093 serum cholesterol, natural log transformed pmol/L fasting insulin, inverse normal transformed nmol/L
1094 total testosterone, inverse normal transformed nmol/L bioavailable testosterone, and inverse normal
1095 transformed nmol/L SHBG. SHBG = sex hormone-binding globulin, IVW = inverse-variance weighted.

1096 **Table 5. Results of multivariable MR mediation analysis examining the effect of BMI and endometrial**
1097 **cancer with previously reported molecular risk factors as potential mediators.**

Mediator	Outcome	Direct Effect of BMI on Outcome	Indirect Effect of Mediator on Outcome	% Mediated (95% CI)	P value
Fasting	Overall	1.75	1.07	11% (1 to 21%)	2.89 x 10 ⁻²
insulin	Endometrial Cancer				
	Endometrioid	1.72	1.11	16% (3 to 28%)	1.24 x 10 ⁻²
	Endometrial Cancer				
Bioavailable	Overall	1.70	1.11	15% (10 to 20%)	1.43 x 10 ⁻⁸
testosterone	Endometrial Cancer				
	Endometrioid	1.72	1.11	15% (9 to 22%)	2.15 x 10 ⁻⁶
	Endometrial Cancer				
SHBG	Overall	1.80	1.04	7% (1 to 12%)	1.81 x 10 ⁻²
	Endometrial Cancer				
	Endometrioid	1.86	1.02	2% (-9 to 14%)	6.87 x 10 ⁻¹
	Endometrial Cancer				

1098 Direct effect is defined as the remaining effect of the exposure (BMI) on the outcome (endometrial
1099 cancer risk) when the effect of the candidate mediator on the outcome has been adjusted for. Indirect
1100 effect is defined as the effect of the exposure (BMI) on the outcome (endometrial cancer risk) through
1101 the candidate mediator. SHBG = sex hormone-binding globulin.

1102	Supporting Information Captions
1103	S1 Appendix. List of unique molecular traits identified from literature search.
1104	S2 Appendix. Supplementary methods.
1105	S3 Appendix. PubMed search terms for literature review.
1106	S4 Table. Additional information on GWAS, including covariates adjusted for.
1107	S5 Table. Sample overlap between GWAS.
1108	S6 Table. Conditional F-statistics for multivariable Mendelian randomization of BMI and mediators on
1109	endometrial cancer risk.
1110	S7 Figure. Leave-one-out analysis for MR examining the effect of adult BMI on overall endometrial
1111	cancer risk.
1112	S8 Table. Results of female BMI sensitivity analysis MR.
1113	S9 Table. Results from sensitivity analyses examining the influence of Winner's curse on GWAS with
1114	overlapping samples.
1115	S10 Figure. Leave-one-out analysis for MR examining the effect of adult BMI on endometrioid
1116	endometrial cancer risk.
1117	S11 Figure. Leave-one-out analysis for MR examining the effect of adult BMI on non-endometrioid
1118	endometrial cancer risk.
1119	S12 Figure. Leave-one-out analysis for MR examining the effect of total testosterone level on
1120	endometrial cancer risk.

1121 **S13 Figure. Leave-one-out analysis for MR examining the effect of bioavailable testosterone level on**
1122 **endometrial cancer risk.**

1123 **S14 Figure. Leave-one-out analysis for MR examining the effect of fasting insulin level on endometrial**
1124 **cancer risk.**

1125 **S15 Figure. Leave-one-out analysis for MR examining the effect of SHBG level on endometrial cancer**
1126 **risk.**

1127 **S16 Figure. Leave-one-out analysis for MR examining the effect of total serum cholesterol level on**
1128 **endometrial cancer risk.**

1129 **S17 Table. Results of female-specific SNP fasting insulin sensitivity analysis MR.**

1130 **S18 Table. Results of female-specific SNP CRP sensitivity analysis MR.**

1131 **S19 Figure. Leave-one-out analysis for MR examining the effect of total testosterone level on**
1132 **endometrioid endometrial cancer risk.**

1133 **S20 Figure. Leave-one-out analysis for MR examining the effect of bioavailable testosterone level on**
1134 **endometrioid endometrial cancer risk.**

1135 **S21 Figure. Leave-one-out analysis for MR examining the effect of fasting insulin level on**
1136 **endometrioid endometrial cancer risk.**

1137 **S22 Figure. Leave-one-out analysis for MR examining the effect of SHBG level on endometrioid**
1138 **endometrial cancer risk.**

1139 **S23 Figure. Leave-one-out analysis for MR examining the effect of adult BMI on fasting insulin level.**

1140 **S24 Figure. Leave-one-out analysis for MR examining the effect of adult BMI on SHBG level.**

1141 **S25 Figure. Leave-one-out analysis for MR examining the effect of adult BMI on bioavailable**
1142 **testosterone level.**

1143 **S26 Figure. Leave-one-out analysis for MR examining the effect of adult BMI on total serum**
1144 **cholesterol level.**

1145 **S27 Figure. Leave-one-out analysis for MR examining the effect of adult BMI on C-reactive protein**
1146 **level.**

1147 **S28 Table. Results of female-specific SNP BMI sensitivity mediation analysis.**

1148 **S29 Table. Results of female-specific SNP fasting insulin mediation analysis.**

1149 **S30 Table. Results of multivariable MR mediation analysis examining the effect of BMI and**
1150 **endometrial cancer with fasting insulin as a potential mediator with BMI-adjusted fasting insulin**
1151 **instrument and 100 SNPs from BMI instrument.**

1152 **S31 Table. Results from sensitivity analyses examining the influence of Winner's curse on GWAS with**
1153 **overlapping samples in analyses determining the mediating role of traits in the relationship between**
1154 **BMI and endometrial cancer risk.**

1155 **S32 Table. Results from sensitivity analyses examining the influence of Winner's curse on GWAS with**
1156 **overlapping samples in analyses determining the interdependent effects of mediators of the**
1157 **relationship between BMI and endometrial cancer risk.**

1158 **S33 Table. Conditional F-statistics for further multivariable Mendelian randomization analyses.**

1159 **S34 Table. Results of multivariable MR mediation analysis examining the effect of endometrial cancer**
1160 **with pairs of confirmed mediating molecular traits without BMI.**

1161 **S35 Table. Results of multivariable MR mediation analysis examining the effect of BMI and**
 1162 **endometrial cancer with fasting insulin as a potential mediator with 100 SNPs from BMI instrument.**

1163 **S36 Table. Results of multivariable MR mediation analysis examining the effect of BMI and**
 1164 **endometrial cancer with fasting insulin as a potential mediator with BMI-adjusted fasting insulin**
 1165 **instrument.**

1166 **S37 Table. Conditional F-statistics for fasting insulin (adjusted for BMI) and BMI with differing**
 1167 **thresholds used to construct fasting insulin.**

1168 **S38 Table. Results of multivariable MR mediation analysis examining the effect of BMI and**
 1169 **endometrial cancer with all confirmed mediating molecular traits with BMI-adjusted fasting insulin.**

1170 **S39 Table. Results of multivariable MR mediation analysis examining the effect of BMI and**
 1171 **endometrial cancer with pairs of confirmed mediating molecular traits.**

1172 **S40 Table. Results of multivariable MR mediation analysis examining the effect of BMI and**
 1173 **endometrial cancer with all confirmed mediating molecular traits including fasting insulin adjusted for**
 1174 **BMI with 100 SNPs from BMI instrument.**

1175 **S41 Table. Results of multivariable MR mediation analysis examining the effect of BMI and**
 1176 **endometrial cancer with pairs of confirmed mediating molecular traits with 100 SNPs from BMI**
 1177 **instrument.**

1178 **S42 Table. Results of multivariable MR mediation analysis examining the effect of endometrial cancer**
 1179 **with all confirmed mediating molecular traits without BMI.**

1180 **S43 Table. Results of sensitivity analysis examining the effect of only including 100 SNPs for the BMI**
 1181 **instrument in MR analyses.**

1182 **S44 Table. Conditional F-statistics with different levels of genetic correlation for SHBG and BMI in**
1183 **multivariable Mendelian randomization analyses.**

1184 **S45 Table. Results of HEIDI test-filtered low-density lipoprotein (LDL) cholesterol and overall**
1185 **endometrial cancer MR.**

20 risk factors identified from
two PubMed literature searches

5 risk factors excluded due to no
GWAS available with any SNPs at
genome-wide significance
($P < 5 \times 10^{-8}$)
30-minute postchallenge insulin,
TNF- α , visfatin, progesterone,
IGFBP-1

1 risk factor excluded due to
inappropriate instrument, as
demonstrated by Steiger filtering
Oestradiol

14 risk factors investigated in
MR analyses
LDL-cholesterol, HDL-cholesterol,
triglyceride, total serum
cholesterol, glucose, fasting
insulin, IGF-1, IL-6, adiponectin,
leptin, CRP, total testosterone,
bioavailable testosterone, SHBG

Fig 1.




Fig 2.




Fig 3.




Fig 4.




Fig 5.




Fig 6.