

1 Global Burden of Disease worldwide statistical analysis
2 of risk factors impacting noncommunicable disease deaths
3 compared with assessments by systematic
4 literature reviews

5
6 David K Cundiff (0000-0002-3206-9665), independent researcher,^{1 3} Chunyi Wu (0000-0002-
7 2186-3433), Research Epidemiologist/Statistician^{2 3}

8 ¹ Long Beach, California, USA

9 ² Area Specialist Lead in Epidemiology and Statistics, Michigan Medicine, Ann Arbor,
10 Michigan, USA

11 ³ Volunteer collaborators with the Institute of Health Metrics and Evaluation, Seattle,
12 Washington, USA

13 Correspondence to: David K Cundiff davidkcundiff@gmail.com Phone: 01-562-438-8805

14 Word count: abstract=501, text=4799

15

16 **Abstract**

17 **Background** Based on systematic literature reviews, the Institute of Health Metrics and
18 Evaluation (IHME) Global Burden of Disease (GBD) landmark risk factor paper attributes about
19 8 million noncommunicable diseases deaths/year to a composite of 15 dietary risk factors.
20 Another 27 million noncommunicable deaths out of 56.5 million total deaths worldwide are
21 attributed to high levels of metabolic risk factors.

22 **Methods** We format and population weight IHME GBD worldwide data resulting in 7886 rows
23 of data from cohorts of about 1 million people each representing about 7.8 billion people in
24 2020. Noncommunicable disease deaths/100k/year (NCD) are correlated with the four metabolic
25 risk factors, 15 dietary risk factors, six dietary covariates, and the composite “dietary risks”
26 variable. We compare NCD groups of cohorts (about 1000 cohorts, representing about 1 billion
27 people) with the lowest and highest NCD and the same for body mass index (BMI). We compare
28 literature reviews attributions of NCD causation with findings from the GBD data in assessing
29 the impact of dietary and metabolic risk factors on NCD.

30
31 **Findings** Literature review derived dietary risks composite risk factor correlated with NCD
32 account for 1.9% of worldwide NCD. Using GBD data to compare groups with the lowest versus
33 the highest rates of NCD: the lowest NCD 990 cohorts average 923.8 NCD and consume 320.4
34 kcal/day from six animal foods. The highest NCD 1054 cohorts average 2064 NCD and on
35 average and consume 117.2 kcal/day of six animal foods. Vitamin A deficiency in children is 75%
36 higher in children in the high NCD cohorts. Evaluating high BMI cohorts, 973 cohorts with
37 $BMI \geq 25$ account for 10.5% of the worldwide NCD. These high BMI cohorts also have higher

38 than average fasting plasma glucose (FPG: FPG mean=4.71 versus FPG mean=4.25) and low
39 density lipoprotein cholesterol (LDLC: LDLC mean=2.79 versus LDLC mean=2.30).

40

41 **Interpretation** The evidence basis of the dietary risks composite risk factor, accounting for only

42 1.9% of NCD worldwide should be questioned. Systematic literature reviews based versus GBD

43 data based estimates of metabolic and dietary risk factor impacts on NCD vary markedly.

44 Systematic literature reviews of studies involving individual subjects are better at ascertaining

45 the impact on NCD of obesity, hyperglycemia, high LDLC, and hypertension. High levels of

46 metabolic risk factors cannot be analysed with GBD data giving only the mean values for NCD

47 and risk factors. Since high mean BMI cohorts (mean BMI \geq 25) account for only 10.5% of

48 worldwide NCD, literature reviews appear to overestimate the worldwide impacts of high

49 metabolic risk factors on NCD (i.e., accounting for nearly 50% of worldwide deaths: 27

50 million/56.5 million worldwide deaths). High NCD, including cardiovascular disease deaths,

51 associated with low animal food intake suggests that fat soluble vitamin deficiencies (vitamins

52 A, D, E and K2, all largely from animal foods that also supply fatty acids for gut absorption)

53 may partly account for high rates of NCD worldwide, especially in developing countries.

54 Literature review based and GBD raw data based approaches should be complementary in

55 assessing dietary and other attributable risks for NCD and other individual health outcomes.

56 **Funding:** None

57

58 **Research in context**

59 **Evidence before this study**

60 The Global Burden of Diseases, Injuries, and Risk Factors Study (GBD 2019) provides the most
61 recent assessment of deaths, attributable to metabolic, environmental and occupational, and
62 behavioural risk factors. Controversies abound in the field of nutritional epidemiology. The lipid
63 hypothesis that dietary saturated fat and cholesterol promote cardiovascular diseases has been
64 disputed with no scientific consensus on the outcome.

65

66 **Added value of this study**

67

68 This study assesses the evidence basis of deaths attributed to dietary risks and metabolic risk
69 factors derived from systematic reviews of the medical literature in GBD 2019. This study uses
70 GBD raw data formatted and population weighted to assess the worldwide deaths from
71 noncommunicable diseases attributable to dietary and metabolic risk factors. With this GBD data
72 analysis methodology, the levels of 20 dietary risk factors are assessed in cohorts with the
73 highest and lowest rates of noncommunicable disease deaths. Worldwide GBD data show that
74 many deaths are associated with low levels of animal foods and saturated fatty acids (kcal/day
75 percapita) and associated with low levels of metabolic risk factors. These findings are at variance
76 with the lipid hypothesis operating uniformly worldwide. In the UK, USA, Mexico, and Japan,
77 with subnational GBD data, many deaths are attributable to high levels of animal foods, saturated
78 fatty acids, and metabolic risk factors, supporting the lipid hypothesis.

79

80

81

82 **Implications of all the available evidence**

83 The findings of systematic literature reviews in assessing the health impacts of high metabolic
84 risk factors are superior than using GBD data analysis because GBD data only assess mean
85 cohort levels of metabolic risk factors. However, GBD data can show effects of dietary risk
86 factors acting directly and mediating metabolic risks leading to noncommunicable disease
87 deaths. This GBD data based methodology can enhance understanding of the complex
88 interrelationships of diet, metabolic risk factors and noncommunicable disease deaths. The
89 findings are consistent with a “fat soluble vitamin hypothesis” that deficiencies of vitamins A, D,
90 E, and K2 and fatty acids from inadequate animal food consumption lead to increased
91 cardiovascular disease and noncommunicable disease deaths.

92

93 **Introduction**

94 Institute of Health Metrics and Evaluation (IHME) Global Burden of Disease (GBD) researchers
95 attribute about 8 million deaths/year worldwide to poor diet quality out of a total of about 56.5
96 million deaths a year worldwide.¹ GBD investigators ascribe high body mass index ($BMI \geq 25$) to
97 poor “diet quality,” excessive caloric intake, and low physical activity. Diet quality is defined as,
98 “the dietary risks that are based on the joint effects of 15 diet quality components.” Each dietary
99 risk factor has an optimal range or theoretical minimum risk exposure level range (TMREL
100 range in g/day) to consume based on systematic reviews of the literature. “Dietary risks” is a
101 composite metric including harmful risk factors: (1) high processed meat (TMREL=0), (2) high
102 red meat (TMREL=0), (3) high sugary beverages (TMREL=Any intake in g/day of
103 beverages with ≥ 50 kcal per 226.8 gram serving), (4) high trans fatty acids (TMREL=0),
104 and (5) high sodium (TMREL=average 24-hour urinary sodium excretion of 1-5 g/day)
105 and protective risk factors: (6) low milk (TMREL=90-100 g/day), (7) low omega-3 fatty acids
106 (TMREL= 430-470 milligrams of eicosapentaenoic acid (EPA) and docosahexaenoic
107 acid (DHA)), (8) low fruits (TMREL= 310-340 grams/day), (9) low vegetables (TMREL=
108 280-320 grams/day), (10) low nuts and seeds (TMREL= 10-19 grams/day) (11) low
109 legumes TMREL= 90-100 g/day, (12) low whole grains (TMREL= 140-160 g/day), (13) low
110 calcium (TMREL=1.06 – 1.11 g/day), (14) low fibre (21-22 g/day), and low (15)
111 polyunsaturated fatty acids (TMREL= 7-9% total energy intake).

112
113 GBD researchers ascribe another 27 million noncommunicable disease deaths to high levels of
114 metabolic risk factors high BMI, high fasting plasma glucose (FPG), high low density
115 lipoprotein cholesterol (LDL-C), and high systolic blood pressure (SBP)). Low levels of these

116 metabolic risk factors are not also included as risk factors for NCD. For example, while the
117 TMREL for BMI is 20 - 25 kg/M², no population attributable risk for BMI < 20 is reported.¹

118

119 This paper will use population weighted, formatted worldwide GBD data to begin to assess the
120 impact of dietary and metabolic risk factors on noncommunicable disease deaths/100k/year
121 (NCD). Findings will be compared with those derived from systematic literature reviews.

122

123

124 **Methods**

125 As volunteer collaborators with the IHME, we utilise raw GBD ecological data (≈ 1.4 Gigabytes)
126 on NCD, metabolic risk factors, dietary risk factors and covariates, and other risk factors of male
127 and female cohorts 15-49 years old and 50-69 years old from each year 1990-2017 from 195
128 countries and 365 subnational locations ($n=1120$ cohorts). We also use 2019 GBD data on the
129 dietary risks composite metric correlated with NCD. To maximally utilise the available GBD
130 data, we average the values for ages 15-49 years old together with 50-69 years old for NCD and
131 each risk factor exposure for each male and female cohort for each year. Finally, for each male
132 and female cohort, data from all 28 years (1990-2017) on mean NCD and on each of the risk
133 factor exposures are averaged using the computer software program R.

134
135 Food risk factors come as g/day consumed percapita. GBD dietary covariate data originally come
136 from Food and Agriculture Organization surveys of animal and plant food commodities available
137 percapita in countries worldwide—as opposed to consumed per capita.² Supplementary Table 1
138 lists the relevant GBD risk factors, covariates, and other available variables with definitions of
139 those risk factor exposures.³

140
141 GBD worldwide citations of over 12,000 surveys constituting ecological data inputs for this
142 analysis are available online from IHME.² The main characteristics of IHME GBD data sources,
143 the protocol for the GBD study, and all risk factor values have been published by IHME GBD
144 data researchers and discussed elsewhere.⁴⁻⁷ These include detailed descriptions of categories of
145 input data, potentially important biases, and methodologies of analysis. We do not clean or pre-
146 process any of the GBD data. GBD cohort risk factor and NCD data from the IHME have no

147 missing records. Because the GBD data for the study come from IHME, no ethics committee
148 approval or institutional review board review is needed for this post hoc statistical analysis. The
149 updated 2019 raw data with all the variables from the GBD 2017 data we use for this analysis
150 may be obtained by volunteer researchers collaborating with IHME.⁸

151
152 To weigh the country and subnational data according to population, internet searches (mostly
153 Wikipedia) yields the most recent population estimates for countries and subnational states,
154 provinces, and regions. The 1120 GBD cohorts available are population weighted by a software
155 program in R into an analysis dataset with 7886 population weighted cohorts representing about
156 7.8 billion people in 2020. Each male or female cohort in the population-weighted analysis
157 dataset represents approximately 1 million people (range: < 100,000 to 1.5 million). For
158 example, India with about 1.234 billion people has 617 rows of the same data for males and 617
159 rows for females. Maldives, with about 445,000 people, has a single row of data for males and
160 another for females. Without population-weighting the data, cohorts in India and Maldives each
161 would have had one row for males and one row for females in the analysis dataset, invalidating
162 the analysis results. World population data from the World Bank or the Organisation for
163 Economic Co-operation and Development could not be used because they do not include all 195
164 countries or any subnational data.

165
166 This report follows the STrengthening the Reporting of OBservational studies in Epidemiology
167 (STROBE) guidelines for reporting global health estimates.⁹

168

169 Supplementary Table 2 details how omega-3 fatty acid g/day is converted to fish g/day using
170 data on the omega-3 fatty acid content of frequently eaten fish from the National Institutes of
171 Health Office of Dietary Supplements (USA).¹⁰ As shown in Supplementary Table 3, we convert
172 all of the animal and plant food data, including alcohol and sugary beverage consumption, from
173 g/day to kcal/day. For the g to kcal conversions, we use the Nutritionix track app,¹¹ which tracks
174 types and quantities of foods consumed. Saturated fatty acids risk factor (0-1 portion of the entire
175 diet) is not available with GBD data from 2017, so GBD saturated fatty acids risk factor data
176 from GBD 2016 is used. Polyunsaturated fatty acid and trans fatty acid GBD risk factor data
177 from 2017 (0-1 portion of the entire diet) are also utilised. These fatty acid data expressed as 0-1
178 portion of the entire diet are converted to kcal/day by multiplying by the kcal/day available for
179 each cohort.

180

181 The principle outcome variable is noncommunicable disease deaths/100k/year (NCD). NCD is a
182 combination variable consisting of the deaths/100k/year of over 100 diseases including cancers,
183 diabetes, and diseases of the cardiovascular, respiratory, hepatic, renal, and central nervous
184 system.

185

186 **Statistical methods**

187 To determine the strengths of the risk factor correlations with mean NCD of population weighted
188 worldwide cohorts (7886 cohorts) or subgroups of cohorts (e.g., highest NCD and lowest NCD
189 cohorts, etc.), we utilise Pearson correlation coefficients: r , 95% confidence intervals (95% CIs),
190 and p values. The subgroup of worldwide data for a sensitivity analysis is the subnational data
191 from the UK, USA, Mexico, and Japan ($n=730$ cohorts). As with the worldwide data analysis,

192 we report NCD correlating with risk factors and NCD trend correlating with risk factor trends for
193 this subgroup.

194

195 We use SAS and SAS Studio statistical software 9.4 (SAS Institute, Cary, NC) for the data
196 analysis.

197 **Results**

198 Table 1 shows that the dietary risks composite risk factor correlated with NCD determined that
199 27.1 / 1426 (1.9%) of worldwide NCD are attributable to dietary risks. The dietary risks
200 composite risk factor is moderately strongly correlated with NCD ($r=0.390$, 95% CI: 0.371 to
201 0.408, $p<0.0001$). In comparison with the worldwide findings, the USA dietary risks related to
202 NCD is higher while the total NCD is lower, giving much higher percentages of NCD from
203 dietary risks (3.62%: 42.90 dietary risks related NCD / 1184 total NCD=0.0362). The correlation
204 of the dietary risks composite with NCD is very strong compared with worldwide and the other
205 regions in this table ($r=0.937$, 95% CI: 0.924 to 0.948, $p<0.0001$).

206

207 While the USA and the UK have similar rates of NCD (NCD=1184 and 1195, respectively), the
208 USA has more than threefold more the dietary risks composite risk factor related to NCD (42.90
209 /13.06=3.27).

210

211 As noted in the IHME GBD risk factors landmark article, the African Sahel region has the lowest
212 level of the dietary risks composite related disability adjusted life years (DALYs) in the world.
213 However, the African Sahel has above the worldwide average number of NCD (1580 versus
214 1426 worldwide). The GBD risk factors landmark article noted Central and Eastern Europe as
215 having among the highest DALYs in the world. However, the range of total NCD in the African
216 Sahel (1115-3229) largely overlaps the range of total NCD in Central and Eastern Europe (875-
217 3193).

218

219 Table 2a shows worldwide NCD correlated with metabolic risk factors. Mean BMI, mean FPG,
220 and mean LDL-C all negatively correlate with NCD ($r=-0.308$, 95% CI: -0.328 to -0.288,
221 $p<0.0001$, $r=-0.141$, 95% CI: -0.163 to -0.119, $p<0.0001$, $r=-0.341$, 95% CI: -0.360 to -0.321,
222 $p<0.0001$, respectively). Only mean SBP worldwide positively correlates with NCD ($r=-0.068$,
223 95% CI: 0.046 to 0.090, $p<0.0001$). On the other hand, Table 2b shows that the literature review
224 derived composite dietary risks related to NCD are positively correlated with all four metabolic
225 risk factors.

226
227 Table 3 compares the NCD in cohorts with the lowest mean BMI (n=952 cohorts) with the NCD
228 in the cohorts with the highest mean BMI (n=987 cohorts). The NCD is much higher in the low
229 BMI cohorts (mean NCD in low BMI cohorts=1821 versus mean NCD in high BMI cohorts
230 1207). The low BMI group with high NCD had a small fraction of the animal food kcal/day of
231 the high BMI group (low BMI six animal foods: 39.32 kcal/day versus high BMI six animal
232 foods: 367.2) and much less intake of fatty acids (169.3 versus 492.9 kcal/day).

233
234 Table 4 compares cohorts with the lowest and highest rates of NCD—each group with about
235 1000 cohorts. The male and female NCD are averaged for each location because of the
236 substantial differences in NCD between male and female cohorts. The mean animal food intake
237 in the low NCD cohorts (animal foods=320.4 kcal/day) is over twice the worldwide average
238 (animal foods=156.1 kcal/day) and nearly three times the mean animal food intake of the highest
239 NCD cohorts (animal foods=117.3 kcal/day). SFA+PUFA+TFA is 66% higher in the lowest
240 NCD cohorts compared with the highest NCD cohorts (406.3 kcal/day versus 244.5 kcal/day:
241 $406.3/244.5=1.662$). The rate of vitamin A deficiency in children is 75% higher in the highest

242 NCD cohorts than the lowest NCD cohorts (24678 / 14082=1.752). Vitamin A is a fat soluble
243 vitamin coming largely from animal foods.

244

245 Table 4 also reports the TMREL levels of the metabolic risk factors, 15 dietary risk factors,
246 alcohol, and vitamin A deficiency. The 990 cohorts with the lowest mean NCD of male-female
247 pairs of cohorts have mean BMI=23.77, but the range goes from below to above the TMREL
248 range (BMI range: 19.61 to 28.57). The 1054 cohorts with the highest NCD (mean NCD=2064)
249 also have wide ranging BMI levels (mean BMI=21.60, range: 17.95 to 27.73). The 576 cohorts
250 within the FPG TMREL range (FPG: 4.8 to 5.4 mmol/L) average 1409 NCD with a very wide
251 range (NCD range: 426 to 3349). The 7307 cohorts with mean FPG < 4.8 mmol/L have mean
252 NCD=1428, again with a very wide range (NCD range: 424 to 4321). Not a single cohort has a
253 mean LDLC or mean SBP within the TMREL.

254

255 Table 5 shows the NCD trend (by least squared linear regression) from 1990 to 2017 correlated
256 with the trends of metabolic and dietary risk factors. Notably, NCD trends downwards (- 2.11
257 NCD) while all the metabolic risk factors and macronutrient risk factors trend up except for
258 LDLC, corn availability, rice availability, sweet potato availability, and trans fatty acids. The
259 NCD trend correlates negatively with BMI, FPG, SBP, and negatively with all macronutrient risk
260 factors except for fish (p=0.86), sugary beverages, rice, sweet potatoes, and trans fatty acids.

261

262 As a contrast to the worldwide GBD data analysis, Table 6a shows NCD correlated with risk
263 factors from subnational data of the USA, UK, Mexico, and Japan (n combined=730 cohorts). In
264 these mostly developed countries (Mexico is classified advanced developing¹²) with NCD much

265 lower than worldwide (NCD four countries=1071 versus worldwide=1426), mean BMI is
266 strongly positively correlated with NCD ($r=0.504$, (95% CI 0.447 to 0.556, $p<0.0001$). As
267 opposed to the worldwide NCD analysis in Table 4, processed meat, red meat, milk, poultry,
268 SFA, PUFA, and kcal available are all at least moderately strongly correlated with NCD. As
269 opposed to the worldwide NCD trend's correlations in Table 5; the four country NCD trend is
270 positively correlated with the trends of processed meat, red meat, fish, poultry, eggs, and SFA.
271
272

273 **Discussion**

274 Table 1 shows that systematic literature review derived dietary risks account for only 1.9%:
275 (27.07 dietary risks related NCD / 1426 total NCD), bringing the dietary risks metric into
276 question. Regarding the USA, Table 1 shows the relatively high percent of NCD deaths
277 accounted for by dietary risks (% NCD due to dietary risks: 3.62%). Additionally, the USA has a
278 relatively strong correlation of dietary risks correlated with NCD with total NCD ($r=0.937$, 95%
279 CI: 0.924 to 0.948, $p<0.0001$). This suggests that the dietary risks metric, based on systematic
280 literature reviews, may be biased toward the NCD data from the USA and other developed
281 countries over developing countries.

282

283 The Sahel region of Africa, which has the world's lowest shares of DALYs attributed to dietary
284 risks of any region,¹ has higher than the worldwide average of NCD (Sahel region NCD: 1580
285 versus worldwide NCD: 1426. This also implies that the dietary risks metric may be biased
286 toward dietary risk factors as reported in medical literature from developed countries. In Central
287 and Eastern Europe, which literature reviews show have among the world's highest shares of
288 DALYs attributed to dietary risks of any region,¹ the much higher than average NCD has a range
289 that greatly overlaps the range of the Sahel region of Africa (Central and Eastern Europe: mean
290 NCD=2067, range: 875-3193 versus Sahel region: mean NCD=1580, range: 1115-3229). This
291 comparison further calls into question the evidence basis of the composite dietary risks metric.

292

293 In Tables 2a and 2b, NCD negatively correlate with BMI, FPG, and LDLC. However, literature
294 review derived dietary risks related to NCD positively correlate with BMI, FPG, and LDLC.

295 This also suggests that the dietary risks metric is biased toward the survey findings of the USA

296 and other developed countries in which high levels of BMI, FPG, and LDLC are more prevalent
297 than in developing countries.

298
299 Tables 3a and 3b detail how NCD in the cohorts lowest in BMI are much higher than in cohorts
300 highest in BMI (BMI<19 kg/M²: mean NCD=1821 versus BMI >24.95 kg/M²: mean
301 NCD=1207). This is further suggestive evidence that literature derived high BMI, high FPG, and
302 high LDLC attributions of NCD and noncommunicable disease DALYs are biased toward
303 developed countries over developing countries.

304
305 Table 3 also shows that the high NCD of cohorts with large numbers of underweight cohorts
306 (BMI< 18.5) are associated with very low mean animal food kcal/day intake that is about one-
307 ninth the animal food kcal/day intake of the highest BMI cohorts (mean animal food intake
308 lowest mean BMI cohorts=39.32 kcal/day versus mean animal food intake of the cohorts with
309 the highest mean BMI=367.2 kcal/day: 367.2/39.32=9.33). These worldwide data are at variance
310 with the lipid hypothesis (saying that dietary saturated fat and cholesterol, largely from animal
311 foods, cause high LDLC and atherosclerosis) as the dominant dietary risk factor for
312 cardiovascular disease (CVD). However, as is shown in Table 6, dietary SFA and cholesterol
313 may still strongly contribute to CVD mostly in developed countries with high intakes of animal
314 foods and the associated high rates of overweight and obesity.

315
316 Compared with analysing these worldwide GBD data, systematic literature reviews with
317 individual participant analysis would likely be much better than using GBD mean cohort values
318 of with about 1 million people per cohort in documenting CVD and NCD associated with

319 obesity, overweight, normal weight and underweight. Flegal, et. al., with an individual
320 participant level systematic review of BMI levels related to longevity, report that, compared to
321 normal weight ($18.5 \leq \text{BMI} < 25$), the summary HRs are the following: overweight ($25 \leq \text{BMI} < 30$):
322 (HR $r=0.94$ (95% CI, 0.91-0.96), stage 1 obesity ($30 \geq \text{BMI} < 35$): HR $r=0.95$ (95% CI, 0.88-1.01),
323 and above stage 1 obesity ($\text{BMI} \geq 35$): HR $r=1.29$ (95% CI, 1.18-1.41).¹³ A systematic review by
324 Aune, et. al. including underweight people finds BMI=20 gives a 15% increase in all cause
325 mortality (HR $r=1.15$ (95% CI, 1.13-1.17), $p < 0.0001$) and BMI=17.5 gives a 47% increase in all
326 cause mortality (HR $r=1.47$ (95% CI, 1.43-1.51), $p < 0.0001$)¹⁴ These are consistent with the
327 Table 3 findings for low BMI.

328

329 While GBD data on mean BMIs of cohorts cannot probe the impact of overweight and obesity on
330 NCD as can literature reviews, GBD data should be very informative for estimating the NCD
331 impact of cohorts with high BMI on average (i.e., mean BMI ≥ 25) in comparison with cohorts
332 that are underweight and normal weight on average (i.e., mean BMI < 25). The 973 cohorts with
333 mean BMI ≥ 25 have mean NCD=1208, while the 6913 cohorts with mean BMI < 25 have average
334 NCD=1457. The BMI ≥ 25 cohorts account for 10.5% of the total NCD and the BMI < 25 account
335 for the remaining 89.5% of the NCD (BMI ≥ 25 cohorts: 973 cohorts * 1208 (mean NCD) =
336 1,175,384 total NCD versus BMI < 25 cohorts: 6913 cohorts * 1457 (mean NCD) = 10,072,241
337 total NCD: 1,175,384 + 10,072,241 = 11,247,625 (total NCD), mean BMI ≥ 25 : 1,175,384 /
338 11,247,625 = 0.1045 and mean BMI < 25 : 10,072,241 / 11,247,625 = 0.8955).

339

340 BMI ≥ 25 cohorts also have higher FPG and LDLC: FPG mean=4.71 and LDLC mean=2.79
341 versus the BMI < 25 cohorts with: FPG mean=4.25 and LDLC mean=2.30. The mean levels of

342 SBP of the BMI \geq 25 cohorts and the BMI<25 cohorts differ by only 0.2 mm Hg (BMI \geq 25
343 cohorts SBP mean=133.71 versus BMI<25 cohorts SBP mean=133.91). These comparisons
344 strongly suggest that high levels of dietary and metabolic risk factors should account for about
345 10% of the worldwide NCD rather than about 62%, which can be derived from what is reported
346 from the analyses based on systematic literature reviews ((35 million worldwide deaths/year
347 attributable to dietary risks and metabolic risk factors / 56.5 million worldwide deaths/year¹)
348 =0.6195).

349
350 These GBD data showing that low consumption of animal foods and fatty acids correlate
351 strongly with low BMI and high NCD suggests that deficiencies of the fat soluble vitamins A, D,
352 E, and K might contribute to CVD and thereby NCD. Animal foods are the primary sources of
353 each of these vitamins and their absorption in the gut depends on adequate fatty acid intake that
354 also comes largely from animals. By different mechanisms, deficiencies of each of these fat
355 soluble vitamins raises the risk of CVD (vitamins A,¹⁵ D,¹⁶ E,¹⁵ and K¹⁷).

356
357 Our “fat soluble vitamin hypothesis” is that the high rates of CVD and, consequently, NCD in
358 countries with low animal food and low fatty acid intake stem to a significant degree from
359 deficiencies of fat soluble vitamins. For instance, over three times the vitamin A deficiency
360 prevalence in children occurs in the low BMI cohorts compared to high BMI cohorts (32.2% of
361 children < 5 years old of low BMI cohorts have vitamin A deficiency versus 10. 3% of children
362 from the high BMI cohorts, see Table 3).

363

364 Table 4, which compares 990 cohorts with the lowest NCD with 1054 cohorts with the highest
365 NCD, gives similar findings as Table 3 with low and high BMI levels. The metabolic and dietary
366 risk factors for the mean NCD of the lowest 1054 cohorts does not perceptively correspond
367 better with TMREL ranges than do the mean NCD of the highest 990 cohorts.

368
369 Again the higher intake of animal foods is associated with lower levels of NCD in Table 4. This
370 is all consistent with the hypothesis that fat soluble vitamin deficiencies significantly contribute
371 to NCD. The 75% higher rate of vitamin A deficiencies of the highest cohorts in NCD (vitamin
372 A deficiency rate =24,687/100k/year) versus the lowest NCD cohorts (14,082/100k/year) and is
373 another example.

374
375 Table 5 shows how the worldwide downward NCD trend from 1990 to 2017 relates to the trends
376 of the metabolic and dietary risk factors. While NCD trends downwards by -2.11 /year on
377 average, BMI, FPG, and SBP are trending upwards and LDLC is neutral. The NCD trend
378 negatively correlating with the BMI and FPG trends reinforces Table 2a data showing negative
379 correlations of NCD with BMI and FPG.

380
381 These metabolic risk factor correlations with NCD and the metabolic risk factor trends with the
382 NCD trend do not exclude that high levels of metabolic risk factors contribute significantly to
383 NCD. We have a vast literature relating individuals with high metabolic risk factor levels with
384 increased NCD. However, these GBD cohort data, now including the trend analysis, suggest that
385 BMI, FPG and perhaps LDLC correlate negatively with NCD.

386

387 In Table 5, all of the 20 food commodities are trending upwards except for corn, trans fatty acids,
388 rice and sweet potatoes. Kilocalories (kcal) /day available, a covariate, goes up by 12
389 kcal/day/year. As might be expected, trends of sugary beverages and trans fatty acid (the only
390 clearly always harmful food risk factors) positively correlate with the NCD trend. With
391 worldwide development, vitamin A deficiency in children is on the decline (-269
392 cases/100k/year from 1990-2017), most likely related to increased animal foods, which contain
393 most of the dietary vitamin A and fatty acids, which are required for gut absorption of vitamin A.
394
395 Despite the major increase in percapita food intake worldwide from 1990 to 2017, the NCD trend
396 and the trends of most macronutrients are still negatively correlated, indicating that more food
397 percapita is needed to drive down NCD, especially in developing countries. Dietary choices need
398 to be improved in places, especially developed countries, where abundant food availability
399 allows for and often promotes high levels of metabolic risk factors.

400
401 Table 4 shows that the cohorts with minimal NCD had higher mean sodium intake (3.67 g/day)
402 than the cohorts with maximum NCD (3.27 g/day). Accordingly, sodium g/day trend correlates
403 negatively with the NCD trend. This suggests that sodium lowering public health campaigns may
404 not be effective to benefit population health. However, individuals with CVD, hypertension, or
405 chronic diseases causing fluid overload still should reduce dietary sodium.

406
407 Table 6 shows how using data primarily from developed countries can easily support the lipid
408 hypothesis that defines saturated fat and cholesterol as the major dietary drivers of CVD and
409 NCD. However Table 6 findings are not generalisable to the world. In the four countries with

410 subnational GBD data in Table 6, the high level animal food consumption (animal foods=445
411 kcal/day) reverses the worldwide negative correlation of animal food with NCD (Four countries
412 cohorts: NCD versus animal foods, $r=0.214$, 95% CI (0.144 to 0.282), $p<0.0001$). Meat is
413 especially strongly positively correlated with NCD (NCD versus processed meats $r=0.576$, 95%
414 CI (0.525 to 0.622), $p<0001$ and NCD versus red meat $r=0.715$, 95% CI (0.677 to 0.748),
415 $p<0001$).

416
417 The unexpected positive correlations of NCD with corn availability (covariate), potato
418 availability (covariate) and fruit in Table 6 may be explained. Corn availability includes high
419 fructose corn syrup used widely in developed countries. Corn availability and sugary beverages
420 are highly positively correlated in these cohorts (corn availability versus sugary beverages,
421 $r=0.944$, 95% CI 0.935 to 0.951, $p<0.0001$). Fruits include 100% fruit juice in the GBD database
422 (see Supplementary Table 1). In the USA, about one-third of fruits consumed are in the form of
423 100% fruit juices.¹⁸ In these four countries combined, fruits are positively correlated with sugary
424 beverages (fruits versus sugary beverages, $r=0.186$, 95% CI 0.115 to 0.255, $p<0.0001$). Potato
425 availability includes highly processed potatoes like potato chips and French fries. According to
426 the International Potato Center, at least 50% of potatoes consumed worldwide are highly
427 processed.¹⁹ Processed potatoes may contain up to 6 g (or 54 kcal) per 100 kcal serving.²⁰ While
428 SFA worldwide averages only 191.3 kcal/day and is negatively correlated with NCD (SFA
429 versus NCD worldwide ($n=7886$ cohorts), $r=-0.253$, 95% CI -0.274 to -0.232, $p<0.0001$), SFA
430 averages 269.4 kcal/day and is positively correlated with NCD in these four mostly developed
431 countries combined (SFA versus NCD four countries with subnational data ($n=730$ cohorts),

432 $r=0.642$, 95% CI 0.597 to 0.682, $p<0.0001$). The lipid hypothesis is strongly supported by data
433 on people from these four countries and other developed countries.

434
435 As mentioned in the IHME GBD 2019 risk factor paper, the assessment of the joint effects of
436 dietary and other risk factors depends on two critical factors: the correlation of risk exposure and
437 the estimation of the joint effects of groups of risks together.¹ While the estimates of exposure to
438 the individual dietary risk factors appears excellent and improving with each iteration of the
439 GBD project, the estimation of the joint effects of the 15 food variables composite dietary risks
440 and the metabolic risk factors are problematic, as discussed regarding Tables 1 and 2 above.

441
442 The IHME GBD methodology assumption that, for each age-sex-location-year, the estimates of
443 the prevalence of exposure of each of the dietary variables are independent from each other is
444 said to be supported by findings from USA data from the National Health and Nutrition
445 Examination Survey. While this may have been the case for the USA where intakes of dietary
446 commodities vary much less than they do worldwide, the assumption of independence of
447 exposures of different dietary variables worldwide is highly dubious. As shown in Tables 3 and
448 4, there are marked differences between animal foods intake (four dietary risk factors and two
449 dietary covariates) between cohorts with the lowest and highest mean BMI and the lowest and
450 highest mean NCD. The GBD 2019 risk factor paper states in the limitations,

451
452 “To avoid over-estimation of the joint effects, we computed the non-mediated
453 relative risks and then assumed that non-mediated relative risks are multiplicative.
454 This approach does not capture potential synergy between relative risks in which

455 some combinations might be super-multiplicative. For some areas such as diet,
456 the joint estimation is very important for public policy. Further, more detailed
457 work is needed to strengthen the evidence base for understanding mediation.”¹

458
459 This GBD formatted data analysis is perhaps the detailed work that is needed to strengthen the
460 evidence base for understanding mediation and for more accurately determining the deaths and
461 DALYs attributable to dietary and other risk factors.

462
463 Also mentioned under limitations in the GBD risk factor paper, “...we assume that relative risks
464 as a function of exposure are universal and apply in all locations and time periods.”¹ Tables 4-6
465 of this GBD data analysis gives a clear example of how the level of exposure of animal foods in
466 cohorts worldwide versus in the four countries with subnational data lead to diametrically
467 opposing correlations between NCD and the six animal foods kcal/day. Likewise, Table 6 shows
468 the worldwide NCD trend negative correlations with trends of animal foods and SFA are
469 reversed in the subnational data analysis (n=730 cohorts) for processed meat, red meat, milk,
470 poultry and SFA.

471
472 **Limitations**

473 The GBD data on animal foods, plant foods, alcohol, sugary beverages, and fatty acids are not
474 comprehensive and comprised only 1191.4 kcal/day on average worldwide. Subnational data are
475 available on only four countries. Because the data formatting and statistical methodology are
476 new, this is necessarily a post hoc analysis and no pre-analysis protocol is possible. This GBD
477 data analysis should be repeated with the most recently released GBD 2019 data when it
478 becomes available to volunteer collaborators.

479 **Generalisability**

480 The new methodology presented for probing the GBD raw data to find correlations between risk
481 factors (e.g., dietary and metabolic risk factors) and health outcomes (e.g., NCD) can be applied
482 to any of the risk factors and health outcomes available with the new GBD 2019. The intake of
483 each of the 20 dietary variables in this analysis that corresponds with the 990 cohorts with the
484 lowest NCD worldwide will not be the optimal healthy diet for everyone. The total kcal/day of
485 animal foods, plant foods and fatty acids are all important. Having comprehensive data on
486 dietary inputs is key to more accurate and reliable analyses. Baseline metabolic risk factor levels
487 are important. The optimal diet to lower NCD risk for someone with BMI=35 will be different
488 than in someone with BMI=20. This analysis is merely a proof of concept the the GBD data
489 analysis methodology can derive plausible results and that the GBD data analysis methodology
490 can be complimentary to the systematic literature review methodology.

491

492 **Conclusion**

493 Statistically analysing GBD data provides a tool to compliment the systematic literature review
494 methodology in forming policy recommendations, clinical practice guidelines, and public health
495 recommendations. Eventually, GBD data analysis can bring more rigor, precision, and consensus
496 to the field of population health, especially in the areas of dietary and metabolic risk factors.

497

498

499 **Table 1 Dietary risks related NCD compared with total NCD worldwide (n=7886 cohorts,**
 500 **representing about 7.5 billion people in 2020)**

Regions or countries	NCD in region or country	NCD r	SD	Min	Max	Dietary risks versus total NCD r	95% CI	p	(Dietary risks related NCD / Total NCD) * 100 = %
World	Dietary risks related NCD	27.07	22.92	2.6	220				
World	Total NCD	1426	463	424	4321	0.390	0.371 to 0.408	<.0001	1.9
USA	Dietary risks related NCD	42.90	19.46	14.17	98.5				
USA	Total NCD	1184	316	697	2312	0.937	0.924 to 0.948	<.0001	3.62
UK	Dietary risks related NCD	13.06	14.67	2.71	61.1				
UK	Total NCD	1195	330	777	1823	0.610	0.428 to 0.740	<.0001	1.09
African Sahel	Dietary risks related NCD	25.03	11.28	11.7	47.1				
African Sahel	Total NCD	1580	340	1115	3229	0.565	0.441 to 0.666	<.0001	1.58
Central and Eastern Europe	Dietary risks related NCD	50.24	42.14	8.79	220				
Central and Eastern Europe	Total NCD	2067	827	875	3193	0.355	0.253 to 0.447	<.0001	2.43

501
 502
 503

504 **Table 2. Worldwide NCD correlated with metabolic risk factors and dietary risks related**
 505 **NCD also correlated with metabolic risk factors**

506 **Table 2a. Worldwide NCD correlated with metabolic risk factors**

NCD versus metabolic risk factors (n=7886 cohorts)	Mean	SD	Max	Min	NCD r	95% CI	p
NCD	1426	462.8	423.8	4321			
BMI kg/M ²	21.79	2.31	17.95	29.39	-0.308	-0.328 to -0.288	<.0001
Fasting plasma glucose mmol/L	4.31	0.35	3.32	5.58	-0.141	-0.163 to -0.119	<.0001
LDL cholesterol mmol/L	2.36	0.40	1.27	3.25	-0.341	-0.360 to -0.321	<.0001
Systolic BP mm Hg	133.9	4.33	123.4	147.9	0.068	0.046 to 0.090	<.0001

507

508 **Table 2b. Dietary risks related NCD correlated with metabolic risk factors**

Dietary risks related NCD versus metabolic risk factors (n=7886 cohorts)	Mean	SD	Min	Max	Diet related NCD r	95% CI	p
Dietary risks related NCD	22.60	20.31	1.7	196.8			
BMI kg/M ²	21.79	2.31	17.95	29.39	0.207	0.186 to 0.228	<.0001
Fasting plasma glucose mmol/L	4.31	0.35	3.32	5.58	0.113	0.091 to 0.134	<.0001
LDL cholesterol mmol/L	2.36	0.40	1.27	3.25	0.147	0.125 to 0.169	<.0001
Systolic BP mm Hg	133.9	4.33	123.4	147.9	0.228	0.207 to 0.249	<.0001

509

510

511 **Table 3. BMI<19.00 versus BMI >24.95 associated with NCD and NCD risk factors**

512 **Table 3a. BMI<19 associated with NCD and NCD risk factors**

NCD with mean BMI<19 kg/M² (n=952 cohorts)	Mean	SD	Min	Max
NCD	1821	215	1242	3229
BMI kg/M²	18.86	0.10	17.95	19.00
FPG mmol/L	4.21	0.21	3.48	4.32
LDLC mmol/L	1.89	0.16	1.41	2.22
SBP mm Hg	130.9	2.39	125.7	139.9
Processed meat kcal/day	1.33	0.32	0.31	1.62
Red meat kcal/day	11.29	13.09	3.84	61.38
Fish kcal/day	1.57	0.77	0.47	7.53
Milk kcal/day	14.99	6.91	1.35	19.94
Poultry kcal/day	5.85	4.53	2.72	24.99
Eggs kcal/day	4.29	1.06	1.23	5.81
Alcohol kcal/day	110.3	40.89	33.37	316.4
Sugary beverages kcal/day	316.2	51.57	199.4	350.0
Corn kcal/day	15.27	14.10	0.47	148.4
potatoes kcal/day	47.21	41.85	10.27	289.1
SFA kcal/day	126.5	24.41	71.02	155.0
PUFA kcal/day	30.46	8.77	2.99	47.74
TFA kcal/day	12.32	4.13	2.64	15.32
Fruits kcal/day	20.89	7.53	3.58	33.95
Vegetables kcal/day	58.24	26.10	11.60	99.86
Nuts and seeds kcal/day	5.40	2.25	1.09	12.55
Whole grains kcal/day	66.99	28.77	4.03	124.3
Legumes kcal/day	82.31	32.00	26.42	180.8
Rice kcal/day	238.4	120.1	1.42	443.6
Sweet potatoes kcal/day	6.03	8.12	2.04	53.93
Kcal /day	2246	189	1579	2347
Vitamin A deficient children/100k/yr	32157	6122	21075	43336
Sodium g/day	3.24	0.50	2.02	4.69
Calcium g/day	0.20	0.05	0.11	0.28
Dietary fiber g/day	8.23	2.94	2.78	16.25
Animal foods kcal/day	39.32	16.23	18.58	97.38
Plant foods kcal/day	540.7	120.6	229.8	809.6
SFA+PUFA+TFA kcal/day	169.3	22.94	79.92	185.0

513

514

515 **Table 3b. BMI>24.95 associated with NCD and NCD risk factors**

NCD with mean BMI >24.95 (n=987 cohorts)	Mean	SD	Min	Max
NCD	1207	326.9	660.9	3021
BMI kg/M²	26.06	0.69	24.97	29.39
Fasting plasma glucose mmol/L	4.71	0.32	3.69	5.58
LDL cholesterol mmol/L	2.79	0.22	1.83	3.20
Systolic BP mm Hg	133.8	5.29	125.9	146.0
Processed meat kcal/day	20.90	17.46	0.60	68.77
Red meat kcal/day	100.7	53.73	18.65	207.8
Fish kcal/day	11.69	4.36	3.07	27.73
Milk kcal/day	65.95	28.52	5.81	146.8
Poultry kcal/day	133.4	62.63	18.53	411.9
Eggs kcal/day	34.56	10.69	7.41	51.53
Alcohol kcal/day	127.3	88.00	6.07	429.8
Sugary beverages kcal/day	285.4	206.5	131.0	1472
Corn kcal/day	40.57	63.45	0.32	300.7
potatoes kcal/day	111.6	57.67	12.36	338.1
SFA kcal/day	275.9	74.39	109.5	479.4
PUFA kcal/day	183.8	97.77	36.04	346.9
TFA kcal/day	33.28	22.40	2.25	77.76
Fruits kcal/day	63.45	19.82	16.23	161.4
Vegetables kcal/day	115.7	44.94	22.06	304.2
Nuts and seeds kcal/day	18.24	11.47	0.07	103.0
Whole grains kcal/day	52.49	31.10	1.60	173.5
Legumes kcal/day	37.70	20.82	2.80	120.3
Rice kcal/day	30.25	33.13	3.59	179.6
Sweet potatoes kcal/day	2.35	3.77	0.02	37.83
Kcal /day	3227	221	2246	3898
Vitamin A deficient children/100k/yr	10320	7607	1400	39657
Sodium g/day	3.52	0.67	1.53	6.47
Calcium g/day	0.58	0.17	0.17	1.04
Dietary fiber g/day	11.01	2.10	5.74	18.15
Animal foods kcal/day	367.2	148.8	73.58	618.3
Plant foods kcal/day	472.3	91.79	227.0	817.9
SFA+PUFA+TFA kcal/day	492.9	163.4	194.6	828.0

516

517 **Table 4. Comparing metabolic and dietary risk factor values in cohorts with the lowest and**
 518 **highest NCD**

519 **Table 4a. Lowest NCD cohorts**↓

NCD and risk factors (n=990 cohorts with minimal NCD)	Mean	SD	Min	Max	TMREL low kcal/day	TMREL high kcal/day
NCD	923.8	267.0	423.8	1403		
BMI kg/M²	23.77	1.49	19.61	28.57	20	25
FPG mmol/L	4.49	0.30	3.54	5.58	4.8	5.4
LDLC mmol/L	2.73	0.30	1.60	3.25	0.7	1.3
SBP mm Hg	134.1	4.05	123.4	144.4	110	115
Processed meat kcal/day	12.46	11.75	0.87	66.78	0	0
Red meat kcal/day	81.85	48.92	12.03	205.4	0	0
Fish kcal/day	49.34	93.27	2.66	370.4	64	70
Milk kcal/day	54.38	32.16	14.60	146.8	180	250
Poultry kcal/day available	89.81	47.20	4.74	240.6		
Eggs kcal/day available	32.61	13.59	4.77	69.64		
Alcohol kcal/day	103.6	65.99	4.25	296.7	0	0
Sugary beverages kcal/day	350.8	257.2	72.91	1472	200	200
Corn kcal/day available	37.20	52.04	1.79	281.5		
potatoes kcal/day available	95.68	48.95	18.09	307.8		
SFA kcal/day	245.0	77.92	129.3	481.1		
PUFA kcal/day	143.3	60.62	34.28	334.7	207	266
TFA kcal/day	17.97	16.28	3.59	73.64	0	0
Fruits kcal/day	63.66	18.63	21.61	113.7	112	122
Vegetables kcal/day	105.8	48.24	9.48	228.7	182	208
Nuts and seeds kcal/day	14.42	11.49	0.26	45.28	58	110
Whole grains kcal/day	50.07	32.34	1.61	165.1	323	370
Legumes kcal/day	49.40	26.95	3.01	134.7	125	139
Rice kcal/day available	58.10	49.08	3.59	171.6		
Sweet potatoes kcal/day available	4.65	6.90	0.04	37.83		
Kcal available/day	2959	403	1948	3572		
Vitamin A deficiency in children	14082	8954	1368	44100	0	0
Sodium g/day	3.67	1.15	1.33	6.70	1	5
Calcium g/day	0.49	0.17	0.19	1.04	1.06	1.1
Dietary fiber g/day	10.84	2.32	5.41	18.15	21	22
Animal foods kcal/day	320.4	141.7	52.52	701.4		
Plant foods kcal/day	479.0	91.16	293.6	730.8		
SFA+PUFA+TFA kcal/day	406.3	117.6	198.2	732.4		

520
 521 ↓ Because NCD levels of males are much higher than for females, for each location the NCD of males
 522 and females are averaged and the average number of NCD for each male-female cohort pair<1070
 523

524 **Table 4b. Highest NCD cohorts**↓

NCD and risk factors (n=1054 cohorts with maximal NCD)	Mean	SD	Min	Max	TMREL low kcal/day	TMREL high kcal/day
NCD	2064	572	1080	4321		
BMI kg/M²	21.60	2.31	17.95	27.73	20	25
FPG mmol/L	4.25	0.45	3.38	5.18	4.8	5.4
LDLC mmol/L	2.27	0.48	1.34	3.20	0.7	1.3
SBP mm Hg	135.6	5.27	123.7	147.9	110	115
Processed meat kcal/day	3.82	5.52	0.22	65.14	0	0
Red meat kcal/day	39.04	35.31	6.56	194.8	0	0
Fish kcal/day	4.37	3.28	0.40	20.32	64	70
Milk kcal/day	25.26	19.90	1.06	100.9	180	250
Poultry kcal/day	30.39	28.21	1.06	207.7		
Eggs kcal/day	14.39	14.69	0.79	49.16		
Alcohol kcal/day	58.14	45.65	5.77	340.4	0	0
Sugary beverages kcal/day	294.7	75.44	131.2	768.5	200	200
Corn kcal/day	31.11	42.66	0.20	268.8		
potatoes kcal/day	112.2	110.8	3.94	533.9		
SFA kcal/day	186.3	60.78	71.02	447.5		
PUFA kcal/day	49.43	40.69	5.93	333.8	207	266
TFA kcal/day	8.72	5.62	2.64	50.09	0	0
Fruits kcal/day	25.70	14.17	3.58	107.5	112	122
Vegetables kcal/day	59.86	41.70	14.43	152.6	182	208
Nuts and seeds kcal/day	4.76	3.88	0.05	49.94	58	110
Whole grains kcal/day	34.76	33.77	1.34	146.9	323	370
Legumes kcal/day	39.53	29.20	0.64	194.7	125	139
Rice kcal/day	79.65	121.8	1.42	461.8		
Sweet potatoes kcal/day	9.94	26.90	0.02	364.7		
Kcal /day available	2392	478	1579	3481		
Vitamin A deficiency in children	24678	13103	1718	50969	0	0
Sodium g/day	3.27	0.88	1.33	6.47	1	5
Calcium g/day	0.30	0.17	0.08	0.86	1.06	1.1
Dietary fiber g/day	9.06	2.94	2.72	20.67	21	22
Animal foods kcal/day	117.3	97.21	14.53	612.9		
Plant foods kcal/day	397.5	199.7	145.0	847.1		
SFA+PUFA+TFA kcal/day	244.5	89.76	79.92	694.5		

525
 526 ↓ Because NCD levels of males are much higher than for females, for each location, the NCD of males
 527 and females are averaged and the average number of NCD for each male-female cohort pair>1763
 528

529 **Table 5. NCD trend versus risk factor trends**

NCD trend and risk factor trends worldwide (n=7886 cohorts)	Mean	SD	Min	Max	NCD trend v risk factor trend r	95% CI	p
NCD trend	-2.11	46.54	-119	355			
BMI trend	0.08	0.03	-0.02	0.22	-0.055	-0.077 to -0.033	<.0001
FPG trend	0.02	0.01	-0.01	0.04	-0.158	-0.179 to -0.136	<.0001
LDLC trend	0.00	0.01	-0.02	0.02	0.043	0.021 to 0.065	0.0001
SBP trend	0.02	0.11	-0.48	0.31	-0.055	-0.077 to -0.033	<.0001
Processed meat kcal/day trend	0.05	0.16	-0.81	1.68	-0.143	-0.165 to -0.121	<.0001
Red meat kcal/day trend	0.55	1.09	-3.29	4.73	-0.114	-0.136 to -0.092	<.0001
Fish kcal/day trend	0.22	0.95	-0.13	9.79	0.002	0.020 to 0.024	0.8647
Milk kcal/day trend	0.19	0.30	-2.02	2.50	-0.072	-0.094 to -0.050	<.0001
Poultry kcal/day trend	1.15	1.04	-0.87	7.66	-0.020	-0.042 to -0.002	0.0701
Eggs kcal/day trend	0.30	0.49	-0.88	1.27	-0.147	-0.169 to -0.126	<.0001
Alcohol kcal/day trend	0.03	0.16	-0.91	1.99	-0.270	-0.290 to -0.249	<.0001
Sugary beverages kcal/day trend	0.29	1.14	-5.40	14.00	0.081	0.059 to 0.103	<.0001
Corn kcal/day trend	-0.03	0.60	-4.36	2.81	-0.056	-0.078 to -0.034	<.0001
Potatoes kcal/day trend	0.24	1.83	-12.5	11.03	-0.110	-0.131 to -0.088	<.0001
SFA trend	0.21	0.92	-3.66	2.09	-0.253	-0.274 to -0.232	<.0001
PUFA trend	1.00	0.96	-3.03	5.47	-0.026	-0.048 to -0.004	0.0195
TFA trend	-0.21	0.45	-2.68	0.01	0.123	0.102 to 0.145	<.0001
Fruits kcal/day trend	0.62	0.60	-2.08	4.10	-0.095	-0.117 to -0.073	<.0001
Vegetables kcal/day trend	1.39	1.61	-7.96	8.47	-0.027	-0.049 to -0.005	0.0151
Nuts and seeds kcal/day trend	0.28	0.27	-0.34	3.12	-0.136	-0.158 to -0.115	<.0001
Whole grains kcal/day trend	0.17	0.58	-2.60	8.46	0.162	0.141 to 0.183	<.0001
Legumes kcal/day trend	0.37	0.98	-3.24	5.40	-0.087	-0.109 to -0.065	<.0001
Rice kcal/day trend	-0.78	1.29	-5.05	3.50	0.191	0.170 to 0.212	<.0001
Sweet potatoes kcal/day trend	-0.46	1.34	-4.82	3.91	0.124	0.102 to 0.146	<.0001
Kcal/day available trend	12.12	10.10	-28.1	47.89	-0.180	-0.201 to -0.158	<.0001
Vitamin A deficient children/100k/yr trend	-269	253	-1425	169.4	-0.057	-0.078 to -0.034	<.0001
Sodium g/day trend	-0.001	0.044	-0.219	0.085	-0.107	-0.128 to -0.085	<.0001
Calcium g/day trend	0.003	0.003	-0.007	0.018	0.155	0.133 to 0.176	<.0001
Dietary fiber g/day trend	0.06	0.07	-0.23	0.54	-0.073	-0.095 to -0.051	<.0001

530
531
532
533

534 **Table 6. NCD and the NCD trend versus risk factors and risk factor trends in USA, UK,**
 535 **Mexico, and Japan combined**

536 **Table 6a NCD correlated with risk factors in USA, UK, Mexico, and Japan combined**

NCD versus risk factors USA, UK, Mexico, and Japan (combined n=730 cohorts)	Mean	SD	Min	Max	r	95% CI	p
NCD	1071	343	424	2312			
BMI kg/M²	25.18	1.92	21.37	27.73	0.504	0.447 to 0.556	<.0001
FPG mmol/L	4.78	0.24	4.03	5.17	-0.214	-0.282 to -0.143	<.0001
LDLC mmol/L	2.79	0.12	2.33	3.09	-0.106	0.177 to 0.034	0.0041
SBP mm Hg	131.7	4.20	126.8	141.6	-0.258	0.325 to 0.189	<.0001
Processed meat kcal/day	28.27	14.81	3.98	68.77	0.576	0.525 to 0.622	<.0001
Red meat kcal/day	104.1	44.09	41.43	207.8	0.715	0.677 to 0.748	<.0001
Fish kcal/day	66.79	103.3	5.89	370.4	-0.479	0.533 to 0.421	<.0001
Milk kcal/day	66.62	28.54	25.35	103.9	0.451	0.391 to 0.507	<.0001
Poultry kcal/day	136.4	59.04	43.47	190.9	0.402	0.339 to 0.460	<.0001
Eggs kcal/day	42.73	6.86	27.43	69.64	-0.363	0.424 to 0.298	<.0001
Alcohol kcal/day	172.6	69.05	4.25	296.7	0.399	0.336 to 0.458	<.0001
Sugary beverages kcal/day	280.7	302.4	72.91	1472	0.160	0.089 to 0.230	<.0001
Corn kcal/day	63.53	86.99	10.34	300.7	0.043	0.030 to 0.115	0.2477
potatoes kcal/day	79.60	36.62	18.09	147.4	0.360	0.295 to 0.421	<.0001
SFA kcal/day	269.4	76.29	129.4	464.2	0.409	0.347 to 0.468	<.0001
PUFA kcal/day	212.4	91.12	88.94	334.7	0.363	0.298 to 0.425	<.0001
TFA kcal/day	38.34	20.36	6.49	76.54	0.257	0.187 to 0.323	<.0001
Fruits kcal/day	62.54	12.78	31.67	86.50	0.132	0.060 to 0.203	0.0003
Vegetables kcal/day	113.1	37.27	49.16	224.9	-0.214	0.282 to 0.144	<.0001
Nuts and seeds kcal/day	18.06	9.02	3.34	38.04	0.376	0.312 to 0.437	<.0001
Whole grains kcal/day	73.64	35.54	31.01	235.1	-0.046	0.118 to 0.027	0.2184
Legumes kcal/day	49.56	20.88	14.93	86.31	-0.244	0.311 to 0.174	<.0001
Rice kcal/day	39.98	44.60	11.73	144.4	-0.525	0.575 to 0.470	<.0001
Sweet potatoes kcal/day	4.58	4.13	0.52	12.12	-0.489	0.542 to 0.421	<.0001
Kcal /day available	3133	324	2456	3463	0.513	0.457 to 0.564	<.0001
Vitamin A deficient children/100k/yr	9705	6865	4319	28421	-0.045	0.117 to 0.028	0.2246
Sodium g/day	4.11	1.18	2.34	6.70	-0.390	0.450 to 0.326	<.0001
Calcium g/day	0.59	0.16	0.30	0.86	0.487	0.429 to 0.540	<.0001
Dietary fiber g/day	11.97	2.31	7.87	17.97	-0.035	0.107 to 0.038	0.3447
Animal foods kcal /day	444.9	115.2	159.8	701.4	0.214	0.144 to 0.282	<.0001
Plant foods kcal/day	504.6	94.57	321.3	758.6	-0.192	0.261 to 0.121	<.0001
SFA+PUFA+TFA kcal/day	520.1	172.3	257.3	828.0	0.404	0.341 to 0.463	<.0001

537
 538
 539
 540
 541
 542

543 **Table 6b NCD trend correlated with risk factor trends in USA, UK, Mexico, and Japan**

NCD trend versus risk factor trends: USA, UK, Mexico, and Japan (n combined=730 cohorts)	Mean	SD	Min	Max	r	95% CI	p
NCD trend	-13.87	14.95	-67.77	30.64			
BMI kg/M² trend	0.069	0.026	0.007	0.116	-0.105	0.176 to 0.032	0.0047
FPG mmol/L trend	0.008	0.006	-0.011	0.034	0.489	0.431 to 0.542	<.0001
LDLC mmol/L trend	-0.008	0.011	-0.022	0.009	0.512	0.456 to 0.564	<.0001
SBP mm Hg trend	-0.140	0.127	-0.475	0.250	0.209	0.138 to 0.277	<.0001
Processed meat kcal/day trend	0.084	0.369	-0.812	0.684	0.380	0.316 to 0.440	<.0001
Red meat kcal/day trend	-0.300	0.943	-2.94	2.66	0.493	0.436 to 0.546	<.0001
Fish kcal/day trend	1.529	2.795	-0.028	9.790	0.371	0.306 to 0.432	<.0001
Milk kcal/day trend	0.220	0.280	-0.438	1.012	-0.242	0.309 to 0.173	<.0001
Poultry kcal/day trend	2.003	0.783	0.802	4.592	0.113	0.040 to 0.184	0.0023
Eggs kcal/day trend	0.105	0.339	-0.143	0.955	0.483	0.426 to 0.537	<.0001
Alcohol kcal/day trend	0.100	0.244	-0.114	1.993	-0.411	0.469 to 0.369	<.0001
Sugary beverages kcal/day trend	0.878	2.88	-5.40	14.00	-0.023	0.096 to 0.049	0.5313
Corn kcal/day available trend	-0.134	0.303	-1.48	0.539	-0.413	0.472 to 0.351	<.0001
Potatoes kcal/day available trend	-0.073	0.170	-0.226	0.409	0.218	0.147 to 0.286	<.0001
Saturated fatty acids trend	0.886	0.425	-0.585	1.453	0.642	0.597 to 0.682	<.0001
Poly unsaturated fatty acids trend	1.217	0.947	-0.906	3.214	-0.342	0.404 to 0.276	<.0001
Trans fatty acids trend	-1.031	0.762	-2.68	-0.023	-0.209	0.277 to 0.138	<.0001
Fruits kcal/day trend	-0.138	0.383	-1.43	1.06	0.239	0.169 to 0.306	<.0001
Vegetables kcal/day trend	0.175	1.250	-2.83	6.61	0.146	0.074 to 0.216	<.0001
Nuts and seeds kcal/day trend	0.442	0.287	0.003	1.23	-0.450	0.506 to 0.390	<.0001
Whole grains kcal/day trend	0.306	1.433	-1.07	8.46	0.195	0.124 to 0.264	<.0001
Legumes kcal/day trend	-0.251	0.208	-0.706	0.156	-0.644	0.684 to 0.599	<.0001
Rice kcal/day available trend	-0.124	0.423	-0.983	0.133	-0.383	0.443 to 0.319	<.0001
Sweet potatoes kcal/day available trend	-0.002	0.030	-0.057	0.022	-0.357	0.419 to 0.219	<.0001
Kcal available/day trend	7.68	6.23	-5.49	13.19	-0.428	0.485 to 0.367	<.0001
Vitamin A deficient children/100k/yr trend	-40.52	69.62	-266.0	80.03	-0.451	0.506 to 0.390	<.0001
Sodium g/day trend	0.008	0.044	-0.106	0.051	-0.404	0.463 to 0.341	<.0001
Calcium g/day trend	0.001	0.002	-0.002	0.007	0.088	0.015 to 0.159	0.0176
Dietary fiber g/day trend	0.034	0.052	-0.143	0.104	-0.336	0.398 to 0.269	<.0001

544

545

546 **Supplementary Table 1. Definitions of IHME GBD risk factors and covariates related to**

547 **BMI**

Variables	Definition
Alcohol	Any alcohol consumption (g/day)
Body-mass index	Body mass index (BMI) (kg/m ²)—the dependent variable of interest
Corn	Corn availability percapita (g/day), a covariate
Eggs	Eggs availability percapita (g/day) a covariate
Fasting plasma glucose	Fasting plasma glucose (mmol/L)
Fish	This variable expressed in g/day is derived by determining the weight of fish in g corresponding to 1 g of omega-3 fatty acids (eicosapentaenoic acid and docosahexaenoic acid) by averaging the fish g per 1 g of omega-3 fatty acids 20 species of fish= 117.04 g/day fish/1 g/day omega-3 fatty acids (Supplementary Table 2)
Fruits	Consumption of fruits (includes fresh, frozen, cooked, canned, or dried fruit but excludes fruit juices and salted or pickled fruits) (g/day)
Kilocalories available /day	The mean number of kilocalories percapita available per day to people in each location (kcal/day available), a covariate
LDL cholesterol	Serum low-density lipoprotein cholesterol (mmol/L)
Legumes	Consumption of beans, lentils, pulses (g/day)
Milk	Consumption of milk including non-fat, low-fat, and full-fat milk but excluding soy milk and other plant derivatives (g/day)
Nuts and seeds	Consumption of nuts and seeds (g/day)
Poultry	Poultry availability percapita (g/day), a covariate
Potatoes	Potatoes availability percapita (g/day), a covariate
Processed meat	Consumption of any processed meat (includes meat preserved by smoking, curing, salting, or addition of chemical preservatives, including bacon, salami, sausages, or deli or luncheon meats like ham, turkey, and pastrami) (g/day)
Red meat	Consumption of red meat (includes beef, pork, lamb, and goat but excludes poultry, fish, eggs, and all processed meats) (g/day)
Rice	Rice availability percapita (g/day), a covariate
Seafood omega-3 fatty acids	Seafood omega-3 fatty acids (eicosapentaenoic acid and docosahexaenoic acid) in tablet or fish form (g/day)
Sugar-sweetened beverages	Consumption of any beverage with ≥50 calories of sugar per one-cup serving, including carbonated beverages, sodas, energy drinks, fruit drinks but excluding 100% fruit and vegetable juices (g/day)
Sweet potatoes	Sweet potato availability percapita (g/day), a covariate
Systolic blood pressure	Systolic blood pressure (mm Hg)
Vegetables	Consumption of frozen, cooked, canned, or dried vegetables (excluding legumes salted or pickled, juices, nuts and seeds, and starchy vegetables such as potatoes or corn) (g/day)
Whole grains	Consumption of whole grains (bran, germ, and endosperm in their natural proportions) from breakfast cereals, bread, rice, pasta, biscuits, muffins, tortillas, pancakes, and others (g/day)

548

549

550 **Supplementary Table 2. Omega-3 Fatty Acid g to fish g calculation¶¶**

Fish	DHA g/3 ounce fish	EPA g/3 ounce fish	Omega-3 Fatty Acids (DHA + EPA) g/3 ounce fish mean	Fish 3 ounces = 85.02 g	Fish (g) per omega-3 Fatty Acids (g)=columns E/F
Salmon Atlantic farmed	1.24	0.59			
Salmon Atlantic wild	1.22	0.35			
Herring Atlantic	0.94	0.77			
Sardines canned in tomato sauce drained	0.74	0.45			
Mackerel Atlantic	0.59	0.43			
Salmon pink canned drained	0.63	0.28			
Trout rainbow wild	0.44	0.40			
Oysters eastern wild	0.23	0.30			
Sea bass	0.47	0.18			
Shrimp	0.12	0.12			
Lobster	0.07	0.10			
Tuna light canned in water drained	0.17	0.02			
Tilapia	0.11				
Scallops	0.09	0.06			
Cod Pacific	0.1	0.04			
Tuna yellowfin	0.09	0.01			
Mean DHA and EPA Omega-3 Fatty Acids g/3 ounce fish	0.4531	0.2733			
Calculations total Omega-3 FA g to fish g			0.7264	85.02	117.043

551 ¶¶ Data on omega-3 fatty acid content of varieties of fish came from the National Institutes of
 552 Health Office of Dietary Supplements (USA)

553

554

555

556 **Supplementary Table 3. Calculations of kcal/day from g/day of animal and plant foods¶**

Foods	Food sub-categories	kcal/serving	g/serving	kcal/g
Milk (2% fat)		122	244	0.5
Fish		218	170	1.28
Eggs		72	50	1.44
Poultry		187	85	2.91
Red meat		247	85	2.91
Processed meat				
	Salami	222	59	3.76
	Pastrami	104	71	1.46
	Ring baloney	86	28	3.07
	Pepperoni	94	100	0.94
Average processed meat		126.5	64.5	1.96
Fruits		97	162	0.60
Vegetables		59	91	0.65
Legumes		249	179	1.39
Nuts		172	28	6.14
Seeds				
	Flax seeds	55	10	5.5
	Chia seeds	58	12	4.83
	Fennel seeds	34.5	10	3.45
	Hemp seeds	55.3	10	5.53
Average of seeds		50.7	10.5	4.83
Average of nuts and seeds		111.4	19.25	5.78
Corn		99	103	0.96
Potatoes		161	173	0.93
Sweet potatoes		115	151	0.76
Rice		205	158	1.3
Whole grains		120	52	2.31

557 ¶ Source: NutritionIX app

558

559

560 **Acknowledgments**

561• **Authors' contributions:** DKC acts as guarantor; conceived and designed the study, acquired
562 and analysed the data, interpreted the study findings, drafted the manuscript, critically reviewed
563 and edited the manuscript and tables, and approved the final version for publication.

564

565 CW designed software programs in R to format and population weight the data,
566 aided with the SAS statistical analysis, critically reviewed the manuscript, and
567 approved the final version for publication.

568

569 The corresponding author attests that all listed authors meet authorship criteria and that no others
570 meeting the criteria have been omitted.

571

572 **Copyright:** The authors retain the copyright to the paper

573

574 The Patient and Public Involvement

575 • When and how were patients/public first involved in the research?

576 IHME acquired, catalogued and extracted information from over 12,000 surveys from
577 government and non government agencies in order to compile the GBD database.

578 • How were the research question(s) developed and informed by their priorities,
579 experience, and preferences?

580 The diverse surveys contributing to the GBD database had many different priorities and

581 intentions unrelated to this post hoc analysis. The current analysis research questions have been

582 developed to give researchers, policymakers, and the public the methodological tool to quantify
583 the impacts of dietary and other risk factor patterns.

584 ● How were patients/public involved in

585 (a) the design and conduct of the study?

586 The way patients/public were involved in the collection of the surveys has not been
587 systematically studied and reported on. Patients/public are the intended beneficiaries of
588 this analysis of the GBD data.

589 (b) choice of outcome measures?

590 The outcome measures chosen all relate to human health.

591 (c) recruitment to the study?

592 The recruitment methods vary by the survey.

593 ● How were (or will) patients/ public be involved in choosing the methods and agreeing
594 plans for dissemination of the study results to participants and linked communities?

595 IHME and all the health surveyors that contributed will make these decisions. This analysis of
596 the data will be available to all by open access.

597

598● **Competing interests statement:** Both authors have completed the ICMJE uniform disclosure
599 form at www.icmje.org/coi_disclosure.pdf and declare: no support from any organisation for the
600 submitted work; no financial relationships with any organisations that might have an interest in
601 the submitted work in the previous three years; no other relationships or activities that could
602 appear to have influenced the submitted work.

603

604 **Contributors:** Martin Sebera, from the Department of Kinesiology, Faculty of Sports Studies

605 Masaryk University, Czech Republic, critiqued statistical aspects of the manuscript and provided
606 useful input. Pavel Grasgruber, from Masaryk University, Czech Republic, provided suggestions
607 after reviewing the manuscript.

608

609 **Transparency declaration:** The manuscript is an honest, accurate, and transparent account of
610 the study being reported; that no important aspects of the study have been omitted; and that any
611 discrepancies from the study as planned have been explained.

612

613 **Ethics:** Studies based solely on data from IHME GBD database do not need approval from any
614 bioethics committee.

615

616• **Funding:** This research received no specific grant from any funding agency in the public,
617 commercial or not-for-profit sectors. The Bill and Melinda Gates Foundation funded the
618 acquisition of the data for this analysis by the IHME. The data were provided to the authors as
619 volunteer collaborators with IHME.

620•

621
622 **Details of the role of the study sponsors:** While IHME GBD faculty and staff by virtue of
623 Gates Foundation grants provided the raw data for this analysis, they did not vet the analysis or
624 sponsor the manuscript.

625

626 **Statement of independence of researchers from funders:** The researchers have received no
627 funding. Gates Foundation funded IHME to collect and analyse the GBD data.

628 **Dissemination declaration:** Dissemination of this manuscript to the participants of the more
629 than 12,000 surveys is not possible individually, but the information will become in the public
630 domain.

631
632 **Disclosures:** We thank Scott Glenn and Brent Bell from IHME who supplied us with the GBD
633 risk factor exposure data for the risk factors and for noncommunicable disease death data.

634•
635 **Ethics:** Studies were based solely on data from the IHME GBD database and do not need
636 approval from any bioethics committee.

637
638 **Participant informed consents:** Not applicable.

639
640 **Author access to data:** As volunteer collaborators with the Institute of Health Metrics and
641 Evaluation, we received about 1.4 gigabytes of raw data on noncommunicable disease deaths and
642 32 relevant risk factors.

643
644 **Data sharing statement:** The raw, unformatted data used in this analysis is now out of date. The
645 2019 GBD data on all the variables in this analysis may be obtained from the IHME by volunteer
646 collaborating researchers. Our data formatting software code in R and SAS and our formatted
647 database are available on request to researchers.

648
649 **Protocol, submitted as a supplementary file:** Not applicable.

650

651 **STROBE checklist:** Submitted.

652

653 **Patient consent:** Not applicable.

654

655

656 **References**

- 657 1. Murray CJL, Aravkin AY, Zheng P, et al. **Global burden of 87 risk factors in 204**
658 **countries and territories, 1990-2019: a systematic analysis for the Global Burden of**
659 **Disease Study 2019.** *The Lancet.* 2020; **396** (10258): 1223-1249.
660 [https://www.thelancet.com/journals/lancet/article/PIIS0140-6736\(20\)30752-2/fulltext](https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(20)30752-2/fulltext)
- 661 2. Global Burden of Disease Study 2017 (GBD 2017) Data Input Sources Tool. Seattle,
662 Washington: Institute of Health Metrics and Evaluation Accessed August 10, 2019.
663 <http://ghdx.healthdata.org/gbd-2017/data-input-sources>
- 664 3. **GBD 2017 Risk Factor Collaborators. Global, regional, and national comparative**
665 **risk assessment of 84 behavioural, environmental and occupational, and metabolic**
666 **risks or clusters of risks for 195 countries and territories, 1990–2017: a systematic**
667 **analysis for the Global Burden of Disease Study 2017: Table of risk factor**
668 **definitions.** *The Lancet.* 2018; **392** (10159): 1923-94.
669 [https://www.thelancet.com/action/showFullTableHTML?isHtml=true&tableId=tbl1&pii=](https://www.thelancet.com/action/showFullTableHTML?isHtml=true&tableId=tbl1&pii=S0140-6736%2818%2932225-6)
670 [S0140-6736%2818%2932225-6](https://www.thelancet.com/action/showFullTableHTML?isHtml=true&tableId=tbl1&pii=S0140-6736%2818%2932225-6)
- 671 4. PROTOCOL FOR THE GLOBAL BURDEN OF DISEASES, INJURIES, AND RISK
672 FACTORS STUDY (GBD) Version 3.0; Issue 26. Seattle, Washington: Institute for
673 Health Metrics and Evaluation Accessed August 2, 2019.
674 http://www.healthdata.org/sites/default/files/files/Projects/GBD/GBD_Protocol.pdf
- 675 5. Naghavi M, Abajobir AA, Abbafati C, et al. **Global, regional, and national age-sex**
676 **specific mortality for 264 causes of death, 1980–2016: a systematic analysis for the**
677 **Global Burden of Disease Study 2016.** *The Lancet.* 2017; **390** (10100): 1151-1210.
678 [https://doi.org/10.1016/S0140-6736\(17\)32152-9](https://doi.org/10.1016/S0140-6736(17)32152-9)

- 679 6. Flaxman AD, Lee YY, Vos T, et al. An Integrative Metaregression Framework for
680 Descriptive Epidemiology. Seattle, WA: University of Washington Press, 2015.
- 681 7. GBD 2017 Risk Factor Collaborators. **Global, regional, and national comparative risk
682 assessment of 84 behavioural, environmental and occupational, and metabolic risks
683 or clusters of risks for 195 countries and territories, 1990–2017: a systematic
684 analysis for the Global Burden of Disease Study 2017.** . *The Lancet*. 2018; **392**
685 (10159): 1923-94. [https://doi.org/10.1016/S0140-6736\(18\)32225-6](https://doi.org/10.1016/S0140-6736(18)32225-6)
- 686 8. Call for Collaborators. . Seattle, Washington: Institute for Health Metrics and Evaluation
687 at the University of Washington 2019; Accessed April 25, 2019.
688 <http://www.healthdata.org/gbd/call-for-collaborators>
- 689 9. Stevens GA, Alkema L, Black PRE, et al. **Guidelines for Accurate and Transparent
690 Health Estimates Reporting: the GATHER statement.** *The Lancet*. 2016; **388**
691 (10062): e19-e23. [https://doi.org/10.1016/S0140-6736\(16\)30388-9](https://doi.org/10.1016/S0140-6736(16)30388-9)
- 692 10. Omega 3 Fatty Acids: Fact Sheet for Health Professionals. The Office of Dietary
693 Supplements 2018; Accessed September 1, 2018.
694 <https://ods.od.nih.gov/factsheets/Omega3FattyAcids-HealthProfessional/>
- 695 11. Nutritionix Track App. Syndigo LLC Accessed April 25, 2019.
696 <https://www.nutritionix.com/>
- 697 12. Mexico: profiled by the IHME GBD project. Seattle, Washington: Institute of Health
698 Metrics and Evaluation, University of Washington Accessed November 25, 2020.
699 <http://www.healthdata.org/mexico>
- 700 13. Flegal KM, Kit BK, Orpana H, Graubard BI. **Association of All-Cause Mortality With
701 Overweight and Obesity Using Standard Body Mass Index Categories: A Systematic**

- 702 **Review and Meta-analysis.** *JAMA*. 2013; **309** (1): 71-82.
- 703 <https://doi.org/10.1001/jama.2012.113905>
- 704 14. Aune D, Sen A, Prasad M, et al. **BMI and all cause mortality: systematic review and**
- 705 **non-linear dose-response meta-analysis of 230 cohort studies with 3.74 million**
- 706 **deaths among 30.3 million participants.** *BMJ*. 2016; **353**: i2156.
- 707 <https://www.bmj.com/content/bmj/353/bmj.i2156.full.pdf>
- 708 15. Jayedi A, Rashidy-Pour A, Parohan M, Zargar MS, Shab-Bidar S. **Dietary and**
- 709 **circulating vitamin C, vitamin E, β -carotene and risk of total cardiovascular**
- 710 **mortality: a systematic review and dose–response meta-analysis of prospective**
- 711 **observational studies.** *Public Health Nutrition*. 2019; **22** (10): 1872-1887.
- 712 [https://www.cambridge.org/core/article/dietary-and-circulating-vitamin-c-vitamin-e-](https://www.cambridge.org/core/article/dietary-and-circulating-vitamin-c-vitamin-e-carotene-and-risk-of-total-cardiovascular-mortality-a-systematic-review-and-doseresponse-metaanalysis-of-prospective-observational-studies/71D0B7C4B76F03C3E8FB00A4925551F0)
- 713 [carotene-and-risk-of-total-cardiovascular-mortality-a-systematic-review-and-](https://www.cambridge.org/core/article/dietary-and-circulating-vitamin-c-vitamin-e-carotene-and-risk-of-total-cardiovascular-mortality-a-systematic-review-and-doseresponse-metaanalysis-of-prospective-observational-studies/71D0B7C4B76F03C3E8FB00A4925551F0)
- 714 [doseresponse-metaanalysis-of-prospective-observational-](https://www.cambridge.org/core/article/dietary-and-circulating-vitamin-c-vitamin-e-carotene-and-risk-of-total-cardiovascular-mortality-a-systematic-review-and-doseresponse-metaanalysis-of-prospective-observational-studies/71D0B7C4B76F03C3E8FB00A4925551F0)
- 715 [studies/71D0B7C4B76F03C3E8FB00A4925551F0](https://www.cambridge.org/core/article/dietary-and-circulating-vitamin-c-vitamin-e-carotene-and-risk-of-total-cardiovascular-mortality-a-systematic-review-and-doseresponse-metaanalysis-of-prospective-observational-studies/71D0B7C4B76F03C3E8FB00A4925551F0)
- 716 16. Wang L, Song Y, Manson JE, et al. **Circulating 25-Hydroxy-Vitamin D and Risk of**
- 717 **Cardiovascular Disease: A Meta-Analysis of Prospective Studies.** *Circulation:*
- 718 *Cardiovascular Quality and Outcomes*. 2012; **5** (6): 819-829.
- 719 <http://circoutcomes.ahajournals.org/content/5/6/819.abstract>
- 720 17. Beulens JWJ, Booth SL, van den Heuvel EGHM, Stoecklin E, Baka A, Vermeer C. **The**
- 721 **role of menaquinones (vitamin K2) in human health.** *British Journal of Nutrition*.
- 722 2013; **110**: 1357–1368. [https://www.cambridge.org/core/journals/british-journal-of-](https://www.cambridge.org/core/journals/british-journal-of-nutrition/article/the-role-of-menaquinones-vitamin-k2-in-human-health/5B9F317B526629D8BA77B6435F1E5509)
- 723 [nutrition/article/the-role-of-menaquinones-vitamin-k2-in-human-](https://www.cambridge.org/core/journals/british-journal-of-nutrition/article/the-role-of-menaquinones-vitamin-k2-in-human-health/5B9F317B526629D8BA77B6435F1E5509)
- 724 [health/5B9F317B526629D8BA77B6435F1E5509](https://www.cambridge.org/core/journals/british-journal-of-nutrition/article/the-role-of-menaquinones-vitamin-k2-in-human-health/5B9F317B526629D8BA77B6435F1E5509)

- 725 18. Clemens R, Drewnowski A, Ferruzzi MG, Toner CD, Welland D. **Squeezing fact from**
726 **fiction about 100% fruit juice.** *Advances in nutrition (Bethesda, Md.)*. 2015; **6** (2):
727 236S-243S. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4352186/>
- 728 19. POTATO PROCESSING AND USES. Lima, Peru: International Potato Center
729 Accessed May 28, 2020. <https://cipotato.org/potato/potato-processing-uses/>
- 730 20. Amount of Saturated Fat in Potatoes. Diet and Fitness Today Accessed July 10, 2020.
731 <http://www.dietandfitnesstoday.com/saturated-fat-in-potatoes.php>
- 732
- 733