

1 **Title: Long-term Exposures to Air Pollutants Affect FeNO in Children: A Longitudinal**
2 **Study**

3
4 **Authors:** Yue Zhang, PhD^{1,2,3}, Sandrah P. Eckel, PhD⁴, Kiros Berhane, PhD⁵, Erika Garcia,
5 PhD⁴, Patrick Muchmore, PhD⁴, Noa Ben-Ari Molshatzki, PhD⁴, Edward B. Rappaport, MS⁴,
6 William S. Linn, MA⁴, Rima Habre, ScD⁴, Frank D. Gilliland, MD, PhD⁴

7
8 **Affiliations:**

9 ¹ Department of Internal Medicine, University of Utah, Salt Lake City, UT, USA;
10 ² Department of Family and Preventive Medicine, University of Utah, Salt Lake City, UT, USA;
11 ³ Veteran Affairs Salt Lake City Health Care System, Salt Lake City, UT, USA;
12 ⁴ Department of Preventive Medicine, University of Southern California, Los Angeles, CA, USA;
13 ⁵ Department of Biostatistics, University of Columbia, New York, NY, USA;

14
15
16 **Corresponding author contact information:**

17 Yue Zhang
18 Division of Epidemiology
19 Department of Internal Medicine
20 University of Utah
21 295 Chipeta Way
22 Salt Lake City, UT, 84018
23 E-mail: zhang.yue@hsc.utah.edu
24 Tel: 801-213-3735
25 Fax: 801-581-3623

26
27 **Funding:** National Institute of Health, Environmental Protection Agency, Hastings Foundation.
28

29 **ABSTRACT**

30 Fractional exhaled nitric oxide (FeNO) is a marker of airway inflammation shown to be
31 responsive to short-term air pollution exposures; however, effects of long-term exposures are
32 uncertain. Using longitudinal assessments of FeNO and air pollutant exposures, we aimed to
33 determine whether FeNO is a marker for chronic effects of air pollution exposures after
34 accounting for short-term exposures effects.

35
36 FeNO was assessed up to six times 2004-2012 in 3607 schoolchildren from 12 communities in
37 the Southern California Children's Health Study. Within-community long-term ambient air
38 pollution exposures (PM_{2.5}, PM₁₀, NO₂, O₃) were represented by differences between
39 community-specific annual averages and the eight-year average spanning the study period.
40 Linear mixed-effect models estimated within-participant associations of annual average air
41 pollution with current FeNO, controlling for previous FeNO, prior seven-day average pollution,
42 potential confounders, and community-level random intercepts. We considered effect
43 modification by sex, ethnicity, asthma, and allergy at baseline.

44
45 We found FeNO was positively associated with annual average air pollution, after accounting for
46 short-term exposures. One standard deviation higher annual PM_{2.5} and NO₂ exposures (PM_{2.5}:2.0
47 µg/m³; NO₂:2.7 ppb) were associated, respectively, with 4.6% (95%CI:2.3%-6.8%) and 6.5%
48 (95%CI:4.1%-8.9%) higher FeNO. These associations were larger among females. We found
49 little evidence supporting association with PM₁₀ or O₃.

50
51 Annual average PM_{2.5} and NO₂ levels were associated with FeNO in schoolchildren, adding new
52 evidence that long-term exposure affects FeNO beyond the well-documented short-term effects.
53 Longitudinal FeNO measurements may be useful as an early marker of chronic respiratory
54 effects of long-term PM_{2.5} and NO₂ exposures in children.

55

56 **Key messages**

- 57 • We show strong evidence that long-term exposures to air pollutants affect FeNO,
58 independent of the well-documented associations with short-term exposures to air
59 pollution
60 • Longitudinal FeNO measurements may be useful as an early marker of chronic
61 respiratory effects of long-term air pollution exposures in children.

62
63 **Capsule summary**

64 Annual average PM_{2.5} and NO₂ were associated with FeNO in schoolchildren, adding new
65 evidence that long-term exposure affects FeNO beyond the well-documented short-term effects.
66

67 **Key words:**

68 FeNO, pollution, particulate matter, nitrogen dioxide, children
69

70 **Abbreviations**

71 ATS: American Thoracic Society

72 BMI: body mass index

73 CHS: Southern California Children's Health study

74 EC: elemental carbon

75 FeNO: Fractional exhaled nitric oxide

76 NO: Nitric oxide

77 NO₂: Nitrogen dioxide

78 O₃: Ozone

79 PM_{2.5}: particulate matter with an aerodynamic diameter of 2.5 or less

80 PM₁₀: particulate matter with an aerodynamic diameter of 10 µm or less

81 SHS: secondhand tobacco smoke
82

83 INTRODUCTION

84 Breathing polluted air has a multitude of impacts on human health, many of which are thought to
85 be driven by inflammatory mechanisms. The primary route of exposure is inhalation, thus the
86 respiratory system is a major target organ for toxic effects as it receives the largest dose. The
87 fractional concentration of exhaled nitric oxide (FeNO) is a marker of airway inflammation
88 which has been shown to be useful for the assessment of acute respiratory effects of short-term
89 exposure (hours to weeks) to ambient air pollutants,¹⁻³ including particulate matter with an
90 aerodynamic diameter of 2.5 or 10 μm or less (PM_{2.5} and PM₁₀, respectively), black smoke,
91 elemental carbon (EC), nitrogen dioxide (NO₂), and ultrafine particles. In the Southern
92 California Children's Health study (CHS), we have reported positive cross-sectional associations
93 of short-term average PM₁₀, PM_{2.5}, and ozone (O₃) exposures in the week(s) before FeNO
94 measurement with FeNO.⁴

95
96 Currently, there is limited and inconsistent evidence for effects of longer-term air pollution
97 exposures on FeNO. For example, a number of cross-sectional studies have related FeNO to
98 either location-based proxies for long-term exposures to traffic/industrial activity or long-term
99 (seasonal to annual average) ambient pollution. Some of these studies found significant positive
100 associations⁵⁻⁷ (two with adjustment for short-term exposures^{8,9}) but others did not.¹⁰⁻¹⁵ An early
101 analysis of CHS data over a short follow-up period is one of the few studies to date relating
102 longitudinally assessed FeNO to long-term ambient air pollution exposures. Using two
103 assessments of FeNO approximately one year apart in ~1200 schoolchildren, we found that
104 increased annual average PM_{2.5} or NO₂ was associated with increased within-participant FeNO,
105 after adjusting for short-term exposures (in the prior week).¹⁶

106
107 We hypothesize that FeNO is affected by both long-term and short-term exposures to ambient air
108 pollution. In this study, the estimand of interest is long-term ambient air pollution exposure
109 effect, and short-term exposure is treated as nuisance parameter. We examined the full
110 longitudinal CHS data—which includes up to six repeated FeNO assessments in more than 3000
111 schoolchildren—to investigate whether long-term (i.e., annual average) ambient air pollution
112 exposures are associated with within-participant changes in FeNO, after accounting for short-
113 term (i.e., weekly average) exposures.

115 MATERIAL AND METHODS

116 Study Population

117 Participants in this study were part of the most recent southern California Children's Health
118 Study (CHS) cohort, which started enrollment in 2002-2003 from kindergarten and first grade
119 classrooms. Detailed information on study design, participant recruitment, and data collection
120 has been reported previously.¹⁷ This analysis includes 3607 school children aged 6-17 years from
121 12 communities in Southern California, who had FeNO assessed up to 6 times in approximately
122 annual visits from 2004-2005 through 2011-2012. Ethical approval of the protocol was obtained
123 from the University of Southern California institutional review board (approval # HS-13-00150,
124 continuing review date April 17, 2019). Informed assent was obtained for each child, and
125 informed consent from a parent or guardian. The research protocol was reviewed and approved
126 by the University of Southern California Health Campus Institutional Review Board.

127 FeNO measurement

129 FeNO was measured at schools following the American Thoracic Society (ATS) guidelines using
130 an offline technique at the first two visits (2004-2005 and 2005-2006 school years) and online
131 technique in the subsequent four visits. For offline measurement, breath was collected in
132 aluminized Mylar bags using commercial breath sampling kits (Sievers Division, GE Analytical
133 Instruments, Boulder, Colorado, USA). Each child took 2-3 initial deep breaths through the kit's
134 NO scrubber followed by a near-vital-capacity breath and immediate controlled exhalation
135 through the sampling kit. The collected bag samples were measured in central laboratory for
136 FeNO level at targeted 100 ml/sec flow rate using a standard chemiluminescent analyzer system
137 (Sievers Division). For online measurement, FeNO was measured directly using EcoMedics
138 CLD-88-SP analyzers with DeNOx accessories to provide NO-free inhaled air (EcoPhysics Inc,
139 Ann Arbor, MI, USA/Duernten, Switzerland). Each child was asked to provide at least three
140 maneuvers at the target 50 ml/sec flow rate. The plateau concentration for each maneuver was
141 taken from the 3 second plateau with the minimum NO coefficient of variation. FeNO was
142 represented by the mean of two or three ATS acceptable plateaus that differed by less than 15%
143 or 1 *ppb* where all ATS acceptable plateaus were less than 10 *ppb*. Offline FeNO data were
144 converted to estimates of FeNO as would be measured online at 50 ml/sec expiratory flow. The
145 conversion was performed using a statistical model determined in the post hoc comparison study,
146 which involved 362 children measured on and off line in the same testing session. We exclude
147 observations with reported use of inhaled corticosteroids. Details of FeNO data collection,
148 quality control, and conversion procedures have been reported previously.¹⁸⁻²⁰

149 150 **Air Pollution Data**

151 Daily 24-hour average PM_{2.5}, PM₁₀, NO₂, O₃ (10am-6pm only), and temperature data were
152 obtained for the study period from central monitoring sites in each community operated by local
153 air pollution agencies in conformance with US Environmental Protection Agency (EPA)
154 requirements. For each pollutant, community-specific study period averages were computed (a
155 seven-year average, from 2004/05-2011/12). For each pollutant and each participants' FeNO test
156 date, annual average pollution was calculated based on the 12 months preceding the test date in
157 that participant's community. Weekly averages were calculated based on the 7 days preceding
158 the test date. Air pollution data was interpolated to fill in data gaps on days with missing
159 exposure using modeled predictions based on data from nearby monitors. While such processes
160 could be potential sources of bias, previous sensitivity analyses on CHS data that limited use to
161 complete data have not altered main study findings.⁴ The primary "long-term" exposure of
162 interest for this study was the annual within-community fluctuation, defined as the difference
163 between annual averages and the seven year average concentration for each community. We
164 refer to seven day average as "short-term" exposures hereafter.

165 166 **Other Covariates**

167 Race/ethnicity, medication use, current wheeze status, exposure to secondhand tobacco smoke
168 (SHS), asthma history, and allergy history were determined from annual self-administered
169 questionnaires. Questionnaires were completed by parents from baseline to year 2006-2007, and
170 by children from follow-up year 2007-2008 onwards. Questionnaires were collected at the school
171 visit during which the child's FeNO test was performed. Ambient air nitric oxide was measured
172 at the time of FeNO collection. Trained technicians measured height and weight at every FeNO
173 test date following a standardized protocol. Age- and sex-specific body mass index (BMI)

174 percentiles were calculated based on the Centers for Disease Control and Prevention BMI growth
175 charts²¹.

176

177 **Statistical Analysis**

178 FeNO had a right-skewed distribution, so we used geometric mean and geometric standard
179 deviation to quantify central tendency and dispersion and performed all analyses with natural
180 log-transformed FeNO. In descriptive data analyses, we summarized the characteristics of the
181 study participants, compared the distribution of baseline year log FeNO by these characteristics,
182 and also summarized the distribution of the community-specific seven-year pollutant averages
183 and the annual within-community fluctuations. We also calculated Pearson correlation
184 coefficients of annual within-community fluctuations between pairs of pollutants.

185

186 We estimated linear mixed effect models to examine the association of log FeNO with annual
187 within-community fluctuations in air pollution, adjusting for short-term exposures, confounders,
188 and design variables as well as log FeNO at the previous visit. More details on the model
189 specification could be found in the Supplementary appendix. Potential confounders included sex,
190 race/ethnicity, and respiratory allergy at baseline, as well as time-varying measures of asthma
191 status, medication use, current wheeze, secondhand tobacco smoke exposure, recent respiratory
192 illness, concurrent room air nitric oxide, month of FeNO test, and average temperature in the
193 week preceding the FeNO test. To assess confounding, we used the analytic change-in-estimate
194 ($\geq 10\%$) criterion. We also evaluated whether long-term exposure associations were modified by:
195 baseline asthma, baseline allergy, sex, and ethnicity. In sensitivity analyses, the final models
196 were refitted by: (a) removing the five communities which contributed to only the first four (of
197 six) FeNO visits, (b) removing one community at a time, (c) adding a community-level random
198 slope on long-term exposure to investigate potential heterogeneity by community, (d) subsetting
199 to only online FeNO collection (the last four of the six visits), or (e) accounting for the serial
200 correlation of FeNO observations from same participant by using an autoregressive within-
201 participant level random component instead of controlling for previous visit log FeNO. In all
202 models, missing data were assumed to be missing at random. All analyses were conducted using
203 the statistical package R version 3.5.1²². Hypothesis tests were performed under a 0.05
204 significance level and a two-sided alternative.

205

206 **RESULTS**

207 The 3607 children, with mean age 9 at baseline, were evenly distributed by sex (male: 49.7%),
208 primarily Hispanic (56.3%) or non-Hispanic White (33.2%), 41.5% had allergies, 15.1% had
209 asthma, and 3% had reported a recent respiratory illness (Table 1). At baseline, the geometric
210 mean FeNO was 12.1 ppb (geometric SD: 1.9 ppb) and baseline FeNO was statistically
211 significantly associated with most participant characteristics. Specifically, boys on average had
212 significantly higher FeNO than girls (p-value=0.015). FeNO differed by race/ethnicity (p-value
213 <0.001), with African American children having higher FeNO than the other race/ethnicity
214 groups. Children with respiratory conditions, such as asthma, wheeze, allergy, and recent
215 respiratory illness, had statistically significantly higher FeNO than those without (p-values all
216 <0.001). FeNO was positively associated with age and height (p-value <0.001), but not BMI (p-
217 value=0.831). FeNO varied by month of collection (p-value<0.001), with the lowest geometric
218 means in cooler months (December – March).

219

220
221
222
223

Table 1: Summary of Participant Characteristics and FeNO at Baseline (N=3607)

Variable	Category	N (%)	Geometric mean FeNO (GSD)*	p-value†
Sex	Male	1792 (49.7)	12.4 (2.0)	0.015
	Female	1815 (50.3)	11.9 (1.9)	
Race/ethnicity	Non-Hispanic White	1193 (33.2)	11.4 (1.8)	<0.001
	Hispanic	2022 (56.3)	12.4 (1.9)	
	African American	122 (3.4)	15.5 (2.2)	
	Asian	173 (4.8)	11.2 (1.9)	
	Other	82 (2.3)	14.2 (2.1)	
ETS	Yes	174 (5.2)	13.1 (2.0)	0.181
	No	3207 (94.8)	12.0 (1.9)	
Allergy	Yes	1292 (41.5)	13.6 (2.0)	<0.001
	No	1820 (58.5)	11.3 (1.8)	
Age	<=9	2443 (67.7)	11.2 (1.8)	<0.001
	>9	1164 (32.3)	14.3 (2.0)	
Height	<=132cm	1947 (54.3)	11.4 (1.8)	<0.001
	>132cm	1638 (45.7)	13.1 (2.0)	
Asthma	Yes	546 (15.1)	16.3 (2.3)	<0.001
	No	3060 (84.9)	11.5 (1.8)	
Medication Use	Yes	299 (9.4)	17.2 (2.3)	<0.001
	No	2896 (90.6)	11.6 (1.8)	
Wheeze	Yes	368 (10.9)	16.6 (2.2)	<0.001
	No	2995 (89.1)	11.6 (1.9)	
BMI	<=75%	1803 (50.3)	12.1 (1.9)	0.831
	>75%	1781 (49.7)	12.1 (1.9)	
Month	Jan	319 (8.8)	10.8 (2.0)	<0.001
	Feb	554 (15.4)	11.0 (1.8)	
	Mar	678 (18.8)	11.7 (1.9)	
	Apr	629 (17.4)	13.6 (1.9)	
	May	323 (9.0)	14.4 (1.9)	
	Jun	367 (10.2)	13.7 (1.9)	
	Oct	117 (3.2)	13.1 (1.8)	
	Nov	250 (6.9)	12.1 (1.8)	
	Dec	370 (10.3)	10.1 (2.0)	
	Room air NO	<=5	1831 (50.8)	
>5		1775 (49.2)	11.0 (1.9)	
Respiratory Illness	Yes	82 (3.0)	15.4 (2.1)	<0.001
	No	2631 (97)	11.8 (1.9)	

*: Results based on geometric means (GM) and geometric standard deviations (GSD) due to skewness of FeNO data.

†: p-values are based on t-tests for comparisons of natural log-transformed FeNO levels across categories (not

including missing category).

224
 225 Community-specific seven-year average air pollution concentrations ranged approximately two
 226 to three-fold across the 12 communities (Table 2). Most communities experienced a downward
 227 trend in annual average PM_{2.5} and NO₂ over the study period (Figure 1), which was a major
 228 source of variation in the annual within-community fluctuations for these pollutants. The
 229 standard deviation of annual within-community fluctuations was 2.0 µg/m³ for PM_{2.5}, 6.7
 230 µg/m³ for PM₁₀, 2.7 ppb for NO₂, and 2.5 ppb for O₃ (Table 2). As shown in Table 3, PM_{2.5} and
 231 PM₁₀ had the highest correlation in seven-year averages across-communities (Pearson’s R=0.81),
 232 while PM_{2.5} and NO₂ had the largest correlation in annual within-community fluctuations
 233 (Pearson’s R=0.64) and annual within-community fluctuations of O₃ were negatively correlated
 234 with the other pollutants. Log FeNO was correlated with log FeNO at the previous visit
 235 (Pearson’s R=0.65), indicating moderately strong first order autocorrelation.

236
 237

Table 2: Distribution of the 7-year Average Air Pollutant Concentrations and Annual Within-Community Fluctuation across 12 Communities

Pollutant [#]	7-year Average		Annual Within-Community Fluctuation	
	Mean (SD)	Min-Max	Mean (SD)	Min-Max
PM _{2.5} , µg/m ³	12.9 (3.8)	6.8-19.4	0.1 (2.0)	-3.8-4.8
PM ₁₀ , µg/m ³	32.0 (8.3)	22.5-52.7	2.1 (6.7)	-10.2-23.3
NO ₂ , ppb	16.7 (6.3)	6.7-25.6	-0.1 (2.7)	-7.2-5.7
O ₃ , ppb	45.1 (8.4)	30.7-56.6	-0.1 (2.5)	-6.1-7.7

[#]: All pollutants are 24 hr average, except for ozone (10am–6pm average).

238
 239
 240


Table 3: Pearson Correlation Coefficients of Annual Within-Community Fluctuation in Air Pollution (lower triangular matrix) and 7-year average air pollutant concentrations (upper triangular matrix).

Pollutant [#]	PM _{2.5}	PM ₁₀	NO ₂	O ₃
PM _{2.5}	1	0.81	0.75	0.2
PM ₁₀	0.55	1	0.29	0.31
NO ₂	0.64	0.37	1	-0.02
O ₃	-0.41	-0.55	-0.43	1

*: P-value < 0.05

[#]: All pollutants are 24 hr average, except for ozone (10am–6pm average).

241
 242



243
 244 **Figure 1: Community-specific annual average air pollution during the study period.**
 245 Note: 1=Alpine; 2=Lake Elsinore; 3=Lake Gregory; 6=Long Beach; 7=Mira Loma; 8=Riverside; 9=San Dimas;
 246 11=Santa Maria; 12=Upland; 13=Glendora; 14=Anaheim; 15=San Bernardino; 16=Santa Barbara. Study
 247 Year 3=2004-2005; Study Year 4=2005-2006; Study Year 5=2006-2007; Study Year 6=2007-2008; Study
 248 Year 7=2008-2009; Study Year 8=2009-2010; Study Year 9=2010-2011; Study Year 10=2011-2012; The dot
 249 points are the annual average air pollution level prior to the dates of FeNO measurement, and triangle

250 points are the annual average air pollution level at each study year. The curve lines are the fitted trends
 251 of annual average air pollution level across the study years for each community.
 252 Long-term exposures to PM_{2.5} and NO₂ were positively associated with within-person changes in
 253 FeNO, after adjustment for covariates and short-term exposure (Table 4). A one SD (2.0 µg/m³)
 254 increase in the annual within-community fluctuation of PM_{2.5} was associated with a 4.6% within-
 255 participant increase in FeNO (95% CI: 2.3%-6.8%), controlling for covariates including FeNO at
 256 the previous visit and seven-day average PM_{2.5}. Similarly, a one SD (2.7 ppb) increase in the
 257 annual within-community fluctuation of NO₂ was associated with a 6.5% within-participant
 258 increase in FeNO (95% CI: 4.1%-8.9%), controlling for covariates. We found no evidence for an
 259 association of long-term PM₁₀ and FeNO. Long-term O₃ was negatively associated with FeNO in
 260 the primary analysis, but this association was attenuated and no longer statistically significant in
 261 sensitivity analyses (Tables 1S and 2S). All models were adjusted for sex, race, respiratory
 262 allergy, asthma, medication use, wheeze, secondhand tobacco smoking, recent respiratory illness,
 263 room air nitric oxide, month, temperature, and short-term exposure of air pollution.
 264

Table 4: Adjusted* percent difference in FeNO associated with a one standard deviation (SD)[†] increase in long-term air pollution exposures[‡]

Pollutant	Percent difference in FeNO (95% CI)
PM _{2.5}	4.55 (2.33, 6.82)
PM ₁₀	0.63 (-1.88, 3.21)
NO ₂	6.46 (4.08, 8.9)
O ₃	-2.62 (-4.49, -0.71)

* Models adjusted for: sex, race, respiratory allergy, asthma, medication use, wheeze, secondhand tobacco smoking, recent respiratory illness, room air nitric oxide, month, temperature, prior 7 day average of the same pollutant, and natural log-transformed FeNO at the previous visit.

[†] Long-term exposures are annual within-community fluctuations, calculated as the difference between annual air pollutant concentration and seven-year average concentration.

[‡]SDs are: 2.0 µg/m³ for PM_{2.5}, 6.7 µg/m³ for PM₁₀, 2.7 ppb for NO₂ and 2.5 ppb for O₃.

265
 266
 267 There was evidence that the associations of long-term PM_{2.5} and NO₂ with FeNO varied by sex,
 268 with larger estimated associations in females (interaction p-values: 0.007 for PM_{2.5} and 0.066 for
 269 NO₂) (Table 5). In sensitivity analyses (Supplementary Table 1), we still observed statistically
 270 significant positive associations of long-term PM_{2.5} and NO₂ with FeNO after: (a) removing the
 271 five communities followed only through four years, (b) removing one community at a time
 272 (Supplementary Table 2), and (c) taking out the adjustment for short-term air pollution, though
 273 we did observe some heterogeneity in the associations.
 274

275

Table 5: Interaction Tests* on the Effects of Annual Within-Community Fluctuation in Air Pollutants# on FeNO level

Effect Modifier		PM _{2.5} % diff/Unit (95% CI)	PM ₁₀ % diff/Unit (95% CI)	NO ₂ % diff/Unit (95% CI)	O ₃ % diff/Unit (95% CI)
Sex	Female	3.1 (1.8,4.3)	0.2 (-0.2,0.6)	2.7 (1.8,3.6)	-1.0 (-1.9,-0.2)
	Male	1.4 (0.1,2.6)	-0.1 (-0.5,0.4)	1.8 (0.8,2.9)	-1.1 (-2.0,-0.1)
Interaction P-value		0.007	0.143	0.066	0.957
Asthma	No	2.4 (1.2,3.5)	0.1 (-0.3,0.5)	2.4 (1.5,3.2)	-1.0 (-1.8,-0.2)
	Yes	1.1 (-1.1,3.4)	-0.2 (-0.9,0.5)	2.3 (0.7,3.9)	-1.1 (-2.8,0.5)
Interaction P-value		0.249	0.288	0.919	0.899
Allergy	No	2.6 (1.3,3.9)	0.1 (-0.3,0.6)	2.5 (1.5,3.5)	-1.3 (-2.2,-0.4)
	Yes	2.1 (0.7,3.6)	0.1 (-0.4,0.5)	2.1 (1.1,3.2)	-1.5 (-2.5,-0.5)
Interaction P-value		0.491	0.753	0.471	0.731
Hispanic	No	2.4 (1.0,3.7)	-0.1 (-0.5,0.4)	2.2 (1.3,3.1)	-1.4 (-2.3,-0.6)
	Yes	2.2 (1.0,3.4)	0.2 (-0.2,0.6)	2.6 (1.5,3.7)	-0.6 (-1.5,0.4)
Interaction P-value		0.799	0.232	0.420	0.083

*: Annual Within-community Fluctuation: The difference between annual air pollutant concentration and 7-year average concentration (within-communities). All models adjusted for sex, race, respiratory allergy, asthma, medication use, wheeze, secondhand tobacco smoking, recent respiratory illness, room air nitric oxide, month, temperature, and air pollution level in the prior week.

#: All pollutants are 24 hr average, except for ozone (10am–6pm average).

276

277

278 DISCUSSION

279 In a large, prospective cohort study of schoolchildren in 12 Southern California communities, we
 280 found statistically significant positive associations of FeNO—assessed longitudinally over an
 281 eight-year period—with within-community fluctuations in annual average PM_{2.5} and NO₂ after
 282 adjusting for prior week exposure, previous visit FeNO, and confounders. This provides the
 283 strong evidence that long-term exposures to air pollutants affect FeNO, independent of the well-
 284 documented associations with short-term exposures to air pollution.

285 Results of this longitudinal study (using all six FeNO visits in the CHS) are consistent with a
 286 preliminary longitudinal analysis of CHS data (using only the first two online FeNO
 287 assessments),¹⁶ which had found *within-participant* associations of long-term PM_{2.5} and NO₂
 288 (but not PM₁₀ or O₃) with FeNO while adjusting for short-term exposure. The preliminary CHS
 289 study had targeted within-participant changes in FeNO using a different but complementary
 290 statistical approach in which the change in FeNO was modeled as a function of the change in
 291 annual average air pollution.¹⁶ When studying FeNO, it is important to consider within-
 292 participant changes over time from longitudinally collected data, given the considerable
 293 unexplained across-participant heterogeneity in FeNO typically observed in cross-sectional
 294 studies (e.g., R²<0.3).²³ Indeed, in our data we observed moderately strong autocorrelation in

295 FeNO indicating relatively stable within-person FeNO. As an additional example, in 158
296 asthmatic adults, the intraclass correlation of FeNO over one year was 0.73, with low baseline
297 FeNO tending to stay low and higher baseline FeNO tending to stay high, but with more
298 variability.²⁴ To our knowledge, the only other study relating longitudinally assessed FeNO to
299 long-term air pollution exposures found positive but non-statistically-significant associations of
300 FeNO (measured twice over six months in 130 children) with seasonal black carbon levels at
301 subjects' residence (estimated by land-use regression), even after adjusting for short-term
302 exposures (2 hour, 24 hour, or 7 day average black carbon from central site monitors).²⁵ This
303 study's statistical analysis did not fully target within-participant associations since it used linear
304 mixed models with a participant-level random intercept but did not decompose time-varying
305 exposures (X_{ij}) into within- and between-person terms (i.e., $X_{ij}-\bar{X}_i$ and \bar{X}_i).²⁶
306 A number of cross-sectional studies have related FeNO to long-term (seasonal to annual average)
307 ambient pollution or location-based proxies for long-term exposures to traffic/industrial activity.
308 By design, these studies can estimate only between-participant associations of exposures with
309 FeNO and hence are subject to the limitations of this approach. Some cross-sectional studies
310 provide evidence supporting long-term exposure effects. In 241 healthy, full-term infants,
311 prenatal average NO₂ from a central monitoring site was associated with increased FeNO
312 measured during sleep at five weeks of age, even after adjusting for mean NO₂ in the 2 days
313 prior to the FeNO test or residential distance to major road.⁹ Two studies relating FeNO to
314 proximity to industrial activity found FeNO to be elevated in children living closer to oil
315 refinery/petrochemical activity⁶ and in children attending schools closer to a steel mill,⁷ though
316 neither study adjusted for short-term air pollution exposures. Three studies^{5, 8, 27} found a similar
317 magnitude positive association between a proxy measure for traffic exposure (length of road in a
318 circular buffer around the home), with the strongest associations for the smallest buffers around
319 the home. While the length of road traffic exposure proxy likely represents both long- and short-
320 term exposures, one of these studies adjusted for short-term exposure (central site 24 hour
321 average ambient pollution) and found the length of road association to be robust to the
322 adjustment,⁸ suggesting a long-term exposure effect. Other cross-sectional studies have found no
323 evidence for long-term exposure effects, perhaps due to exposure metrics relying on spatial
324 variation which are not sensitive to traffic sources on smaller, local roads (e.g., land-use
325 regression predicted seasonal or annual averages,^{14, 15, 25} traffic counts,^{10, 13} or distance to major
326 road^{11, 13}) or too few locations with high/low exposure to separate the source of interest from
327 potential confounders.¹²
328 Many study communities observed declines in annual average PM_{2.5} and NO₂ during the study
329 period. These declines were a key source of the variation in our long-term exposure metric
330 (within-community annual fluctuations, calculated as the difference between annual averages
331 and the seven-year average concentration for each community). Our standard interpretation of
332 the observed positive associations as “percent *increase* in FeNO per SD *increase* in exposure”
333 (e.g., 6.5% increase in FeNO (95% CI: 4.1%-8.9%) per 2.7 ppb increase in long-term NO₂),
334 could alternatively be reformulated as a “percent *decrease* in FeNO per SD *decrease* in
335 exposure”. For example, scaled to an observed declines, in Glendora annual average NO₂
336 decreased by 11.3 ppb over the study period, and we estimated a 30% *decrease* in FeNO (95%
337 CI: 18.2% to 42.9%) per 11.3 ppb *decrease* in long-term NO₂). A number of studies have used
338 natural or designed experiments to demonstrate that FeNO decreased following decreasing
339 (primarily short-term) exposures. For example, following the adoption of clean fuel/technology
340 school buses, 10-30% reductions in FeNO were observed in 275 schoolchildren.²⁸ Following

341 dramatic drops in air pollution in response to anti-air pollution measures enacted for the Beijing
342 Olympics, FeNO decreased ~57% in 125 health young adults (and subsequently rose 135% after
343 the games)³ and ~27% in 36 children.²⁹ Within-participant FeNO also declined in a group of 42
344 retired teachers relocated from a higher air pollution city to a lower pollution city for 9 days³⁰
345 and in 37 untreated allergic children moved from an urban to a rural environment for 7 days,
346 with the reductions in FeNO restricted to participants with the highest FeNO at baseline in the
347 urban environment.³¹

348 Several additional results in our study merit further discussion. We found evidence for larger
349 effects of long-term PM_{2.5} and NO₂ in females as compared to males (p=0.007 and 0.07,
350 respectively), but no evidence for effect modification by asthma or allergy. In the preliminary
351 CHS analysis of long-term air pollution effects on FeNO, we had found no evidence for effect
352 modification by sex.¹⁶ Sex/gender differences have been observed previously in air pollution
353 epidemiology analyses, generally with larger effects for boys in early life and girls/women in
354 later childhood and older age.³² We also found a statistically significant negative association for
355 long-term O₃, which was not robust to our various sensitivity analyses. Within-community
356 annual average O₃ was negatively correlated with PM_{2.5}, PM₁₀, and NO₂ and unlike those
357 pollutants, there were trends of increasing O₃ during the study period. Finally, the association of
358 long-term NO₂ with FeNO was larger in magnitude when leaving out one community (Upland)
359 or including a community-level random slope. Upland had the highest annual average NO₂ at the
360 beginning of the study period, and one of the largest declines over the study period.

361 This study has several strengths, including: a large ethnically diverse population-based cohort of
362 schoolchildren; prospective longitudinal assessment of FeNO (up to 6 measurements over an
363 eight year period); a Southern California study location with substantial within- and between-
364 community variation in ambient air pollution exposures during the study period, representative
365 of the range observed nationally in the United States; as well as a study design and statistical
366 analyses approach that allowed us to target within-person associations of long-term air pollution
367 and FeNO (automatically controlling for time-constant confounders) while adjusting for short-
368 term air pollution exposures.

369 This study has several limitations. The method for FeNO assessment changed during the study
370 period, as technology improved (offline assessment used for the first two FeNO visits and online
371 assessment used for the next four FeNO visits). Although the offline values were converted to
372 predicted online values based on a published internally-developed prediction model with an $R^2 =$
373 0.94³³, there remains a possibility that this change in breath collection technique may have
374 impacted our effect estimates. In a sensitivity analysis restricting to only online FeNO visits, the
375 association of long-term NO₂ with FeNO remained similar and statistically significant (though
376 with wider confidence intervals because of smaller sample size) and the association of long-term
377 PM_{2.5} with FeNO was still positive, but attenuated and no longer statistically significant. A
378 limitation of our study design is that FeNO was assessed in only the most recent CHS cohort,
379 allowing only for investigation of associations with within-community fluctuations in annual
380 average air pollution. We were unable to investigate associations of within-community
381 fluctuations in multi-year averages. Previous high impact CHS publications have taken
382 advantage of multiple CHS cohorts (with staggered recruitment over time from the same
383 communities) to demonstrate that, in the same communities, declines in long-term (multi-year)
384 average air pollution have been associated with improved lung function growth,³⁴ reduced
385 bronchitic symptoms,³⁵ and reduced asthma incidence.³⁶ However, the most recent CHS cohort

386 was still recruited from multiple cities, with a wide range of annual average ambient air pollution
387 levels at baseline and differing trends over the course of the study period.

388
389 In conclusion, our findings provide the strongest evidence to date that long-term PM_{2.5} and NO₂
390 exposures affect within-participant FeNO, independent of the effects of short-term exposures,
391 even during a study period with declining pollution levels. Longitudinal FeNO measurements
392 may be useful as an early marker of chronic respiratory effects of long-term air pollution
393 exposures in children.

394
395 **Acknowledgements**

396 This work was supported by the Office of the Director of the National Institutes of Health
397 (UG3OD023287 and UG3OD023249); the National Institutes of Environmental Health Sciences
398 (grant # R01ES023262, R01ES027860); the Southern California Environmental Health Sciences
399 Center (grant # P30ES007048) funded by the National Institute of Environmental Health
400 Sciences; the Children's Environmental Health Center (grant #s P01ES009581, R826708-01 and
401 RD831861-01) funded by the National Institute of Environmental Health Sciences and the
402 Environmental Protection Agency; the Maternal and Developmental Risks from Environmental
403 and Social Stressors (MADRES) Center (grant #s P50ES026086, 83615801-0) funded by the
404 National Institute of Environmental Health Sciences, the National Institute for Minority Health
405 and Health Disparities and the Environmental Protection Agency and the Hastings Foundation.
406 We are indebted to the school principals, teachers, students and parents in each of the study
407 communities for their cooperation and especially to the members of the field team for their
408 efforts.

409

410 **REFERENCES**

- 411
- 412 1. La Grutta S, Ferrante G, Malizia V, Cibella F, Viegi G. Environmental effects on
413 fractional exhaled nitric oxide in allergic children. *J Allergy (Cairo)* 2012; 2012:916926.
 - 414 2. Scarpa MC, Kulkarni N, Maestrelli P. The role of non-invasive biomarkers in detecting
415 acute respiratory effects of traffic-related air pollution. *Clin Exp Allergy* 2014; 44:1100-
416 18.
 - 417 3. Huang W, Wang G, Lu SE, Kipen H, Wang Y, Hu M, et al. Inflammatory and oxidative
418 stress responses of healthy young adults to changes in air quality during the Beijing
419 Olympics. *Am J Respir Crit Care Med* 2012; 186:1150-9.
 - 420 4. Berhane K, Zhang Y, Linn WS, Rappaport EB, Bastain TM, Salam MT, et al. The effect
421 of ambient air pollution on exhaled nitric oxide in the Children's Health Study. *Eur*
422 *Respir J* 2011; 37:1029-36.
 - 423 5. Eckel SP, Berhane K, Salam MT, Rappaport EB, Linn WS, Bastain TM, et al. Residential
424 traffic-related pollution exposures and exhaled nitric oxide in the children's health study.
425 *Environ Health Perspect* 2011; 119:1472-7.
 - 426 6. Rusconi F, Catelan D, Accetta G, Peluso M, Pistelli R, Barbone F, et al. Asthma
427 symptoms, lung function, and markers of oxidative stress and inflammation in children
428 exposed to oil refinery pollution. *J Asthma* 2011; 48:84-90.
 - 429 7. Acat M, Aydemir Y, Turğut M, Çörtük M, Çakar M, Yaşar Z, et al. Comparison of the
430 Fractional Exhaled Nitric Oxide Levels in Adolescents at Three Schools Located Three
431 Different Distances from a Large Steel Mill. *Can Respir J* 2017; 2017.
 - 432 8. Dales R, Wheeler A, Mahmud M, Frescura AM, Smith-Doiron M, Nethery E, et al. The
433 influence of living near roadways on spirometry and exhaled nitric oxide in elementary
434 schoolchildren. *Environ Health Perspect* 2008; 116:1423-7.
 - 435 9. Latzin P, Röösli M, Huss A, Kuehni CE, Frey U. Air pollution during pregnancy and
436 lung function in newborns: a birth cohort study. *Eur Respir J* 2009; 33:594-603.
 - 437 10. Cakmak S, Mahmud M, Grgicak-Mannion A, Dales RE. The influence of neighborhood
438 traffic density on the respiratory health of elementary schoolchildren. *Environ Int* 2012;
439 39:128-32.
 - 440 11. Brown MS, Sarnat SE, DeMuth KA, Brown LAS, Whitlock DR, Brown SW, et al.
441 Residential proximity to a major roadway is associated with features of asthma control in
442 children. *PLoS One* 2012; 7:e37044.
 - 443 12. Spencer-Hwang R, Soret S, Ghamsary M, Rizzo N, Baum M, Juma D, et al. Gender
444 Differences in Respiratory Health of School Children Exposed to Rail Yard–Generated
445 Air Pollution: The ENRRICH Study. *J Environ Health* 2016; 78:8.
 - 446 13. Graveland H, Van Roosbroeck SA, Rensen WM, Brunekreef B, Gehring U. Air pollution
447 and exhaled nitric oxide in Dutch schoolchildren. *Occup Environ Med* 2010.
 - 448 14. Liu C, Flexeder C, Fuertes E, Cyrus J, Bauer C-P, Koletzko S, et al. Effects of air
449 pollution on exhaled nitric oxide in children: Results from the GINIplus and LISApplus
450 studies. *Int J Hyg Environ Health* 2014; 217:483-91.
 - 451 15. Yang A, Janssen NA, Brunekreef B, Cassee FR, Hoek G, Gehring U. Children's
452 respiratory health and oxidative potential of PM_{2.5}: the PIAMA birth cohort study.
453 *Occup Environ Med* 2016; 73:154-60.

- 454 16. Berhane K, Zhang Y, Salam MT, Eckel SP, Linn WS, Rappaport EB, et al. Longitudinal
455 effects of air pollution on exhaled nitric oxide: the Children's Health Study. *Occup*
456 *Environ Med* 2014; 71:507-13.
- 457 17. McConnell R, Berhane K, Yao L, Jerrett M, Lurmann F, Gilliland F, et al. Traffic,
458 susceptibility, and childhood asthma. *Environ Health Perspect* 2006; 114:766-72.
- 459 18. Linn WS, Rappaport EB, Berhane KT, Bastain TM, Salam MT, Gilliland FD. Extended
460 exhaled nitric oxide analysis in field surveys of schoolchildren: a pilot test. *Pediatr*
461 *Pulmonol* 2009; 44:1033-42.
- 462 19. Linn WS, Rappaport EB, Eckel SP, Berhane KT, Zhang Y, Salam MT, et al. Multiple-
463 Flow Exhaled Nitric Oxide, Allergy, and Asthma in a Population of Older Children.
464 *Pediatr Pulmonol* 2013.
- 465 20. Islam T, Urman R, Gauderman WJ, Milam J, Lurmann F, Shankardass K, et al. Parental
466 stress increases the detrimental effect of traffic exposure on children's lung function. *Am*
467 *J Respir Crit Care Med* 2011; 184:822-7.
- 468 21. 2009. CfDCaPCASpftCgc. A SAS program for the CDC growth charts. . 2009.
- 469 22. Team RDC. R: a language and environment for statistical computing. R Foundation for
470 Statistical Computing, Vienna, Austria. 2014.
- 471 23. Garcia-Marcos P, Soriano-Pérez M, Perez-Fernández V, Valverde-Molina J. Exhaled
472 nitric oxide in school children: Searching for the lost variability. *Allergol immunopathol*
473 2016; 44:206-13.
- 474 24. Silkoff P, Laviolette M, Singh D, FitzGerald J, Kelsen S, Backer V, et al. Longitudinal
475 stability of asthma characteristics and biomarkers from the Airways Disease Endotyping
476 for Personalized Therapeutics (ADEPT) study. *Respir Res* 2016; 17:43.
- 477 25. De Prins S, Dons E, Van Poppel M, Panis LI, Van de Mierop E, Nelen V, et al. Airway
478 oxidative stress and inflammation markers in exhaled breath from children are linked
479 with exposure to black carbon. *Environ Int* 2014; 73:440-6.
- 480 26. Enders CK, Tofighi D. Centering predictor variables in cross-sectional multilevel models:
481 a new look at an old issue. *Psychological methods* 2007; 12:121.
- 482 27. Holguin F, Flores S, Ross Z, Cortez M, Molina M, Molina L, et al. Traffic-related
483 exposures, airway function, inflammation, and respiratory symptoms in children. *Am J*
484 *Respir Crit Care Med* 2007; 176:1236-42.
- 485 28. Adar SD, D'Souza J, Sheppard L, Kaufman JD, Hallstrand TS, Davey ME, et al.
486 Adopting clean fuels and technologies on school buses. *Pollution and health impacts in*
487 *children. Am J Respir Crit Care Med* 2015; 191:1413-21.
- 488 29. Lin W, Huang W, Zhu T, Hu M, Brunekreef B, Zhang Y, et al. Acute respiratory
489 inflammation in children and black carbon in ambient air before and during the 2008
490 Beijing Olympics. *Environ Health Perspect* 2011; 119:1507.
- 491 30. Hao Y, Zhang G, Han B, Xu X, Feng N, Li Y, et al. Prospective evaluation of respiratory
492 health benefits from reduced exposure to airborne particulate matter. *Int J Environ Health*
493 *Res* 2017; 27:126-35.
- 494 31. Renzetti G, Silvestre G, D'Amario C, Bottini E, Gloria-Bottini F, Bottini N, et al. Less air
495 pollution leads to rapid reduction of airway inflammation and improved airway function
496 in asthmatic children. *Pediatrics* 2009; 123:1051-8.
- 497 32. Clougherty JE. A growing role for gender analysis in air pollution epidemiology.
498 *Environmental health perspectives* 2009; 118:167-76.

- 499 33. Linn WS, Berhane KT, Rappaport EB, Bastain TM, Avol EL, Gilliland FD. Relationships
500 of online exhaled, offline exhaled, and ambient nitric oxide in an epidemiologic survey of
501 schoolchildren. *J Expo Sci Environ Epidemiol* 2009; 19:674-81.
- 502 34. Gauderman WJ, Urman R, Avol E, Berhane K, McConnell R, Rappaport E, et al.
503 Association of improved air quality with lung development in children. *N Engl J Med*
504 2015; 372:905-13.
- 505 35. Berhane K, Chang C-C, McConnell R, Gauderman WJ, Avol E, Rappaport E, et al.
506 Association of changes in air quality with bronchitic symptoms in children in California,
507 1993-2012. *JAMA* 2016; 315:1491-501.
- 508 36. Garcia E, Berhane KT, Islam T, McConnell R, Urman R, Chen Z, et al. Association of
509 changes in air quality with incident asthma in children in California, 1993-2014. *Jama*
510 2019; 321:1906-15.