Modern contraceptive use among adolescent girls and young women in Benin: a mixed-methods study.

**AUTHORS**:

Noudéhouénou Crédo Adelphe AHISSOU<sup>1,2</sup>

Lenka BENOVA<sup>3</sup>

Thérèse DELVAUX <sup>3</sup>

Charlotte GRYSEELS <sup>3</sup>

Jean-Paul DOSSOU 2,3

Sourou GOUFODJI<sup>2</sup>

Lydie KANHONOU<sup>2</sup>

Christelle BOYI<sup>2</sup>

Armelle VIGAN<sup>2</sup>

Koen PEETERS 1,3

Miho SATO  $^1$ 

Mitsuaki MATSUI<sup>1</sup>

# AFFILIATIONS:

1 Department of Global Health, Nagasaki University School of Tropical Medicine and Global Health

Nagasaki 852-8523, Japan

- 2 Centre de Recherche en Reproduction Humaine et en Démographie Cotonou, Benin
- 3 Department of Public Health, Institute of Tropical Medicine Antwerp Antwerp, Belgium

# CORRESPONDING AUTHOR

Mitsuaki MATSUI

e-mail: mmatsui@nagasaki-u.ac.jp

c/o Nagasaki University School of Tropical Medicine and Global Health Sakamoto 1-12-4, Nagasaki 852-8523, Japan

Word count - 5102 words

# 1 ABSTRACT

## 2 Objectives

3 The study aimed to assess the determinants of modern contraceptive method use

4 among young women in Benin.

5 Design

6 A mixed-methods design.

## 7 Setting and participants

8 We used the Benin 2017-18 Demographic and Health Survey datasets for 9 quantitative analysis. Data collection was conducted using multiple-cluster sampling 10 method and through household survey. Qualitative part was conducted in the city of 11 Allada, one of the Fon cultural capitals. The participants were purposively selected.

## 12 Outcomes

13 Contraceptive prevalence rate, unmet need for modern method, and percentage of 14 demand satisfied by a modern method for currently married and sexually active 15 unmarried women were measured in the quantitative part. Access barriers and 16 utilization of modern methods were assessed in the qualitative part.

# 17 Results

Overall, 8.5% (95%CI: 7.7-9.5%) among young women ages 15 to 24 were using modern contraceptives and 13% (95%CI: 12.1-14.0%) among women ages 25 or more. Women 15-24 had a higher unmet need, and a lower demand satisfied by modern contraceptive methods compared to women aged 25 or more. 60.8% (56.9-64.7%) of all unmarried young women had unmet need for modern contraceptives. Young women were more likely to use male condoms which they

obtain mainly from for-profit outlets, pharmacies, and relatives. The factors associated with demand satisfied by a modern method were literacy, being unmarried, knowing a greater number of modern contraceptive methods, and experiencing barriers in access to health services. On the other hand, the qualitative study found that barriers to using modern methods include community norms about pre-marital sexual intercourse, perceptions about young women's fertility, spousal consent, and the use of non-modern contraceptives.

31 Conclusion

32 Contraceptive use is low among young women in Benin. The use of modern 33 contraceptives is influenced by socio-demographic factors and social norms. 34 Appropriate interventions might promote comprehensive sexuality education, 35 increase community engagement, provide youth-friendly services, and address 36 gender inequalities.

37

# 38 KEYWORDS

39 Contraception, modern method, demand satisfied, unmet need, adolescent, Benin40

# 41 Strengths and limitations of this study

- 42 This study used a nationally representative survey data which allows
- 43 generalizability at the national level.
- We used a mixed methods design which enables us to compare and discuss the
- 45 large-scale quantitative data with qualitative information from the ground.
- The quantitative information could have been affected by recall errors and
- 47 biases.
- 48

# 50 INTRODUCTION

51 Despite efforts toward making access and use of contraceptive services a basic reproductive right for all women, many countries still face high rates of unintended and 52 53 unwanted pregnancies, [1, 2]. In sub-Saharan Africa alone, about 14 million unintended (unwanted or mistimed) pregnancies occur every year, [3]; and adolescent girls and young 54 55 women 15 to 24 years old are the most vulnerable group.[4]. The reasons behind this situation include the high prevalence (about 70%) of sexually active young women with low 56 57 utilization of effective contraceptive methods (less than 10%),[4]. Additionally, unmarried sexually active adolescents are likely to have a high unmet need for contraception, which 58 59 increases their risk of unintended pregnancies, [5, 6]. In developing countries, nearly half 60 (49%) of pregnancies are unintended among adolescent girls of 15 to 19 years old, [7]. 61 Adolescence is viewed as the starting point in the continuum of care for reproductive, 62 maternal, neonatal, and child health; and is a phase when poor access and utilization of 63 contraception are likely to result in poor health outcomes across the continuum of care, [8, 64 9]. Early and unintended pregnancies result in increased risks of maternal mortality and 65 morbidity, premature births, low birth weight, unsafe abortions, and social consequences 66 such as stigmatization, school dropout, and poverty, [10, 11].

In Benin, although the use of modern contraceptives has been slowly increasing since 2006, it is still relatively low. The modern contraceptive prevalence among all women reached 12% in 2018 compared to 6% in 2006,[12]. At the same time, almost half (48%) of all adolescent girls age 15 to 19 are sexually active, and one in five girls has already had a child or is pregnant,[12]. According to the government of Benin (2017), only 5.4% of women ages 15 to 24 were using modern contraceptive methods in 2017,[13]. Recent data

showed a total fertility rate of 5.7 among all women of reproductive ages 15 to 49 years old, and the modern contraceptive prevalence rate (mCPR) was estimated at 12%,[12]. Of all pregnancies in the country, 19% were unintended,[14]; and in 2017, both the maternal mortality ratio and infant mortality rate remained high at 397 per 100,000 live births, and 30 per 1,000 live births, respectively,[6, 12].

78 Existing studies on the use of modern contraceptive methods in Benin largely reported 79 on women of reproductive age as a whole, rather than focusing on specific age 80 groups, [15-20]. MacQuarrie suggested that young women should be studied 81 separately, as they do not have the same needs for or access to contraception as 82 adult women, [21]. However, for Benin, the literature is sparse on the use of 83 reproductive health services among adolescent girls and young women. There two studies 84 which focused on young women's contraceptive use in African countries including 85 Benin, [22, 23]. However, these studies employed quantitative method and did not 86 explore in-depth factors that affect decision-making such as adolescent girls and 87 young women's perceptions, preferences, and interactions with peers and providers. 88 Moreover, little is known about the contraceptive practices and needs of sexually 89 active and inactive young women specifically.

90 Therefore, this study aimed to assess contraceptive use, perceptions, and practices 91 among adolescent girls and young women in Benin. Specifically, we used secondary 92 quantitative data to examine the rates of modern method use, unmet needs for 93 modern contraception, and to evaluate the association between various factors and 94 the demand satisfied for a modern contraceptive among sexually active young 95 women. We then used qualitative methods to further explore the socio-cultural

- 96 factors influencing the use of modern contraceptive methods among young women
- 97 of ages 18 to 24 years.
- 98
- 99 METHODS
- 100 Study design

101 The study was conducted using a sequential explanatory mixed-methods design 102 where the collection and analysis of quantitative data were followed by the 103 collection and analysis of qualitative data. The qualitative results thus aim to explain 104 and interpret the findings of the quantitative study,[24].

105

### 106 Quantitative strand

107 Data source

We used the most recent Benin Demographic and Health Survey (BDHS), conducted between November 2017 and February 2018,[12]. The BDHS2017-18 was a nationally representative cross-sectional survey conducted by the National Institute of Statistics and Economic Analysis and funded by the USAID. The BDHS 2017-18 employed a two-stage stratified cluster sampling during which data were collected with standardized questionnaires.

114 Study population

Included in the study were women ages 15 to 49 years at the time of the survey, residing in sampled households, and who agreed to respond to the BDHS 2017-18. We defined adolescent girls as those 15 to 19 years old and young women as those 20 to 24 years. In descriptive analyses, we compared indicators of contraceptive use

in each of the age groups to all women 25 years and older. In the analysis of the determinants of modern contraceptive methods use, we focused on women who reported to be sexually active (married and unmarried) because they are exposed to the risk of pregnancy, and we used the combined age groups 15 to 24 years due to sample size considerations.

124 Definitions

Some definitions of indicators from the BDHS 2017-18 were adopted, while others
were generated or recategorized as necessary for our study-specific objectives.

127 *Modern method*: Respondents were asked which method of contraception they were 128 using at the time of the survey. If they selected more than one method (for example, 129 implant and condom), only the most effective method was noted in the dataset. 130 Based on the definition of Hubacher and Trussell (2015), we considered modern contraceptive methods to include: oral contraceptive pills, intrauterine devices 131 132 (IUD), injectables, diaphragm, male/female condoms, female/male sterilization, 133 implants/Norplant, foam, or jelly, and emergency contraception,[25]. All other 134 methods were grouped as traditional methods.

135 Current modern method users: Women who reported using any modern
136 contraceptive method at the time of the survey were considered modern method
137 users.

138 Contraceptive prevalence rate: The proportion of women who reported using a 139 method at the time of the survey. This was estimated for modern contraceptive 140 methods among different groups of women based on their marital status.

Sexual activity: Women who reported sexual intercourse 30 days preceding the survey were
categorized as sexually active.

143 Source of modern method: The source of the modern method is the most recent 144 source of provision among current modern methods users. We grouped sources of 145 modern methods into public sources (government hospital, government health center, mobile clinic, government community health worker, and other public 146 147 sectors), non-public sources (private hospital/clinic, private clinic or doctor, and private mobile clinic), for-profit outlets (pharmacy, family planning clinic, and shop), 148 relatives and other sources (friends/relatives). A second categorization in the 149 150 analysis was done based on the theoretical capacity model of providers proposed by 151 Radovich et al., [22]. It divides the providers into two groups, 'comprehensive' or 152 'limited'. Comprehensive providers are public or private providers that can provide both short- and long-acting methods. Limited providers include service locations, 153 154 for-profit outlets, partners, and friends, which do not generally provide long-acting 155 methods.

156 *Unmet need for modern methods:* Unmet need for contraception describes women who are at risk of pregnancy (married or sexually active), do not wish to be pregnant, 157 158 but are not using a modern contraceptive method. We applied an algorithm 159 proposed by Bradley et al. (2012) to identify 'unmet need for limiting' and 'unmet 160 need for spacing' among married women and sexually active unmarried women, [26]. 161 Also, we adopted the definition of FP2020 (2021) to compute the 'unmet need for 162 modern methods', [4]. Because women using a traditional method are less protected 163 against pregnancy, they were included with the non-users and considered to have an

unmet need for modern contraception,[27]. Unmet need was estimated among the
denominator of all women regardless of whether they were in need of family
planning or not.

167 *Percentage of demand satisfied by a modern method:* This is the percentage of women at 168 risk of pregnancy and who do not wish to become pregnant, who are using modern 169 methods. The numerator is current modern method users, and the denominator is all 170 women in need of family planning (current users of any contraceptive methods and women 171 with unmet need for family planning).

Existence of barriers to access to health services: Women were asked whether getting permission to go, getting money to pay for services, the distance to travel, or going alone to a health center were barriers in their access to health services. We created a binary variable categorizing women who said that they had none of these issues versus those who had one or more.

177 Core indicators: The core indicators include 'contraceptive prevalence rate', 'unmet 178 need for modern method', and 'percentage of demand satisfied by a modern method' 179 for currently married and sexually active unmarried women. Demand satisfied by a 180 modern method among women ages 15 to 24 in need of family planning was the main 181 outcome variable in the logistic regression.

182 *Explanatory variables:* We included in the logistic regression the respondents' 183 socio-demographic characteristics (literacy, ethnicity, religion, marital status 184 [married/cohabiting at the time of the survey or no], parity, and household wealth 185 quintile); knowledge of modern contraceptive methods; and the existence of 186 barriers to access health services.

#### 187 Data analysis

188 Data analysis was carried out using Stata (version 13, Stata Corp., Texas, USA). We 189 conducted a descriptive analysis to describe the socio-demographic characteristics 190 of the study population within three age groups (15 to 19, 20 to 24, 25 or older). We 191 used logistic regression to conduct bivariate and multivariable analyses to assess the 192 associations between the explanatory variables and modern method use among 193 sexually active women of ages 15 to 24 years in need of contraception. We retained 194 all explanatory variables in multivariable logistic regression. We used the svyset 195 command to adjust for clustering, stratification, and the sampling weights of 196 individual women in all analyses.

197 Missing data

There was no missing data in any variables used except for the source of the modern method. Of condom users 15 to 49 years old, 26 out of 1805 (1.4%) had a missing source of method; and 19 of them were 15 to 24 years old. We grouped them within the "pharmacy" category, which is the most common source for condom users.

202

203 Qualitative strand

204 Study site and population

205 Data collection was conducted in the city of Allada, one of the Fon cultural capitals – 206 the largest ethnic group in Benin - and located in southern Benin about 50 km from 207 Cotonou, the largest city of Benin. The region offers the advantage of being 208 semi-rural with diverse religions and care practices such as spiritual care, 209 alternative care, and biomedicine,[28]. Spiritual healing involves using mystical

210 practices, advice prescribed by traditional healers which are viewed as effective in 211 treating health problems. Such practices could include consulting deities, 212 performing rituals, using leaves, or consuming herbal teas. Alternative care refers to 213 the use of herbal infusions and powders, as well as the use of certain leaves and 214 barks to address health issues.

We restricted the target population of the qualitative study to women ages 18 to 24 years old due to consent issues with minors. We collected data from two main groups: young pregnant women from whom we wanted to learn about access barriers and utilization of modern methods based on their experiences, particularly in cases where pregnancy might be unintended. They were recruited after pre or postnatal visits, or consultation about family planning services in the main public health center of Allada.

222 Healthcare providers helped us identify women that met our inclusion criteria 223 (pregnant young women). We later approached them face-to-face outside the health 224 facilities after consultations. Non-pregnant adolescent girls and young women were 225 recruited in schools through purposive sampling, and in the community using 226 convenience and snowball sampling techniques. In the community, we visited public 227 places (markets, school yards, parks) and approached young women face-to-face and 228 were selected if they met our inclusion criteria and were willing to participate in the 229 study. New potential participants were then referred by previous informants.

230 Since the use of modern contraceptive methods may also be affected by other actors 231 in decision-making and service provision, we collected data from young men ages 18 232 to 24, health care providers in public and private local health centers, relatives of

233 adolescent girls, and the leaders of a non-governmental organization (NGO) which 234 promotes contraceptive methods in the region. Those informants were all recruited 235 by purposive sampling techniques. 236 Data collection 237 Data were collected through in-depth interviews and focus group discussions with 238 young people, females, and males separately. The group discussions were conducted 239 in isolated public places (parks, youth gathering places) to ensure quietness and that 240 participants speak with ease and without fear of being seen by relatives. We 241 conducted the discussions mainly in the Fon language. The number and 242 characteristics of the participants are shown in Table 1.

#### 243 Table 1. Numbers and characteristics of participants in informant interviews and focus

244 group discussions.

	Informant interview	Focus group discussion
Pregnant women	12	-
Non-pregnant women	13	7
Young men	3	7
Relatives of adolescent girls	2	-
Service providers, NGO leaders	6	-
Total	36	14

245

Based on an adapted theoretical framework from the PASS Health Seeking Behaviour model of Hausmann-Muela et al., [29], and the first findings from the quantitative analysis, we designed interview guides specific to each target group. The question guides focused on perceptions of ideal family size, knowledge of contraceptive methods, and decision-making for contraceptive use.

251 Data analysis

All recorded sessions were transcribed into French. Those that were not conducted in French were translated. A content analysis of transcripts was done using the software MAXQDA 11. In the inductive analysis, narrative data was parsed and categorized into units of meaningful information and the categories were linked to understanding young women's decision-making for contraception.

257

258 **RESULTS** 

#### 259 Quantitative strand

Overall, 15,928 women of reproductive age (15 to 49 years old) were surveyed on the DHS; 39.4% (n=6,251) were 15 to 24 years old. The sociodemographic characteristics of the respondents by three age groups (15 to 19 years old; 20 to 24 years old; and 25 years or older) are shown in Table 2.

264	Table 2. Socio-demographic characteristics	of the quantitative	analysis sample (Benin	DHS 2017-18); N=15,	928 (women aged 15-49)
-----	--	---------------------	------------------------	---------------------	------------------------

2	6	5
_	_	_

	Age groups	15 - 19 years				20 - 24 yea	rs	2	25 years or older			
			(n=3,335)			(n=2,916)			(n=9,677)			
Variables		n	%	95%CI	n	%	95%Cl	n	%	95% CI		
Residence	Urban	1468	42.2	39.4-45.1	270	41,6	39.0-44.4	4307	42.8	40.8-44.8		
	Rural	1867	57.8	54.9-60.7	1646	58.4	55.7-61.0	5370	57.2	55.2-59.2		
Ethnicity	Fon	1227	37.4	34.4-40.4	1004	34.6	32.0-37.3	3514	37.5	35.4-39.7		
	Others	2108	62.6	59.6-65.6	1912	65.4	62.7-68.0	6163	62.5	60.3-64.6		
Religion	No religion	165	4.6	3.8-5.7	140	4.4	3.6-5.4	528	5.3	4.5-6.3		
	Traditional	235	6.9	5.8-8.3	224	7.8	6.5-9.3	1010	10.4	9.1-11.8		
	Islam	892	26.9	24.2-29.8	932	32.7	29.8-35.8	2853	29.7	27.5-32.0		
	Christianity	2018	60.8	57.9-63.6	1585	53.8	50.8-56.7	5181	53.5	51.2-55.8		
	Others	25	0.8	0.5-1.3	35	1.3	0.8-2.3	105	1.1	0.89-1.5		
Literacy	Can read	1899	57.0	54.5-59.6	1301	44.3	41.6-47.0	2290	23.0	21.5-24.6		
	Cannot read	1436	43.0	40.1-45.5	1615	55.7	53.0-58.4	7387	77.0	75.4-78.5		
Marital status	Not married	2712	81.5	79.7-83.3	972	32.7	30.4-35.2	1074	11.1	10.3-12.0		
and	Sexually inactive	2344	70.3	68.2-72.3	688	23.4	21.5-25.3	822	8.6	7.9-9.4		
sexual activity	Sexually active	368	11.2	10.0-12.6	284	9.3	8.1-10.8	252	2.5	2.2-2.9		
	Married	623	18.5	16.7-20.3	1944	67.3	64.8-69.6	8603	88.9	88.0-89.7		
Parity	0	2826	84.8	83.2-86.3	974	32.5	30.5-34.7	372	3.7	3.3-4.1		
	1	426	12.8	11.5-14.3	925	32.6	30.7-34.5	690	7.2	6.6-7.8		
	2	79	2.2	0.2-2.9	663	23.2	21.5-25.0	1212	12.4	11.7-13.2		
	3 or more	4	0.1	0.0-0.3	354	11.7	10.4-13.1	7403	76.7	75.6-77.9		
	Median [IQR]	0	[0,0]		1	[0, 2]		4	[3,6]			
Wealth	Poorest	538	15.7	13.2-17.6	464	15.0	12.9-17.3	1854	18.5	16.6-20.6		
quintile	Poorer	658	20.3	18.3-22.5	536	18.9	17.1-20.9	1782	18.6	17.3-20.1		
	Middle	631	19.8	17.8-22.0	564	19.9	17.9-22.0	1790	19.3	18.0-20.6		
	Richer	691	21.2	19.1-23.6	630	22.5	20.4-24.8	1960	21.1	19.6-22.7		
	Richest	817	23.4	20.6-26.4	722	23.8	21.2-26.6	2291	22.5	20.6-24.6		

267	Modern contraceptive use was 11.2% (95%CI: 10.6-12.0%) among all women of
268	reproductive age (15 to 49 years old) versus 8.5% (95%CI: 7.7-9.5%) among women aged 15
269	to 24, and 13.0% (95%CI: 12.1-14.0%) among 25 years old women or older.
270	Contraceptive practices and knowledge in the study population are shown in Table 3,
271	where we compared four groups: adolescents and young women (15-24 years) and
272	women 25 years or older, (separately married and unmarried sexually active). Among
273	all women, the unmet need for modern contraception was significantly higher among
274	young unmarried sexually active women (60.8%) compared to all other groups.
275	Specifically, among women in need of family planning, the percentage of demand
276	satisfied by a modern method was lowest among 15 to 24 years old married women
277	(17.3%).

# Table 3. Need for, demand for, and knowledge of modern contraception among sexually active women in Benin (n=12,074)

	Age 15 – 24 years old							25 years old or more							
	Marital status	Unmarried, sexually active				Marrie	d	Unma	rried, sexu	ally active		Married			
			(n=652	2)		(n=2567)			(n=652	:)		(n=2567	')		
Indicator		n	%	95% Cl	n	%	95% Cl	n	%	95% Cl	n	%	95% Cl	p-value	
Modern contraceptive	Total	152	23.2	19.8-26.9	202	8.1	7.0 – 9.4	58	22.5	17.6-29.2	1132	13.0	12.1 - 14.0	< 0.001	
prevalence rate	for limiting	1	0.19	0.0 - 1.3	8	0.3	0.2 - 0.7	18	7.7	4.7 – 12.4	498	5.7	5.1-6.3		
	for spacing	151	23.0	19.7-26.7	194	7.8	6.7 – 9.0	40	14.8	10.7-20.2	634	7.4	6.7 - 8.1		
Unmet need for modern	Total	393	60.8	56.9-64.7	989	38.8	36.7 – 41.0	113	44.5	38.0 - 51.0	3027	35.1	33.9 – 36.3	< 0.001	
contraception	for limiting	5	0.72	0.3 - 1.9	28	1.1	0.7 – 1.7	33	13.4	9.4 – 18.7	1186	13.6	12.7 – 14.5		
	for spacing	388	60.1	56.1-64.0	961	37.7	35.5 – 55.6	80	31.1	25.3 – 37.5	1841	21.5	20.5 – 22.5		
Total need for modern	Total	545	84.0	80.6-86.9	1191	46.9	44.4 - 49.3	171	67.0	60.2 - 73.2	4159	48.1	46.8-49.4	< 0.001	
contraception	for limiting	6	0.9	0.4 – 2.2	1155	1.5	1.0 - 2.1	51	21.1	16.2 - 27.2	1684	19.2	18.2 – 20.4		
	for spacing	539	83.1	79.5 - 86.2	1376	45.4	43.0 – 47.8	120	45.9	39.1-52.9	2475	28.9	27.8 – 30.0		
Existence of barriers to	None (0)	274	40.2	36.0-44.6	1,022	40.3	37.4 - 43.1	113	45.2	38.2-52.5	3,405	39.6	37.7-41.5	0.44	
access health services	Any (>=1)	378	59.8	55.4-64.0	1,545	59.7	56.8 - 62.5	139	54.8	47.5-61.9	5, 198	60.4	58.5 – 62.3		
Number of modern	Fewer than four	87	13.8	10.9 - 17.2	470	18.0	15.9 – 20.3	13	5.8	3.2-10.2	958	10.9	9.8 - 12.1	< 0.001	
methods known by	Four to six	316	50.0	45.2 - 54.8	1,208	46.9	44.3 – 49.5	92	35.8	29.4-42.9	3,674	43.6	41.7–45.4		
women	More than six	249	36.2	31.8-40.9	889	35.0	32.6 – 37.6	147	58.4	51.4-65.2	3, 97 1	45.6	43.6 - 47.5		
	Median [IQR]	6 [4, 7]			5 [4, 7]			7 [5, 8]			6 [5, 8]				
Percentage of demand sat method (as of % of wome	tisfied by a modern n in need)		27.6	23.8-31.7		17.3	15.0 - 19.8		33.6	26.1-42.1		27.1	25.4 – 28.9	<0.001	

medRxiv preprint doi: https://doi.org/10.1101/2021.06.07.21257758; this version posted October 17, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

281	Table 4 shows the method mix among users of modern methods. Among unmarried
282	sexually active women ages 15 to 24 years, 63.4% of modern contraceptive users
283	were using short-term methods, and more than 60% of these being male condom
284	users. The use of modern contraceptives was lower among married young women
285	(27.1%), sexually active unmarried women ages 25 or above (25.7%), and among
286	married women 25 years old or more (19.3%). Although there was no substantial
287	difference in long-term method use among married women (72.9% vs 80.7%) of 15 to
288	24 years old and 25 or older, the use of implants and IUDs was lower among young
289	women (49.4% vs 61.0%). Regarding the source of contraceptives, unmarried
290	sexually active young women obtained their contraceptives from for-profit outlets or
291	relatives (62.1%) or comprehensive providers (36.2%). The public sector was the
292	main source of methods among young and adult married women (77.0% and 78.9%).

293	Table 4: Modern contraceptive practices among unmarried and married sexually active modern contraceptive users, by age group (n=1544)
294	

	Age	15 – 24 years old						25 years old or more							
	Marital status and	Unmarried, sexually active				Married			Unmar	ried, sexua	Illy active		Married		
	Sexual activity		(n=15	2)		(n=202)				(n=58)			(n=1132)		
Variables		n	%	95% Cl		n	%	95% Cl	n	%	95% Cl	n	%	95% Cl	
Methods	Short-term	100	63.4	54.4-71.6		53	27.1	20.6-34.7	17	25.7	15.3-39.8	229	19.3	16.8-22.1	
of contraception	Oral contraceptive pills	6	2.6	1.1-6.0		26	12.3	8 3-18 0	7	11.4	5.1-23.6	151	12.7	10.6-15.1	
	Male condoms	94	60.8	51.8-69.1		27	14.8	9.93-21.4	10	14.3	7.3-26.2	78	6.6	5.2-8.4	
	Long-term	52	36.6	28.4-45.6	1	49	72.9	65.3-79.4	41	74.3	60.2-84.7	903	80.7	77.9-83.2	
	Implants & IUD	36	24.7	17.8-33.3		99	49.4	41.3-57.6	30	54.4	39.4-68.7	688	61.0	57.7-64.2	
	Injectables	16	11.9	6.92-19.6		48	22.9	17.3-29.7	11	19.9	10.3-34.9	191	17.6	15.4-20.0	
	Others	0	-	-		2	0.6	0.2-2.47	0	-	-	24	2.2	1.4-3.3	
Source of method	Public	49	33.9	25.8-43.1	1	43	70.8	62.9-77.6	30	61.1	47.3-73.3	824	72.4	68.8-75.8	
	Non-public medical	6	3.9	1.5-9.7		16	7.23	4.2-12.3	11	13.2	7.0-23.5	111	10.4	8.4-12.8	
	For-profit outlets and		62.1	53.0-70.5			22.0	16.2-29.2	17	25.7	15.3-39.8		17.2	14.6-20.1	
	relatives	97				43						197			
Capacity of provider	Comprehensive	52	36.2	27.9-45.4	1	56	77.0	69.8-82.9	38	70.7	56.6-81.7	892	78.9	75.9-81.6	
	Limited	100	63.8	54.6-72.1		46	23.0	17.1-30.2	20	29.3	18.3-43.4	240	21.1	18.4-24.1	

297 Factors associated with demand satisfied by a modern method

298	We examined the demand satisfied by a modern method among women ages 15 to 24
299	years in need of contraception (n=1736). Table 5 shows the percentages of modern
300	method users and the crude and adjusted odds ratios of each factor on the use of
301	modern methods. The multivariable analysis showed that being unmarried, higher
302	parity, existence of barriers in access to health services, and a higher number of
303	modern methods known by a woman were positively associated with demand
304	satisfied by a modern method.

Variables		Modern method use (95% CI)	Crude	OR [95% CI]	P-value	Adjuste	ed OR [95% CI]	P-value
Residence	Urban	21.9 (18.8-25.2)	Ref			Ref		
	Rural	19.5 (16.9-22.5)	0.87	[0.67-1.12]	0.283	1.00	[0.72-1.38]	0.999
Ethnicity	Others	21.6 (19.1-24.4)	Ref			Ref		
	Fon	18.1 (15.0-21.7)	0.80	[0.61-1.05]	0.111	0.76	[0.55-1.05]	0.095
Religion	Christians	20.6 (18.0-23.4)	Ref			Ref		
	Islam	21.6 (18.0-25.8)	1.07	[0.80-1.42]	0.666	1.04	[0.73-1.47]	0.828
	Traditional and others	17.1 (12.0-23.9)	0.80	[0.52-1.24]	0.312	0.83	[0.53-1.30]	0.406
Literacy	Cannot read	17.8 (15.2-20.7)	Ref			Ref		
	Can read	23.6 (20.5-27.0)	1.43	[1.10-1.85]	0.007	1.20	[0.89-1.61]	0.233
Marita	Married	17.3 (15.1-19.8)	Ref			Ref		
status	Not married	27.6 (23.8-31.7)	1.82	[1.42-2.35]	<0.001	3.33	[2.21-5.01]	<0.001
Parity	None	23.4 (19.9-27.3)	Ref			Ref		
	One	19.6 (16.3-23.3)	0.80	[0.59-1.07]	0.133	2.11	[1.39-3.22]	0.001
	Two or more	18.6 (18.5-22.4)	0.75	[0.54-1.02]	0.071	2.22	[1.39-3.55]	0.001
Wealth	Poorest	21.9 (15.6-29.8)	Ref			Ref		
quintiles	Poorer	18.2 (13.9-23.4)	0.79	[0.46-1.35]	0.388	0.78	[0.45-1.33]	0.358
	Middle	20.3 (16.0-25.5)	0.91	[0.55-1.48]	0.701	0.79	[0.49-1.28]	0.333
	Richer	18.5 (14.7-23.1)	0.81	[0.49-1.35]	0.420	0.69	[0.41-1.16]	0.161
	Richest	24.0 (20.0-28.5)	1.13	[0.70-1.81]	0.625	0.94	[0.56-1.57]	0.814
Barriers to access to	None	17.6 (14.6-21.1)	Ref			Ref		
the services	Any	22.2 (19.6-25.1)	1.34	[1.02-1.76]	0.038	1.52	[1.15-2.02]	0.003
Number of modern	Fewer than four	9.7 (6.30-14.8)	Ref			Ref		
methods known	Four to six	21.1 (18.2-24.3)	2.48	[1.48-4.15]	<0.001	2.43	[1.45-4.07]	0.001
	More than six	23.7 (20.2-27.5)	2.88	[1.70-4.85]	< 0.001	3.00	[1.73-5.21]	<0.001

# 305 Table 5. Percentages and factors associated with modern method use among 15-24-year-old women in need of contraception (n=1,736)

# 307 Qualitative strand

We identified three main barriers in access to and utilization of contraception services among adolescents, which are (i) socio-cultural perceptions related to sexuality and fertility and (ii) perceptions about the purpose and effectiveness of contraception. A conceptual model of the barriers is shown in Figure 1.

312

313 Figure 1. A conceptual model of the barriers to access and utilization of contraception

314 services from qualitative analyses

315

316 Sociocultural norms

Sociocultural norms including gender norms, cultural, and religious perceptions affect people's perceptions about young women's fertility and the need for contraception in Allada. These norms include views about pregnancy before marriage, male partners' role in women's contraceptive use, local perceptions of women's fertility, resulting in different perceptions of contraception such as the misuse of biomedicine, perceived effectiveness of natural methods, mistrust in modern methods, and negative views about modern method users.

324 To be pregnant before marriage

There is a common negative perception about sexual intercourse among adolescents. Becoming pregnant as an adolescent and before marriage is not well viewed in the community as judged as promiscuity. Therefore, unmarried young women avoid such judgment through the use of traditional methods such as abstinence or fertility awareness. This allows young women to avoid exposing their pre-marital sexual

activity publically. This situation prevents unmarried young women from using
contraceptives.

- 332 *"To avoid pregnancy, one must abstain from sex. If a girl really wants to avoid shame in*
- front of her classmates, she must refrain from sex." (less than 20 years old/pregnant)
- 334 Perceptions regarding the ideal family size
- 335 In the study area, there was cultural pressure on having a certain number of children based
- on social norms about the ideal family size. It implies that when a young woman gets
- married, she may not be allowed to use contraception until she has several children.
- 338 *"If we say that someone has enough children, it will be from seven. That is when the*
- 339 person will say, 'Yes, I do have children.'" (less than 20 years old/male)
- 340 None of the respondents mentioned cultural pressures that might force young unmarried
- 341 women to have children early.
- 342 Decision-making dynamics
- 343 Women do not often have the decision-making power to use contraception.
- 344 "One day, some people went door-to-door to inform us of the presence of an NGO at
- 345 the health center for family planning services. I then informed my husband and did the
- 346 injection for 3 months. But after he forbade me to continue it, because of rumors about
- 347 other women's bad experiences. When your husband refuses you have no choice."
- 348 (20-24 years old pregnant)
- 349 Perceptions of contraception
- 350 Contraception perceived as abortion
- Religious beliefs, especially Catholicism, also affect the utilization of contraception.

- 352 *"In my opinion, 'planning' is an abortion, a deadly sin. We are killing babies without*
- knowing and it is against the ten commandments of God. We are also killing ourselves
- because we do not know the consequences of those things in our body." (less than 20
- 355 years old/non-pregnant)
- 356 Perceptions of modern method users
- 357 Respondents believed that the use of contraception facilitates sexual promiscuity.
- 358 Contraception was also considered to be a eugenic procedure.
- 359 "When she was 12 years old, people told me to have her adopt a contraceptive method.
- 360 I thought it was asking her to prostitute herself and I did not. Here now, as you see, she
- 361 came home with pregnancy while she is still having her training." (Relative of pregnant)
- 362 Perceived effectiveness of natural methods
- 363 Faith in natural methods contributes to reluctance to use modern contraceptives among
- 364 young women.
- 365 *"I do not plan to use any of those things [modern methods]. My family members told*
- 366 me they will prepare an herbal tea for me and I will not get pregnant for the period I
- 367 want they know how to do it." (less than 20 years old/pregnant)
- Beliefs were led by fear of side effects of modern methods, and these opinions were in part
  based on rumor or hearsay.
- 370 "It happened to one of my relatives. She chose the one that is inserted in the arm.
- 371 Three months later, she had no more periods. Then when she had menstrual periods
- 372 again, it was too much. It was when it became excessive that she had to go where she
- 373 had it inserted for removal. Finally, she died." (20-24 years old/female)
- 374 Misuse of biomedicine

Some also perceived 'aspirin' to be effective in preventing pregnancy and used it as a typeof modern method.

377 *"I often use Sedaspir [commercial name of acetylsalicylic acid] and it works well. I buy*378 the box at the pharmacy for 1500 franc CFA [equivalent to 3 US dollars]." (20-24 years
379 old/pregnant).

380

# 381 DISCUSSION

We investigated modern method use, unmet need for contraception, and the reasons why modern methods were not used; focusing on adolescent girls and young adult women (15 to 24 years old) using the most recent Benin Demographic and Health Survey (BDHS 2017-18) combined with a qualitative study conducted in the Allada area in Benin.

387 Our quantitative analysis revealed that characteristics of modern method use 388 differed between unmarried sexually active and married young women ages 15 to 24. 389 Among all women, young unmarried sexually active women had both the highest 390 modern contraceptive use and the highest unmet need for modern contraceptives. 391 Modern method users tended to adopt more short-term than long-term methods, 392 especially the male condoms which they acquired from providers with a limited 393 range of methods. Young married women had the lowest percentage of demand 394 satisfied by a modern method among women who wished to delay or limit births. 395 Factors associated with modern method use among young women in need of 396 contraception were being unmarried, having at least one child, experiencing barriers 397 in access to health services, and knowing a greater number of modern

medRxiv preprint doi: https://doi.org/10.1101/2021.06.07.21257758; this version posted October 17, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

contraceptives. As suggested by the qualitative analysis, the reasons underlying the
high unmet need for modern contraceptives in young women are related to the
access and use of modern methods which can be categorized as 'sociocultural norms'
and 'perceptions of contraception'.

402 Young unmarried women had the highest unmet need for modern contraceptives as a 403 percentage of all women, which may explain the high rate of unintended pregnancy 404 in Benin. Very few unmarried young women want to have a child due to community 405 negative perceptions about pre-marital sexual intercourse and pregnancy, therefore 406 in need of contraception. However, as unmarried young women are considered in 407 the communities not to be ready for sexual intercourse they are reluctant to request 408 contraceptives despite their wish to avoid a pregnancy, [19, 20]. Moreover, the fear 409 of adverse effects contributes to young women's non-use or discontinuation of 410 modern contraceptives, [20, 30, 31].

411 Among unmarried young women in need of family planning, those with demand 412 satisfied by a modern contraceptive tend to use male condoms, the most adopted 413 modern method in Benin (45.3%), which can be easily obtained from for-profit outlets, partners, and relatives. This finding is similar to that of Ali and Cleland, [32], 414 415 secondary data analysis from 36 African countries which showed that male condoms 416 constitute 51.3% of modern methods among unmarried women ages 15 to 24 years. Our qualitative analysis suggested that while access to male condoms has fewer 417 418 barriers, male partners are in most cases the ones to obtain the method for use. 419 Consequently, the decision to use a male condom is likely to be made by the male 420 partner whose failure to acquire the method or unwillingness to use could be a

421 barrier to the young woman's contraceptive use. Male condoms are obtained in 422 shops, pharmacies, οr from non-governmental organizations during 423 awareness-raising campaigns. Also, young women may assume that male partners 424 obtain contraceptives mainly from pharmacies which could explain why pharmacies and relatives are the most frequent sources of methods reported in the DHS2017-18. 425 426 Young people's choice of providers including shops and pharmacies is based on their 427 values for discretion, respect of confidentiality, and the quickness of the service 428 delivery. This is in line with the findings of Murray et al., [4], which suggested that 429 the preference of pharmacies by young women in requesting contraceptive services 430 may be driven by their perception that the providers are youth friendly.

431 Also, young married women in need of family planning had a lower percentage of 432 demand satisfied by a modern contraceptive compared to unmarried young women 433 and older married women. Most young women want to have a child once they get 434 married which explains the low unmet need as a percentage of all women. However, 435 married young women who wished to delay the next pregnancies face several 436 challenges in decision-making. For instance, the qualitative study found that women 437 are often under pressure to have children once they get married due to traditional 438 beliefs about fertility that prevent married women from using contraceptives. 439 Women are expected to demonstrate their fertility after marriage, which may be even 440 stronger for younger women as suggested by the literature, [33-37]. Moreover, in a 441 social context such as Allada, where large families of seven children are the average, the 442 family size is often decided by male partners and women feel less empowered to decide on 443 their own to adopt contraceptives. This may explain the high total fertility rate of 5.7 in

444 Benin. To get around the barrier of spousal consent, some married women request 445 contraceptives discretely, after giving several births and as their households face 446 economic challenges to raise the children. Most of the women would request long-term 447 methods such as implants to be inserted in the leg or other unusual and less exposed parts 448 of the body.

Our study revealed that factors associated with modern method use among those in 449 450 need included literacy, not being married, having children, knowing a greater 451 number of modern methods, and experiencing barriers to access health services. 452 Although 'literacy' was not often used as a variable in previous studies, among young 453 women there was evidence of a positive association between educational attainment 454 and the demand satisfied, [34, 38]. Similarly, the number of modern methods known 455 was positively associated with use. The strongest association was found between 456 women's marital status and the demand satisfied by a modern method. The 457 multivariate analysis revealed that young unmarried women were four times more 458 likely to use modern methods. This finding strengthens the evidence that married 459 young women are being left behind in terms of demand satisfied. Also, a similar 460 positive association between being unmarried and using modern contraceptives was 461 found by Dansou et al. in a study among all women of reproductive age (15 to 49 462 years old) in Benin, [20]. Several other studies conducted in other sub-Saharan 463 African countries such as Burkina Faso, Ethiopia, Nigeria, and Ghana found that 464 modern method prevalence was higher among unmarried adolescent girls compared 465 to married,[39, 40]. Young unmarried women used contraceptives to avoid pregnancy 466 because of the social stigma associated with early pregnancy in Benin, and because

467 they have no income to support a child on their own, [10, 20]. This finding is also 468 supported by our qualitative data. Having a child was associated with modern 469 method use. Other studies revealed that women with children were more likely to 470 use modern methods, [41, 42]. Health education from care providers during 471 pregnancy and in the postpartum period might be a contributing factor because the 472 BDHS 2017-8 showed that coverage of antenatal care provided by skilled personnel 473 and delivery in a health facility were 83% and 84%, respectively. Counterintuitively, 474 we found that having barriers to access to health services was associated with a higher 475 likelihood of modern methods use. We hypothesize that women who had accessed 476 contraceptive methods were more likely to report barriers, while those who did not report 477 such barriers did not theoretically exist.

478 Finally, our qualitative analysis revealed aspirin or paracetamol was used in Allada 479 to prevent pregnancy, despite the lack of a biomedical evidence-based for such an 480 effect. Boko et al. (2015) noted the same practice in Benin,[43]. The practice may be 481 driven by the fact that requesting such medicine does not expose a young woman to 482 any social stigma as the drug is not viewed as a contraceptive method by providers. 483 This common practice may imply that the use of oral contraception use among young 484 women in the DHS2017-18 is an over-estimate. In a context such as in Allada where 485 other ineffective drugs are used as oral contraceptives, it is important to promote 486 accurate knowledge as well as practices through health education activities.

487

488 Limitations

medRxiv preprint doi: https://doi.org/10.1101/2021.06.07.21257758; this version posted October 17, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

489 Our study has several limitations. We used secondary data, our analysis could have 490 been affected by recall errors and biases in some variables collected on the DHS. For 491 example, several assumptions were made in the DHS to assess the need for 492 contraception among women, which were criticized to have possibly induced an 493 underestimation in unmet needs for family planning, [44]. The assumptions 494 concerned the frequency of sexual activity of women, women's capacity to conceive (fecundity criteria, length of lactational amenorrhea which is 24 months in DHS 495 496 versus 6 months clinically accepted standard), and the validity of their reports on 497 whether recent births were desired. In the qualitative study, having a male 498 interviewer might have affected responses from young women. Meanwhile, this was 499 addressed as a young female interviewer continued interviews with female 500 participants. Moreover, we could not ascertain whether a respondent in the 501 qualitative study was sexually active whereas the analysis of determinants in the 502 quantitative study focused on sexually active young women.

503

#### 504 Conclusion

Access to and use of modern contraceptive methods is low among women in Benin, especially among adolescent girls and young women. Adolescent girls and young women tend to use short-term methods such as condoms which they obtain from easily accessible providers and relatives. Unmarried and married women face different barriers to use modern contraceptives, which need to be carefully addressed. Most importantly young married women who wish to delay births should be specifically targeted in interventions to address challenges related to decision-making to use contraceptives.

512	Raising awareness in communities with strong culture towards encouraging early
513	marriage and high fertility, as well as promoting comprehensive sexuality education
514	and providing youth-friendly services may help reduce the incidence of unintended
515	pregnancies among adolescent girls and young women,[45, 46].
516	

# 517 Acknowledgments

518 We thank the Demographic Health Survey Program (http://dhsprogram.com/) for

519 granting us access to the BDHS2017-18 dataset. We acknowledge TOTIN Mahoutondji

520 Gilberte for her work as a research assistant during the qualitative data collection.

521 We would like to thank the study participants for sharing their experiences as well as

522 the local administrations of Allada for facilitating our study implementation.

523

# 524 Contributors

NCAA, MM, MS, LB, TD, J-P D, and SG were involved in conception and study design.
LB and MM provided statistical expertise. NCAA, AV, and CB collected qualitative
data in the field. MM, MS, TD, CG, LK, and KP provided qualitative data analysis
expertise. NCAA, MM, and MS were involved in the drafting of the manuscript. LB, TD,
J-P D, and CG were involved in the critical revision of the manuscript for important
intellectual content. All authors were involved in the final approval of the

532

533 Funding

- 534 This study was supported by core funding from Nagasaki University School of
- 535 Tropical Medicine and Global Health to MM (grant number N/A).

536

- 537 Competing interests
- 538 None declared.

539

540 Patient and public involvement

541 Patients and/or the public were not involved in the design, conduct, reporting, or

542 dissemination plans of this research.

543

- 544 Patient consent for publication
- 545 Not required.

546

#### 547 Data availability statement

548 Quantitative data (BDHS 2017-18) is available from the DHS program 549 (https://dhsprogram.com/). Qualitative data including interview audios and 550 transcripts cannot be shared publicly because it contains information which enables 551 one to identify each individual.

552

#### 553 Ethical statement

554 The BDHS 2017-18 received approvals from the National Statistics Council and the 555 National Ethics Committee for Health Research in Benin. As DHS data are secondary 556 data, the quantitative strand did not require ethical approvals. The qualitative strand

557	was conducted as part of the "Continuum des Soins de Santé Reproductive au Bénin"
558	study which received approvals from the Local Ethics Committee for Biomedical
559	Research of the University of Parakou in Benin (approval number:
560	0092/CLERB-UP/P/SP/R/SA), the Institute of Tropical Medicine Antwerp in Belgium
561	(approval number: IRB/AB/AC/044), and Nagasaki University School of Tropical
562	Medicine and Global Health in Japan (approval number: NU-TMGH_064). The
563	participants in the interviews and the focus group discussions were informed of the
564	research objectives, their voluntary participation in the study, and there was no
565	disadvantage in refusal or withdrawal from the study. Written consent was obtained
566	from each participant.
567	
568	References
500	
569	1. United Nations Population Fund (UNFPA). Programme of Action. New York: UNFPA 1994.
569 570	<ol> <li>United Nations Population Fund (UNFPA). Programme of Action. New York: UNFPA 1994.</li> <li>Singh S, Sedgh G, Hussain R. Unintended pregnancy: worldwide levels, trends, and</li> </ol>
569 570 571	<ol> <li>United Nations Population Fund (UNFPA). Programme of Action. New York: UNFPA 1994.</li> <li>Singh S, Sedgh G, Hussain R. Unintended pregnancy: worldwide levels, trends, and outcomes. <i>Stud Fam Plann</i> 2010;41(4):241-50. doi:</li> </ol>
569 570 571 572	<ol> <li>United Nations Population Fund (UNFPA). Programme of Action. New York: UNFPA 1994.</li> <li>Singh S, Sedgh G, Hussain R. Unintended pregnancy: worldwide levels, trends, and outcomes. <i>Stud Fam Plann</i> 2010;41(4):241-50. doi: 10.1111/j.1728-4465.2010.00250.x [published Online First: 2011/04/06]</li> </ol>
569 570 571 572 573	<ol> <li>United Nations Population Fund (UNFPA). Programme of Action. New York: UNFPA 1994.</li> <li>Singh S, Sedgh G, Hussain R. Unintended pregnancy: worldwide levels, trends, and outcomes. <i>Stud Fam Plann</i> 2010;41(4):241-50. doi: 10.1111/j.1728-4465.2010.00250.x [published Online First: 2011/04/06]</li> <li>Hubacher D, Mavranezouli I, McGinn E. Unintended pregnancy in sub-Saharan Africa:</li> </ol>
569 570 571 572 573 574	<ol> <li>United Nations Population Fund (UNFPA). Programme of Action. New York: UNFPA 1994.</li> <li>Singh S, Sedgh G, Hussain R. Unintended pregnancy: worldwide levels, trends, and outcomes. <i>Stud Fam Plann</i> 2010;41(4):241-50. doi: 10.1111/j.1728-4465.2010.00250.x [published Online First: 2011/04/06]</li> <li>Hubacher D, Mavranezouli I, McGinn E. Unintended pregnancy in sub-Saharan Africa: magnitude of the problem and potential role of contraceptive implants to alleviate</li> </ol>
569 570 571 572 573 574 575	<ol> <li>United Nations Population Fund (UNFPA). Programme of Action. New York: UNFPA 1994.</li> <li>Singh S, Sedgh G, Hussain R. Unintended pregnancy: worldwide levels, trends, and outcomes. Stud Fam Plann 2010;41(4):241-50. doi: 10.1111/j.1728-4465.2010.00250.x [published Online First: 2011/04/06]</li> <li>Hubacher D, Mavranezouli I, McGinn E. Unintended pregnancy in sub-Saharan Africa: magnitude of the problem and potential role of contraceptive implants to alleviate it. Contraception 2008;78(1):73-8. doi: 10.1016/j.contraception.2008.03.002</li> </ol>
569 570 571 572 573 574 575 576	<ol> <li>United Nations Population Fund (UNFPA). Programme of Action. New York: UNFPA 1994.</li> <li>Singh S, Sedgh G, Hussain R. Unintended pregnancy: worldwide levels, trends, and outcomes. Stud Fam Plann 2010;41(4):241-50. doi: 10.1111/j.1728-4465.2010.00250.x [published Online First: 2011/04/06]</li> <li>Hubacher D, Mavranezouli I, McGinn E. Unintended pregnancy in sub-Saharan Africa: magnitude of the problem and potential role of contraceptive implants to alleviate it. Contraception 2008;78(1):73-8. doi: 10.1016/j.contraception.2008.03.002 [published Online First: 2008/06/17]</li> </ol>
569 570 571 572 573 574 575 576 577	<ol> <li>United Nations Population Fund (UNFPA). Programme of Action. New York: UNFPA 1994.</li> <li>Singh S, Sedgh G, Hussain R. Unintended pregnancy: worldwide levels, trends, and outcomes. Stud Fam Plann 2010;41(4):241-50. doi: 10.1111/j.1728-4465.2010.00250.x [published Online First: 2011/04/06]</li> <li>Hubacher D, Mavranezouli I, McGinn E. Unintended pregnancy in sub-Saharan Africa: magnitude of the problem and potential role of contraceptive implants to alleviate it. Contraception 2008;78(1):73-8. doi: 10.1016/j.contraception.2008.03.002 [published Online First: 2008/06/17]</li> <li>FP2020. Indicators 3-4 : Unmet Need and Demand Satisfied.</li> </ol>
569 570 571 572 573 574 575 576 577 578	<ol> <li>United Nations Population Fund (UNFPA). Programme of Action. New York: UNFPA 1994.</li> <li>Singh S, Sedgh G, Hussain R. Unintended pregnancy: worldwide levels, trends, and outcomes. <i>Stud Fam Plann</i> 2010;41(4):241-50. doi: 10.1111/j.1728-4465.2010.00250.x [published Online First: 2011/04/06]</li> <li>Hubacher D, Mavranezouli I, McGinn E. Unintended pregnancy in sub-Saharan Africa: magnitude of the problem and potential role of contraceptive implants to alleviate it. <i>Contraception</i> 2008;78(1):73-8. doi: 10.1016/j.contraception.2008.03.002 [published Online First: 2008/06/17]</li> <li>FP2020. Indicators 3-4 : Unmet Need and Demand Satisfied. <i>http://2016-2017progressfamilyplanning2020org/en/measurement-section/unmet-n</i></li> </ol>
569 570 571 572 573 574 575 576 577 578 579	<ol> <li>United Nations Population Fund (UNFPA). Programme of Action. New York: UNFPA 1994.</li> <li>Singh S, Sedgh G, Hussain R. Unintended pregnancy: worldwide levels, trends, and outcomes. Stud Fam Plann 2010;41(4):241-50. doi: 10.1111/j.1728-4465.2010.00250.x [published Online First: 2011/04/06]</li> <li>Hubacher D, Mavranezouli I, McGinn E. Unintended pregnancy in sub-Saharan Africa: magnitude of the problem and potential role of contraceptive implants to alleviate it. Contraception 2008;78(1):73-8. doi: 10.1016/j.contraception.2008.03.002 [published Online First: 2008/06/17]</li> <li>FP2020. Indicators 3-4 : Unmet Need and Demand Satisfied. http://2016-2017progressfamilyplanning2020org/en/measurement-section/unmet-n eed-and-demand-satisfied-indicators-3-4 [accessed November 2020]</li> </ol>

580	5. Okigbo CC, Speizer IS. Determinants of Sexual Activity and Pregnancy among Unmarried
581	Young Women in Urban Kenya: A Cross-Sectional Study. PLoS One
582	2015;10(6):e0129286. doi: 10.1371/journal.pone.0129286 [published Online First:
583	2015/06/06]
584	6. World Health Organization (WHO). Family planning evidence brief: family planning
585	financing. Geneva: WHO, 2018.
586	https://apps.who.int/iris/bitstream/handle/10665/255863/WHO-RHR-18.26-eng.pdf
587	[accessed August 2019]
588	7. Darroch JE, Woog V, Bankole A, et al. Costs and Benefits of Meeting the Contraceptive
589	Needs of Adolescents. New York: Guttmacher Institute 2016.
590	http://www.guttmacher.org/sites/default/files/report_pdf/adding-it-up-adolescents
591	-report.pdf [accessed May 2020]
592	8. Fescina R, B DM, JL DR, et al. Guides for the PHC focused continuum of care of women
593	and newborns. Montevideo: Latin American Center for Perinatology, Women and
594	Reproductive Health, PAHO, 2009.
595	9. World Health Organization (WHO). Health for the world's adolescents: A second chance
596	in the second decade. Geneva: WHO,2014
597	10. Sedgh G, Ashford LS, Hussain R. Unmet Need for Contraception in Developing
598	Countries <sup>1</sup> 2: Examining Women's Reasons for Not Using a Method. <i>New York:</i>
599	Guttmacher Institute 2016.
600	https://www.guttmacher.org/sites/default/files/report_pdf/unmet-need-for-contrac
601	eption-in-developing-countries-report.pdf [accessed May 2020]

- 602 11. McCurdy RJ, Jiang X, Schnatz PF. Long-acting reversible contraception in adolescents in
- 603 Sub-Saharan Africa: evidence from demographic and health surveys. Eur J
- 604 *Contracept Reprod Health Care* 2018;23(5):357-64. doi:
- 605 10.1080/13625187.2018.1519535 [published Online First: 2018/11/23]
- 606 [dataset] 12. Institut National de la Statistique et de l'Analyse Économique (INSAE) and ICF.
- 607 Enquête Démographique et de Santé au Bénin, 2017-2018. Cotonou, Bénin and
- 608 Rockville, Maryland, USA : INSAE and ICF. 2019.
- 609 13. Government of Benin. FAMILY PLANNING 2020 COMMITMENT. 2017.
- 610 http://www.familyplanning2020.org/news/benin-fp2020-commitment [accessed
- 611 March 2020]
- 612 14. Chae S, Woog V, Zinsou C, et al. Obstacles à la pratique contraceptive des femmes au
- 613 Bénin. En Bref, New York: Guttmacher Institute 2015.
- 614 https://www.guttmacher.org/sites/default/files/report\_pdf/ib-benin-contraception-
- 615 fr.pdf [accessed March 2020]
- 616 15. USAID, Communities AP. Benin's Community-Based Access to Injectable Contraceptives
- 617 Pilot Project. 2015.
- 618 https://www.advancingpartners.org/sites/default/files/sites/default/files/resources
- 619 /apc\_benin\_cba2i\_pilot\_brief\_march\_2015\_english\_final.pdf [accessed July 2020]
- 620 16. Okegbe T, Affo J, Djihoun F, et al. Introduction of Community-Based Provision of
- 621 Subcutaneous Depot Medroxyprogesterone Acetate (DMPA-SC) in Benin:
- 622 Programmatic Results. *Glob Health Sci Pract* 2019;7(2):228-39. doi:
- 623 10.9745/ghsp-d-19-00002 [published Online First: 2019/06/07]

624	17. Igras S,	Diakité M, Lund	gren R. Moving	; from theory	to practice: A	participatory	/ social
-----	--------------	-----------------	----------------	---------------	----------------	---------------	----------

- 625 network mapping approach to address unmet need for family planning in Benin.
- 626 Glob Public Health 2017;12(7):909-26. doi: 10.1080/17441692.2016.1147589
- 627 [published Online First: 2016/03/08]
- 628 18. Ahovey EC. Besoins non satisfaits en planification familiale au sein du couple 🛛:
- 629 caractéristiques socio-démographiques et cadre de vie au Benin. 2002
- 630 <http://www.cicred.org/Eng/Seminars/Details/Seminars/Bangkok2002/001Bangkok
- 631 Ahovey.pdf> [Accessed July 2020].
- 632 19. Sedgh G, Hussain R. Reasons for contraceptive nonuse among women having unmet
- 633 need for contraception in developing countries. *Stud Fam Plann* 2014;45(2):151-69.
- 634 doi: 10.1111/j.1728-4465.2014.00382.x [published Online First: 2014/06/17]
- 635 20. Dansou J, Adekunle AO, Arowojolu AO. Factors behind the preference in contraceptives
- 636 use among non-pregnant and sexually active women in Benin Republic. Cent Afr J

637 Public Health 2017;3(5):80-89. doi: 10.11648/j.cajph.20170305.15

638 21. MacQuarrie KLD. Unmet need for family planning among young women: levels and

- 639 trends. DHS Comparative Reports No 34. Rockville, Maryland, USA: ICF International,
- 640 2014.

642

641 22. Radovich E, Dennis ML, Wong KLM, et al. Who Meets the Contraceptive Needs of Young

Women in Sub-Saharan Africa? J Adolesc Health 2018;62(3):273-80. doi:

- 643 10.1016/j.jadohealth.2017.09.013 [published Online First: 2017/12/19]
- 644 23. Ahinkorah BO. Predictors of modern contraceptive use among adolescent girls and
- 645 young women in sub-Saharan Africa: a mixed effects multilevel analysis of data from

- 647 10.1186/s40834-020-00138-1.
- 648 24. Maxwell J. Designing a Qualitative Study. *Qualitative Research Design An Interactive*
- 649 Approach (Third Edition) 2012
- 650 25. Hubacher D, Trussell J. A definition of modern contraceptive methods. *Contraception*
- 651 2015;92(5):420-1. doi: 10.1016/j.contraception.2015.08.008 [published Online First:
- 652 2015/08/16]
- 653 26. Bradley SEK, Croft T, Fishel J, et al. Revising Unmet Need for Family Planning. DHS

654 Analytical Studies No 25 Calverton, Maryland, USA: ICF International 2012

655 27. Festin MP, Kiarie J, Solo J, et al. Moving towards the goals of FP2020 - classifying

656 contraceptives. *Contraception* 2016;94(4):289-94. doi:

- 657 10.1016/j.contraception.2016.05.015 [published Online First: 2016/06/12]
- 658 28. Vigan A, Boyi C, Ahissou C, et al. Déterminants de l'utilisation des services le long du
- 659 continuum des Soins de Santé Reproductive au Bénin: une exploration qualitative.
- 660 Rapport d'étude. Ministère de la Santé, Centre de Recherche en Reproduction
- 661 Humaine et Démographie (CERRHUD), Institut de Médecine Tropicale (IMT), 2020
- 662 29. Hausmann-Muela S, Muela Ribera J, Toomer E, et al. The PASS-model: a model for
- 663 guiding health-seeking behavior and access to care research. *Malaria Reports*
- 664 2012;2:e3. doi: 10.4081/malaria.2012.e3
- 665 30. Ochako R, Mbondo M, Aloo S, et al. Barriers to modern contraceptive methods uptake
- among young women in Kenya: a qualitative study. *BMC Public Health* 2015;15:118.
- 667 doi: 10.1186/s12889-015-1483-1 [published Online First: 2015/04/18]

668	31. Campbell OM, Benova L, Macleod D, et al. Who, What, Where: an analysis of private
669	sector family planning provision in 57 low- and middle-income countries. Trop Med
670	Int Health 2015;20(12):1639-56. doi: 10.1111/tmi.12597 [published Online First:
671	2015/09/29]
672	32. Ali MM, Cleland J. Long term trends in behaviour to protect against adverse
673	reproductive and sexual health outcomes among young single African women.
674	Reprod Health 2018;15(1):136. doi: 10.1186/s12978-018-0576-6 [published Online
675	First: 2018/08/16]
676	33. Asiimwe JB, Ndugga P, Mushomi J. Socio-demographic factors associated with
677	contraceptive use among young women in comparison with older women in Uganda.
678	DHS Working Papers No 95, 2013.
679	34. Asiimwe JB, Ndugga P, Mushomi J, et al. Factors associated with modern contraceptive
680	use among young and older women in Uganda; a comparative analysis. BMC Public
681	Health 2014;14:926. doi: 10.1186/1471-2458-14-926 [published Online First:
682	2014/09/10]
683	35. Ngome E, Odimegwu C. The social context of adolescent women's use of modern
684	contraceptives in Zimbabwe: a multilevel analysis. Reprod Health 2014;11:64. doi:
685	10.1186/1742-4755-11-64 [published Online First: 2014/08/12]
686	36. Shahabuddin AS, Nöstlinger C, Delvaux T, et al. What Influences Adolescent Girls'
687	Decision-Making Regarding Contraceptive Methods Use and Childbearing? A
688	Qualitative Exploratory Study in Rangpur District, Bangladesh. PLoS One
689	2016;11(6):e0157664. doi: 10.1371/journal.pone.0157664 [published Online First:
690	2016/06/24]

691	37. Mardi A, Ebadi A, Shahbazi S, et al. Factors influencing the use of contraceptives
692	through the lens of teenage women: a qualitative study in Iran. BMC Public Health
693	2018;18(1):202. doi: 10.1186/s12889-018-5116-3 [published Online First:
694	2018/02/01]
695	38. Tamang L, Raynes-Greenow C, McGeechan K, et al. Factors associated with
696	contraceptive use among sexually active Nepalese youths in the Kathmandu Valley.
697	Contracept Reprod Med 2017;2:13. doi: 10.1186/s40834-017-0040-y [published
698	Online First: 2017/12/05]
699	39. Hounton S, Barros AJ, Amouzou A, et al. Patterns and trends of contraceptive use
700	among sexually active adolescents in Burkina Faso, Ethiopia, and Nigeria: evidence
701	from cross-sectional studies. Glob Health Action 2015;8:29737. doi:
702	10.3402/gha.v8.29737 [published Online First: 2015/11/13]
703	40. Marrone G, Abdul-Rahman L, De Coninck Z, et al. Predictors of contraceptive use among
704	female adolescents in Ghana. <i>Afr J Reprod Health</i> 2014;18(1):102-9. [published
705	Online First: 2014/05/07]
706	41. de Vargas Nunes Coll C, Ewerling F, Hellwig F, et al. Contraception in adolescence: the
707	influence of parity and marital status on contraceptive use in 73 low-and
708	middle-income countries. <i>Reprod Health</i> 2019;16(1):21. doi:
709	10.1186/s12978-019-0686-9 [published Online First: 2019/02/23]
710	42. Ahinkorah BO. Predictors of unmet need for contraception among adolescent girls and
711	young women in selected high fertility countries in sub-Saharan Africa: A multilevel
712	mixed effects analysis. PLoS One 2020;15(8):e0236352. doi:
713	10.1371/journal.pone.0236352 [published Online First: 2020/08/08]

- 43. Boko I, Baxerres C, Ouattara F, et al. Interroger au Bénin les usages populaires d'un
- 715 médicament abortif, le misoprostol. *Rev med perinat* 2017;9(1):20-24. doi:
- 716 10.1007/s12611-016-0388-2
- 717 44. Bradley SE, Casterline JB. Understanding unmet need: history, theory, and measurement.
- 718 Stud Fam Plann 2014;45(2):123-50. doi: 10.1111/j.1728-4465.2014.00381.x
- 719 [published Online First: 2014/06/17]
- 45. International Planned Parenthood Federation (IPPF). Provide: Strengthening youth
- 721 friendly services. London: IPPF, 2008.
- 46. UNESCO. UN urges Comprehensive Approach to Sexuality Education.
- 723 <a href="https://en.unesco.org/news/urges-comprehensive-approach-sexuality-education">https://en.unesco.org/news/urges-comprehensive-approach-sexuality-education</a>
- 724 [Accessed November 2020].



medRxiv preprint doi: https://doi.org/10.1101/2021.06.07.21257758; this version posted October 17, 20° (which was not certified by peer review) is the author/funder, who has granted medRxiv a licen It is made available under a CC-BY-NC-ND 4.0 International liv