

Frontline Healthcare workers suffering from psychosomatic disorders during COVID-19 (a pandemic) – A Systematic review

Rhythm Joshi^a, Nidhi.B.Agarwal^a, Dinesh Bhurani^b, Mohd Ashif Khan^{a*}

^aCentre for Translational & Clinical Research, School of Chemical and Life Sciences, Jamia Hamdard, New Delhi 110062, India

^bDepartment of Hemato-Oncology & Bone Marrow Transplantation
Rajiv Gandhi Cancer Institute & Research Centre, Rohini, New Delhi 110085, India

***Corresponding author**

Dr. Mohd Ashif Khan
Centre for Translational & Clinical Research
School of Chemical and Life Sciences
Jamia Hamdard
Hamdard Nagar, New Delhi 110062, India
Email: mhdashifkhan@gmail.com, makhan@jamiahamdard.ac.in,
Mobile no: +91-989102639

Abstract

Purpose: The emergence of SARS CoV-2, has imposed high pressure on the healthcare system worldwide. As a consequence, frontline healthcare workers were impacted widely. The aim of this systematic review is to examine the impact of COVID-19 on mental status of FHW during pandemic.

Methods: Databases such as PubMed, Scopus, google scholar were searched extensively from the date of inception till April 2021. All cross-sectional studies published in English assessing the mental condition and well-being of frontline caregivers during COVID-19 were included in the study. The quality assessment was done by Newcastle Ottawa scale.

Results: Ten thousand eight hundred sixty-nine articles were found. After conscientious literature search, total 78 articles were included satisfying the objective of the review. The highest and lowest values for the rates of depression, anxiety and insomnia was found to be 99.51% & 6.07%, 85.7% & 73.6%, and 5.3% & 11.4%, respectively.

Conclusion: It has been found that FHW were psychologically impacted by the pandemic. This could be due to lack of resources such as PPE, organizational support, inefficient relevant knowledge regarding the novel virus, its extremely indelible transmission rates, fear of contamination, stigmatization, and/or due to prevalence of ignorance by government and health policy makers.

Prospero registration no- CRD42021244612

Key words; Mental disorders, Frontline healthcare workers, Psychosomatic disorders, psychological morbidities, COVID-19.

List of abbreviation

COVID-19- Coronavirus disease 2019

HCWs- Healthcare workers

FL- Frontline

GAD-7- Generalized anxiety disorder assessment scale

PHQ-9- patient health questionnaire

HADS- Hospital anxiety depression scale

ISI- 7- Insomnia severity index

PLC-5- post traumatic stress disorder checklist

IES-R- Impact event scale revised

PHQ-5- Patient health questionnaire

Y-BOCS- Yale brown obsessive compulsive scale

ESS- Epworth sleepiness scale

HADS- Hospital anxiety depression scale

BDI- IA- Beck Depression Inventory

PSQI- Pittsburgh sleep quality index

HARS- Hamilton Anxiety rating scale

NASA- TLX- NASA task load index

STAI-Y1- State- Trait Anxiety Inventory (STAI)

BDI-II- Beck depression inventory- II

DASS-21- Depression, anxiety and stress scale-21 items

PSS-14- The Perceived stress scale 14- item

PSS-10- The perceived stress scale 10- items

BAI- Beck anxiety inventory

PDEQ- Peritraumatic dissociative Experience questionnaire

PSI WHOQL-BREF- World Health organization quality of life instrument

GHQ-12- General Health questionnaire 12- item scale

PHQ-2- Patient health questionnaire 2- item scale

GSI- Global severity index scale 53 item

GAD-2- Generalized anxiety disorder 2- item scale

Pro QoL- Professional Quality of life scale

SAS- Self- rating anxiety scale

EQ-5D-3L- Health status assessment questionnaire of the patient.

ED- Emergency department

ALP- Assistant Laboratory personnel

AN- Assistant nurse

N- Nurse

MD- Medical doctor

Psy- Clinical Psychologist

MLT- Medical laboratory technician

HT- Health Technician

1. INTRODUCTION

Coronavirus belong to the family Coronaviridae and order Nidovirales. These are non-segmented, enveloped and single stranded RNA viruses classified under genus coronavirus (P. Chen et al., 2020). Coronavirus disease (COVID-19), whose first case reported in Hubei Province of Wuhan, China in November 2019, has created havoc in lives of people all around the world.

COVID-19 pandemic has led to the worst crisis globally affecting more than 5,378 million people and mortality rate of 1.3 million people as of 14 Nov 2020 (C et al., 2020).

Healthcare workers (HCWs) are at risk of developing mental distress and other psychological symptoms due to high work demand (Fauzi et al., 2020). Moreover, recently various studies have reported the prevalence of depression, anxiety, and insomnia among HCW due to current pandemic (de Burgos-Berdud et al., 2021; Santabárbara et al., 2021).

High transmission and mortality rate of COVID-19 among HCW may be one of the reasons for the prevalence of psychiatric morbidity among them. According to reports, by the end of Feb, 2020 more than 3000 HCW in China got transmitted with the virus (Gan et al., 2020) with the prevalence rate of depression, anxiety, and stress-related symptoms to be 50%, 44.7%, and 73.4%, respectively (Elbay et al., 2020a). Also, Italy faced the death of almost 100 physicians and 26 nurses and over 12,680 HCW were infected from COVID-19 till 7th April 2020 (Barranco and Ventura, 2020). France reported high infection rates among HCWs and more than 50 deaths (McKay et al., 2020; Shaukat et al., 2020) Challenges were knocked together for HCWs due to rapid spread of the disease and delayed onset of symptoms in infected patients or asymptomatic carriers (Hall, 2020; McKay et al., 2020; Shaukat et al., 2020).

Conditions similar to that of battle field are being seen by HCWs in these unprecedented challenges. High levels of anxiety and uncertainties are being experienced and thus persistent

risks of development of stress exposure syndrome and professional burnout are prevailing among those working in frontline]. (Albott et al., 2020; Antonijevic et al., 2020a; Kannampallil et al., 2020; Keubo et al., 2020).

In this phase of COVID-19 attack, HCW being the frontline warriors are the ones who are least talked about.

In this century world has not faced a pandemic as big as COVID-19, which is also new for HCWs, thus, it is need of today to explore the psychological crisis they are being facing (Thamer H. Alenazi et al., 2020). Hence, this systematic review aims to report the mental health condition of HCW during COVID-19.

2. METHOD

2.1 Data research strategy

This study was conducted following PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-analysis) guidelines using suitable keywords such as [Mental disorders OR Depression OR Anxiety OR Bipolar disorder OR Cognitive impairment OR Neurodegenerative disorders] AND [Healthcare workers OR Medical professional OR Doctors OR Nurses] AND [COVID-19] in PubMed, Embase/Elsevier, Cochrane, Science direct and Google scholar from inception till April 2021. Additionally, citations of the included studies were screened manually for the identification of relevant studies. Language of the articles was restricted to English.


2.2 Selection criteria for included studies

Articles fulfilling the following criteria were included: 1. Studies which examined the impact of SARS-CoV-2 on mental health of healthcare professionals; 2. Studies which investigated mental health of FHW; 3. Use of at least one validated quantitative scoring scale to measure mental health outcomes; 4. Available in English Language; 5. Hospital based studies; 6. Conducted from 31st December 2019 (when China reported the first case of SARS-CoV-2 in

Wuhan) till April 2021; 7. Original research (i.e., commentaries, editorials and reviews were excluded), be published in peer reviewed journals. Inclusion of studies in the present systematic review is depicted using PRISMA flowchart (**Figure 1**).

Figure 1

PRISMA Flow diagram showing study selection process



Studies investigating non-hospital-based HCWs, published in languages other than English or having fewer than 20 participants, and editorials or protocols were excluded.

2.3 Data derivation and evaluation of study trait

To extract the desirable data, structured format was used and information such as study country, population size, type of mental health, measurement tool used for mental health, severity of outcome, and prevalence rate were tabulated.

2.4 Outcomes

The objective of this systematic review was to assess the prevalence of depression, anxiety, insomnia, stress and distress among FHW during COVID-19..

3. RESULTS

3.1 Literature search

Initially, 10,919 records were found. Out of these, 27 duplicates were removed and 4253 potential studies (title and abstract) were screened. Further, after the exclusion of 4071 articles, full texts of 182 articles were assessed.. Finally, 78 articles were included in the study (Fig. 1) (AlAteeq et al., 2020; Thamer H Alenazi et al., 2020; Alshekaili et al., 2020a; Altintas, 2017; An et al., 2020; Antonijevic et al., 2020a; Ayhan Başer et al., 2020; Azoulay et al., n.d.; Bhargava et al., 2020; Buselli et al., 2020; Caillet et al., 2020a; X. Chen et al., 2020; Kannampallil et al., 2020; Keubo et al., 2020; Liu et al., 2020) (Altintas, 2017)(AlAteeq et al., 2020; Thamer H Alenazi et al., 2020; Alshekaili et al., 2020a)(An et al., 2020; Antonijevic et al., 2020b; Ayhan Başer et al., 2020; Azoulay et al., n.d.; Bhargava et al., 2020; Buselli et al., 2020; Caillet et al., 2020a; X. Chen et al., 2020; Liu et al., 2020)(W. Cai et al., 2020; J. Chen et al., 2020; Q. Chen et al., 2020; X. Chen et al., 2020; Cheng et al., 2020; Chew et al., 2020; Di Tella et al., 2020; Elbay et al., 2020b; Elhadi et al., 2020b; Fadhli et al., n.d.; Gallopeni et al., 2020; Herrero San Martin et al., 2020; Hong et al., 2020; Huang

and Zhao, 2020; Janati Idrissi et al., 2020; Johnson et al., 2020; Kang et al., 2020a; Kannampallil et al., 2020; Keubo et al., 2020; Koksai et al., 2020a; Korkmaz et al., 2020; Labrague and De los Santos, 2020; J. Lai et al., 2020; G. Li et al., 2020; R. Li et al., 2020; Liang et al., 2020; M. Liu et al., 2020; Y. Liu et al., 2020a; Lu et al., 2020; Luceño-Moreno et al., 2020; Nie et al., 2020; Ning et al., 2020; Prasad et al., 2020; Que et al., 2020; Rossi et al., 2020; Şahin et al., 2020a; Salopek-Žiha et al., 2020; Sandesh et al., 2020; Shah et al., 2020; Shahrour and Dardas, 2020; Shechter et al., 2020; Si et al., 2020; Skoda et al., 2020; Stojanov et al., 2020; Suryavanshi et al., 2020; Teo et al., 2020; Tian et al., 2020; Wang et al., 2020, 2020; L. Q. Wang et al., 2020; Wańkiewicz et al., 2020; Wilson et al., 2020; Xiao et al., 2020; Xing et al., 2020; Yáñez et al., 2020; Yang et al., 2020a; Yin et al., 2020; W. R. Zhang et al., 2020)

3.2 Study characteristics

Seventy-eight studies met the inclusion criteria for this systematic review. All studies quantitatively synthesize data of the psychological illness such as anxiety, depression, stress disorders, insomnia and PTSD affecting FHW due to COVID-19.

Out of 78 studies, 35 were from China; 4 studies were conducted in USA and Turkey; 3 studies were from Saudi Arabia and Italy; 2 studies were from Serbia, Singapore, Pakistan and India; 1 study each from Oman, Malaysia, Kosovo, Morocco, Norway, Cameroon, Philippines, Croatia, Jordan, Germany, Poland, Peru, Iran and United Kingdoms. One study included the dermatologists from all the continents- Asia, North America, Europe and others. (Altintas, 2017)(An et al., 2020)(Q. Chen et al., 2020; X. Chen et al., 2020; Cheng et al., 2020)(G. Li et al., 2020; Wang et al., 2020)(Huang and Zhao, 2020; Juan et al., 2020; Wang et al., 2020)(Kang et al., 2020b; J. Lai et al., 2020; X. Lai et al., 2020; Liang et al., 2020; Liu et al., 2020; Y. Liu et al., 2020b)(Liu et al., 2020; Nie et al., 2020; Ning et al., 2020; L. Q. Wang et al., 2020), (Tian et al., 2020)(W. Cai et al., 2020; Xiao et al., 2020)(Si et al., 2020;

Xiaoming et al., 2020)(Y. Liu et al., 2020b; Xiaoming et al., 2020; Xing et al., 2020; Yang et al., 2020b)(Yin et al., 2020; M. Zhan et al., 2020; S. X. Zhang et al., 2020; Yifang Zhou et al., 2020)(B. Zhou et al., 2020; Zhou et al., 2020, 2020; Yifang Zhou et al., 2020; Z. Zhu et al., 2020b).(Civantos et al., 2020; Kannampallil et al., 2020; Prasad et al., 2020; Shechter et al., 2020)(Ayhan Başer et al., 2020; Koksall et al., 2020b; Korkmaz et al., 2020)(Şahin et al., 2020b)(Buselli et al., n.d.; Di Tella et al., 2020; Rossi et al., 2020)(AlAteeq et al., 2020; Thamer H Alenazi et al., 2020)

Sample size of the study ranged from 107 (Altintas, 2017) – 8817 (Xiaoming et al., 2020) with an average age of 19- 60< years.

3.3 Risk of Bias assessment/ Quality assessment

The quality of included articles was assessed using Newcastle-Ottawa Scale (NOS). To remove the risk of bias, quality assessment was done by two authors (RJ and MAK) independently. The quality was based on three categories i.e., selection, comparability and outcome consisting of 7 questions. In this, a study can be awarded a maximum of one star for each numbered item within the Selection and Exposure categories. A maximum of two stars can be given for Comparability. Later, stars of specific question can be added later or total score would be the number of stars given to the specific set of questions in a category. Scores ranging 7-10 were deemed to be of high quality, 6-7 represent medium quality and 1-6 were considered to be of low quality (**Table-1**).

Table 1

Quality of assessment table

Sno.	Author and year of study	Selection	Comparability	Outcome	Total	Quality of the study
1	(Altintas, 2017), 2017, Saudi Arabia	4	2	3	9	High
2	(Alshekaili et al., 2020b)Oman	4	2	3	9	High
3	(AlAteeq et al., 2020),2020, Saudi Arabia	4	2	3	9	High
4	(Thamer H Alenazi et al., 2020)2020, Saudi Arabia	5	2	2	9	High
5	(An et al., 2020)2020 China	5	2	3	10	High

6	(Antonijevic et al., 2020a)2020, Serbia	4	2	2	9	High
7	(Ayhan Başer et al., 2020)2020, TURKEY	4	1	2	7	Low
8	(Azoulay et al., n.d.)2020, France	5	1	3	9	High
9	(Bhargava et al., 2020)2020, Asia, North America, Central/ south America, Europe, Other.	4	2	2	8	High
10	(Buselli et al., n.d.)2020, ITALY	4	1	2	7	Low
11	(W. Cai et al., 2020), 2020, China	5	1	2	8	High
12	(Caillet et al., 2020a)2020, France	5	2	2	9	High

13	(Liu et al., 2020), 2020, China	5	2	2	9	High
14	(Civantos et al., 2020)2020, US	4	2	2	8	High
15	(Cheng et al., 2020).2020, China	4	1	3	8	High
16	(Chew et al., 2020)2020, Singapore	5	2	2	9	High
17	(Di Tella et al., 2020)2020, Italy	5	1	3	9	High
18	(Elhadi et al., 2020a)2020, Libya	4	2	2	8	High
19	(Fauzi et al., 2020)2020, Malaysia.	5	1	2	8	High
20	(Galopeni et al., 2020)2020, Kosovo	4	2	2	8	High
21	(G. Li et al., 2020)2020, China	4	1	3	8	High

22	(Herrero San Martin et al., 2020)2020, Spain	5	1	2	8	High
23	(Wang et al., 2020), 2020, China	5	2	3	10	High
24	(Hong et al., 2020)2020, China	5	2	2	9	High
25	(J. Zhu et al., 2020)2020, China	5	2	2	9	High
26	(Huang and Zhao, 2020)2020, China	5	1	2	8	High
27	(Janati Idrissi et al., 2020)2020, Morocco	5	1	1	7	Low
28	(Juan et al., 2020), 2020, China	4	2	2	8	High
30	(Kang et al., 2020a)China	5	1	2	8	High

31	(Kannampallil et al., 2020)USA	4	2	2	8	High
32	(Keubo et al., 2020), 2020, Cameroon	5	1	3	9	High
33	(Koksal et al., 2020a), 2020, Turkey	4	2	3	9	High
34	(Korkmaz et al., 2020)Turkey	4	2	2	8	High
35	(Labrague and De los Santos, 2020) Philippines	4	2	3	9	High
36	(J. Lai et al., 2020) , 2020, China	4	2	2	8	High
37	(Lee et al., 2020) Singapore	4	1	2	7	Low
38	(Liang et al., 2020), 2020 China	4	2	3	9	High
39	(Lu et al., 2020), 2020, China	4	1	2	7	Low

40	(Luceño-Moreno et al., 2020) 2020, Spain	4	2	2	8	High
41	(L. Q. Wang et al., 2020)2020, China	4	1	2	7	Low
42	(Nie et al., 2020) 2020, China	4	1	2	7	Low
43	(Ning et al., 2020)2020, China	5	1	2	8	High
44	(Pouralizadeh et al., 2020)2020, Iran	5	2	3	10	
45	(Prasad et al., 2020)2020, USA	4	1	3	8	High
46	(Q. Chen et al., 2020),2020, China	5	2	3	10	High
47	(Que et al., 2020) 2020, China	5	1	2	8	High

48	(R. Li et al., 2020), 2020, China	5	1	3	9	High
49	(Rossi et al., 2020)2020, Italy	4	1	2	7	Low
50	(Şahin et al., 2020a)2020, Turkey	5	1	2	8	High
51	(Salopek-Žiha et al., 2020)2020, Croatia	5	2	2	9	High
52	(Sandesh et al., 2020)2020, Pakistan	4	2	2	8	High
53	(Shah et al., 2020)2020, UK	5	1	3	9	High
54	(Shahrour, 2020)2020, Jordan	4	2	2	8	High
55	(Shechter et al., 2020) 2020, NYC	5	1	2	8	High

56	(Si et al., 2020), 2020, China	5	1	3	9	High
57	(Skoda et al., 2020)2020., Germany	5	2	3	10	High
58	(Stojanov et al., 2020)2020, Serbia	5	2	2	9	High
59	(Suryavanshi et al., 2020)2020, India	4	2	2	8	High
60	(Wang et al., 2020) 2020, China	4	1	2	7	Low
61	(Tian et al., 2020)2020, China	5	1	3	9	High
62	(Wasim et al., 2020), 2020, Pakistan.	4	2	2	8	High

63	(Wańkiewicz et al., 2020)2020, Poland	4	1	3	8	High
64	(Wilson et al., 2020), 2020, India	5	2	2	9	High
65	(X. Chen et al., 2020) 2020, China	5	2	2	9	High
66	(Xiao et al., 2020), 2020, China	4	1	2	7	Low
67	(Xiaoming et al., 2020)2020 China	5	2	3	10	High
68	(Xing et al., 2020), 2020, China	4	2	2	8	High
69	(Yang et al., 2020a)2020, China	5	2	2	9	High
70	(Yáñez et al., 2020), 2020, Peru	4	1	2	7	Low
71	(Y. Liu et al., 2020b), 2020, China	5	2	2	9	High

72	(Yin et al., 2020), 2020, China	4	1	2	7	Low
73	(Zhan et al., 2020), 2020, China	5	2	3	10	High
74	(Y. Wang et al., 2020a), 2020, China	5	1	3	9	High
75	(Yifang Zhou et al., 2020)China	4	2	2	8	High
76	(W. R. Zhang et al., 2020), 2020, China	5	1	3	9	High
77	(Z. Zhu et al., 2020a),2020, China	4	2	3	8	High
78	(Zhou et al., 2020)2020, China	4	2	2	8	High

3.4 Depression and Anxiety

Out of 78 studies, 56 articles discussed about the anxiety and depression, suffered by the FHW staff (**Table 2**).

The mental health data assessed by (An et al., 2020) reported prevalence of depression in junior nurses and tertiary hospital to be 66.1% and 89.8%, respectively. Another study reported that nurses suffered from depression and anxiety more as compared to doctors 71.8%, 16.7% and 70%, 19.1%, respectively. (Zhu et al, 2020). A similar study was conducted by Xiao et al., 2020 reported prevalence of depression/anxiety in nurses (62.4%/58.5%) more as compared to doctors (53.7%/49.5%). As per (Wankovicz et al., 2020), 99.51% FHW had PHQ-9 ≥ 4 and 99.03% staff had GAD-7 ≥ 4 . Another study used same scale and reported prevalence rate of depression and anxiety to be 90.92% and 73.46% (Pouralizadeh et al., 2020). Cross-sectional study conducted by Sahin et al., 2020 estimated that 77.6% of HCW had depression 60.2% had anxiety. Another study used DASS-21 scale to measure anxiety and depression and reported that 72.3% HCWs had depression whereas 85.7% had moderately- extreme severe anxiety (Sandesh et al., 2020). A study reported that 82.31% of HCW suffered from depression and 61.40 % had anxiety (Liang et al., 2020). Lee et al, 2020 reported HADS-D >11 was 39.3% and < 11 was 18.1% and HADS-A > 11 was 19.2% and <11 was 38.1% in physicians, HADS-D <11 was 11% and > 11 was 30.7% for HADS-A >11 was 11% and < 11 was 31.4% (Lee et al., 2020) (**Table 2**).

There were 7 studies which solely analyzed the occurrence of anxiety in FHW due to COVID-19. Alenazi et al., 2020 reported the overall prevalence of anxiety in nursing group three times (27.90%) than medicine group (9.91%). A similar study showed the prevalence of anxiety in nurses (35.98%) to be more as compared to doctors (25.26%) by having 35.98% of anxiety (X. Yang et al, 2020). Another study showed that overall 39.81% healthcare employees had anxiety, from which 53.3% suffered severe, 40.4% moderate and 38.2% mild

anxiety (Ayhan baser et al., 2020). Another study conducted in Philippines reported that 37.8% nurses had anxiety (Labrague and De los Santos, 2020). (Korkmaz et al., 2020) revealed that 70.71% healthcare workers experienced anxiety out of which 13% had severe anxiety, 20% had moderate and 38% mild levels of anxiety. According to Liang et al., 2020 and liu et al, 2020 HCWs experienced 12.5% of anxiety. (**Table 2**).

Table 2

Summary of characteristics of included studies

S. No	Study characteristics		Outcome characteristics				Findings		
	Study country (Year)	population size (Sex)	Type of mental health	Measurement tool (range of score)	Definition of outcomes	Measures	Total	By severity of outcome	% of morbidity in FL HCWs
1	(Altintas, 2017), 2020, Saudi Arabia	107 (M+F) Residents- 61.7% Fellow- 9.3% Registrar and consultants- 29.0%	Anxiety, depression, insomnia, stress	PHQ-9, GAD-7, ISI, PSS	PHQ-9 (0-27) GAD-7 (0-27) ISI-7 (0-21) PSS-10 (0-40)	Prevalence (%)	All healthcare workers Depression- 50.5% Anxiety- 46.5% Insomnia- 44.9%	PSS (Stress) Low- 28% Moderate- 68.2% High- 3.7% ISI (Insomnia) Sub-threshold- 29.9% moderate- 13.1%	Frontline healthcare workers PHQ-9- 31.9% GAD-7- 43.5% ISI- 18.8% PSS- 24.6%

medRxiv preprint doi: <https://doi.org/10.1101/2021.11.09.21266105>; this version posted November 11, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

								severe- 1.9% PHQ-9 (Depression) mild- 21.5% moderate- 17.8% moderately severe- 7.5% severe- 3.7% GAD-7 (Anxiety) mild- 25.2% moderate- 15.9% severe- 5.6%	
2	(Alshekaili et al., 2020b) Oman	1139 (M+F) Physicians- 384 Nurses- 449	Anxiety, depression, stress,	DASS-21, ISI	DASS-21 (0-42) ISI (0-28)	Prevalence (%)	NR	NR	Healthcare workers Depression-

		Allied health- 305	insomnia						32.3% Anxiety- 34.1% Stress- 23.8% Insomnia- 18.5%
3	(AlAteeq et al., 2020),2020, Saudi Arabia	502 (M+F) Physician- 111 Nurse- 132 Administrator- 143 Pharmacist- 12 Non-physician specialist- 70 Technician- 34	Depression, anxiety	GAD-7. PHQ-9	GAD-7 (0-21) PHQ-9 (0-27)	Prevalence (%)	NR	Depression Mild- 24.9% moderate- 14.5% moderately severe- 10% severe- 5.8% Anxiety mild- 25.1% moderate- 11% severe- 15.3%	Depression 55.20% Anxiety 51.40%

4	(Thamer H Alenazi et al., 2020)2020, Saudi Arabia	4920 (M+F) Medicine- 734 Nursing- 1913 Pharmacy- 580 Medical laboratories- 365 Dentistry- 299 Radiology- 193 Physical therapy- 107 Respiratory therapy- 52 Others- 677	Anxiety	GAD-7, PHQ-9	GAD-7 (0-21) PHQ (0-27)	Prevalence (%)	Total anxiety- 68.45%	Medicine group Anxiety Low- 15.9% Medium – 15.3% High- 13.6% Nursing Anxiety low- 34.8% medium – 36.2% high- 45.9% Dentistry Anxiety low- 6.1% medium – 6.2% high- 5.6%	Medicine group Anxiety- 9.91% Nursing group Anxiety- 27.90%
---	---	--	---------	--------------	-------------------------	----------------	---------------------------------	---	--

5	(An et al., 2020) China	1103 (M) ED nurses- 43.61%	Depression	PHQ-9, QoL	PHQ-9 (0-27) QOL (16-112)	Prevalence (%)		Depression Mild- 27.7% Moderate- 8.6% Moderate-severe- 5.3% Severe- 2.1%	Junior nurses Depression- 66.1% No depression 69.0% Participation in treating patients- Depression- 16.4% No depression 16.9% Working in tertiary hospitals- Depression- 89.8% no depression
---	-----------------------------------	---	------------	------------	------------------------------	----------------	--	---	---

									85.0% working inpatient services- Depression- 36.8% No depression- 32.2%
6	(Antonišević et al., 2020a) Serbia	1678 (M+F) Medical personnel- 684 Others- 994	Anxiety, depression, stress, resilience	PSS, GAD-7, BDI-IA	GAD-7 (0-21) PSS (0-40) BDI-IA (0-63)	Prevalence (%)	Stress- 62.51% Anxiety- 52.26% Depression- 24.97% Resilience- 55.36%	Frontline medical personnel Stress low- 25.15% moderate- 41.72% high- 33.13% Anxiety minimal- 28.39% mild- 33.35% moderate-	Frontline medical personnel Stress- 15.36% Anxiety- 14.44% Depression- 6.07% Resilience- 10.33%

								<p>16.13%</p> <p>severe- 21.94%</p> <p>Depression</p> <p>minimal- 68.24%</p> <p>mild- 18.24%</p> <p>moderate- 10.14%</p> <p>severe- 3.38%</p> <p>Resilience</p> <p>low- 44.83%</p> <p>moderate- 40.00%</p> <p>high- 15.17%</p>	
7	(Ayhan Başer et al., 2020) 2020, TURKEY	1070 (M+F) Health employees-426 Others-644	Anxiety	BAI	BAI (0-63) Min- 0-7 Mild- 8-15 Moderate- 16-	Prevalence (%)		<p>Healthcare employees</p> <p>ANXIETY</p> <p>Mild- 38.2%</p> <p>Moderate- 40.4%</p>	<p>Healthcare employees</p> <p>Anxiety- 39.8%</p>

medRxiv preprint doi: <https://doi.org/10.1101/2021.11.09.21266105>; this version posted November 11, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

					25 Severe- 26- 63			Severe- 53.3%	
8	(Azoulay et al., n.d.) 2020, France	1058 (F) Nurses- 498 Nursing Assistants- 223 Senior Physicians- 204 Residents- 78 Medical students- 25 other professionals- 27	Anxiety, Depression, Peritraumatic dissociation	HADS, PDEQ	PDEQ (0-4) HADS (0-21)	Prevalence (%)	All Healthcare workers Anxiety- 50.4% Depression- 30.4% peritraumatic dissociation- 32%	NR	Nurses Anxiety- 50% Depression- 31.6% Peritraumatic dissociation- 40% Senior physicians Anxiety- 46.6% depression- 25% Peritraumatic dissociation- 20% residents Anxiety- 41%

									Depression- 19.2% Peritraumatic dissociation- 15.4%
9	(Bhargava et al., 2020)2020, Asia, North America, Central/ south America, Europe, Other.	733 (M+F) Dermatologists-733	Anxiety, depression, insomnia, stress, irritation	Survey questions were made and survey instrument was sent electronically. Participants accessed the survey via the link.	NR	Prevalence (%)	Total psychological outcomes- 77.2%	NR	Dermatologists Stress- 73.9% Irritability- 30.7% Insomnia- 30.8% Depression- 27.6%
10	(Buselli et al., n.d.)2020,	265 (M+F) Medical doctors- 85	Depression, anxiety	ProQOL-5, GAD-7, PHQ-	GAD-7 (0-21) PHQ-9	Prevalence (%)	Prevalence (%)	NR	Frontline HCWs Anxiety- 23.2%

	ITALY	Nurses- 133 Healthcare assistants- 47		9	(0-27)				Depression- 22.8%
11	(W. Cai et al., 2020), 2020, China	1521 (M+F) Doctor- 511 Nurses- 546 Technician- 45 Pharmacist- 72	Somatization, obsessive-compulsion, depression, anxiety etc.	SLC-90, CD-RISC, SSRS.	SCL-90 (0-4) CD-RISC (0-100) SSRS (0-50)	Prevalence (%)		NR	Frontline healthcare workers Prevalence of psychological abnormality Doctor- 17.1% Nurse- 13.2%

12	(Caillet et al., 2020a)2020, France	208 (M+F) Nurse anesthetics- 15 Care assistant- 41 Surface technician- 6 Anesthesia nurse student- 1 Nurse student- 12 Medical student- 8 Nurse- 84 Intern- 11 Physiotherapist- 5 Physicians- 17 Secretary- 4	Anxiety, Depression, PTSD	HADS, IES-R	IES-R (0-88) HADS (0-21)	Prevalence (%)	All Healthcare workers Anxiety- 48% Depression- 16% PTSD- 27% Burnout- 10.0%	NR	Nurse Anxious- 55% Non-anxious- 45% Physicians- Anxious- 41% Non-anxious- 59%
----	-------------------------------------	---	---------------------------	-------------	-----------------------------	----------------	---	----	--

It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/).

13	(Liu et al., 2020), 2020, China	512 (M+F) Clinical department- 369 Fever clinic- 68 Service/managerial- 75	Anxiety	SAS	SAS (20-80)	Prevalence (%)		Anxiety Normal- 87.5% Mild- 10.35% Moderate- 1.36% Severe- 0.78%	Healthcare workers Anxiety- 12.5%
14	(Civantos et al., 2020)2020, US	349 (M+F) Residents= 165 Attending physicians= 184	Anxiety, depression, burnout	Mini Z burnout scale, GAD-7, IES-R	GAD-7 (0-21) IES-R (0-23) PHQ-2 (0-6) Mini Z assessment (0-5)	Prevalence (%)		(Residents and Attending physicians) Mini Z burnout symptoms Negative- 78.2% Positive- 21.8% GAD-7 Normal- 52.1% Mild- 28.9%	Residents Burnout symptoms- 29.7% Anxiety- 50.3% Distress- 58.2% Depression- 10.3% Attending physicians Burnout

								<p>Moderate- 11.5%</p> <p>Severe- 7.4%</p> <p>IES</p> <p>Subclinical- 39.8%</p> <p>Mild- 32.7%</p> <p>Moderate- 20.9%</p> <p>Severe- 6.6%</p> <p>PHQ-2</p> <p>Positive- 10.6%</p> <p>Negative- 89.4%</p>	<p>symptoms- 14.7%</p> <p>Anxiety- 45.7%</p> <p>Distress- 62.0%</p> <p>Depression- 10.9%</p>
--	--	--	--	--	--	--	--	--	--

15	(Cheng et al., 2020). 2020, China	534 (M+F) Doctors- 289 Nurses- 245	Anxiety	SAS, PSQI	SAS (20-75), PSQI (0-21)	Prevalence (%)		Healthcare workers ANXIETY Mild- 10.7% Moderate- 2.6% Severe- 0.7%	Healthcare workers Anxiety- 14.0% Poor sleep 30.0%
16	(Chew et al., 2020) 2020, Singapore	906 (M+F)	Anxiety, depression, stress, psychological distress, PTSD	DASS-21	DASS-21 (0-42)	Prevalence (%)		Depression Moderate-severe- 5.3% anxiety Moderate-severe- 8.7% stress Moderate-severe- 2.2% psychological distress Moderate-	Healthcare workers Depression- 10.6% Anxiety- 15.7% Stress- 5.2% PTSD- 7.4%

								severe- 3.8%	
17	(Di Tella et al., 2020) Italy	145 (M+F)	PTSS, Depression	STAIY1, BD-II, PCL-5	STAI Y1 (20-80) BD-II (0-60)	Prevalence (%)	Healthcare workers STAI Y1- 71.00% BDI-II- 31.00% PCL-5- 26.2%		COVID Workers Psychological outcomes STAI Y1- 5.5% BDI-II- 13.7% PCL-5- 29.2%
18	(Elhadi et al., 2020a) Libya	745 (M+F) Medical- 265 Surgical- 150 Emergency medicine- 136 Obstetrics and	Anxiety, depression	HADS	HADS (0-21)	Prevalence (%)		NR	Healthcare workers Anxiety symptoms- 46.7% Depressive symptoms- 56.3%

		gynecology- 49 Pediatrics- 76 ICU- 69							
19	(Fauzi et al., 2020)2020, Malaysia.	1050 (M+F)	Depression, Anxiety, Stress	NASA-TLX, DASS-21	NASA-TLX (1-20) DASS-21 (0-42)	Prevalence (%)		NR	Healthcare workers Depression- 30.95% Anxiety- 29.7% Stress- 23.52%

20	(Galopeni et al., 2020)2020, Kosovo	592 (M+F) Nurses- 51.4%	Anxiety, depression	HADS	HADS (0-21) Normal- 0-7 Borderline- 8-10 Abnormal- 11-21	prevalence (%)		Physicians anxiety symptoms normal- 3.0% borderline- 20.8% abnormal- 18.9% Depressive symptoms Normal- 3.2% borderline- 20.4% abnormal- 49.3% Nurses Anxiety symptoms normal- 4.9% borderline- 23.3% abnormal- 23.1%	Healthcare workers Anxiety - 44.6% Depression- 38.7%
----	-------------------------------------	----------------------------	---------------------	------	---	----------------	--	--	---

								<p>Depressive symptoms</p> <p>Normal- 5.4%</p> <p>borderline- 28.0%</p> <p>abnormal- 17.9%</p>	
21	(G. Li et al., 2020) 2020, China	<p>4396 (F)</p> <p>Medical technician- 407</p> <p>Nurse- 3380</p> <p>Doctor- 582</p>	Stress, anxiety, Depression	GAD-7, PHQ-9, IES-R	<p>GAD-7(0-21) PHQ-27 (0-27)</p> <p>IES-R (0-88)</p>	Prevalence (%)	NR	NR	<p>Doctors</p> <p>Stress- 27.31%</p> <p>Non-stress- 72.69%</p> <p>Nurses</p> <p>Stress- 33.34%</p> <p>Non-stress-</p>

									66.66%
22	(Herrero San Martin et al., 2020) 2020, Spain	170 (M+F) Healthcare workers- 100 Non- Healthcare workers- 70	Insomnia	ESS, PSQI, ISI, HARS	ESS (0-24) PSQI (0-21) ISI (0-28) HARS (0-56)	Prevalence (%)	Total Insomnia- 47.65%	NR	Healthcare workers Insomnia- 57%
23	(Wang et al., 2020), 2020, China	1045 (M+F) Doctors- 149 Nurse- 773 Others- 123	Anxiety, depression, insomnia	HADS, PSS, ISI	HADS (0-21) PSS ISI (0-28)	Prevalence (%)	Healthcare workers (high risk group) Anxiety- 55.4% Depression- 43.6% Insomnia-	NR	Doctors Anxiety (HADS-A>10)- 21.1% depression (HADS-D >10)- 17.4% Stress (PSS>29)- 24.2%

							13.5%		<p>Insomnia (ISI>14)- 11.4%</p> <p>Nurses</p> <p>Anxiety (HADS-A>10)- 8.6%</p> <p>Depression (HADS-D)- 13.6%</p> <p>Stress (PSS>29)- 21.7%</p> <p>Insomnia (ISI>14)- 11.4%</p>
24	(Hong et al., 2020) China	4692 (M+F) Nurses' isolation ward- 381 non-isolation	depression, anxiety, somatic symptoms, suicidal	PHQ-9, GAD-7, PHQ-15	PHQ-9 (0-4) GAD-7 (0-3) PHQ-15	Prevalence (%)	NR	NR	<p>Healthcare workers</p> <p>Depression- 9.4%</p> <p>Anxiety- 8.1%</p>

		ward- 4311 direct care of the patients- 506	ideation						Somatic symptoms- 42.7% Suicidal ideation- 6.5%
25	(J. Zhu et al., 2020) China	165 (M+F) Doctors- 70 Nurses- 86	anxiety, depression	SAS, SDS		Prevalence %	NR	NR	Doctors Anxiety- 11.4% Depression- 45.6% Nurses Anxiety- 27.5% Depression- 43.0%
26	(Huang and Zhao, 2020) China	7,236 (M+F) HW-2250(31.1%) Teachers or students- 1404(19.4%) Enterprise or	Anxiety symptoms, depression symptoms, sleep quality	GAD-7, CES-D, PSQI Scale,	GAD-7 (0-21) CES-D (0-60 points) PSQI scale (0-21)	Prevalence (%)	Total psychological outcomes Anxiety symptoms -	NR	Based occupation (group-1) Depression- 35.8%

		institution workers- 1809(25.0%) Another occupation- 1773(24.5%)					35.1% depressive symptoms- 20.1% sleep quality- 18.2%		Anxiety- 19.8% Poor sleep quality- 73.6%
27	(Janati Idrissi et al., 2020) 2020, Morocco	827(M+F) Healthcare workers- 282 institution workers- 112 teachers or students- 305 others- 115	sleep disorders, anxiety, depression	DBAS-16, AIS, ESS, HARS, BDI	ESS (0-24), HAS (0-56) BDI(0-16)	Prevalence (%)	NR	NR	Healthcare workers Insomnia- 56.0% Anxiety- 29.5% Depression- 35.6%
28	(Juan et al., 2020), 2020, China	456 (M+F) Doctors- 195 (42.8) Nurses- 261 (57.2)	depression, anxiety, somatization obsessive compulsive	IES-R (0-23) PHQ-9 (0-27) GAD-7 (0-21) PHQ-15 (0-30) Y-BOCS (0-	IES-R (0- 23) PHQ-9 (0- 27)	prevalence (%)		Doctors IES-R- Sub-clinical- 54.9%	Doctors IES-R- 19.29% GAD-7- 13.81%

			disorder	40)	GAD-7 (0-7)			mild- 34.4%	PHQ-9- 12.93%
					PHQ-5 (0-30)			moderate-severe- 10.8%	Y-BOCS- 15.78%
					Y-BOCS (0-40)			GAD-7 None- 67.7%	PHQ-15- 15.13%
								mild- 26.2%	Nurses
								moderate-severe- 6.2%	IES-R- 23.9%
								PHQ-9 None- 69.7%	GAD-7- 17.7%
								mild- 23.1%	PHQ-9- 16.6%
								moderate-severe- 7.2%	Y-BOCS- 28.28%
								Y-BOCS None- 63.1%	PHQ-15- 18.2%
								mild- 31.8%	
								moderate-	

29	(Johnson et al., 2020) Norway	1773 (M+F)	PTSD, depression, anxiety	PCL-5, PHQ-9, GAD-7, CAS-1	PCL-5 (0-80) GAD-7 (0-21) PHQ-9 (0-27)	Prevalence (%)	NR	Direct contact group PCL-5 non-clinical- 63.4% subclinical- 22.5% PTSD- 14.1% PHQ-9 Normal- 35.6% Mild- 38.3% Moderate- 15.4% Moderate severe- 8.1% Severe- 2.7% GAD-7 Normal- 51.0% Mild- 25.2% Moderate- 17.8% Severe- 6.0%	Healthcare workers PTSD- 28.9% Anxiety- 21.2% Depression- 20.5%
----	--------------------------------------	-------------------	---------------------------	----------------------------	--	----------------	----	--	---

30	(Kang et al., 2020a) China	994 (M+F) Physicians- 183 Nurses- 811	Depression, Anxiety, Stress	GAD-7, PHQ-9, ISI, IES-R	PHQ-9 (0-27) GAD-7 (0-21) ISI-7 (0-21) IE-22 (0-88)	Prevalence (%)	NR	NR	Healthcare workers Mental Health disturbances Sub-threshold 36.9% mild- 34.4% moderate- 22.4% severe- 6.2%
31	(Kannampallil et al., 2020) USA	393 (F) Fellow-132 Residents- 261	Depression, anxiety, stress, Burnout,	DASS-21, PFI	DASS-21 (0-42)	prevalence (%)	NR	NR	Physician trainees Depression- 27.2% Anxiety- 18.6% Stress- 24.7% Burnout- 40.7%

32	(Keubo et al., 2020), 2020, Cameroon	292 ALP- AN- N- MD- Psy- MLT- HT- 65	(M+F) 2 19 168 74 5 18	Anxiety, Depression	HADS	HADS (0-21)	Prevalence (%)	NR	Medical doctors (MD) Anxiety Absent- 40.5% Low- 23.0% moderate- 24.3% Severe- 12.2% Depression Absent- 33.8% Low- 27.0% Moderate- 24.3% Severe- 12.2% Nurses (n) Anxiety Absent- 34.5% Low- 26.3%	All healthcare workers Mild to severe anxiety- 42.20% Depression- 43.50%
----	--------------------------------------	---	--	------------------------	------	-------------	----------------	----	--	---

								<p>Moderate- 42.1%</p> <p>Severe- 15.8%</p> <p>Depression</p> <p>Absent- 22.6%</p> <p>Low- 21.1%</p> <p>Moderate- 36.8%</p> <p>Severe- 15.8%</p>	
33	(Koksal et al., 2020a), 2020, Turkey	<p>702 (M+F)</p> <p>Anesthetic technician- 363</p> <p>Operating theater nurse- 339</p>	Anxiety, depression	HADS	HADS (0-21)	prevalence (%)	Total Prevalence- 49.32%	NR	<p>Nurse Depression</p> <p>Yes- 50.2%</p> <p>No- 47.2%</p> <p>Anxiety</p> <p>Yes- 48.8%</p> <p>No- 47.7%</p>

34	(Korkmaz et al., 2020) Turkey	140 (M+F) Physicians- 30 nurses- 70 Assistant healthcare staff- 40	Anxiety	PSQI, BAI, PSI WHOQOL-BREF	PSQI (0-21) BAI (0-63) PSI WHOQOL-BREF (0-100)	Prevalence (%)	Total Anxiety Total- 33%	Healthcare workers Anxiety mild- 38% moderate- 20% severe- 13%	Healthcare workers Anxiety- 70.7%
35	(Labrague and De los Santos, 2020) Philippines	325 (F) Nurses- 325	Anxiety	BRCS, PSSQ, POS	BRCS (4-20)	Prevalence (%)	NR	NR	Nurses Anxiety- 37.8%

36	(J. Lai et al., 2020) , 2020, China	1257 Physicians- Nurses-	(M+F) 493 764	Depression, anxiety, insomnia, distress.	PHQ-9, GAD-7, ISI, IES-R	PHQ-9 (0-27) GAD-7 (0-21) ISI (0-28) IES-R (0-23)	Percentage (%)	Healthcare workers PHQ-9- 50.43% GAD-7- 44.55% ISI- 33.96% IES-R- 71.51%	Frontline Healthcare workers PHQ-9 Normal- 41.5% mild- 40.4% moderate- 11.3% severe- 6.7% GAD-7 Normal- 48.4% mild- 35.4% moderate- 9.1% severe- 6.8% ISI Absence- 59.3% subthreshold-	Frontline Healthcare workers PHQ-9- 24.26% GAD-7- 21.40% ISI- 16.86% IES-R- 31.66%
----	-------------------------------------	--------------------------------	---------------------	---	--------------------------	--	----------------	---	---	---

								28.3% moderate- 10.5% severe- 1.7% IES-R Normal- 23.7 mild- 34.0% moderate- 27.9% severe- 14.1%	
37	(Lee et al., 2020) Singapore	270 (M+F) Antitheologist- 42.6% Nurses- 57.4%	Anxiety, depression	GHQ-12, HADS	GHQ-12 (0-36) HADS (0-21)	Prevalence (%)	Healthcare workers HADS-A <11- 69.6% >11- 30.4% HADS-D <11- 70.0%	NR	Physicians HADS-A <11- 31.4% >11- 11.1% HADS-D >11- 30.7% <11- 11.9%

							>11- 30.0%		<p>Nurse</p> <p>HADS-A</p> <p><11- 38.1%</p> <p>>11- 19.2%</p> <p>HADS-D</p> <p>>11- 39.3%</p> <p><11- 18.1%</p>
38	(Liang et al., 2020), 2020 China	<p>899 (M+F)</p> <p>Medical workers- 899</p> <p>Hubei province- 138</p> <p>Other regions- 761</p>	Depression, anxiety, insomnia, resilience	PHQ-9, GAD-7, ISI, CD-RISC-10	PHQ-9 (0-27) GAD-7 (0-21) ISI (0-28) CD-RISC-10 (0-40)	Prevalence (%)	NR	<p>Medical workers</p> <p>Depression</p> <p>Mild- 58.28%</p> <p>Moderate- 24.24%</p> <p>Anxiety</p> <p>Mild- 47.16%</p> <p>Moderate-</p>	<p>Medical workers</p> <p>Depression- 82.31%</p> <p>Anxiety- 61.40%</p> <p>Insomnia- 43.04%</p>

								14.23%	
								Insomnia	
								Mild- 31.81%	
								Moderate- 11.23%	
39	(Lu et al., 2020), 2020, China	2299(M+F) Medical staff- 2042 Administrative staff- 257	Fear, anxiety, depression	NRS, HAMA, HAMD	NRS (0-10) HAMA (0-70) HAMD (0-85)	prevalence (%)	NR	Medical staff Proportion of fear No/mild- 29.4% Moderate- 43.9% Severe/extreme- 26.7% proportion of anxiety no- 74.5% mild/ moderate- 22.6%	Medical staff Fear scale 62.67% HAMA 22.66% HAMD- 10.74%

								severe/ extreme- 2.9%	
								Proportion of depression	
								No- 87.9%	
								mild/ moderate- 11.8%	
40	(Luceño-Moreno et al., 2020) 2020, Spain	1422 (M+F) Medical staff- 143 Nursing staff- 486 Assistant nurse- 443 Caregiver- 117 Others- 233	PTSD, Depression, Anxiety	HADS, IES-R	HADS (0-21) IES-R (0-23)	Prevalence (%)	NR	NR	Healthcare workers Anxiety (HADS)- 79.32% Depression (HADS)- 51.33% Post-traumatic stress (IES-R)- 82.98%

41	(L. Q. Wang et al., 2020) 2020, China	274 (M+F) Doctors- 40.8% Nurses- 59.2%	Anxiety, Depression, Sleep quality, stress	GAD-7, PSQI, PSS-14, CD- RISC-10	GAD-7 (0-21) PSQI (0-21) PSS-14 (0-56) CD-RISC (0-100)	Prevalence (%)	NR	NR	Healthcare workers Anxiety- 13.9% Depression- 16.1% Insomnia- 19.7%
42	(Nie et al., 2020) 2020, China	263 (M+F) Emergency department- 196 non-emergency department-67	Psychological distress	GHQ-12, SCSQ, IES-R, PSSS		Prevalence (%)			Frontline nurses Psychological distress- 25.4%
43	(Ning et al., 2020) 2020, China	612 (M+F) Doctors- 317	anxiety, depression	SAS, SDS	SAS (20-80) SDS (≥ 53)	Prevalence (%)	NR	NR	Nurses anxiety- 20.3% Depression-

		Nurses- 295							30.2% Doctors Anxiety- 12.6% Depression- 20.2%
44	(Pouralizadeh et al., 2020) Iran	441 (M+F) Certified nursing assistant- 2 licensed practical nurse- 6 registered nurse- 340 head nurse- 49 supervisor- 40	anxiety, depression	GAD-7, PHQ-9	GAD-7 (0-4)	Prevalence (%)	NR	Healthcare workers GAD-7 no anxiety- 26.5% mild- 34.7% moderate- 19.7%	Healthcare workers GAD-7- 73.4% PHQ-9- 90.9%

45	(Prasad et al., 2020)2020, USA	347 (M+F) Nurses- 248 Administrative staff-63 Advanced practice provider- 36	Distress, burnout, anxiety, depression	Mini Z assessment, IES, PHQ-2	IES-R (0-23) PHQ-2 (0-6) Mini Z assessment (0-5)	Prevalence (%)	NR	Nurse Distress Sub-clinical- 14.1% Mild- 22.2% Moderate- 38.3% Severe- 25.4% Burnout Negative- 66.5% Positive- 33.5% Anxiety Normal- 25.8% Mild- 39.9% Moderate- 21.4% Severe- 12.9%	Nurses Burnout- 30% Anxiety- 69.5% Distress- 84.1% Depression- 22.8%
----	--------------------------------	---	--	-------------------------------	--	----------------	----	---	---

								<p>Depression</p> <p>Negative- 78.2%</p> <p>Positive- 21.8%</p>	
46	(Q. Chen et al., 2020),2020, China	<p>902 (M+F)</p> <p>Doctors- 543</p> <p>Nurses- 311</p> <p>Others-48</p>	Anxiety, stress, depression	CMBI, TCSQ, GAD-7, PHQ-9	GAD-7(0-21) PHQ-27 (0-27)	Prevalence (%)	<p>Healthcare workers moderate/severe anxiety and Depression- 24.50%</p>	<p>Doctors Anxiety</p> <p>None- mild- 82.82%</p> <p>Moderate-severe- 17.68%</p> <p>depression</p> <p>None- mild- 81.40%</p>	<p>Healthcare workers</p> <p>Moderate-severe- 18.99%</p> <p>Anxiety- 18.99%</p> <p>Moderate-severe- depression- 16.63%</p>

								<p>Moderate-severe- 18.60%</p> <p>Nurse Anxiety</p> <p>none- mild- 85.53%</p> <p>Moderate-severe- 14.47%</p> <p>depression</p> <p>None- mild- 82.96%</p> <p>Moderate - severe- 17.04%</p>	
47	(Que et al., 2020) 2020, China	2285 (M+F) Medical residents- 913 Nurses- 208 physicians- 860 Technicians- 179	anxiety, depression, insomnia	GAD-7, PHQ-9, ISI	GAD-7 (0-21) PHQ-9 (0-27) ISI (0-28)	Prevalence (%)	Overall psychological problems- 56.58%	<p>Healthcare workers</p> <p>Anxiety</p> <p>No- 53.96%</p>	<p>Healthcare workers</p> <p>Anxiety- 46.04%</p> <p>Depression-</p>

		public health practitioners- 125						<p>Mild- 34.44%</p> <p>Moderate/severe- 11.60%</p> <p>Depression No- 55.62%</p> <p>Mild- 31.55%</p> <p>Moderate/severe- 12.82%</p> <p>Insomnia No- 71.25%</p> <p>Sub-threshold- 21.97%</p> <p>Moderate/severe- 6.78%</p>	<p>4437%</p> <p>Insomnia- 28.75%</p> <p>Other psychological problems- 56.59%</p>
--	--	----------------------------------	--	--	--	--	--	--	--

48	(R. Li et al., 2020), 2020, China	176 (M+F) Senior nurse- 120 Nurse-in-charge- 56	Anxiety	HAMA	HAMA (0-10)	Prevalence (%)		Anxiety mild- 27.3% moderate- 25% severe- 25%	Frontline Nurses Anxiety- 77.3%
49	(Rossi et al., 2020)2020, Italy	1379 (M+F) Nurse- 472 Physicians- 433 General practitioner- 86 Others- 275 HCA- 112	anxiety, depression, insomnia	GPS, PHQ-9, GAD-7, ISI PSS	PHQ-9 (0-27) GAD-7 (0-21) ISI (0-28) PSS (0-40)	Prevalence (%)	NR	NR	Healthcare workers GPS-PTSD Score ≥ 3 - 49.38% PHQ-9 Score 15- 24.73% GAD-7 Score 15- 19.80% ISI Score ≥ 27 - 8.27% PSS Score, first quartile- 21.90%

50	(\$ahin et al., 2020a)2020, Turkey	620 (M+F) Physicians- 569 Nurses- 254 Others- 105 Frontline- 569 Second line- 370	Depression, anxiety, insomnia and distress	PHQ-9, GAD-7, ISI, IES-R	PHQ-9 (0-27) GAD-7 (0-21) ISI (0-28) IES-R (0-88)	Prevalence (%)	NR	Frontline Healthcare workers PHQ-9 Normal- 19.7% Mild- 38.5% Moderate- 25.7% Moderately severe- 9.5% Severe- 6.7% GAD-7 Normal- 37.6% Mild- 44.1% Moderate- 10.5% severe- 7.7% ISI	Healthcare workers Depression- 77.6% Anxiety- 60.2% Insomnia- 50.4% Distress- 76.4%
----	------------------------------------	---	--	--------------------------	--	----------------	----	---	--

								<p>Normal- 46.7%</p> <p>Sub- threshold- 38.5%</p> <p>Moderate- 12.7%</p> <p>Severe- 2.1%</p> <p>IES-R</p> <p>Normal- 22.1%</p> <p>Mild- 45.2%</p> <p>Moderate- 19.9%</p> <p>Severe- 12.8%</p>	
51	(Salopek-Žiha et al., 2020)2020, Croatia	124 (M+F) Physicians- 27 Nurses- 97	depression, stress, anxiety	DASS-21	DASS-21 (0-42)	Prevalence (%)	NR	NR	<p>Healthcare workers</p> <p>Anxiety</p> <p>moderate- ver</p> <p>severe- 11%</p> <p>moderate- extremely severe</p>

medRxiv preprint doi: <https://doi.org/10.1101/2021.11.09.21266105>; this version posted November 11, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

									17% Stress moderate- extremely severe 10% 67% Worry
52	(Sandesh et al., 2020) 2020, Pakistan	112 (M+F)	Depression, anxiety, stress	DASS-21	DASS-21 (0-42)	Prevalence (%)	NR	NR	Healthcare workers Depression Moderate/ severe 72.3% Anxiety moderate- extremely severe 85.7% stress Moderate- extremely severe

									90.1%
53	(Shah et al., 2020) 2020, UK	207 (M+F) Consultant- 58 Registrar- 99 SHO- 50	Anxiety, Depression	GAD-2, PHQ-2	PHQ-2 (0-6) GAD-2 (0-21)	Prevalence (%)	total GAD-2 >3- 24.64% PHQ-2 ≤2- 15.94%	NR	Consultant GAD-2≤2- 69.70% GAD-2≥2- 30.30% PHQ-2≤ 80.81% PHQ-2≥2- 19.19%
54	(Shahrour, 2020) 2020, Jordan	448 (M+F) Nurses- 73%	Acute stress disorder, PTSD	SASRQ, BSI-18, GSI	BSI (0-72) SASRQ (0-150) GSI (0-45)	Prevalence (%)	NR	NR	Nurses Acute stress disorder- 64%

55	(Shechter et al., 2020) 2020, NYC	657 (M+F) Attending physicians- 141 Resident/fellow- 141 Nurses/AP providers- 375	Acute stress disorder, depression, anxiety	PC-PTSD, PHQ-2, GAD-2	PC-PTSD (0- 4) PHQ-2 (0- 6) GAD-2 (0- 21)	Prevalence (%)	All participants Acute stress- 56.6% Depression- 47.64% Anxiety- 29.52%	NR	<p>Attending physicians</p> <p>Acute stress disorder- 40% Depression- 38% Anxiety- 15%</p> <p>Residents/fellows</p> <p>Acute stress disorder- 54% Depression- 44% Anxiety- 17%</p> <p>Nurses/ap providers</p> <p>Acute stress disorder- 64% Depression- 53% Anxiety- 40%</p>
----	---	---	--	-----------------------	---	----------------	--	----	---

56	(Si et al., 2020), 2020, China	863 (M+F) Doctors- 377 Nurse- 211 Other HCWs- 275	PTSD, Depression, anxiety, stress	IES-R, DASS-21	DASS-21 (0-42) IES-R (0-23)	Prevalence (%)	NR	NR	Healthcare workers Depression- 13.6% Anxiety- 13.9% Stress- 8.6% Post-traumatic stress- 40.2%
57	(Skoda et al., 2020)2020., Germany	2224 (M+F) physicians- 492 nursing staff- 1511 paramedics- 221 non-healthcare professionals- 10639	Anxiety, Depression, fear	GAD-7, PHQ-2, EQ-5D-3L	GAD-7 (0-21) PHQ-2 (0-6) EQ-5D-3L (0-100)	Prevalence (%)	NR	NR	Physicians GAD-7<10- 94.10% GAD-7 ≥10- 5.89% Nurses GAD-7 <10- 88.41% GAD-7 ≥10-

									11.38%
58	(Stojanov et al., 2020)2020, Serbia	201 (M+F)	Anxiety, depression, sleep quality	GAD-7, SF36, PSQI	GAD-7 (0-21) PSQI (0-21)	Prevalence (%)	NR	NR	Healthcare workers Anxiety GAD-7≥10- 31.8% Depression SDS Score 66 17.6%
59	(Suryavanshi et al., 2020)2020, India	197 (M+F) Physicians- 66 Nurses- 47	Depression, anxiety	PHQ-9, GAD-7, QoL-1	PHQ-9 (0-27) GAD-7 (0-21)	Prevalence (%)	In all healthcare workers	NR	Physicians Depression- 8% Anxiety- 30% Depression and Anxiety combined- 15% Depression- 47% Anxiety- 50%

									<p>Nurses</p> <p>Depression- 28%</p> <p>Anxiety- 25%</p> <p>Depression and Anxiety combined- 12%</p>
60	(Wang et al., 2020) 2020, China	123 (M+F) Pediatric healthcare workers- 123		PSQI, SAS, SDS	PSQI (0-21), SAS (0-4) SDS (>53)	Prevalence (%)	Sleep disturbances - 38%	NR	<p>Doctors</p> <p>PSQI >7- 62%</p> <p>PSQI ≤ 7- 25%</p> <p>Nurses</p> <p>PSQI >7- 38%</p> <p>PSQI ≤ 7- 75%</p>
61	(Tian et al., 2020) 2020, China	845 (M+F)	depression, anxiety, stress,	PSS-10, GAD-7, ISI-7,	PSS-10 (0-40) GAD-7 (0-	prevalence (%)	NR	NR	<p>Frontline healthcare workers</p>

			insomnia		21) ISI (0-28)				Depression- 45.6% Anxiety- 20.7% Insomnia- 27.0% Moderate- severe stress- 60.8%
62	(Wasim et al., 2020), 2020, Pakistan.	356 (M+F) Doctors- 214 Nurses-78 Paramedics- 17.97%	insomnia, depression, anxiety, stress	ISI, DASS-21	ISI (0-28) DASS-21 (0-42)	Prevalence (%)	Depression- 62.35% anxiety- 63.33% insomnia- 53.37% stress- 55.33%	Healthcare workers Insomnia Depression none/min- 8.42% mild- 76.9% moderate- 13.15% severe- 1.6% Anxiety None/min-	DOCTORS Insomnia- 50.52% non- insomnia- 71.1% NURSES Insomnia- 24.21% non- insomnia- 19.3%

								16.9%	
								Mild- 63.15%	
								Moderate- 14.21%	
								Severe- 5.8%	
								Stress	
								None/min- 15.8%	
								Mild- 54.78%	
								Moderate- 26.9%	
								Severe- 2.63%	
63	(Wańkiewicz et al., 2020) 2020, Poland	441 (M+F) Frontline HCWs- 206 Second-line HCWs- 135	Anxiety, depression, insomnia	GAD-7, PHQ-9, ISI	GAD-7 (0-27) PHQ-9 (0-21) ISI (0-28)	Prevalence (%)	NR	NR	Frontline Healthcare workers GAD-7 GAD-7≤4- 0.97% GAD-7 ≥5

medRxiv preprint doi: <https://doi.org/10.1101/2021.11.09.21266105>; this version posted November 11, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

									99.03% PHQ-9 PHQ-9 ≤4- 0.49% PHQ-9 >4- 99.51% ISI ISI ≤8- .00% ISI >8- 100.00%
64	(Wilson et al., 2020), 2020, India	350 (M+F) Doctor- 295 Nurses- 55	Stress, depression, anxiety	GAD-7, PHQ-9, PSS-10	GAD (0-21) PHQ-9 (0-21) PSS-10 (0-47)	Prevalence (%)	NR	Healthcare workers STRESS Low- 17.4% moderate- 78.9% high- 3.7% DEPRESSION none-min- 50.6%	Healthcare workers Perceived stress- 82.57% Depressive symptoms- 49.42% Anxiety symptoms-

								mild- 38.0% moderate- 8.6% moderately severe- 2.0 % severe- 0.8% ANXIETY none- min- 33.7% mild- 48.6% moderate- 13.7% severe- 4.0%	66.28%
65	(X. Chen et al., 2020) 2020, China	252 (M+F)	Distress, anxiety	GAD-7	GAD-7 (0-21)	Prevalence (%)			Healthcare workers Psychological Distress- 32.5% Anxiety disorder 28.2%

66	(Xiao et al., 2020), 2020, China	<p>958 (F+M)</p> <p>Clinical physician- 378</p> <p>Nurse- 359</p> <p>Working in CT room- 323</p> <p>Work in clinical laboratory- 87</p> <p>Others- 102</p>	Stress, anxiety, depression	PSS-14 HAD	PSS (0-56) HAD (0-21)	Prevalence (%)	<p>Total anxiety and depression</p> <p>Clinical physicians- 40.70%</p> <p>Nurses- 45.30%</p>	NR	<p>Clinical physicians</p> <p>Anxiety- 45.5%</p> <p>Depression- 53.7%</p> <p>Nurses</p> <p>Anxiety- 50.5%</p> <p>Depression- 62.4%</p>
----	----------------------------------	---	-----------------------------	------------	-----------------------	----------------	---	----	--

medRxiv preprint doi: <https://doi.org/10.1101/2021.11.09.21266105>; this version posted November 11, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/).

67	(Xiaoming et al., 2020) China	8817 Doctor- Nurse- Others- 920	(M+F) 3212 4685	Depression, anxiety, somatic symptoms	PHQ-9, GAD-7, PHQ-15, SSI	GAD-7 (0-21) PHQ-9 (0-27) PHQ-15 (0-30)	Prevalence (%)	NR	<p>PHQ-9</p> <p>No symptoms- 69.8%</p> <p>min- 20.8%</p> <p>mild- 6.2%</p> <p>moderate- 2.1%</p> <p>severe- 1.1%</p> <p>GAD-7</p> <p>no symptom- 79.3%</p> <p>Low- 15.6%</p> <p>medium- 3.2%</p> <p>high- 1.9%</p> <p>PHQ-15</p> <p>no symptom- 53.8%</p>	<p>Healthcare workers</p> <p>PHQ-9- 30.23%</p> <p>GAD-7- 20.68%</p> <p>PHQ-15- 46.18%</p> <p>SSI- 6.53%</p>
----	---	---	------------------------------	--	---------------------------	---	----------------	----	--	--

								Low- 26.4% medium- 13.7% high- 6.1% SSI no symptom- 93.5% Low- 6.5%	
68	(Xing et al., 2020), 2020, China	548 (M+F)	Somatization, Obsessive compulsive disorder, anxiety, phobic anxiety	SCL-90	SCL-90 (0-90)	Prevalence (%)	NR	NR	Doctors and nurses Depression- 29.74% Anxiety- 34.2%
69	(Yang et al., 2020a)2020, China	449 (M+F) Doctors= 285 Nurses= 164	Anxiety otolarygology healthcare	SAS	SAS (20-75)	Prevalence (%)	Total anxiety- 29.18%	NR	Doctors Anxiety Yes- 25.26%

			workers						No- 74.74% Nurse Yes- 35.98% No- 64.02%
70	(Yáñez et al., 2020), 2020, Peru	303 (M+F) Physicians- 53 Nurses- 63 Pharmacists- 63 technicians- 80 volunteers- 20 others- 24	Anxiety, distress	GAD-7, K-6	GAD-7 (0-21) K-6 (6-30)	Prevalence (%)	NR	NR	Healthcare workers Anxiety- 2.7% Mental distress- 26.1%

71	(Y. Liu et al., 2020b), 2020, China	2031 (M+F) Pediatric healthcare workers- 2031	Anxiety, depression, stress	DASS-21	DASS-21 (0-42)	Prevalence (%)	All healthcare workers depression- 14.81%, stress- 9.98%, anxiety- 18.3%	Doctors Depression Mild- 8.86% Moderate- 6.42% Severe- 0.93% Extremely severe- 0.82% Anxiety Mild- 7.93% Moderate- 7.93% Severe- 8.97% extremely severe- 1.17% Stress mild- 5.71% Moderate- 4.66%	Doctors Depression- 7.18% Anxiety- 8.21% Stress- 5.02% Nurses Depression- 7.63% Anxiety- 10.7% Stress- 4.97%
----	-------------------------------------	--	-----------------------------	---------	----------------	----------------	---	--	---

medRxiv preprint doi: <https://doi.org/10.1101/2021.11.09.21266105>; this version posted November 11, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/).

								Severe- 0.93% Extremely severe- 0.58%	
72	(Yin et al., 2020), 2020, China	377 (M+F) Doctors- 67 Nurses- 264 others- 40	PTSS post-traumatic stress symptoms	DSM-5 (PCL-5), PSQI	PCL-5 (0-80) PSQI (0-21)	Prevalence (%)	NR	NR	Healthcare workers, PCL-5- 3.8% Very bad sleep satisfaction- 7% Frequent sleep disturbances- 11.3% Difficulty falling asleep

medRxiv preprint doi: <https://doi.org/10.1101/2021.11.09.21266105>; this version posted November 11, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/).

									6.7%
									Had less than 6hrs of sleeping time- 6.2%
73	(Zhan et al., 2020), 2020, China	1794 (M+F)	Insomnia	AIS, FS-14, CPSS	AIS (1-75) FS-14 (0-63) CPSS (0-80)	Prevalence (%)	NR	NR	Nurses Insomnia- 52.8%
74	(Y. Wang et al., 2020a), 2020, China	1897 (M+F) Doctors- 563	Depression, Anxiety, PTSD	PHQ-9, GAD-7, IES-R	PHQ-9 (0-27) GAD-7 (0-21) IES-R (0-28)	Prevalence (%)	Healthcare workers Depression Yes- 15.0% No- 85.0% Anxiety Yes- 27.1% No- 72.9% PTSD	NR	Frontline healthcare workers Depression Yes- 28.7% No- 78.3% Anxiety Yes- 38.5% No- 61.5%

							Yes- 9.8% No- 90.2% Yes- 15.0%		PTSD Yes- 15.4%
75	(Yifang Zhou et al., 2020)China	1931 (M+F)	Poor sleep quality	PSQI	PSQI (0-21)	Prevalence (%)	NR	NR	Frontline Healthcare workers Poor sleep quality- 18.4%
76	(W. R. Zhang et al., 2020), China	2182 (M+F) Medical healthcare workers- 927 Non-medical health workers- 1255	Insomnia, anxious and depressive symptoms, somatic symptoms, obsessive-compulsive symptoms and phobic anxiety.	ISI, SCL-90-R, PHQ-4, PHQ-2, GAD-2	PHQ-9 (0-27) GAD-7 (0-21) IES-R (0-28)	Prevalence (%)	Medical HW + non-medical HW Insomnia- 33.9% Anxiety- 10.4% Depression- 10.6% Somatization symptoms-	NR	Medical Healthcare workers Insomnia- 38.4% Anxiety- 13.0% Depression- 12.2% Somatization symptoms- 1.6% Obsessive

							0.9% Obsessive compulsive symptoms- 3.5% Phobic anxiety- 2.9%		compulsive symptoms- 5.3% Phobic anxiety 3.6%
77	(Z. Zhu et al., 2020a),2020, China	5062 (M+F) Physicians- 1004 Nurses- 3417 Technicians- 641	Stress, depression, anxiety	PHQ-9, GAD- 7, IES-R	PHQ-9 (0- 27) GAD-7 (0-21) IES- R (0-28)	Prevalence (%)	Healthcare workers Stress- 29.8% Depression- 13.5% Anxiety- 24.1%	NR NR	Doctors Depression- 16.7% Anxiety- 19.1% Psychological stress- 16.1% Nurses Depression- 71.8% Anxiety- 70.9% Psychological

									stress- 74.9%
78	(Zhou et al., 2020) China	1705 (M+F) Frontline- 606 General population- 1099	Depression, anxiety, somatization, insomnia, suicide.	PHQ-9, GAD-7, SCL-90, ISI	PHQ-9(0-27) GAD-7 (0-21) SCL-90 (12-60) ISI (0-28)	Prevalence (%)	NR	NR	Frontline staff Depressive symptoms- 56% Anxiety symptoms- 44% Somatic symptoms- 10% Insomnia- 32% Suicide 13.0%

COVID-19- Coronavirus disease 2019, SARS-CoV-19- severe acute respiratory syndrome coronavirus 2, MERS- Middle East Respiratory syndrome ,HCWs- Healthcare workers , FL- Frontline , GAD-7- Generalized anxiety disorder assessment scale, PHQ-9- patient health questionnaire, HADS- Hospital anxiety depression scale , ISI- 7- Insomnia severity index, PLC-5- post traumatic stress disorder checklist, IES-R- Impact event scale revised , PHQ-5- Patient health questionnaire , Y-BOCS- Yale brown obsessive compulsive scale , ESS- Epworth

sleepiness scale , HADS- Hospital anxiety depression scale , BDI- IA- Beck Depression Inventory , PSQI- Pittsburgh sleep quality index, HARS- Hamilton Anxiety rating scale , NASA- TLX- NASA task load index , STAI-Y1- State- Trait Anxiety Inventory (STAI), BDI-II- Beck depression inventory- II, DASS-21- Depression, anxiety and stress scale-21 items , PSS-14- The Perceived stress scale 14- item , PSS-10- The perceived stress scale 10-items , BAI- Beck anxiety inventory, PDEQ- Peritraumatic dissociative Experience questionnaire , PSI WHOQL-BREF- World Health organization quality of life instrument , GHQ-12- General Health questionnaire 12- item scale , PHQ-2- Patient health questionnaire 2- item scale , GSI- Global severity index scale 53 item , GAD-2- Generalized anxiety disorder 2- item scale , Pro QoL- Professional Quality of life scale , SAS- Self- rating anxiety scale , CES-D- Centre for epidemiologic studies depression scale , AIS- Abbreviated injury scale , DBAS-16- Dysfunctional beliefs and attitudes about sleep scale, 16 item self-reported scale, CAS-I- Cognitive attention syndrome , PFI- Professional fulfillment index scale, BRCS- Brief Cognitive Rating Scale , PSSQ- Perceived social support questionnaire, POS- The palliative outcome scale , CD-RISC-10- Connor-Davidson Resilience Scale (CD-RISC),, NRS- Numeric rating scale , HAMA- Hamilton Anxiety Rating scale , HAMD- Hamilton Rating scale for depression , TCSQ- Tayside Childre's Sleep Questionnaire , CMBI- Chinese Maslach Burnout Inventory , SASRQ- Stanford Acute Stress Reaction Questionnaire , BSI- 18- Brief symptom Inventory 18 , PC-PTSD- Primary care- Post Traumatic stress disorder , EQ-5D-3L- Health status assessment questionnaire of the patient, SF36- 36 item short form survey, SDS- Self- Rating Depression scale, SCL-90- The symptom Checklist- 90-R, K-6- Kessler 6 scale , DSM-5- The diagnostic and statistical Manual of Mental disorders, fifth Edition , FS-14- The Chadler fatigue scale , CPSS- Child PTSD symptom scale, ED- Emergency department, ALP- Assistant laboratory personnel, AN- Assistant nurse, N- nurse, MD- Medical doctor, Psy- Clinical, psychologist, MLT- Medical laboratory technician, HT- Health technicians

3.5 Insomnia

Eighteen cross-sectional studies estimated the prevalence of insomnia and poor sleep quality among the FHW (**Table 2**). Various studies reported prevalence of insomnia among 32-57% FHWs (Herrero San martin et al., 2020; Zhou et al; Zhang et al, 2020; Janato iddrissi et al, 2020). Another study showed that 6.2% HCW had less than 6 hours of sleep, 11.3% frequently had sleep disturbances, 2.7% had very bad sleep satisfaction, and 6.7% had difficulty in falling asleep (Yin et al., 2020). One study disclosed that 50.4% HCW had insomnia, out of which 2.1% had severe insomnia, 38.5% had sub-threshold and 9.55% had moderately severe insomnia (Sahin et al., 2020). About 8.27% HCW had ISI score ≥ 22 , showing sleeplessness (Rossi Rodolfo et al., 2020). A similar study showed prevalence of 43.04% of insomnia in FHW, out of which 11.2% had moderate insomnia and 31.81% had

mild symptoms of insomnia (Liang et al., 2020). Insomnia severity index of 11% nurses and 11.4% doctors came out to be more than 14 i.e. $ISI > 14$. (Bhargava et al., 2020) and (Cheng et al., 2020) evaluated the prevalence of 30% poor sleep quality and insomnia in HCW. Another study showed that nurses (60.97%) had more insomnia than doctors (39.02%) (S.Wang et al., 2020). Another study reported that 1.6% HCW suffering from insomnia had severe depression, 5.8% had severe anxiety and 2.63% suffered from severe stress. Contrary to other studies this showed that doctors (50.52%) had more insomnia than nurses (24.21%) (Wasim et al., 2020) (**Table 2**).

3.6 Stress

Twenty studies assessed the stress level of FHW. Various studies reported the prevalence of stress among HCW in range of 60.8%-90.1% (Zhu et al., 2020; Luseno moreno et al, 2020; (Sandesh et al., 2020). Another study demonstrated that 82.57% of HCW had stress, out of which 3.7% had high stress levels and 78.9% had moderate stress (Wilson et al., 2020). Luseno moreno et al, 2020 revealed that 82.98% HCW were dealing with PTSD whereas Wang et al., 2020 reported only 15.4% FHW had PTSD. From one study it was disclosed that high stress levels were indicated in nurses (81.5%) than doctors (11.5%) (Li et al, 2020). Other studies also showed prevalence of stress and PTSD among HCW in the range of 21.7%-28.9% (Almater et al., 2020; Alshekaili et al., 2020a; Fauzi et al., 2020; Johnson et al., 2020; Kannampallil et al., 2020; Wang et al., 2020) (**Table 2**).

3.7 Distress

Seven studies evaluated the prevalence of distress among HCW. Chen et al, 2020 estimated that besides anxiety disorders suffered by frontline workers which came out to be 32.5%, 28.2% participants had psychological distress. Another study analyzed that physicians (62.0%) suffered more distress than residents (58.2%) (Civantos et al., 2020). **Lai et al., 2020** examined distress using IES-R scoring tool and reported that 31.66% FHW were in distress

along with 24.26% depression and 21.40% anxiety. Prasad et al., 2020 highlighted that 84.1% nurses tolerated distress during pandemic whereas Nie et al., 2020 reported prevalence rate of 25.1% of distress among nurses. In a similar article, 76.4% distress prevalence was observed in HCW (Şahin et al., 2020a). Yanez et al., 2020 used K-6 scoring tool and found that HCW suffered more with distress (26.1%) than anxiety (2.7%) (Table-2).

3.8 Other psychological symptoms

(Antonijevic et al., 2020b) evaluated resilience in HCW to be 10.33% whereas somatization, peritraumatic dissociation estimations were assessed by (Azoulay et al., n.d.) (Bhargava et al., 2020). Burnout (Civantos et al., 2020), suicidal ideation, (Hong et al., 2020)(Azoulay et al., n.d.), somatization and obsessive-compulsive disorder in FHW was assessed by various studies (H. Cai et al., 2020; Juan et al., 2020; B. Zhou et al., 2020)(Xiaoming et al., 2020). Zhang et al., 2020 in his study showed the phobic anxiety encountered by medical healthcare workers (Table 2).

4. DISCUSSION

Healthcare workers has acknowledged the outspread of COVID-19 conscientiously (Wasim et al., 2020). Medical staff during the course of pandemic went through exorbitant levels of anxiety, depression, insomnia , somatization and obsessive compulsive disorders and were also prone to grow high grade mental disorders (W. R. Zhang et al., 2020). Thus, in the context of the COVID-19 pandemic, this timely review is both relevant and urgent.

HCWs are presented with unique set of challenges and stressors which include uncertainties such as duration, lack of adequate PPE, gowns, respiratory mask, and face shields, magnitude of crisis, unpreparedness, health threats, transmission risk to loved ones and colleagues (Albott et al., 2020; Anmella et al., 2020; Hall, 2020; Shaukat et al., 2020; Walton et al., 2020). Family safety is the major concerns thus limiting their physical contact with family members to minimize the risk of viral transmission (Albott et al., 2020; Hall, 2020; Jun et al.,

2020; Walton et al., 2020). Difficulty in falling asleep, lethargy, irritability generalized anxiety, dejection and feeling submerged because of daily demands served as factors for inconvenience (W. R. Zhang et al., 2020). Also, continuous influx of challenges in treating patients leads to mental exertion, boredom and decreased interest (Walton et al., 2020).

The reasons for their mental health disturbance are due to ambiguity in job role and poor shifts management, leading to lessened efficiency and burnout (Selamu et al., 2017)(Anmella et al., 2020)(AlAteeq et al., 2020). There is need to monitor these psychological morbidities on regular basis.

One study showed that, anxiety and depression levels among medical staff were higher than the national norms (Cheng et al., 2020). Unmarried or single FHWs, due to lower professional enervation and household liability, indicated surege in mental health problem (Z. Zhu et al., 2020c). According to a survey, anxiety levels showed the drift from first increasing then decreasing and was associated with the age, profession and education. Constant distress, worry, irritability, disturbed concentration, work alteration, difficulties in interpersonal relationships, somatization and depression, imposed uncertainty, and burdened HCW ((Albott et al., 2020)(Elhadi et al., 2020a).

Levels of depression and anxiety were also seen among the ICU professionals. A clinical study showed that female nurses were found to be more anxious than doctors. Also, young age appeared to be the risk factors for anxiety and depression (Caillet et al., 2020a). In other studies, incidences of anxiety levels were less among the trained staff working in close proximity with COVID patients. The psychological end results of the pandemic should make us question about the individualized care to HCWs (Caillet et al., 2020b)(Buselli et al., n.d.)2020; (Gallopini et al., 2020); (Elhadi et al., 2020a).

Also, FHWs had reduced confidence than non-FHW in handling virus. Experiences HCW had less mental health issues as compared to interns (M. Zhang et al., 2020) Good training

and relevant knowledge of the disease can effectively lessens the expanse of anxiety, consternation and precariousness (Yifang Zhou et al., 2020). Less educated workers were more likely to get insomnia (Wasim et al., 2020). A survey of 534 pediatric medical staff estimated that outpatient emergency departments and those having bachelor's degree had high anxiety (Cheng et al., 2020).

Results from the studies conducted in China and other pandemics in the past indicated that FHW had higher distress levels in pandemic (J. Lai et al., 2020); (Prasad et al., 2020). Clinical PTSD symptoms were developed in those who had sub-clinical symptoms due to work setting and continuous challenges and traumatic occurrences, thus there is a need to monitor the sub-clinical symptoms among the persons treating COVID-19 patients (Johnson et al., 2020)(Caillet et al., 2020a). High level of stress lead to burnout which accentuate the exploration of interdependence between burnout, stress and PTSD (Johnson et al., 2020). Supplications have been made in several papers that mental aids should be called for HCWs and particular measures should be taken on psychiatric illness so as to reduce PTSD (Johnson et al., 2020). Women were seen more vulnerable in developing PTSD and required adequate social support by partners (Di Tella et al., 2020). Clinical study showed that, the prevalence of psychological distress among FHW was more than in general population in China (Nie et al., 2020).

In another study it was also reported that people were afraid to meet the nurses and doctors as they feared that they might carry virus with them and can transmit disease to them. FHWs also had to handle the stigmatization from the public which lead to psychological stress eventually (Yifang Zhou et al., 2020). Another considerable apprehension was mistreatment and abuse (either physical or verbal). About 52.3% physicians disclosed minimum one episode of certain kind of abuse. About 31% of the physicians experienced stigmatization by patients (Elhadi et al., 2020a).

Sleep disturbances among HCWs are individualistically associated with depression. About, 38% HCWs suffered from sleep disturbances (Wang et al., 2020). In accordance with a cross-sectional survey, conducted in China among HCW who were directly involved with COVID-19 infected patients, 50.4% suffered from depression, 44.6% from anxiety, 34% from insomnia and 71% from distress (Wańkiewicz et al., 2020). The collaborative pervasiveness of sleep disturbances according to a recent systematic review and meta-analysis established that prevalence of sleep disorders among Chinese FHWs was found to be 39.2%. US doctors working in emergency departments had 31% of poor sleep quality and among Pakistani physicians it was 36.8% (Qiu et al., 2020)(Yifang Zhou et al., 2020).

Belief in conspiracy theory in regard to COVID-19 origin were considered to be related to the weak mental health (X. Chen et al., 2020). Service factors which include lack of quality training being given to HCW in providing mental healthcare in isolation units in hospitals, as in this situational setting of pandemic when transmission rates are high and people are dying, there are common mental health problems which are being faced by both patients and care providers. Lack of organizational support is also the major reason for lack of confidence and burnout and psychological morbidities. Inexperienced doctors with inadequate training, unqualified hand washing and improper knowledge of infection control are the factors responsible for the increment of psychological disturbances among medical staff contributing to the development of risks in their life (Xiang et al., 2020);(Kisely et al., 2020); Shaukat et al., 2020; (Albott et al., 2020); (Jun et al., 2020); (Y. Wang et al., 2020b)(Keubo et al., 2020); (Krishnamoorthy et al., 2020).

5. CONCLUSION

Medical staff has suffered from high degrees of mental disturbances due to miscellaneous inducements. Governing the robustness among FHW during this time of universal calamity caused due to outbreak of SARS-CoV-2 is very crucial. Many avoidable particulars can be

contemplated as the components which are being used to control the current situation of the widespread are burdensome to recast. The prevailing phobia, can be cut down by the proper propagation of knowledge, information and resources of the current situation. Stigma, intolerance and prejudices can be subjugated by educating the healthcare workers, their families and general population. Health policy makers must weigh up the seriousness of the preventive approaches to attenuate the psychological manifestations. HCWs need to be emancipated and their roles and efforts are needed to be acknowledged. Psychological support and potent course of action should be available promptly at multifarious modalities as they play a condemnatory part at odds with COVID-19.

Acknowledgements

None.

Declaration of competing interest

The authors have no relevant affiliation or financial involvement with any organization or entity with a financial interest in or financial conflict with the subject matter or materials discussed in the manuscript. This includes employment, consultancies, honoraria, stock ownerships or options, expert testimony, grants or patents received or pending, or royalties.

Contributors

Rhythm Joshi: Conceptualization, Methodology, Data Curation, Writing-Original Draft, Visualization. **Nidhi.B.Agarwal:** Validation, Writing-Review and Editing. **Dinesh Bhurani:** Writing-Review and Editing. **Mohd Ashif Khan:** Conceptualization, Methodology, Data Curation, Writing-Original Draft, Writing-Review and Editing, Visualization, Validation

Funding

This research received no grants from any funding agency- commercial, public or non- profit sectors.

References

- AlAteeq, D.A., Aljhani, S., Althiyabi, I., Majzoub, S., 2020. Mental health among healthcare providers during coronavirus disease (COVID-19) outbreak in Saudi Arabia. *J. Infect. Public Health* 13, 1432–1437. <https://doi.org/10.1016/j.jiph.2020.08.013>
- Albott, C.S., Wozniak, J.R., McGlinch, B.P., Wall, M.H., Gold, B.S., Vinogradov, S., 2020. Battle Buddies: Rapid Deployment of a Psychological Resilience Intervention for Health Care Workers during the COVID-19 Pandemic. *Anesth. Analg.* 131, 43–54. <https://doi.org/10.1213/ANE.0000000000004912>
- Alenazi, Thamer H., BinDhim, N.F., Alenazi, M.H., Tamim, H., Almagrabi, R.S., Aljohani, S.M., H Basyouni, M., Alzubair, R.A., Althumiri, N.A., Alqahtani, S.A., 2020. Prevalence and predictors of anxiety among healthcare workers in Saudi Arabia during the COVID-19 pandemic. *J. Infect. Public Health* 13, 1645–1651. <https://doi.org/10.1016/j.jiph.2020.09.001>
- Alenazi, Thamer H., Bindhim, N.F., Alenazi, M.H., Tamim, H., Althumiri, N.A., Alqahtani, S.A., 2020. Prevalence and predictors of anxiety among healthcare workers in Saudi Arabia during the COVID-19 pandemic. *J. Infect. Public Health*. <https://doi.org/10.1016/j.jiph.2020.09.001>
- Almater, A.I., Tobaigy, M.F., Younis, A.S., Alaqeel, M.K., Abouammoh, M.A., 2020. Effect of 2019 Coronavirus Pandemic on Ophthalmologists Practicing in Saudi Arabia: A Psychological Health Assessment. *Middle East Afr. J. Ophthalmol.* 27, 79–85. https://doi.org/10.4103/meajo.MEAJO_220_20
- Alshekaili, M., Hassan, W., Al Said, N., Al Sulaimani, F., Jayapal, S.K., Al-Mawali, A., Chan, M.F., Mahadevan, S., Al-Adawi, S., 2020a. Factors associated with mental health outcomes across healthcare settings in Oman during COVID-19: Frontline versus non-

frontline healthcare workers. *BMJ Open* 10, 1–7. <https://doi.org/10.1136/bmjopen-2020-042030>

Alshekaili, M., Hassan, W., Said, N. Al, Sulaimani, F. Al, Chan, M.F., Jayapal, S.K., Al-, A., 2020b. Factors associated with mental health outcomes across healthcare settings in Oman during COVID-19: frontline frontline healthcare workers versus non- - 1–7. <https://doi.org/10.1136/bmjopen-2020-042030>

Altintas, A.K., 2017. Allen RC. Orbital metastases: When to suspect? When to biopsy?. *Middle East Afr J Ophthalmol* 2018;25:60–4. *Middle East Afr J Ophthalmol* 24, 43–47. <https://doi.org/10.4103/meajo.MEAJO>

An, Y., Yang, Y., Wang, A., Li, Y., Zhang, Q., Cheung, T., Ungvari, G.S., Qin, M.Z., An, F.R., Xiang, Y.T., 2020. Prevalence of depression and its impact on quality of life among frontline nurses in emergency departments during the COVID-19 outbreak. *J. Affect. Disord.* 276, 312–315. <https://doi.org/10.1016/j.jad.2020.06.047>

Anmella, G., Fico, G., Roca, A., Gómez-Ramiro, M., Vázquez, M., Murru, A., Pacchiarotti, I., Verdolini, N., Vieta, E., 2020. Unravelling potential severe psychiatric repercussions on healthcare professionals during the COVID-19 crisis. *J. Affect. Disord.* 273, 422–424. <https://doi.org/10.1016/j.jad.2020.05.061>

Antonijevic, J., Binic, I., Zikic, O., Manojlovic, S., Tosic-golubovic, S., Popovic, N., 2020a. Mental health of medical personnel during the COVID-19 pandemic 1–9. <https://doi.org/10.1002/brb3.1881>

Antonijevic, J., Binic, I., Zikic, O., Manojlovic, S., Tosic-Golubovic, S., Popovic, N., 2020b. Mental health of medical personnel during the COVID-19 pandemic. *Brain Behav.* 10, 1–9. <https://doi.org/10.1002/brb3.1881>

Ayhan Başer, D., Çevik, M., Gümüştakim, Ş., Başara, E., 2020. Assessment of individuals' attitude, knowledge and anxiety towards COVID-19 at the first period of the outbreak in

- Turkey: A web-based cross-sectional survey. *Int. J. Clin. Pract.* 74, 1–8.
<https://doi.org/10.1111/ijcp.13622>
- Azoulay, E., Cariou, A., Bruneel, F., Demoule, A., Kouatchet, A., Souppart, V., Combes, A., Klouche, K., Argaud, L., Barbier, F., Jourdain, M., Reignier, J., Papazian, L., Guidet, B., Géri, G., Guisset, O., Labbé, V., Mégarbane, B., Der, G. Van, Guitton, C., Friedman, D., Pochard, F., Darmon, M., Klouche, K., Argaud, L., Barbier, F., Jourdain, M., Reignier, J., Labbé, V., Megarbane, B., Meersch, G. Van Der, Guitton, C., Azoulay, E., Darmon, M., Souppart, V., Pochard, F., Kenstish-, N., n.d. Clinicians Managing COVID-19 Patients □: A Cross-Sectional Study 1–46. <https://doi.org/10.1164/rccm.202006-2568OC>
- Barranco, R., Ventura, F., 2020. Covid-19 and infection in health-care workers: An emerging problem. *Med. Leg. J.* 002581722092369. <https://doi.org/10.1177/0025817220923694>
- Bhargava, S., Sarkar, R., Kroumpouzou, G., 2020. Mental distress in dermatologists during COVID-19 pandemic: Assessment and risk factors in a global, cross-sectional study. *Dermatol. Ther.* 33, 1–6. <https://doi.org/10.1111/dth.14161>
- Buselli, R., Corsi, M., Baldanzi, S., Chiumiento, M., Lupo, E. Del, Dell'oste, V., Bertelloni, C.A., Massimetti, G., Dell'osso, L., Cristaudo, A., Carmassi, C., 2020. Professional quality of life and mental health outcomes among health care workers exposed to SARS-CoV-2 (COVID-19). *Int. J. Environ. Res. Public Health* 17, 1–12.
<https://doi.org/10.3390/ijerph17176180>
- Buselli, R., Corsi, M., Baldanzi, S., Chiumiento, M., Lupo, E. Del, Oste, V.D., Bertelloni, C.A., Massimetti, G., Osso, L.D., Cristaudo, A., Carmassi, C., n.d. Professional Quality of Life and Mental Health Outcomes among Health Care Workers Exposed to 2, 1–12.
- C, J.M., Blais-rochette, C., Kokou-kpolou, C.K., Noorishad, P., Mukunzi, J.N., Mcintee, S., Dalexis, R.D., Labelle, P., C, J.M., 2020. Journal Pre-proof.
- Cai, H., Tu, B., Ma, J., Chen, L., Fu, L., Jiang, Y., Zhuang, Q., 2020. Psychological impact

- and coping strategies of frontline medical staff in Hunan between January and March 2020 during the outbreak of coronavirus disease 2019 (COVID) in Hubei, China. *Med. Sci. Monit.* 26, 1–16. <https://doi.org/10.12659/MSM.924171>
- Cai, W., Lian, B., Song, X., Hou, T., Deng, G., Li, H., 2020. A cross-sectional study on mental health among health care workers during the outbreak of Corona Virus Disease 2019. *Asian J. Psychiatr.* 51, 102111. <https://doi.org/10.1016/j.ajp.2020.102111>
- Caillet, A., Coste, C., Sanchez, R., Allaouchiche, B., 2020a. Psychological Impact of COVID-19 on ICU Caregivers. *Anaesth. Crit. Care Pain Med.* 39, 717–722. <https://doi.org/10.1016/j.accpm.2020.08.006>
- Caillet, A., Coste, C., Sanchez, R., Allaouchiche, B., 2020b. na of. *Anaesth. Crit. Care Pain Med.* <https://doi.org/10.1016/j.accpm.2020.08.006>
- Chen, J., Liu, X., Wang, D., Jin, Y., He, M., Ma, Y., Zhao, X., Song, S., Zhang, L., Xiang, X., Yang, L., Song, J., Bai, T., Hou, X., 2020. Risk factors for depression and anxiety in healthcare workers deployed during the COVID-19 outbreak in China. *Soc. Psychiatry Psychiatr. Epidemiol.* <https://doi.org/10.1007/s00127-020-01954-1>
- Chen, P., Lei, J., Chen, F., Zhou, B., 2020. Experiences and perceptions risk of health-care workers from coronavirus: A protocol for systematic review. *Medicine (Baltimore)*. 99, e20308. <https://doi.org/10.1097/MD.00000000000020308>
- Chen, Q., Liang, M., Li, Y., Guo, J., Fei, D., Wang, L., He, L., Sheng, C., Cai, Y., Li, X., Wang, J., Zhang, Z., 2020. Mental health care for medical staff in China during the COVID-19 outbreak. *The Lancet Psychiatry* 7, e15–e16. [https://doi.org/10.1016/S2215-0366\(20\)30078-X](https://doi.org/10.1016/S2215-0366(20)30078-X)
- Chen, X., Zhang, S.X., Jahanshahi, A.A., Alvarez-Risco, A., Dai, H., Li, J., Ibarra, V.G., 2020. Belief in a COVID-19 conspiracy theory as a predictor of mental health and well-being of health care workers in Ecuador: Cross-sectional survey study. *JMIR Public*

Heal. Surveill. 6, 1–7. <https://doi.org/10.2196/20737>

Cheng, F.F., Zhan, S.H., Xie, A.W., Cai, S.Z., Hui, L., Kong, X.X., Tian, J.M., Yan, W.H.,
2020. Anxiety in Chinese pediatric medical staff during the outbreak of Coronavirus
Disease 2019: A cross-sectional study. *Transl. Pediatr.* 9, 231–236.
<https://doi.org/10.21037/TP.2020.04.02>

Chew, N.W.S., Lee, G.K.H., Tan, B.Y.Q., Jing, M., Goh, Y., Ngiam, N.J.H., Yeo, L.L.L.,
Ahmad, A., Ahmed Khan, F., Napoleon Shanmugam, G., Sharma, A.K., Komalkumar,
R.N., Meenakshi, P.V., Shah, K., Patel, B., Chan, B.P.L., Sunny, S., Chandra, B., Ong,
J.J.Y., Paliwal, P.R., Wong, L.Y.H., Sagayanathan, R., Chen, J.T., Ying Ng, A.Y., Teoh,
H.L., Tsivgoulis, G., Ho, C.S., Ho, R.C., Sharma, V.K., 2020. A multinational,
multicentre study on the psychological outcomes and associated physical symptoms
amongst healthcare workers during COVID-19 outbreak. *Brain. Behav. Immun.* 88,
559–565. <https://doi.org/10.1016/j.bbi.2020.04.049>

Civantos, A.M., Byrnes, Y., Chang, C., Prasad, A., Chorath, K., Poonia, S.K., Jenks, C.M.,
Bur, A.M., Thakkar, P., Graboyes, E.M., Seth, R., Trosman, S., Wong, A., Laitman,
B.M., Harris, B.N., Shah, J., Stubbs, V., Choby, G., Long, Q., Rassekh, C.H., Thaler, E.,
Rajasekaran, K., 2020. Mental health among otolaryngology resident and attending
physicians during the COVID-19 pandemic: National study. *Head Neck* 42, 1597–1609.
<https://doi.org/10.1002/hed.26292>

de Burgos-Berdud, I., Valdés-Florido, M.J., López-Díaz, Á., 2021. Are healthcare workers
during the COVID-19 pandemic at risk of psychosis? Findings from a scoping review.
Gen. Hosp. Psychiatry. <https://doi.org/10.1016/j.genhosppsy.2020.06.015>

Di Tella, M., Romeo, A., Benfante, A., Castelli, L., 2020. Mental health of healthcare
workers during the COVID-19 pandemic in Italy. *J. Eval. Clin. Pract.* 26, 1583–1587.
<https://doi.org/10.1111/jep.13444>

Elbay, R.Y., Kurtulmuş, A., Arpacioğlu, S., Karadere, E., 2020a. Depression, anxiety, stress levels of physicians and associated factors in Covid-19 pandemics. *Psychiatry Res.* 290. <https://doi.org/10.1016/j.psychres.2020.113130>

Elbay, R.Y., Kurtulmuş, A., Arpacioğlu, S., Karadere, E., 2020b. Depression, anxiety, stress levels of physicians and associated factors in Covid-19 pandemics. *Psychiatry Res.* 290, 1–5. <https://doi.org/10.1016/j.psychres.2020.113130>

Elhadi, M., Msherghi, A., Elgzairi, M., Alhashimi, A., Bouhuwaish, A., Biala, M., Abuelmeda, S., Khel, S., Khaled, A., Alsou, A., Elmabrouk, A., Bin, F., Alhadi, B., Alhaddad, S., Elmabrouk, O., Ben, T., Alameen, H., Zaid, A., Elhadi, A., Albakoush, A., 2020a. Psychological status of healthcare workers during the civil war and COVID- 19 pandemic□: A cross-sectional study 137. <https://doi.org/10.1016/j.jpsychores.2020.110221>

Elhadi, M., Msherghi, A., Elgzairi, M., Alhashimi, A., Bouhuwaish, A., Biala, M., Abuelmeda, S., Khel, S., Khaled, A., Alsoufi, A., Elmabrouk, A., Alshiteewi, F. Bin, Alhadi, B., Alhaddad, S., Gaffaz, R., Elmabrouk, O., Hamed, T. Ben, Alameen, H., Zaid, A., Elhadi, A., Albakoush, A., 2020b. Psychological status of healthcare workers during the civil war and COVID-19 pandemic: A cross-sectional study. *J. Psychosom. Res.* 137. <https://doi.org/10.1016/j.jpsychores.2020.110221>

Fadhli, M., Fauzi, M., Mohd, H., Robot, R.M., n.d. Doctors ' Mental Health in the Midst of COVID-19 Pandemic□: The Roles of Work Demands and Recovery Experiences 19.

Fauzi, M.F.M., Yusoff, H.M., Robot, R.M., Saruan, N.A.M., Ismail, K.I., Haris, A.F.M., 2020. Doctors' mental health in the midst of covid-19 pandemic: The roles of work demands and recovery experiences. *Int. J. Environ. Res. Public Health* 17, 1–16. <https://doi.org/10.3390/ijerph17197340>

Gallopeni, F., Bajraktari, I., Selmani, E., Tahirbegolli, I.A., Sahiti, G., Muastafa, A., Bojaj,

- G., Muharremi, V.B., Tahirbegolli, B., 2020. Anxiety and depressive symptoms among healthcare professionals during the Covid-19 pandemic in Kosovo: A cross sectional study. *J. Psychosom. Res.* 137, 110212.
<https://doi.org/10.1016/j.jpsychores.2020.110212>
- Gan, W.H., Lim, J.W., Koh, D., 2020. Preventing Intra-hospital Infection and Transmission of Coronavirus Disease 2019 in Health-care Workers. *Saf. Health Work.*
<https://doi.org/10.1016/j.shaw.2020.03.001>
- Hall, H., 2020. The effect of the COVID-19 pandemic on healthcare workers' mental health. *JAAPA* 33, 45–48. <https://doi.org/10.1097/01.JAA.0000669772.78848.8c>
- Herrero San Martin, A., Parra Serrano, J., Diaz Cambriles, T., Arias Arias, E.M., Muñoz Méndez, J., del Yerro Álvarez, M.J., González Sánchez, M., 2020. Sleep characteristics in health workers exposed to the COVID-19 pandemic. *Sleep Med.* 75, 388–394.
<https://doi.org/10.1016/j.sleep.2020.08.013>
- Hong, S., Ai, M., Xu, X., Wang, W., Chen, J., Zhang, Q., Wang, L., Kuang, L., 2020. Immediate psychological impact on nurses working at 42 government-designated hospitals during COVID-19 outbreak in China: A cross-sectional study. *Nurs. Outlook* 0, 1–7. <https://doi.org/10.1016/j.outlook.2020.07.007>
- Huang, Y., Zhao, N., 2020. Mental health burden for the public affected by the COVID-19 outbreak in China: Who will be the high-risk group? *Psychol. Heal. Med.* 00, 1–12.
<https://doi.org/10.1080/13548506.2020.1754438>
- Janati Idrissi, A., Lamkaddem, A., Benouajjit, A., Ben El Bouazzaoui, M., El Houari, F., Alami, M., Labyad, S., Chahidi, A., Benjelloun, M., Rabhi, S., Kissani, N., Zarhbouch, B., Ouazzani, R., Kadiri, F., Alouane, R., Elbiaze, M., Boujraf, S., El Fakir, S., Souirti, Z., 2020. Sleep quality and mental health in the context of COVID-19 pandemic and lockdown in Morocco. *Sleep Med.* 74, 248–253.

<https://doi.org/10.1016/j.sleep.2020.07.045>

Johnson, S.U., Ebrahimi, O. V., Hoffart, A., 2020. PTSD symptoms among health workers and public service providers during the COVID-19 outbreak. *PLoS One* 15, 1–13.

<https://doi.org/10.1371/journal.pone.0241032>

Juan, Y., Yuanyuan, C., Qiuxiang, Y., Cong, L., Xiaofeng, L., Yundong, Z., Jing, C., Peifeng, Q., Yan, L., Xiaojiao, X., Yujie, L., 2020. Psychological distress surveillance and related impact analysis of hospital staff during the COVID-19 epidemic in Chongqing, China. *Compr. Psychiatry* 103, 152198. <https://doi.org/10.1016/j.comppsy.2020.152198>

Compr. Psychiatry 103, 152198. <https://doi.org/10.1016/j.comppsy.2020.152198>

Jun, J., Tucker, S., Melnyk, B.M., 2020. Clinician Mental Health and Well-Being During Global Healthcare Crises: Evidence Learned From Prior Epidemics for COVID-19 Pandemic. *Worldviews Evidence-Based Nurs.* 17, 182–184.

<https://doi.org/10.1111/wvn.12439>

Kang, L., Li, Y., Hu, S., Chen, M., Yang, C., Yang, B.X., Wang, Y., Hu, J., Lai, J., Ma, X., Chen, J., Guan, L., Wang, G., Ma, H., Liu, Z., 2020a. The mental health of medical workers in Wuhan, China dealing with the 2019 novel coronavirus. *The Lancet Psychiatry* 7, e14. [https://doi.org/10.1016/S2215-0366\(20\)30047-X](https://doi.org/10.1016/S2215-0366(20)30047-X)

Psychiatry 7, e14. [https://doi.org/10.1016/S2215-0366\(20\)30047-X](https://doi.org/10.1016/S2215-0366(20)30047-X)

Kang, L., Ma, S., Chen, M., Yang, J., Wang, Y., Li, R., Yao, L., Bai, H., Cai, Z., Xiang Yang, B., Hu, S., Zhang, K., Wang, G., Ma, C., Liu, Z., 2020b. Impact on mental health and perceptions of psychological care among medical and nursing staff in Wuhan during the 2019 novel coronavirus disease outbreak: A cross-sectional study. *Brain. Behav. Immun.* 87, 11–17. <https://doi.org/10.1016/j.bbi.2020.03.028>

Immun. 87, 11–17. <https://doi.org/10.1016/j.bbi.2020.03.028>

Kannampallil, T.G., Goss, C.W., Evanoff, B.A., Strickland, J.R., McAlister, R.P., Duncan, J., 2020. Exposure to COVID-19 patients increases physician trainee stress and burnout. *PLoS One* 15, 1–12. <https://doi.org/10.1371/journal.pone.0237301>

PLoS One 15, 1–12. <https://doi.org/10.1371/journal.pone.0237301>

Keubo, F.R.N., Mboua, P.C., Tadongfack, T.D., Tchoffo, E.F., Tatang, C.T., Zeuna, J.I.,

- Noupoue, E.M., Tsoplifack, C.B., Folefack, G.O., 2020. Psychological distress among health care professionals of the three COVID-19 most affected Regions in Cameroon: Prevalence and associated factors. *Ann. Médico-psychologiques, Rev. Psychiatr.* <https://doi.org/10.1016/j.amp.2020.08.012>
- Kisely, S., Warren, N., McMahon, L., Dalais, C., Henry, I., Siskind, D., 2020. Occurrence, prevention, and management of the psychological effects of emerging virus outbreaks on healthcare workers: rapid review and meta-analysis. *BMJ* 369, m1642. <https://doi.org/10.1136/bmj.m1642>
- Koksal, E., Dost, B., Terzi, Ö., Ustun, Y.B., Özdin, S., Bilgin, S., 2020a. Evaluation of Depression and Anxiety Levels and Related Factors Among Operating Theater Workers During the Novel Coronavirus (COVID-19) Pandemic. *J. Perianesthesia Nurs.* 35, 472–477. <https://doi.org/10.1016/j.jopan.2020.06.017>
- Koksal, E., Dost, B., Ustun, Y.B., 2020b. Evaluation of Depression and Anxiety Levels and Related Factors Among Operating Theater Workers During the Novel Coronavirus (COVID-19) Pandemic 35, 472–477. <https://doi.org/10.1016/j.jopan.2020.06.017>
- Korkmaz, S., Kazgan, A., Çekiç, S., Tartar, A.S., Balcı, H.N., Atmaca, M., 2020. The anxiety levels, quality of sleep and life and problem-solving skills in healthcare workers employed in COVID-19 services. *J. Clin. Neurosci.* 80, 131–136. <https://doi.org/10.1016/j.jocn.2020.07.073>
- Krishnamoorthy, Y., Nagarajan, R., Saya, G.K., Menon, V., 2020. Prevalence of psychological morbidities among general population, healthcare workers and COVID-19 patients amidst the COVID-19 pandemic: A systematic review and meta-analysis. *Psychiatry Res.* 293, 113382. <https://doi.org/10.1016/j.psychres.2020.113382>
- Labrague, L.J., De los Santos, J.A.A., 2020. COVID-19 anxiety among front-line nurses: Predictive role of organisational support, personal resilience and social support. *J. Nurs.*

Manag. 28, 1653–1661. <https://doi.org/10.1111/jonm.13121>

Lai, J., Ma, S., Wang, Y., Cai, Z., Hu, J., Wei, N., Wu, J., Du, H., Chen, T., Li, R., Tan, H.,

Kang, L., Yao, L., Huang, M., Wang, H., Wang, G., Liu, Z., Hu, S., 2020. Factors Associated With Mental Health Outcomes Among Health Care Workers Exposed to Coronavirus Disease 2019. *JAMA Netw. open* 3, e203976.

<https://doi.org/10.1001/jamanetworkopen.2020.3976>

Lai, X., Wang, M., Qin, C., Tan, L., Ran, L., Chen, D., Zhang, H., Shang, K., Xia, C., Wang,

S., Xu, S., Wang, W., 2020. Coronavirus Disease 2019 (COVID-2019) Infection Among Health Care Workers and Implications for Prevention Measures in a Tertiary Hospital in Wuhan, China. *JAMA Netw. open* 3, e209666.

<https://doi.org/10.1001/jamanetworkopen.2020.9666>

Lee, M.C.C., Thampi, S., Chan, H.P., Khoo, D., Chin, B.Z.B., Foo, D.P.X., Lua, C.B.,

Lewin, B., Jacob, R., 2020. Psychological distress during the COVID-19 pandemic amongst anaesthesiologists and nurses. *Br. J. Anaesth.* 125, e384–e386.

<https://doi.org/10.1016/j.bja.2020.07.005>

Li, G., Miao, J., Wang, H., Xu, S., Sun, W., Fan, Y., Zhang, C., Zhu, S., Zhu, Z., Wang, W.,

2020. Psychological impact on women health workers involved in COVID-19 outbreak in Wuhan: A cross-sectional study. *J. Neurol. Neurosurg. Psychiatry* 91, 895–897.

<https://doi.org/10.1136/jnnp-2020-323134>

Li, R., Chen, Y., Lv, J., Liu, L., Zong, S., Li, Hanxia, Li, Hong, 2020. Anxiety and related

factors in frontline clinical nurses fighting COVID-19 in Wuhan. *Medicine (Baltimore)*. 99, e21413. <https://doi.org/10.1097/MD.00000000000021413>

Liang, Y., Wu, K., Zhou, Yongjie, Huang, X., Zhou, Yueyue, Liu, Z., 2020. Mental health in

frontline medical workers during the 2019 novel coronavirus disease epidemic in China: A comparison with the general population. *Int. J. Environ. Res. Public Health* 17, 1–12.

<https://doi.org/10.3390/ijerph17186550>

Liu, C.Y., Yang, Y.Z., Zhang, X.M., Xu, X., Dou, Q.L., Zhang, W.W., 2020. The prevalence and influencing factors for anxiety in medical workers fighting COVID-19 in China: A cross-sectional survey. medRxiv. <https://doi.org/10.1101/2020.03.05.20032003>

Liu, C.Y., Yang, Y.Z., Zhang, X.M., Xu, X., Dou, Q.L., Zhang, W.W., Cheng, A.S.K., 2020. The prevalence and influencing factors in anxiety in medical workers fighting COVID-19 in China: A cross-sectional survey. *Epidemiol. Infect.* 148.

<https://doi.org/10.1017/S0950268820001107>

Liu, M., Cheng, S.Z., Xu, K.W., Yang, Y., Zhu, Q.T., Zhang, H., Yang, D.Y., Cheng, S.Y., Xiao, H., Wang, J.W., Yao, H.R., Cong, Y.T., Zhou, Y.Q., Peng, S., Kuang, M., Hou, F.F., Cheng, K.K., Xiao, H.P., 2020. Use of personal protective equipment against coronavirus disease 2019 by healthcare professionals in Wuhan, China: Cross sectional study. *BMJ* 369, 6–11. <https://doi.org/10.1136/bmj.m2195>

Liu, Y., Wang, L., Chen, L., Zhang, X., Bao, L., Shi, Y., 2020a. Mental Health Status of Paediatric Medical Workers in China During the COVID-19 Outbreak. *Front. Psychiatry* 11, 1–7. <https://doi.org/10.3389/fpsy.2020.00702>

Liu, Y., Wang, L., Chen, L., Zhang, X., Bao, L., Shi, Y., 2020b. Mental Health Status of Paediatric Medical Workers in China During the COVID-19 Outbreak 11, 1–7. <https://doi.org/10.3389/fpsy.2020.00702>

Liu, Y.E., Zhai, Z.C., Han, Y.H., Liu, Y.L., Liu, F.P., Hu, D.Y., 2020. Experiences of front-line nurses combating coronavirus disease-2019 in China: A qualitative analysis. *Public Health Nurs.* 1–7. <https://doi.org/10.1111/phn.12768>

Lu, W., Wang, H., Lin, Y., Li, L., 2020. Psychological status of medical workforce during the COVID-19 pandemic: A cross-sectional study. *Psychiatry Res.* 288, 1–5. <https://doi.org/10.1016/j.psychres.2020.112936>

- Luceño-Moreno, L., Talavera-Velasco, B., García-Albuérne, Y., Martín-García, J., 2020. Symptoms of posttraumatic stress, anxiety, depression, levels of resilience and burnout in spanish health personnel during the COVID-19 pandemic. *Int. J. Environ. Res. Public Health* 17, 1–29. <https://doi.org/10.3390/ijerph17155514>
- McKay, D., Heisler, M., Mishori, R., Catton, H., Kloiber, O., 2020. Attacks against health-care personnel must stop, especially as the world fights COVID-19. *Lancet* 395, 1743–1745. [https://doi.org/10.1016/S0140-6736\(20\)31191-0](https://doi.org/10.1016/S0140-6736(20)31191-0)
- Nie, A., Su, X., Zhang, S., Guan, W., Li, J., 2020. Psychological impact of COVID-19 outbreak on frontline nurses: A cross-sectional survey study. *J. Clin. Nurs.* 29, 4217–4226. <https://doi.org/10.1111/jocn.15454>
- Ning, X., Yu, F., Huang, Qin, Li, X., Luo, Y., Huang, Qing, Chen, C., 2020. The mental health of neurological doctors and nurses in Hunan Province, China, during the COVID-19 outbreak 1–9. <https://doi.org/10.21203/rs.3.rs-22061/v1>
- Pouralizadeh, M., Bostani, Z., Maroufizadeh, S., Ghanbari, A., Khoshbakht, M., Alavi, S.A., Ashrafi, S., 2020. Anxiety and depression and the related factors in nurses of Guilan University of Medical Sciences hospitals during COVID-19: A web-based cross-sectional study. *Int. J. Africa Nurs. Sci.* 13, 100233. <https://doi.org/10.1016/j.ijans.2020.100233>
- Prasad, A., Civantos, A.M., Byrnes, Y., Chorath, K., Poonia, S., Chang, C., Graboyes, E.M., Bur, A.M., Thakkar, P., Deng, J., Seth, R., Trosman, S., Wong, A., Laitman, B.M., Shah, J., Stubbs, V., Long, Q., Choby, G., Rassekh, C.H., Thaler, E.R., Rajasekaran, K., 2020. Snapshot Impact of COVID-19 on Mental Wellness in Nonphysician Otolaryngology Health Care Workers: A National Study. *OTO Open* 4, 2473974X2094883. <https://doi.org/10.1177/2473974x20948835>
- Qiu, D., Yu, Y., Li, R.Q., Li, Y.L., Xiao, S.Y., 2020. Prevalence of sleep disturbances in

- Chinese healthcare professionals: a systematic review and meta-analysis. *Sleep Med.* 67, 258–266. <https://doi.org/10.1016/j.sleep.2019.01.047>
- Que, J., Shi, L., Deng, J., Liu, J., Zhang, L., Wu, S., Gong, Y., Huang, W., Yuan, K., Yan, W., Sun, Y., Ran, M., Bao, Y., Lu, L., 2020. Psychological impact of the covid-19 pandemic on healthcare workers: A cross-sectional study in China. *Gen. Psychiatry* 33, 1–12. <https://doi.org/10.1136/gpsych-2020-100259>
- Rossi, R., Socci, V., Pacitti, F., Di Lorenzo, G., Di Marco, A., Siracusano, A., Rossi, A., 2020. Mental Health Outcomes Among Frontline and Second-Line Health Care Workers During the Coronavirus Disease 2019 (COVID-19) Pandemic in Italy. *JAMA Netw. open* 3, e2010185. <https://doi.org/10.1001/jamanetworkopen.2020.10185>
- Şahin, M.K., Aker, S., Şahin, G., Karabekiroğlu, A., 2020a. Prevalence of Depression, Anxiety, Distress and Insomnia and Related Factors in Healthcare Workers During COVID-19 Pandemic in Turkey. *J. Community Health* 45, 1168–1177. <https://doi.org/10.1007/s10900-020-00921-w>
- Şahin, M.K., Aker, S., Şahin, G., Karabekiroğlu, A., 2020b. Prevalence of Depression, Anxiety, Distress and Insomnia and Related Factors in Healthcare Workers During COVID-19 Pandemic in Turkey. *J. Community Health* 45, 1168–1177. <https://doi.org/10.1007/s10900-020-00921-w>
- Salopek-Žiha, D., Hlavati, M., Gvozdanovi, Z., Gaši, M., Placento, H., Jaki, H., Klapan, D., Šimi, H., 2020. Differences in distress and coping with the covid-19 stressor in nurses and physicians. *Psychiatr. Danub.* 32, 287–293. <https://doi.org/10.24869/PSYD.2020.287>
- Sandesh, R., Shahid, W., Dev, K., Mandhan, N., Shankar, P., Shaikh, A., Rizwan, A., 2020. Impact of COVID-19 on the Mental Health of Healthcare Professionals in Pakistan. *Cureus* 12, 3–7. <https://doi.org/10.7759/cureus.8974>

- Santabárbara, J., Bueno-Notivol, J., Lipnicki, D.M., Olaya, B., Pérez-Moreno, M., Gracia-García, P., Idoiaga-Mondragon, N., Ozamiz-Etxebarria, N., 2021. Prevalence of anxiety in health care professionals during the COVID-19 pandemic: A rapid systematic review (on published articles in Medline) with meta-analysis. *Prog. Neuro-Psychopharmacology Biol. Psychiatry* 107. <https://doi.org/10.1016/j.pnpbp.2021.110244>
- Selamu, M., Thornicroft, G., Fekadu, A., Hanlon, C., 2017. Conceptualisation of job-related wellbeing, stress and burnout among healthcare workers in rural Ethiopia: A qualitative study. *BMC Health Serv. Res.* 17, 1–11. <https://doi.org/10.1186/s12913-017-2370-5>
- Shah, N., Raheem, A., Sideris, M., Velauthar, L., Saeed, F., 2020. Mental health amongst obstetrics and gynaecology doctors during the COVID-19 pandemic: Results of a UK-wide study. *Eur. J. Obstet. Gynecol. Reprod. Biol.* 253, 90–94. <https://doi.org/10.1016/j.ejogrb.2020.07.060>
- Shahrour, G., 2020. Acute stress disorder , coping self-efficacy and subsequent psychological distress among nurses amid COVID-19 1–10. <https://doi.org/10.1111/jonm.13124>
- Shahrour, G., Dardas, L.A., 2020. Acute stress disorder, coping self-efficacy and subsequent psychological distress among nurses amid COVID-19. *J. Nurs. Manag.* 28, 1686–1695. <https://doi.org/10.1111/jonm.13124>
- Shaukat, N., Ali, D.M., Razzak, J., 2020. Physical and mental health impacts of COVID-19 on healthcare workers: A scoping review. *Int. J. Emerg. Med.* 13, 1–8. <https://doi.org/10.1186/s12245-020-00299-5>
- Shechter, A., Diaz, F., Moise, N., Anstey, D.E., Ye, S., Agarwal, S., Birk, J.L., Brodie, D., Cannone, D.E., Chang, B., Claassen, J., Cornelius, T., Derby, L., Dong, M., Givens, R.C., Hochman, B., Homma, S., Kronish, I.M., Lee, S.A.J., Manzano, W., Mayer, L.E.S., McMurry, C.L., Moitra, V., Pham, P., Rabbani, L.R., Rivera, R.R., Schwartz, A., Schwartz, J.E., Shapiro, P.A., Shaw, K., Sullivan, A.M., Vose, C., Wasson, L.,

- Edmondson, D., Abdalla, M., 2020. Psychological distress, coping behaviors, and preferences for support among New York healthcare workers during the COVID-19 pandemic. *Gen. Hosp. Psychiatry* 66, 1–8.
<https://doi.org/10.1016/j.genhosppsych.2020.06.007>
- Si, M., Su, X., Jiang, Y., Wang, W., Gu, X., Ma, L., Li, J., Zhang, S., Ren, Z.-F., Liu, Y.-L., Qiao, Y.-L., 2020. The Psychological Impact of COVID-19 on Medical Care Workers in China. *SSRN Electron. J.* 1–13. <https://doi.org/10.2139/ssrn.3592642>
- Skoda, E.M., Teufel, M., Stang, A., Jöckel, K.H., Junne, F., Weismüller, B., Hetkamp, M., Musche, V., Kohler, H., Dörrie, N., Schweda, A., Bäuerle, A., 2020. Psychological burden of healthcare professionals in Germany during the acute phase of the COVID-19 pandemic: Differences and similarities in the international context. *J. Public Heal. (United Kingdom)* 42, 688–695. <https://doi.org/10.1093/pubmed/fdaa124>
- Stojanov, J., Malobabic, M., Stanojevic, G., Stevic, M., Milosevic, V., Stojanov, A., 2020. Quality of sleep and health-related quality of life among health care professionals treating patients with coronavirus disease-19. *Int. J. Soc. Psychiatry.*
<https://doi.org/10.1177/0020764020942800>
- Suryavanshi, N., Kadam, A., Dhumal, G., Nimkar, S., Mave, V., Gupta, A., Cox, S.R., Gupte, N., 2020. Mental health and quality of life among healthcare professionals during the COVID-19 pandemic in India. *Brain Behav.* 10, 1–12. <https://doi.org/10.1002/brb3.1837>
- Teo, W.Z.Y., Soo, Y.E., Yip, C., Lizhen, O., Chun-Tsu, L., 2020. The psychological impact of COVID-19 on ‘hidden’ frontline healthcare workers. *Int. J. Soc. Psychiatry.*
<https://doi.org/10.1177/0020764020950772>
- Tian, T., Meng, F., Pan, W., Zhang, S., Cheung, T., Ng, C.H., Li, X.H., Xiang, Y.T., 2020. Mental health burden of frontline health professionals in screening and caring the imported COVID-19 patients in China during the pandemic. *Psychol. Med.*

<https://doi.org/10.1017/S0033291720002093>

- Walton, M., Murray, E., Christian, M.D., 2020. Mental health care for medical staff and affiliated healthcare workers during the COVID-19 pandemic. *Eur. Hear. J. Acute Cardiovasc. Care* 9, 241–247. <https://doi.org/10.1177/2048872620922795>
- Wang, H., Huang, D., Huang, H., Zhang, J., Guo, L., Liu, Y., Ma, H., Geng, Q., 2020. The Psychological Impact of COVID-19 Pandemic on Medical Staff in Guangdong, China: A Cross-Sectional Study. *Psychol. Med.* <https://doi.org/10.1017/S0033291720002561>
- Wang, H., Liu, Y., Hu, K., Zhang, M., Du, M., Huang, H., Yue, X., 2020. Healthcare workers' stress when caring for COVID-19 patients: An altruistic perspective. *Nurs. Ethics.* <https://doi.org/10.1177/0969733020934146>
- Wang, L.Q., Zhang, Meng, Liu, G.M., Nan, S.Y., Li, T., Xu, L., Xue, Y., Zhang, Min, Wang, L., Qu, Y.D., Liu, F., 2020. Psychological impact of coronavirus disease (2019) (COVID-19) epidemic on medical staff in different posts in China: A multicenter study. *J. Psychiatr. Res.* 129, 198–205. <https://doi.org/10.1016/j.jpsychires.2020.07.008>
- Wang, S., Wen, X., Dong, Y., Liu, B., Cui, M., 2020. Psychological Influence of Coronavirus Disease 2019 (COVID-19) Pandemic on the General Public, Medical Workers, and Patients With Mental Disorders and its Countermeasures. *Psychosomatics* 61, 616–624. <https://doi.org/10.1016/j.psym.2020.05.005>
- Wang, S., Xie, L., Xu, Y., Yu, S., Yao, B., Xiang, D., 2020. Sleep disturbances among medical workers during the outbreak of COVID-2019. *Occup. Med. (Chic. Ill)*. 70, 364–369. <https://doi.org/10.1093/occmed/kqaa074>
- Wang, Y., Ma, S., Yang, C., Cai, Z., Hu, S., Zhang, B., Tang, S., Bai, H., Guo, X., Wu, J., Du, H., Kang, L., Tan, H., Li, R., Yao, L., Wang, G., 2020a. Acute psychological effects of Coronavirus Disease 2019 outbreak among healthcare workers in China: a cross-sectional study. *Transl. Psychiatry.* <https://doi.org/10.1038/s41398-020-01031-w>

- Wang, Y., Ma, S., Yang, C., Cai, Z., Hu, S., Zhang, B., Tang, S., Bai, H., Guo, X., Wu, J., Du, H., Kang, L., Tan, H., Li, R., Yao, L., Wang, G., Liu, Z., 2020b. Acute psychological effects of Coronavirus Disease 2019 outbreak among healthcare workers in China: a cross-sectional study. *Transl. Psychiatry* 10. <https://doi.org/10.1038/s41398-020-01031-w>
- Wańkiewicz, P., Szylińska, A., Rotter, I., 2020. Assessment of mental health factors among health professionals depending on their contact with covid-19 patients. *Int. J. Environ. Res. Public Health* 17, 1–8. <https://doi.org/10.3390/ijerph17165849>
- Wasim, T., Raana, G., Bushra, N., Riaz, A., 2020. Effect of COVID-19 Pandemic on Mental Wellbeing of Healthcare Workers in Tertiary Care Hospital. *Ann. King Edward Med. Univ.* 26, 140–144.
- Wilson, W., Raj, J.P., Rao, S., Ghiya, M., Nedungalaparambil, N.M., Mundra, H., Mathew, R., 2020. Prevalence and Predictors of Stress, anxiety, and Depression among Healthcare Workers Managing COVID-19 Pandemic in India: A Nationwide Observational Study. *Indian J. Psychol. Med.* 42, 353–358. <https://doi.org/10.1177/0253717620933992>
- Xiang, Y.T., Yang, Y., Li, W., Zhang, L., Zhang, Q., Cheung, T., Ng, C.H., 2020. Timely mental health care for the 2019 novel coronavirus outbreak is urgently needed. *The Lancet Psychiatry* 7, 228–229. [https://doi.org/10.1016/S2215-0366\(20\)30046-8](https://doi.org/10.1016/S2215-0366(20)30046-8)
- Xiao, X., Zhu, X., Fu, S., Hu, Y., Li, X., Xiao, J., 2020. Psychological impact of healthcare workers in China during COVID-19 pneumonia epidemic: A multi-center cross-sectional survey investigation. *J. Affect. Disord.* 274, 405–410. <https://doi.org/10.1016/j.jad.2020.05.081>
- Xiaoming, X., Ming, A., Su, H., Wo, W., Jianmei, C., Qi, Z., Hua, H., Xuemei, L., Lixia, W., Jun, C., Lei, S., Zhen, L., Lian, D., Jing, L., Handan, Y., Haitang, Q., Xiaoting, H.,

- Xiaorong, C., Ran, C., Qinghua, L., Xinyu, Z., Jian, T., Jing, T., Guanghua, J., Zhiqin, H., Nkundimana, B., Li, K., 2020. The psychological status of 8817 hospital workers during COVID-19 Epidemic: A cross-sectional study in Chongqing. *J. Affect. Disord.* 276, 555–561. <https://doi.org/10.1016/j.jad.2020.07.092>
- Xing, J., Sun, N., Xu, J., Geng, S., Li, Y., 2020. Study of the mental health status of medical personnel dealing with new coronavirus pneumonia. *PLoS One* 15, 1–10. <https://doi.org/10.1371/journal.pone.0233145>
- Yáñez, J.A., Jahanshahi, A.A., Alvarez-Risco, A., Li, J., Zhang, S.X., 2020. Anxiety, distress, and turnover intention of healthcare workers in Peru by their distance to the epicenter during the COVID-19 crisis. *Am. J. Trop. Med. Hyg.* 103, 1614–1620. <https://doi.org/10.4269/ajtmh.20-0800>
- Yang, X., Zhang, Y., Li, S., Chen, X., 2020a. Risk factors for anxiety of otolaryngology healthcare workers in Hubei province fighting coronavirus disease 2019 (COVID □ 19). *Soc. Psychiatry Psychiatr. Epidemiol.* 2019. <https://doi.org/10.1007/s00127-020-01928-3>
- Yang, X., Zhang, Y., Li, S., Chen, X., 2020b. Risk factors for anxiety of otolaryngology healthcare workers in Hubei province fighting coronavirus disease 2019 (COVID-19). *Soc. Psychiatry Psychiatr. Epidemiol.* 2019. <https://doi.org/10.1007/s00127-020-01928-3>
- Yin, Q., Sun, Z., Liu, T., Ni, X., Deng, X., Jia, Y., Shang, Z., Zhou, Y., Liu, W., 2020. Posttraumatic stress symptoms of health care workers during the corona virus disease 2019. *Clin. Psychol. Psychother.* 27, 384–395. <https://doi.org/10.1002/cpp.2477>
- Zhan, M., Qin, Y., Xue, X., Zhu, S., 2020. Death from COVID-19 of 23 health care workers in China. *N. Engl. J. Med.* 382, 2267–2268. <https://doi.org/10.1056/NEJMc2005696>
- Zhan, Y., Liu, Y., Liu, H., Li, M., Shen, Y., Gui, L., Zhang, J., Luo, Z., Tao, X., Yu, J., 2020.

Factors associated with insomnia among Chinese frontline nurses Fighting against COVID-19 in Wuhan: A cross-sectional survey. *J. Nurs. Manag.*

<https://doi.org/10.1111/jonm.13094>

Zhang, M., Zhou, M., Tang, F., Wang, Y., Nie, H., Zhang, L., You, G., 2020. Knowledge, attitude, and practice regarding COVID-19 among healthcare workers in Henan, China. *J. Hosp. Infect.* 105, 183–187. <https://doi.org/10.1016/j.jhin.2020.04.012>

Zhang, S.X., Liu, J., Afshar Jahanshahi, A., Nawaser, K., Yousefi, A., Li, J., Sun, S., 2020. At the height of the storm: Healthcare staff's health conditions and job satisfaction and their associated predictors during the epidemic peak of COVID-19. *Brain. Behav. Immun.* 87, 144–146. <https://doi.org/10.1016/j.bbi.2020.05.010>

Zhang, W.R., Wang, K., Yin, L., Zhao, W.F., Xue, Q., Peng, M., Min, B.Q., Tian, Q., Leng, H.X., Du, J.L., Chang, H., Yang, Y., Li, W., Shangguan, F.F., Yan, T.Y., Dong, H.Q., Han, Y., Wang, Y.P., Cosci, F., Wang, H.X., 2020. Mental Health and Psychosocial Problems of Medical Health Workers during the COVID-19 Epidemic in China. *Psychother. Psychosom.* 89, 242–250. <https://doi.org/10.1159/000507639>

Zhou, B., She, J., Wang, Y., Ma, X., 2020. *cept cr ip t Ac ce pt us cr t 1–14.*

Zhou, Y., Cai, S., Jin, M., Jiang, C., Xu, N., Duan, C., Peng, X., Zhao, J., Ma, X., 2020. Economic burden for retinoblastoma patients in China. *J. Med. Econ.* 6998. <https://doi.org/10.1080/13696998.2020.1831518>

Zhou, Y., Wang, W., Sun, Y., Qian, W., Liu, Z., Wang, R., Qi, L., Yang, J., Song, X., Zhou, X., Zeng, L., Liu, T., Li, Z., Zhang, X., Wang, W., Sun, Y., Qian, W., Liu, Z., Wang, R., Qi, L., Yang, J., Song, X., Zhou, X., Zeng, L., Liu, T., Li, Z., Zhang, X., 2020. *Journal Pre-proof.*

Zhou, Yifang, Yang, Y., Shi, T., Song, Y., Zhou, Yuning, Zhang, Z., Guo, Y., Li, X., Liu, Y., Xu, G., Cheung, T., Xiang, Y.T., Tang, Y., 2020. Prevalence and Demographic

Correlates of Poor Sleep Quality Among Frontline Health Professionals in Liaoning Province, China During the COVID-19 Outbreak. *Front. Psychiatry* 11, 1–6.

<https://doi.org/10.3389/fpsyt.2020.00520>

Zhu, J., Sun, L., Zhang, L., Wang, H., Fan, A., Yang, B., Li, W., Xiao, S., 2020. Prevalence and Influencing Factors of Anxiety and Depression Symptoms in the First-Line Medical Staff Fighting Against COVID-19 in Gansu. *Front. Psychiatry* 11, 1–6.

<https://doi.org/10.3389/fpsyt.2020.00386>

Zhu, Z., Ph, D., Xu, S., Ph, D., Wang, H., Med, M., Liu, Z., Ph, D., 2020a. COVID-19 in Wuhan □ : Immediate Psychological Impact on 5062 Health Workers.

Zhu, Z., Xu, S., Wang, H., Liu, Z., Wu, J., Li, G., Miao, J., Zhang, C., Yang, Y., Sun, W., Zhu, S., Fan, Y., Chen, Y., Hu, J., Liu, J., Wang, W., 2020b. COVID-19 in Wuhan: Sociodemographic characteristics and hospital support measures associated with the immediate psychological impact on healthcare workers. *EClinicalMedicine* 24.

<https://doi.org/10.1016/j.eclinm.2020.100443>

Zhu, Z., Xu, S., Wang, H., Liu, Z., Wu, J., Li, G., Miao, J., Zhang, C., Yang, Y., Sun, W., Zhu, S., Fan, Y., Hu, J., Liu, J., Wang, W., 2020c. COVID-19 in Wuhan: Immediate Psychological Impact on 5062 Health Workers. *EClinicalMedicine*.

<https://doi.org/10.1101/2020.02.20.20025338>