

**TRANSGENERATIONAL EFFECT OF EARLY CHILDHOOD FAMINE EXPOSURE
IN THE COHORT OF LENINGRAD SIEGE SURVIVORS' OFFSPRING.**

Kristina Tolkunova^{1*}, Dmitrii Usoltsev^{1,2*}, Ekaterina Moguchaia¹, Maria Boyarinova¹,
Ekaterina Kolesova¹, Anastasia Erina¹, Trudy Voortman³, Elena Vasilyeva¹, Anna
Kostareva¹, Evgeny Shlyakhto¹, Alexandra Konradi^{1,2}, Oxana Rotar^{1,#}, Mykyta
Artomov^{1,2,4,5,#}

1 - Almazov National Medical Research Centre, Saint Petersburg, Russia

2 - ITMO University, St. Petersburg, Russia

3 –Department of Epidemiology, Erasmus MC, University Medical Center, Rotterdam, The
Netherlands

4 - Institute for Genomic Medicine, Nationwide Children's Hospital, Columbus, OH, USA

5 - Department of Pediatrics, College of Medicine, Ohio State University, Columbus, OH,
USA

* - contributed equally

- Correspondence: mykyta.artomov@nationwidechildrens.org, oxana.rotar@gmail.com

The authors declare no conflict of interest

1 **Abstract**

2 **Introduction:** Environmental exposure during early life development can affect
3 disease risk in late-life period. There is little data on the transmission of phenotypic features
4 from famine-exposed individuals to the next generations.

5 **Objective:** The purpose of our study was to investigate the association of parental
6 starvation in the perinatal period and the period of early childhood with the phenotypic
7 features observed in two generations of descendants of Leningrad Siege survivors (DLSS).

8 **Design and methods:** We examined 54 children and 30 grandchildren of 58 besieged
9 Leningrad residents who suffered from starvation in early childhood and prenatal age during
10 the Second World War. Controls from the population-based national epidemiological ESSE-
11 RF study (n=175) were matched on sex, age and body mass index (BMI). Anthropometry,
12 blood pressure (BP) measurement, and blood tests were performed for both groups.
13 Sociodemographic and lifestyle information was collected via questionnaires. Phenotypes of
14 controls and descendants (combined and children and grandchildren separately) were
15 compared, taking into account multiple testing.

16 **Results:** Comparison of two generations descendants with corresponding control
17 groups revealed significantly higher creatinine level and lower glomerular filtration rate
18 (GFR), both in meta-analysis and in both generations independently. The mean values of GFR
19 for all groups were detected within the normal range. Additionally, independent of the
20 creatinine level, differences in the eating pattern were detected: insufficient fish consumption
21 and excessive red meat consumption were significantly more frequent in the children of the
22 Leningrad siege survivors compared with controls. BP and blood lipids did not differ between
23 the groups.

24 **Conclusions:** Ancestral famine exposure in the prenatal period and early childhood
25 may contribute to a decrease of in kidney filtration capacity and altered eating pattern in at
26 least two generations of besieged Leningrad offspring.

27 **Introduction**

28 The theory of fetal origin of health and diseases suggests that parental environmental
29 exposure during fetal development may be an important risk factor for adult-onset diseases, in
30 particular, for cardiovascular diseases (CVD) (1). Susceptibility to diseases resulting from
31 fetal programming can be observed directly after birth in exposed individuals or transmitted
32 from the exposed generation to the descendants.

33 Major famine studies of the Dutch Famine 1944-1945 and the Chinese Famine 1959-
34 1962 have shown that insufficient nutrition during prenatal and early life periods have critical
35 impacts on fetal programming. Importantly, not only were individuals directly exposed to
36 famine during the prenatal period shown to have phenotypic consequences of exposure in
37 later life. It was suggested that the effect of fetal programming can be transmitted to the
38 descendants of exposed individuals, potentially through epigenetic mechanisms. For example,
39 the Dutch Hunger study observed a higher frequency of neonatal obesity in children of
40 women exposed to malnutrition prenatally and a higher body mass index (BMI) at an older
41 age in the offspring of prenatally malnourished men (2,3). The China Health and Nutrition
42 Survey reported significantly lower glomerular filtration rate (GFR), and a faster than
43 expected increase in BMI with age, waist circumference, and blood pressure (BP) in the first
44 generation of descendants of exposed individuals (4,5). Therefore, transgenerational
45 transmission of the famine exposure phenotype could be observed across at least one
46 generation of descendants.

47 The Siege of Leningrad was another tragic example of catastrophic malnutrition of the
48 city's residents having experienced a major supply deficit for two and a half years during the
49 Second World War (6). There are some similarities but also difference with the two other
50 major famine studies: The Siege of Leningrad was longer than Dutch Famine (28 months
51 compared to about 6 months, respectively), the weather was significantly colder, affecting the

52 calorie demand and potential hypothermia, and daily rations were not restored after breaking
53 the blockade (in the Netherlands daily rations quickly rose within a week) (7). The Chinese
54 famine, which is comparable in duration to the famine during the Siege of Leningrad, was
55 characterized by pandemic nature and uneven intensity distribution within the country (8).
56 And both previously mentioned studies involved only one generation of descendants.

57 The purpose of our study was to investigate the association of starvation in the
58 prenatal period and the period of early childhood with phenotypic differences later in life in
59 two generations of descendants of Leningrad Siege survivors (DLSS).

60 **Materials and methods**

61 In 2009-2011, 305 Leningrad siege survivors who starved in early childhood and/or in
62 utero during the Second World War (1941-1944) were examined in the Almazov National
63 Medical Research Centre as part of an observational cohort study, the design and general
64 characteristics were described in previous publications (6). In 2020-2021, we examined 87
65 adult DLSS (57 children and 30 grandchildren from 55 families) of 58 Siege survivors, with
66 the current pregnancy being the only exclusion criteria (**Figure 1A**). A subsample of a
67 population-based cohort (N=1,600) of St. Petersburg residents aged 25-64 years within the
68 national epidemiological study ESSE-RF (9) were selected as controls by matching gender,
69 age and BMI distribution to each generation of descendants. In the current study we refer to
70 Leningrad Siege survivors exposed to starvation in early life as generation zero (F_0), their
71 children as the first generation of DLSS (F_1) and grandchildren as the second generation (F_2).
72 The final control cohorts included 127 individuals for F_1 and 48 for F_2 .

73 The study was approved by the Local Ethics Committee of the Almazov National
74 Medical Research Centre (Protocol #2401-21 from the local ethics committee meeting #01-21
75 dated January 18, 2021). All participants signed an informed consent.

76 Body weight measurement was carried out on the scales of the brand VEM-150- Mass-
77 K (Russia), height - using the height meter RM-1 Diacoms (Russia), waist and hip
78 circumference - using a standard flexible centimeter tape. Abdominal obesity was determined
79 according to the JIS 2009 metabolic syndrome criteria 2009 (10) as: waist circumference ≥ 94
80 cm for males and ≥ 80 cm for females. The BMI was calculated as the ratio of body weight in
81 kilograms to height in meters squared. All respondents were classified as obese (BMI ≥ 30
82 kg/m²) or non-obese (BMI < 30 kg/m²).

83 Double measurement of blood pressure (BP) on the right arm was performed with the
84 automatic tonometer "Omron" (Japan) with intervals of 2 minutes, after resting for 5 minutes
85 in a sitting position, then a single measurement of BP was made 3 minutes after
86 verticalization was made. Hypertension was diagnosed if the mean BP level was $\geq 140/90$
87 mmHg and/or if patient was receiving antihypertensive therapy.

88 The creatinine, blood glucose, lipids study (total cholesterol – TC, low-density
89 lipoproteins – LDL, high-density lipoproteins – HDL, triglycerides – TG) measurement was
90 performed in fasting state (AbbotArchitect 8000 (USA)). The levels of leptin (Leptin ELISA,
91 DBC (Canada)) and adiponectin (Adiponectin ELISA, Mediagnost (Germany)) levels were
92 determined by enzyme immunoassay. We assessed glucose metabolism disorders: participants
93 were divided into fasting hyperglycemia group (glucose level ≥ 5.6 mmol/L) and diabetes
94 mellitus group (the patients self-reported about diabetes). The GFR was calculated using the
95 CKD-EPI formula (11). Groups of participants with high total cholesterol (> 4.9 mmol/L),
96 LDL (> 3.0 mmol/L), triglycerides (≥ 1.7 mmol/L), and reduced HDL (in male < 1.0 and in
97 female < 1.2 mmol/L) were also formed; hypolipidemic therapy was taken into account (12).

98 Behavioral and socioeconomic risk factors were evaluated with questionnaires and
99 interviews: diet pattern, smoking status, alcohol consumption, duration of sleep, level of
100 education, physical activity. The consumption of fresh vegetables and fruits was assessed

101 (respondents were divided into those who had fresh fruits and vegetables in their daily diet
102 and those who consumed irregularly). Excess salt intake was considered to be salting of
103 ready-made foods. Consumption of fish (200 grams) less than 1-2 times a week was
104 considered insufficient. Daily consumption of red meat (150 grams or more) was considered
105 excessive (13). If the respondent consumed ≥ 6 pieces/spoons of sugar per day or daily/almost
106 daily intake of sweets/confectionery, it was considered excessive intake of sweets. The
107 duration of sleep of less than 6 hours a day was considered insufficient. Moderate physical
108 activity ≥ 150 minutes per week was sufficient. Inactivity was recorded in the case of being in
109 a sitting position for more than 9 hours on a weekday. The assessment of smoking status was
110 also carried out, respondents were divided into groups: smoking at the moment, in the past,
111 and those who have never smoked. The number of alcohol doses and the frequency of alcohol
112 consumption were assessed and respondents were divided into those who regularly took
113 alcoholic beverages and those who did not consume at all.

114 Mathematical and statistical data analysis was implemented using the R-4.0
115 programming language (14) and libraries dplyr (v1.0.7) (15), ggplot2 (v3.3.5) (16), tidyr
116 (v1.1.4) (17). Quantitative parameters were described using median values and lower and
117 upper quartiles (Q1-Q3). Nominal data were described with absolute values and percentages.

118 The prevalence of 44 phenotypic risk factors in both generations of DLSS was
119 simultaneously assessed using a logistic model adjusted for sex, generation and BMI. In case
120 of absence of a phenotype in one of the comparison groups, a simple Fisher test was used
121 instead of logistic regression. We report both effect size (beta) and odds ratio for each
122 phenotype. If a phenotype was obtained from the other phenotype through calculations, we
123 considered the pair as a single phenotype for multiple hypothesis correction with Bonferroni
124 approach. In total, 34 independent phenotypes were analyzed and the significance threshold
125 for the p-value was $0.05/34=0.0015$.

126

127

128 **Results**

129 87 DLSS (mean age: 43.8 years; sd: +/- 13.4 years; range: 18-63 years; 44% men)
130 were observed during an ambulatory visit at Almazov National Medical Research Centre (St.
131 Petersburg, Russia). The prevalence of abdominal and obesity did not differ significantly
132 among BMI-matched participants. The BP levels and the prevalence of hypertension was non-
133 remarkable compared to population average. The DLSS group had nominally higher HDL
134 levels. The differences are described in more details in the supplementary and Tab. S1.

135 The DLSS cohort had 57 and 30 individuals from the first and F₂, respectively (**Fig.**
136 **1A**). Initially we performed matching of the DLSS cohort to a group of 1,600 controls from
137 the ESSE population cohort from St. Petersburg. We excluded individuals with known
138 cardiovascular diseases and outliers (more than 6 standard deviations) based on blood
139 biochemical analyses and blood pressure measurements from the control pool. The final
140 control pool available for matching had 1,142 individuals.

141 Controls of the same gender from the ESSE populational cohort were selected for each
142 individual so that the age and BMI of the controls did not differ for more than two units. No
143 controls were found for 8 DLSS individuals (7 youngest individuals from F₂) and (1
144 individual with high BMI from F₂), therefore they were excluded from further analyses. 3
145 individuals from F₁ of DLSS were excluded because they had cardiovascular disease. The
146 final cohort included a total of 175 controls and 76 DLSS (127 controls and 54 DLSS for F₁;
147 48 controls and 22 DLSS for F₂, **Sup. Fig. S1**).

148 Of all the biomarker phenotypes that were analyzed and compared, only creatinine
149 ($p=1.367 \times 10^{-7}$, $\beta=0.092$, $se=0.017$) and GFR ($p=3.94 \times 10^{-8}$, $\beta=-0.08$, $se=0.015$) passed
150 the Bonferroni significance threshold (**Fig. 1B, Sup. Tab. S1**). Additionally, insufficient fish

151 consumption was more frequently observed in the DLSS group compared to controls ($p =$
152 3.15×10^{-6} ; $\beta = 1.361$; $\beta \text{ se} = 0.292$), and also excessive red meat consumption was more
153 frequently observed in the DLSS group ($p = 1.8 \times 10^{-4}$; $\beta = 1.168$; $\beta \text{ se} = 0.312$) (Sup. Tab.
154 S1). Among cardiometabolic risk factors, nominal differences were observed for leptin and
155 HDL levels that were higher in the DLSS group, but still within normal ranges (leptin: $p =$
156 3.12×10^{-3} , $\beta = 0.037$; HDL: $p = 9.47 \times 10^{-3}$, $\beta = 1.08$) and not significant after multiple
157 hypothesis testing correction.

158 Further, we investigated whether this effect could be observed in each generation
159 separately. We analyzed the phenotypic data within each generation independently, adjusting
160 the model for age, BMI, sex. Interestingly, the effect of creatinine and GFR was observed in
161 both generations - F₁: creatinine ($p = 9.71 \times 10^{-5}$, $\beta = 0.071$, $\text{se} = 0.018$) and GFR ($p = 1.09 \times 10^{-5}$,
162 $\beta = -0.076$, $\text{se} = 0.017$) and F₂: creatinine ($p = 3.56 \times 10^{-4}$, $\beta = 0.208$, $\text{se} = 0.058$) and GFR
163 ($p = 4.56 \times 10^{-4}$, $\beta = -0.179$, $\text{se} = 0.051$) (Sup. Tab. S1). The differences in fish and red meat
164 intake habits passed Bonferroni correction only for F₁ (red meat: $p = 3.10 \times 10^{-4}$; $\beta = 1.39$;
165 $\beta \text{ se} = 0.385$. Fish: $p = 5.23 \times 10^{-4}$; $\beta = 1.197$; $\beta \text{ se} = 0.345$), but estimates for F₂ were in
166 similar direction and magnitude (red meat: $p = 0.237$; $\beta = 0.653$; $\beta \text{ se} = 0.55$. Fish: $p =$
167 0.0015 ; $\beta = 1.897$; $\beta \text{ se} = 0.598$) (Sup. Tab. S1).

168 In addition to the consumption of red meat and fish, several other nutritional
169 phenotypes were nominally significant, therefore, we used the principal component analysis
170 (PCA) based on food consumption-related phenotypes (fish, red meat, salt, sugar, vegetable
171 intake, and leptin) to investigate differences in food pattern between DLSS and controls (**Sup.**
172 **Fig. S2A**). We found that PC2, which was characterized by a low fish intake, leptin levels,
173 and high salt and sugar intake, and differed significantly between descendants and controls for
174 both the meta-analysis ($p = 3.77 \times 10^{-9}$; $\beta = -1.06$; $\beta \text{ se} = 0.18$) and for each generation

175 separately: F_1 ($p = 2.15 \times 10^{-6}$; $\beta = -1.00$; $\beta \text{ se} = 0.211$); F_2 ($p = 5.29 \times 10^{-4}$; $\beta = -1.243$;
176 $\beta \text{ se} = 0.359$) (**Sup. Tab. S2**).

177 Subsequently, we investigated whether the food pattern was related to the observed
178 difference in creatinine level. From the control population that was not used in the previous
179 comparisons, we selected individuals to match sex, age, and PC2 distributions obtained from
180 the PCA of food intake in the DLSS cohort (**Sup. Fig. S2B**). Comparison of this cohort with
181 the previously selected control cohort for DLSS comparison, we did not observe significant
182 differences in creatinine and GFR ($p = 0.75$; $p = 0.61$, respectively, **Sup. Fig. S2C**). Therefore,
183 the pattern of food consumption and biochemical markers of renal function can be interpreted
184 as independent associations.

185

186 **Discussion**

187 Our results demonstrate a significantly higher creatinine level and a lower glomerular
188 filtration rate in the descendant generations of individuals exposed to prenatal or early
189 childhood starvation. It is known that most human organs have a critical development period
190 at the prenatal stage. In the case of serious nutritional deficiency, the number of functioning
191 glomeruli decreases, leading to increased filtration load, increased GFR throughout postnatal
192 life, and subsequent damage to functioning glomeruli (18). Reports on the Northern China
193 famine cohort study also presented significantly lower calculated GFR in prenatally exposed
194 participants to starvation and their children compared to controls. Data on the second
195 generation were not available in this study. It has been hypothesized that the starvation of the
196 ancestors in early childhood contributes to a decrease in the renal filtration capacity in their
197 offspring. This may be due to transgenerational changes in the epigenome through the
198 programming of nephron deficiency in germ cells (4). In the long term, renal filtration
199 capacity can cause glomerulosclerosis, and finally hypertension in adulthood (18).

200 Observed differences in the dietary pattern between the descendants and controls do
201 not have a clear direction of effect. On the one hand, increased red meat and decreased fish
202 consumption increase the risk of cardiometabolic disorders (19,20). On the other hand, the
203 decreased salt consumption and higher leptin levels might indicate a more healthy approach to
204 food selection. Comparison of the subsample of the population cohort that matched in dietary
205 behavior with DLSS with the control population indicated that the observed differences in
206 creatine levels and food selection were independent of each other. Therefore, it is likely that
207 there is an independent effect of dietary behavior imprinting observed in DLSS.

208 The absence of statistically significant differences in the prevalence of cardiometabolic
209 diseases can be explained on the one hand by the small size of the cohort studied, and on the
210 other hand by the relatively young age of most of the participants. Also, it is important to note
211 that in this work we studied the offspring, among whom most of the parents are still alive,
212 which may have resulted in survivorship bias. In addition, both a decrease in fertility and an
213 increase in perinatal mortality during famine may have led to selective births of healthier or
214 more resilient individuals. The besieged Leningrad residents included in the study could
215 therefore represent an artificially "healthy" sample.

216 Conclusively, observation of phenotypic effects associated with early-in-life
217 exposures, separated not only in time, but also by generations serve yet another example of
218 Barker's hypothesis of fetal origins of chronic diseases of adult life. Our study suggests that
219 not only individuals directly exposed to traumatic events, but their offspring will benefit from
220 additional health screenings, in particular assessment of kidney function, for earlier detection
221 of disease symptoms and their successful treatment.

222

223 **Figure Caption**

224 **Figure 1.** Comparison of descendants of Leningrad Siege survivors (DLSS) with
225 population controls from St. Petersburg. **(A)** Graphical representation of all family trees in the
226 DLSS cohort (Siege survivors -F₀, the 1st generation of descendants - F₁, the 2nd generation of
227 descendants - F₂); **(B)** Differences in GFR and creatinine levels between each generation of
228 DLSS and the corresponding controls from the populational cohort of St. Petersburg.

Author Contributions

K.T., D.U., O.R., M.A. designed and conceived the study.

K.T., D.U., E.M., M.B., E.K., A.E., T.V., O.R., M.A. collected and analyzed the data.

K.T., D.U., O.R., M.A. designed statistical analysis approach.

K.T., E.M., M.B., E.K., A.E., E.V., An.K., Al.K., O.R. recruited and supervised patients.

Al.K., E.S., M.A. – obtained funding.

O.R., M.A. supervised the study

K.T., D.U., O.R., M.A. wrote the manuscript

All authors edited and reviewed the manuscript text

Funding

K.T., D.U., E.M., M.B., E.K., A.E., E. V., An.K., E.S., Al.K., O.R. were supported by the Ministry of Science and Higher Education of the Russian Federation (Agreement #075-15-2022-301).

M.A. was supported by Nationwide Foundation Pediatric Innovation Fund.

References

1. Barker DJ. Fetal origins of coronary heart disease. *BMJ*. 1995;311(6998):171-4. doi: 10.1136/bmj.311.6998.171

2. Painter RC, Osmond C, Gluckman P, Hanson M, Phillips DI, Roseboom TJ. Transgenerational effects of prenatal exposure to the Dutch famine on neonatal adiposity and health in later life. *BJOG*. 2008;115:1243–9. doi:10.1111/j.1471-0528.2008.01822
3. Veenendaal MVE, Painter RC, de Rooij SR, Bossuyt PM, van der Post JA, Gluckman PD, et al. Transgenerational effects of prenatal exposure to the 1944-45 Dutch famine. *BJOG*. 2013;120:548–54. doi:10.1111/1471-0528.12136pmid
4. Jiang W, Han T, Duan W, Dong Q, Hou W, Wu H, et al. Prenatal famine exposure and estimated glomerular filtration rate across consecutive generations: association and epigenetic mediation in a population-based cohort study in Suihua China. *Aging (Albany NY)*. 2020;12(12):12206-12221. doi:10.18632/aging.103397
5. Li J, Yang Q, An R, Sesso HD, Zhong VW, Chan KHK, et al. Famine and Trajectories of Body Mass Index, Waist Circumference, and Blood Pressure in Two Generations: Results From the CHNS From 1993-2015. *Hypertension*. 2022;79(3):518-531. doi: 10.1161/HYPERTENSIONAHA.121.18022
6. Rotar O, Moguchaia E, Boyarinova M, Kolesova E, Khromova N, Freylikhman O, et al. Seventy years after the siege of Leningrad: does early life famine still affect cardiovascular risk and aging? *J Hypertens*. 2015;33(9):1772-9. doi: 10.1097/HJH.0000000000000640
7. Fredrick J, Stare MD, Nutritional Conditions in Holland. *Nutrition Reviews*. 1945;8(3):225–227. doi:10.1111/j.1753-4887.1945.tb08647.x
8. Susser E, St Clair D. Prenatal famine and adult mental illness: interpreting concordant and discordant results from the Dutch and Chinese Famines. *Soc Sci Med*. 2013;97:325-30. doi: 10.1016/j.socscimed.2013.02.049
9. Orlov AV, Rotar OP, Boyarinova MA, Alieva AS, Dudorova EA, Kolesova EP, et al. Gender characteristics of the prevalence of behavioral risk factors in St. Petersburg residents. *Bulletin*

of the Russian Academy of Medical Sciences. 2015;70(5):585-591. (In Russ.) doi: 10.15690/vramn.v70.i5.1446.

10. Alberti KG, Eckel RH, Grundy SM, Zimmet PZ, Cleeman JI, Donato KA, et al. International Diabetes Federation Task Force on Epidemiology and Prevention; National Heart, Lung, and Blood Institute; American Heart Association; World Heart Federation; International Atherosclerosis Society; International Association for the Study of Obesity. Harmonizing the metabolic syndrome: a joint interim statement of the International Diabetes Federation Task Force on Epidemiology and Prevention; National Heart, Lung, and Blood Institute; American Heart Association; World Heart Federation; International Atherosclerosis Society; and International Association for the Study of Obesity. *Circulation*. 2009;120(16):1640–1645, doi: 10.1161/CIRCULATIONAHA.109.192644
11. Moiseev VC, Mukhin NA, Smirnov AV, Kobalava JD, Bobkova IN, Villevalde SV, et al. Cardiovascular risk and chronic kidney disease: cardio-nephroprotection strategies. *Russian Journal of Cardiology*. 2014;(8):7-37. (In Russ.) doi: 10.15829/1560-4071-2014-8-7-37
12. Kobalava ZD, Konradi AO, Nedogoda SV, Shlyakhto EV, Arutyunov GP, Baranova EI, et al. Arterial hypertension in adults. Clinical guidelines 2020. *Russian Journal of Cardiology*. 2020;25(3):3786. (In Russ.) doi:10.15829/1560-4071-2020-3-3786
13. Visseren FLJ, Mach F, Smulders YM, Carballo D, Koskinas KC, Bäck M, et al; ESC National Cardiac Societies; ESC Scientific Document Group. 2021 ESC Guidelines on cardiovascular disease prevention in clinical practice. *Eur Heart J*. 2021;42(34):3227-3337. doi: 10.1093/eurheartj/ehab484
14. R Core Team (2021). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.
15. Wickham H, François R, Henry L, Müller K. (2021). dplyr: A Grammar of Data Manipulation. R package version 1.0.7. <https://CRAN.R-project.org/package=dplyr>

16. Wickham H. *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York, 2016.
ISBN: 978-3-319-24277-4
17. Wickham H., Henry L. (2021). *tidyr: Tidy Messy Data*. R package version 1.1.4.
<https://CRAN.R-project.org/package=tidyr>
18. Lejarraga H. Perinatal origin of adult diseases. *Arch Argent Pediatr*. 2019;117(3):e232-e242.
English, Spanish. doi: 10.5546/aap.2019.eng.e232
19. Bechthold A, Boeing H, Schwedhelm C, Hoffmann G, Knüppel S, Iqbal K, et al. Food groups and risk of coronary heart disease, stroke and heart failure: A systematic review and dose-response meta-analysis of prospective studies. *Crit Rev Food Sci Nutr*. 2019;59(7):1071-1090. doi: 10.1080/10408398.2017.1392288
20. Pan L, Chen L, Lv J, Pang Y, Guo Y, Pei P, et al. Association of Red Meat Consumption, Metabolic Markers, and Risk of Cardiovascular Diseases. *Front Nutr*. 2022;9:833271. doi: 10.3389/fnut.2022.833271

