- 1 Trends in STI testing, diagnoses, and use of online chlamydia self-sampling services
- 2 among young people during the first year of the COVID-19 pandemic in England
- 3
- 4 Tamilore Sonubi¹, Dahir Sheik-Mohamud¹, Natasha Ratna¹, James Bell¹, Alireza
- 5 Talebi¹, Catherine H Mercer^{2,3}, Katy Sinka¹, Stephanie J Migchelsen¹, Kate Folkard¹
- 6 and Hamish Mohammed^{1,3}
- 7 1. Blood Safety, Hepatitis, Sexually Transmitted Infections (STI) and HIV
- 8 Division UK Health Security Agency, London, UK
- 9 2. Institute for Global Health, University College London, London, UK
- 103. The National Institute for Health Research Health Protection Research Unit in
- Blood Borne and Sexually Transmitted Infections at University College
- London in partnership with UK Health Security Agency, London, UK
- 13 Corresponding author:
- 14 Tamilore Sonubi,
- 15 UK Health Security Agency,
- 16 61 Colindale Avenue,
- 17 London, NW9 5EQ
- 18 Tami.Sonubi@kent.gov.uk
- 19
- 20 Contributors: TS, DSM, NR, JB, SJM and HM planned the analysis. TS, DSM, NR,
- JB and AT reviewed and analysed the results of the analysis. TS and DSM wrote the
- 22 first draft of the manuscript. All co-authors reviewed and edited the manuscript.
- Funding: The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.
- 25 Conflict of interest: None declared.
- Acknowledgments: The authors thank all laboratories and sexual health services that
- 27 report CTAD and GUMCAD surveillance data to UKHSA. We also thank colleagues
- in the CTAD, GUMCAD and NCSP teams at UKHSA. We acknowledge members of
- the National Institute for Health Research Health Protection Research Unit (NIHR
- 30 HPRU) in Blood Borne and Sexually Transmitted Infections (BBSTI) Steering
- 31 Committee: Professor Caroline Sabin (HPRU Director), Dr John Saunders (UKHSA
- 32 Lead), Professor Catherine Mercer, Dr Hamish Mohammed (previously Professor
- 33 Gwenda Hughes), Professor Greta Rait, Dr Ruth Simmons, Professor William
- Rosenberg, Dr Tamyo Mbisa, Professor Rosalind Raine, Dr Sema Mandal, Dr
- 35 Rosamund Yu, Dr Samreen Ijaz, Dr Fabiana Lorencatto, Dr Rachel Hunter, Dr Kirsty
- 36 Foster and Dr Mamooma Tahir. NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

- 1 Ethics statement: This study was undertaken for health protection purposes under
- 2 the permissions granted to UKHSA to collect and process pseudonymised
- 3 surveillance data under Regulation 3 of The Health Service (Control of Patient
- 4 Information) Regulations 2020 and under Section 251 of the NHS Act 2006.
- 5 Some findings from this research have been presented at the British Association for
- 6 Sexual Health and HIV annual conference in 2020 and 2021.

7

1 Abstract (242/250)

2 Purpose:

Measures to control COVID-19 reduced face-to-face appointments and walk-ins at
sexual health services (SHSs). Remote access to SHSs through online self-sampling
for STIs was increased. This analysis assesses how these changes affected service
use and STI testing among young people in England.

7 Methods:

8 Data on all chlamydia, gonorrhoea and syphilis tests from 2019-2020 amongst 9 English-resident 15-24 year olds (hereafter referred to as 'young people') were obtained from national STI surveillance datasets. We calculated proportional 10 differences in tests and diagnoses for each STI, by demographic characteristics 11 including age and socioeconomic deprivation, between 2019 and 2020. Among 12 13 those tested for chlamydia, we used binary logistic regression to determine crude and adjusted odds ratios (OR) between demographic characteristics and being 14 tested for chlamydia by an online service. 15

16 <u>Results:</u>

Compared to 2019, there were declines in testing (30% for chlamydia, 26% for
gonorrhoea, 36% for syphilis) and diagnoses (31% for chlamydia, 25% for
gonorrhoea and 23% for syphilis) among young people in 2020. These reductions
were greater amongst 15-19 year-olds (*vs.* 20-24 year-olds). Among young people
tested for chlamydia, those living in the least deprived areas were more likely to be
tested using an online self-sampling kit compared to those living in the most deprived
areas (males; OR=1.24[1.22-1.26], females; OR=1.28[1.27-1.30]).

24 Conclusion:

- 1 The first year of the COVID-19 pandemic in England saw declines in STI testing and
- 2 diagnoses in young people and disparities in the use of online chlamydia self-
- 3 sampling which risk widening existing health inequalities.
- 4 Key words: Young people, COVID-19 pandemic, STIs, Chlamydia, Gonorrhoea,
- 5 Syphilis, Socioeconomic deprivation, online STI testing, COVID-19 impact, STI service
- 6 provision
- 7
- 8 Implications and contributions (49/50)
- 9 There was a decrease in STI testing of young people during the first year of the
- 10 COVID-19 pandemic in England with larger reductions among teenagers. There was
- an increase in use of online STI self-sampling services but with inequalities in
- 12 provision which risk widening existing inequalities in sexual health.

1 Main text word count/limit: 2444/3500

2

3 Background

Sexually transmitted infections (STIs) present a public health challenge in England.
The burden of STIs is greatest amongst young people aged 16-24 years and they
are most likely to access sexual health services (SHSs) ^[2-6]. Between 2010-12, they
had the highest prevalence of chlamydia (3.1% of women and 2.3% of men in this
age group). The prevalence of gonorrhoea was highest amongst people 20-24 years
of age^[7].

The majority of chlamydia and gonorrhoea infections are asymptomatic, particularly
amongst women, and may result in poor sexual and reproductive health if left
untreated^[8]; this includes a higher risk of pelvic inflammatory disease (PID), and
ectopic pregnancy, which are preventable with early diagnosis and treatment^[9, 10].
Consequently, opportunistic chlamydia screening has been offered to sexually active
15-24 year olds in England since 2003 through the National Chlamydia Screening
Programme (NCSP).

There have been increases in attendances at SHSs between 2016-2020^[3]. However, 17 SHS delivery was greatly disrupted in March 2020 after the introduction of public 18 health measures to reduce COVID-19 transmission such as national lockdowns, the 19 requirement to stay at home, and social distancing. Moreover, the introduction of 20 legislation restricting social interaction may have resulted in some people with STI-21 related symptoms avoiding attending SHSs due to fear of being judged for breaking 22 these rules, creating a false sense of reduced demand on STI testing services^[11]. 23 However, to ensure continued provision of STI testing, SHSs across England were 24 25 rapidly reconfigured to provide more remote care via online consultations^[3].

To understand how STI testing, diagnoses and service use among 15-24 year olds in
England changed during the first year of the COVID-19 pandemic, we compared the
demographic characteristics of young people tested for STIs in 2019 and 2020 then,
among all young people tested for chlamydia, determined the correlates of being
tested via an online service instead of a face to face appointment.

6 <u>Methods</u>

7 Data description

8 We conducted a retrospective cohort study of 15-24 year olds (hereafter referred to as 'young people') residing in England who received an STI test or diagnosis utilising 9 10 data from the GUMCAD STI Surveillance System and the CTAD Chlamydia 11 Surveillance System. GUMCAD is a pseudonymised and depersonalised dataset of all attendances at SHSs in England and was used to obtain data on chlamydia, 12 gonorrhoea and syphilis tests and diagnoses in young people attending this setting; 13 syphilis diagnoses included primary, secondary and early latent stages^[12, 13]. CTAD 14 is a pseudonymised and depersonalised dataset of all publicly provided chlamydia 15 tests and diagnoses, including those made through the NCSP, and was the source 16 of data for chlamydia tests and diagnoses from community-based settings (those 17 18 offering non-specialist STI-related care such as general practices and pharmacies)^[14, 15]. All tests and diagnoses are coded by healthcare practitioners in 19 keeping with surveillance reporting specifications. To avoid double counting of tests 20 21 or diagnoses in each surveillance system, only one test or diagnosis of each STI for each person with a unique person identifier was counted within a 6-week episode^[15]. 22 Neither GUMCAD nor CTAD include personal identifiers so individuals cannot be 23 matched between datasets; individuals are identified using a clinic-specific patient 24 identification code in GUMCAD^[16] and a unique patient identifier number in CTAD^[14]. 25

1 The study period was from 1st January 2019 to 31st December 2020 (inclusive), where data from 2019 relates to the pre-COVID period and 2020 relates to the first 2 3 year of the COVID-19 pandemic. To be considered in the analysis data were restricted to people aged 15-24 years, at the time of the test or diagnosis, residing in 4 England. Residential location was defined using the lower super output area (LSOA), 5 small geographical areas for the reporting of small area statistics with an average 6 7 size of 1,620 residents^[17]. To obtain a measure of area-level socioeconomic deprivation, LSOAs were used to match to guintiles of the 2019 Index of Multiple 8 9 Deprivation (IMD) dataset^[18]. Additionally, the LSOAs were matched to the 2011 census area classification to categorise young people as living either in an urban or 10 rural setting ^[19-21]. Ethnicity was categorised using the national Census classification, 11 as follows: Asian (including Bangladeshi, Chinese, Indian, Pakistani and any other 12 Asian background), Black African, Black Caribbean, Other Black ethnicity, Mixed 13 ethnicity (including White and Black Caribbean, White and Black African, White and 14 Asian and any other Mixed or Multiple ethnic background), Other, and White^[22]. 15

16 Statistical analysis

We determined the proportional change in the characteristics of young people tested 17 for STIs between 2019 and 2020. Demographic and clinical characteristics included 18 age group (15-19 or 20-24 years), area of residence (rural or urban), residential 19 area-level deprivation, (as defined by IMD guintile, where guintile 1 is the most 20 deprived and quintile 5 is the least deprived), ethnicity, gender and public health 21 region of residence (categorised as: East Midlands, East of England, London, North-22 East, North-West, South-East, South-West, West Midlands and Yorkshire and 23 Humber) these characteristics were compared for all three STIs as they can be 24 25 assessed in both CTAD and GUMCAD surveillance systems. Sexual orientation

1 (including heterosexual males, men who have sex with men [MSM], heterosexual females and women who have sex with women [WSW]), and HIV status (categorised 2 3 as; HIV diagnosed, HIV undiagnosed or unknown) were compared for gonorrhoea and syphilis as they are only collected in the GUMCAD surveillance system. Testing 4 services (categorised as physical or online services) were compared for chlamydia 5 only, as the CTAD surveillance system comprehensively captures all chlamydia 6 7 testing from all publicly-commissioned testing services. The Pearson's chi-square test was used to compare these characteristics across both years. 8

9 Subsequently, to assess any inequalities in the access to online self-sampling services for chlamydia testing (hereafter: 'online chlamydia testing'), we restricted 10 11 the sample to young people tested for chlamydia then used binary logistic regression to determine the crude and adjusted associations with being tested via an online 12 service (yes vs. no); all models were stratified by gender. Bivariate models were 13 14 created to determine the unadjusted odds ratios (ORs) for being tested online and residential area-level deprivation, as defined by IMD quintile (the primary 15 independent variable), and each potential confounder (year of test, age group, area 16 of residence and region of residence). All associations with a p-value less than 0.05 17 were considered to be statistically significant and all variables that had significant 18 crude associations were included in the multivariable model. Adjusted odds ratios 19 (aORs) were calculated using hierarchical modelling and covariates were added 20 using a forward building approach. Firstly, Model 1 was constructed with year of test 21 included a priori due to the scale up of online service provision during 2020^[3]. The 22 remaining covariates were added sequentially as follows: Model 2 was based on 23 Model 1 with age group included as a confounder. Model 3 was based on Model 2 24 25 with the addition of area of residence. Lastly, Model 4 comprised Model 3 with the

1 inclusion of region of residence. Ethnicity was excluded from the regression analysis due to a high degree of item non-response: 29% of young people tested for 2 3 chlamydia were reported with an unspecified ethnic group in CTAD. All data analyses were performed using version Stata v15 (College Station, TX, USA)^[23]. 4 5 Results 6 Trends in STI tests and diagnoses 7 There were 26-36% decreases in tests and diagnoses for chlamydia, gonorrhoea 8 and syphilis among young people between 2019 and 2020 (Tables 1-3). However, 9 there were greater proportional decreases among 15-19 compared to 20-24 year 10 olds. By ethnicity, testing and diagnoses of all 3 STIs decreased for all ethnic groups 11 with larger proportional declines among young people of Asian and Black non-12 African/non-Caribbean ethnicities. The number of chlamydia tests fell across all the

different types of services offering testing (47%; 1,041,553 to 554,299), with the

exception of online services where there was a 33% increase in testing between

15 2019 (271,684 tests) and 2020 (361,622 tests). Comparisons by sexual orientation

16 could only be done for gonorrhoea and syphilis and, in both cases, testing and

diagnoses fell to the largest extent (33-46%) among heterosexual men.

18 Correlates of being tested for chlamydia via an online service

19 Amongst all young people tested for chlamydia, those living in the least deprived

areas were more likely to be tested online (unadjusted odds ratios - males: 1.24

21 [1.22-1.26]; females: 1.28 [1.27-1.30]) compared to young people living in the most

22 deprived areas. This association remained after adjusting for confounders in the final

23 model (males: 1.29 [1.27-1.32]; females 1.32 [1.30-1.34]) (Table 4). In the final

24 model, there was a greater likelihood of being tested for chlamydia via an online

- 1 service in 2020 [(males: 2.81 [2.77-2.84]; females: 2.45 [2.44-2.47]) vs. 2019] and a
- 2 similarly increased likelihood amongst 20-24 year olds [(males: 1.47 [1.45-1.49];
- 3 females: 1.63 [1.61-1.64]) vs 15-19 year olds]. Online testing was also more likely
- 4 among residents of urban areas [(males: 1.17 [1.15-1.20]; females: 1.16 [1.15-1.17])
- 5 vs rural] and was generally less likely among all regions of residence compared to
- 6 London (*Appendix B*).

1 <u>Discussion</u>

There was a decrease in STI testing and diagnoses among young people during the
first year of the COVID-19 pandemic, with up to 50% larger decreases in teenagers.
In keeping with the reconfiguration of SHSs in 2020 to offer more remote
consultations, we found a 33% increase in chlamydia testing of young people via
online services, but there was evidence of inequalities in access to testing via this
modality.

8 Among young people tested for chlamydia, those living in the least deprived areas 9 were more likely to be tested for chlamydia online, compared to those living in the 10 most deprived areas. Further inequalities in chlamydia online testing were found, with young people living in rural areas or regions outside London and those aged 15-11 19 being less likely to be tested for chlamydia using an online service. This suggests 12 that there may be socioeconomic or structural barriers to online testing, which may 13 include lack of online access. 15-19 year olds may find it more difficult to be tested 14 for chlamydia using an online service if they are still living with their parents and are 15 unable to discreetly receive the self-sampling kit. The greater likelihood of being 16 tested for chlamydia online for young people living in London reflects the fact that 17 there is a pan-London online sexual health service for all London residents^[24]. 18

The reductions in STI testing between 2019 and 2020 are partly due to the extensive public health measures to help reduce the transmission of COVID-19^[25]. Moreover, individuals may have delayed their visits to SHSs due to fear of COVID infection^[26] and with lockdown restrictions it would have been difficult to meet and interact with new people, reducing the possibility of new sexual encounters^[26]. All these factors may have contributed to the decline in STI testing in 2020.

1 Our findings are consistent with international literature highlighting the negative impact the COVID-19 pandemic had on STI testing. A report from the EuroTEST 2 3 COVID-19 impact assessment consortium found that, among 34 countries in the World Health Organization European Region and in different test settings, 95% of 4 them tested less than half the expected number of people between March and May 5 2020, this decline continued until August 2020^[27]. Research in the USA found a 6 7 reduction in STI testing and case detection resulting in more than, 27,000 missed cases of chlamydia and 5,500 cases of gonorrhoea between March and June 8 9 2020^[25]. The implications of these missed cases are likely to be increased community transmission due to the asymptomatic nature of these STIs and 10 associated long-term sexual and reproductive health complications^[25]. Studies have 11 found that testing via online services is preferred over physical services, particularly 12 amongst young people^[28], but this may not be the case for teenagers who are living 13 14 at home. The advantages of online services include privacy and the ability to selfsample^[29]. Previous research has found that online testing is more likely to be used 15 by women and those between the ages of 20-30 compared to younger age groups. 16 17 Consistent with our findings, research conducted amongst online services and SHSs in London found those living in less deprived areas are more likely to use online 18 services when testing for an STI even when adjusting for confounders^[30]. 19

Our analysis benefitted from a very large sample from national surveillance datasets which included patient-level data with key demographic factors so we could robustly assess differences in testing patterns within different subgroups. However, our analysis is not without limitations. Urban and rural area classifications were based on the 2011 census (the most up to date dataset at the time of writing) and these may not be accurate for all areas of England in 2020. We were unable to adjust for

ethnicity in the regression analysis predicting being tested online for chlamydia due 1 to a high proportion of missing values for ethnicity in the CTAD surveillance system. 2 3 The regression analysis was restricted to chlamydia because we were only able to reliably identify all sources of online testing for chlamydia by using a combination of 4 CTAD and GUMCAD data at the time of writing. While GUMCAD is a rich source of 5 data on STI testing, it underestimates online testing for gonorrhoea and syphilis as it 6 7 could only identify online testing by standalone online providers, and not online 8 testing provided as an alternate service by physical SHSs, in 2019 and 2020. 9 Similarly, we did not perform a regression analysis with count data to determine 10 correlates of being tested for chlamydia online vs not being tested – this is because of a lack of underlying population data for all key variables (e.g. age-group, gender 11 and residential area-level deprivation). However, as we have comprehensive data on 12 all young people tested for chlamydia, we were able to assess the correlates of 13 14 being tested online. Whilst we included deprivation quintile in our analyses, this in an area-level, rather than individual-level, measure of deprivation and is subject to the 15 ecologic fallacy. Additionally, the larger proportional drop in STI tests among 16 17 teenagers may be explained by residual confounding as our analyses could not take risk behaviours such as multiple condomless sex partners into account, and it is 18 unclear how this varied between 15-19 and 20-24 year olds between 2019 and 2020. 19 Reduced testing, missed infections and late diagnoses may have potential 20 consequences such as the increase in PID and infertility^[31]. This will impact the 21 22 guality of life of young people with STIs and increase costs to the healthcare system with the need for treatments for STI-related complications or sequelae. Additionally, 23 the difference in the means of testing between those in the least and most deprived 24 25 areas suggests barriers to access to online services, which should not occur, as STI

- 1 testing is free at the point of delivery in England. Given the increasing shift to online
- 2 service provision, there remains a need to assess how equitably they are provided
- 3 and to reduce the risk of differential access widening existing inequalities in sexual
- 4 health.

1 <u>References</u>

2	[1]	Mapp F, Hickson F, Mercer CH, Wellings K. How social representations of
3		sexually transmitted infections influence experiences of genito-urinary
4		symptoms and care-seeking in Britain: mixed methods study protocol. BMC
5		public health, 2016. 16 (1): p. 1-9. DOI: 10.1186/s12889-016-3261-0.
6	[2]	Clifton S, Mercer CH, Sonnenberg P, et al, STI risk perception in the British
7		population and how it relates to sexual behaviour and STI healthcare use:
8		findings from a cross-sectional survey (Natsal-3). EClinicalMedicine, 2018. 2:
9		p. 29-36. DOI: 10.1016/j.eclinm.2018.08.001.
10	[3]	Mitchell H, Allen H, Sonubi T, et al. Sexually transmitted infections and
11		screening for chlamydia in England, 2019. Public Health England (PHE),
12		London. 2020.
13	[4]	Mercer CH, Tanton C, Prah P, et al. Changes in sexual attitudes and lifestyles
14		in Britain through the life course and over time: findings from the National
15		Surveys of Sexual Attitudes and Lifestyles (Natsal). The Lancet, 2013.
16		382 (9907): p. 1781-1794. DOI: 10.1016/S0140-6736(13)62035-8.
17	[5]	Sonnenberg P, Menezes D, Freeman L, et al. Intimate physical contact
18		between people from different households during the COVID-19 pandemic: a
19		mixed-methods study from a large, quasi-representative survey (Natsal-
20		COVID). BMJ open, 2022. 12(2): p. e055284. DOI: 10.1136/bmjopen-2021-
21		055284.
22	[6]	Tanton C, Geary RS, Clifton S, et al. Sexual health clinic attendance and non-
23		attendance in Britain: findings from the third National Survey of Sexual
24		Attitudes and Lifestyles (Natsal-3). Sexually Transmitted Infections, 2018.
25		94(4): p. 268-276. DOI: 10.1136/sextrans-2017-053193.

1	[7]	Sonnenberg P, Clifton S, Beddows S, et al. Prevalence, risk factors, and
2		uptake of interventions for sexually transmitted infections in Britain: findings
3		from the National Surveys of Sexual Attitudes and Lifestyles (Natsal). The
4		Lancet. 2013. 382 . DOI: 10.1016/S0140-6736(13)61947-9.
5	[8]	Tiller, CM. Chlamydia during pregnancy: implications and impact on perinatal
6		and neonatal outcomes. Journal of Obstetric, Gynecologic, & Neonatal
7		Nursing, 2002. 31 (1): p. 93-98. DOI: 10.1111/j.1552-6909.2002.tb00027.x.
8	[9]	Centers for Disease Control and Prevention (CDC). and Prevention, Sexually
9		transmitted disease surveillance 2012. Centers for Disease Control and
10		Prevention, Atlanta, GA. 2014.
11	[10]	Detels R, Green AM, Klausner JD, et al. The incidence and correlates of
12		symptomatic and asymptomatic Chlamydia trachomatis and Neisseria
13		gonorrhoeae infections in selected populations in five countries. Sexually
14		transmitted diseases, 2011. 38(6): p. 503.
15	[11]	João AL, Lencastre A, Calvão J, et al. COVID-19, fear and sexual behaviour:
16		a survey in a tertiary STI clinic in Lisbon. Sexually Transmitted Infections,
17		2021. 97(7): p. 549-549. DOI: 10.1136/sextrans-2020-054834.
18	[12]	Bardsley M, Wayal S, Blomquist P, et al. Improving our understanding of the
19		disproportionate incidence of STIs in heterosexual-identifying people of black
20		Caribbean heritage: findings from a longitudinal study of sexual health clinic
21		attendees in England. Sexually Transmitted Infections, 2022. 98(1): p. 23-31.
22		DOI: 10.1136/sextrans-2020-054784.
23	[13]	Savage EJ, Mohammed H, Leong G, et al. Improving surveillance of sexually
24		transmitted infections using mandatory electronic clinical reporting: the
25		genitourinary medicine clinic activity dataset, England, 2009 to 2013. Euro

1 Surveillance, 2014. **19**(48): p. 20981. DOI: 10.2807/1560-

2 7917.es2014.19.48.20981.

- [14] Public Health England (PHE). *Chlamydia Testing Activity Dataset: Dataset* specification and technical guidance. Public Health England, London. 2016.
- 5 [15] Chandra NL, Soldan K, Dangerfield C, et al. Filling in the gaps: estimating
- 6 numbers of chlamydia tests and diagnoses by age group and sex before and
- 7 during the implementation of the English National Screening Programme,
- 8 2000 to 2012. Euro Surveillance, 2017. 22(5): p. 23-33. DOI: 10.2807/1560-
- 9 7917.ES.2017.22.5.30453.
- 10 [16] UK Health Security Agency (UKHSA). GUMCAD STI Surveillance System.
- 11 Data specification and technical guidance. UK Health Security Agency
- 12 (UKHSA), London 2021.
- 13 [17] Office for National Statistics (ONS). Super Output Area (SOA). Available at:
- 14 https://webarchive.nationalarchives.gov.uk/ukgwa/20160106001702/http://ww
- 15 w.ons.gov.uk/ons/guide-method/geography/beginner-s-guide/census/super-
- 16 <u>output-areas--soas-/index.html</u> [accessed 13.07.22].
- Inistry of Housing, Communities and Local Government (MHCLG). *The English Indices of Deprivation 2019 (IoD2019).* 2019.
- 19 [19] Office for National Statistics (ONS). 2011 Census: Characteristics of Built-Up
- 20 Areas. 2013. Available at:
- 21 <u>https://www.ons.gov.uk/peoplepopulationandcommunity/housing/articles/char</u>
- 22 <u>acteristicsofbuiltupareas/2013-06-28</u> [accessed 12.05.22].
- 23 [20] Office for National Statistics (ONS). 2011 Built-up Areas Methodology and
- 24 *Guidance*. Office for National Statistics: London. 2013.

- 1 [21] Office for National Statistics (ONS). 2011 rural/urban classification. 2016.
- 2 Available at:
- 3 https://www.ons.gov.uk/methodology/geography/geographicalproducts/ruralur
- 4 <u>banclassifications/2011ruralurbanclassification</u> [accessed: 12.05.22].
- 5 [22] Office for National Statistics (ONS). *Ethnic group, national identity and*
- 6 *religion*. Available at:
- 7 <u>https://www.ons.gov.uk/methodology/classificationsandstandards/measuringe</u>
- 8 <u>quality/ethnicgroupnationalidentityandreligion</u> [accessed 07.07.22].
- 9 [23] StataCorp, LP., Stata data analysis and statistical Software. Special Edition
- 10 Release, 2007. **10**: p. 733.
- 11 [24] Sexual Health London (SHL). *Home STI testing, regular and emergency*
- 12 contraception. Free NHS sexual health services online. Available at:
- 13 https://www.shl.uk/ [accessed 20.07.22].
- 14 [25] Pinto CN, Niles JK, Kaufman HW, et al. Impact of the COVID-19 Pandemic on
- 15 Chlamydia and Gonorrhea Screening in the US. American journal of
- 16 preventive medicine, 2021. **61**(3): p. 386-393. DOI:
- 17 10.1016/j.amepre.2021.03.009.
- 18 [26] Latini A, Magri F, Dona MG, et al. Is COVID-19 affecting the epidemiology of
- 19 STIs? The experience of syphilis in Rome. Sexually Transmitted Infections,
- 20 2021. **97**(1): p. 78. DOI: 10.1136/sextrans-2020-054543.
- 21 [27] Simões D, Stengaard AR, Combs L, Rabenet D. Impact of the COVID-19
- 22 pandemic on testing services for HIV, viral hepatitis and sexually transmitted
- 23 infections in the WHO European Region, March to August 2020.
- 24 Eurosurveillance, 2020. 25(47): p. 2001943. DOI: 10.2807/1560-
- 25 7917.ES.2020.25.47.2001943.

1	[28]	Eaton S, Biggerstaff D, Petrou S, et al. Young people's preferences for the
2		use of emerging technologies for asymptomatic regular chlamydia testing and
3		management: a discrete choice experiment in England. BMJ Open, 2019.
4		9 (1). DOI: 10.1136/bmjopen-2018-023663.
5	[29]	Spence T, Kander I, Walsh J, et al. Perceptions and Experiences of Internet-
6		Based Testing for Sexually Transmitted Infections: Systematic Review and
7		Synthesis of Qualitative Research. Journal of Medical Internet Research,
8		2020. 22 (8). DOI: 10.2196/17667.
9	[30]	Barnard S, Free C, Bakolis I, et al. Comparing the characteristics of users of
10		an online service for STI self-sampling with clinic service users: a cross-
11		sectional analysis. Sexually Transmitted Infections, 2018. 94(5): p. 377-383.
12		DOI: 10.1136/sextrans-2017-053302.
13	[31]	O'Connell CM, Ferone ME., Chlamydia trachomatis Genital Infections.
14		Microbial Cell, 2016. 3 (9): p. 390-403. DOI: 10.15698/mic2016.09.525.
15		

Table 1. Number of gonorrhoea test and diagnoses among 15–24 year olds residing in England, by demographic characteristics: 2019 to 2020

	Tests			Diagnoses		
	2010	2020	Percentage	2010	2020	Percentage
	2013	2020	difference	2013	2020	difference
Age						
15 to 19	238,554	157,303	-34%	8,278	5,815	-30%
20 to 24	559,757	433,457	-23%	17,397	13,447	-23%
Ethnicity						
Asian	31,993	20,516	-36%	985	641	-35%
Black African	35,797	26,257	-27%	1,396	1,259	-10%
Black Caribbean	29,370	21,071	-28%	1,892	1,345	-29%
Other Black ethnicity	9,130	5,755	-37%	513	353	-31%
Mixed ethnicity	47,461	35,606	-25%	2,175	1,729	-21%
White	554,275	382,189	-31%	16,393	11,327	-31%
Other	9,572	6,292	-34%	409	303	-26%
Unknown ethnicity	80,713	93,074	15%	1,912	2,305	21%
Sexual orientation						
Heterosexual Males	196,172	131,192	-33%	6,557	4,554	-31%
MSM	47,550	39,356	-17%	6,525	4,782	-27%
Heterosexual Females	465,906	363,502	-22%	11,037	8,419	-24%
WSW	4,561	5,244	15%	61	78	28%
Unknown sexual orientation	84,122	51,466	-39%	1,495	1,429	-4%
Area of residence						
Rural	79,697	60,021	-25%	1,590	1,100	-31%
Urban	704,883	518,388	-26%	23,643	17,773	-25%
Unknown area of residence	13,731	12,351	-10%	442	389	-12%
Residential area-level deprivation						
(Deprivation Quintile)*						
1 (most deprived)	179,282	130,529	-27%	7,756	5,944	-23%
2	193,177	144,229	-25%	7,027	5,463	-22%
3	157,740	118,779	-25%	4,651	3,437	-26%
4	134,641	98,672	-27%	3,296	2,323	-30%
5 (least deprived)	119,740	86,200	-28%	2,503	1,706	-32%
Unknown deprivation quintile	13,731	12,351	-10%	442	389	-12%
Region of residence						
East Midlands	57,710	43.033	-25%	2.102	1.570	-25%
East of England	74,590	58,536	-22%	1.851	1.535	-17%
London	204,435	151,651	-26%	8,554	6,676	-22%
North–East	30,391	19,930	-34%	965	713	-26%
North–West	88,276	54,658	-38%	3,094	1,915	-38%
South–East	122,652	86,955	-29%	2,509	1,620	-35%
South–West	69,963	56,043	-20%	1,314	862	-34%
West Midlands	80,152	59,789	-25%	2,788	2,341	-16%
Yorkshire and Humber	59,577	48,957	-18%	2,144	1,676	-22%
Unknown region of residence	10,565	11,208	6%	354	354	0%
HIV status						
HIV diagnosed	1,574	969	-38%	316	198	-37%
HIV negative or unknown	796,737	589,791	-26%	25,359	19,064	-25%
Total	700 244	500 760	260/	05 G75	10.000	250/
1	190,311	590,760	-20%	20,070	19,202	-23%
1						

*Deprivation quintile is an area-level measure of deprivation and socioeconomic status based on the 2019 Index of Multiple Deprivation (IMD) quintiles 2

Table 2. Number of syphilis test and diagnoses among 15–24 year olds residing in England, by demographic characteristics: 2019 to 2020

	Tests			Diagnoses		
	2019	2020	Percentage difference	2019	2020	Percentage difference
Age						
15 to 19	136,686	78,620	-42%	223	139	-38%
20 to 24	382,761	255,340	-33%	948	766	-19%
Ethnicity						
Asian	24,157	14,354	-41%	61	40	-34%
Black African	25,799	17,443	-32%	19	24	26%
Black Caribbean	19,560	12,608	-36%	29	36	24%
Other Black ethnicity	6,165	3,547	-42%	17	9	-47%
Mixed ethnicity	31,476	21,445	-32%	72	50	-31%
White	351,999	211,583	-40%	861	626	-27%
Other	6,880	4,455	-35%	21	24	14%
Unknown ethnicity	53,411	48,525	-9%	91	96	5%
Sexual orientation						
Heterosexual Males	146,927	78,987	-46%	192	121	-37%
MSM	44,921	35,974	-20%	693	568	-18%
Heterosexual Females	283,049	185,791	-34%	220	154	-30%
WSW	2,866	2,767	-3%	3	3	0%
Unknown sexual orientation	41,684	30,441	-27%	63	59	-6%
Area of residence						
Rural	50,909	30,084	-41%	74	55	-26%
Urban	459,186	294,991	-36%	1,070	837	-22%
Unknown area of residence	9,352	8,885	-5%	27	13	-52%
Residential area-level deprivation						
(Deprivation Quintile)*	114,267	70,837	-38%	357	275	-23%
1 (most deprived)	128,123	84,643	-34%	327	245	-25%
2	102,827	66,940	-35%	219	168	-23%
3	87,324	54,707	-37%	149	111	-26%
4	77,554	47,948	-38%	92	93	1%
5 (least deprived)						
Unknown deprivation quintile	9,352	8,885	-5%	27	13	-525
Region of residence						
East Midlands	38,033	19,792	-48%	72	43	-40%
East of England	48,082	28,046	-42%	59	59	0%
London	142,416	103,316	-27%	382	315	-18%
North–East	20,684	11,430	-45%	91	78	-14%
North–West	55,936	30,863	-45%	205	127	-38%
South–East	82,258	56,890	-31%	137	99	-28%
South–West	45,034	28,029	-38%	61	63	3%
West Midlands	42,700	23,298	-45%	76	60	-21%
Yorkshire and Humber	37,252	24,131	-35%	72	48	-33%
Unknown region of residence	7,052	8,165	16%	16	13	-19%
HIV status		_			_	_
HIV diagnosed	949	618	-35%	58	43	-26%
HIV negative or unknown	518,498	333,342	-36%	1,113	862	-23%
Total	540 447	222.000	260/	1 1 71	005	000/
IUlai	519,447	<u> </u>	-30%	1,171	905	-23%

1

*Deprivation quintile is an area-level measure of deprivation and socioeconomic status based on the 2019 Index of Multiple Deprivation (IMD) quintiles 2

Table 3. Number of chlamydia test and diagnoses among 15–24 year olds residing in England, by demographic characteristics: 2019-2020

	Tests			Diagnoses		
	2019	2020	Percentage difference	2019	2020	Percentage difference
Age						
15 to 19	418,019	258,419	-38%	48,138	31,140	-35%
20 to 24	914,914	670,879	-27%	80,288	57,165	-29%
Ethnicity						
Asian	38,185	25,549	-33%	3,077	2,046	-34%
Black African	41,229	31,450	-24%	5,587	3,977	-29%
Black Caribbean	35,566	26,380	-26%	5,449	3,653	-33%
Other Black ethnicity	9,914	6,710	-32%	1,408	887	-37%
Mixed ethnicity	55,686	44,242	-21%	6,479	5,031	-22%
White	740,482	533,857	-28%	72,369	51,307	-29%
Other	11,440	7,974	-30%	1,145	794	-31%
Unknown ethnicity	400,431	253,136	-37%	32,912	20,610	-37%
Gender						
Female	940,083	669,050	-29%	82,920	57,636	-30%
Male	380,647	252,121	-34%	44,173	29,476	-33%
Unknown gender	12,203	8,127	-33%	1,333	1,193	-11%
Online vs. Physical services						
Online services	271,684	361,622	33%	22,838	31,726	39%
Physical services	1,041,553	554,299	-47%	104,343	55,607	-47%
Unknown testing service	19,696	13,377	-32%	1,245	972	-22%
Area of residence						
Rural	147,884	106,001	-28%	12,899	9,017	-30%
Urban	1,105,688	774,434	-30%	107,641	74,482	-31%
Unknown area of residence	79,361	48,863	-38%	7,886	4,806	-39%
Residential area-level deprivation						
(Deprivation Quintile)*				33,041		
1 (most deprived)	290,480	202,207	-30%	29,881	22,991	-30%
2	299,190	209,304	-30%	23,280	20,884	-30%
3	255,757	180,401	-29%	18,732	16,095	-31%
4	217,828	154,037	-29%	15,606	12,967	-31%
5 (least deprived)	190,317	134,486	-29%		10,562	-32%
Unknown deprivation quintile	79,361	48,863	-38%	7,886	4,806	-39%
Region of residence						
East Midlands	104,710	77,763	-26%	11,149	7,829	-30%
East of England	137,273	102,422	-25%	11,886	9,302	-22%
London	298,401	199,182	-33%	28,481	18,347	-36%
North–East	61,670	43,572	-29%	5,921	4,818	-19%
North–West	162,097	102,017	-37%	16,571	10,129	-39%
South–East	182,138	125,763	-31%	17,257	11,864	-31%
South–West	134,229	98,483	-27%	11,587	7,833	32%
West Midlands	111,664	77,602	-31%	11,812	8,373	-29%
Yorkshire and Humber	140,751	102,494	-27%	13,762	9,810	-29%
Total	1,332,933	929,298	-30%	128,426	88,305	-31%
1						

2

*Deprivation quintile is an area-level measure of deprivation and socioeconomic status based on the 2019 Index of Multiple Deprivation (IMD) quintiles

		Crude Odds Ratio (95% CI)	Model 1, Adjusted Odds Ratio [∵] (95% CI)	Model 2, Adjusted Odds Ratio [±] (95% CI)	Model 3, Adjusted Odds Ratio [¥] (95% CI)	Model 4, Adjusteo Odds Ratio [‡] (95% CI) 3
	Residential area-level deprivation (Deprivation Quintile)*					4
	1 (most deprived)	1	_	_	_	- 6
Male	2	1.30 (1.28 – 1.33)	1.32 (1.29 – 1.34)	1.30 (1.28 – 1.32)	1.31 (1.29 – 1.34)	1.18 (1.16 – 1.20)
	3	1.40 (1.38 – 1.43)	1.42 (1.40 – 1.45)	1.41 (1.39 – 1.43)	1.46 (1.43 – 1.48)	1.35 (1.32 – 1.37)
	4	1.36 (1.34 – 1.39)	1. 39 (1.36 – 1.41)	1.39 (1.36 – 1.41)	1.45 (1.42 – 1.48)	1.37 (1.35 – 1.40)
	5 (least deprived)	1.24 (1.22 – 1.26)	1.25 (1.23 – 1.28)	1.26 (1.23 – 1.28)	1.31 (1.29 – 1.34)	1.29 (1.27 – 1.32) 10
	Residential area-level deprivation (Deprivation Quintile)*					11 12
	1 (most deprived)	1	_	_	-	- 13
Female	2	1.34 (1.32 – 1.35)	1.35 (1.34 – 1.37)	1.34 (1.32 – 1.35)	1.35 (1.34 – 1.37)	1.20 (1.19 – 1.21) 14
	3	1.40 (1.39 – 1.42)	1.42 (1.40 – 1.43)	1.40 (1.39 – 1.42)	1.46 (1.44 – 1.47)	1.33 (1.32 – 1.35)
	4	1.39 (1.38 – 1.41)	1.40 (1.39 – 1.42)	1.40 (1.38 – 1.41)	1.46 (1.45 – 1.48)	1.38 (1.36 – 1.39)
	5 (least deprived)	1.28 (1.27 – 1.30)	1.29 (1.27 – 1.30)	1.29 (1.28 – 1.31)	1.35 (1.33 – 1.37)	1.32 (1.30 – 1.34)

Table 4. Unadjusted and adjusted logistic regression analysis of the association between deprivation quintile^{*} and chlamydia testing via an online service among 15–24 year olds in England: 2019–2020, stratified by gender

17

18 *Deprivation quintile is an area-level measure of deprivation and socioeconomic status based on the 2019 Index of Multiple Deprivation (IMD) quintiles

19 ** Model 1 adjusted for year of test

20 ± Model 2 adjusted for year of test and age group

21 ¥ Model 3 adjusted for year of test, age group and area of residence

22 ^{‡ Model 4} adjusted for year of test, age group, area of residence and region

1 Appendix A

2 Table A1) Demographic characteristics of 15–24 year old males tested for chlamydia in England:

3 2019-2020

	Number and proportion tested in online services	Number and proportion tested in physical services
Age group (years)		
15 10	37,627	124,890
15-19	(23.1%)	(76.8%)
00.04	152,859	317,392
20–24	(32.5%)	(67.5%)
Area of residence		
Rural	20,521	52,145
	(28.2%)	(71.8%)
Urban	165,813	356,975
	(89.0%)	(68.3%)
Unknown area of residence	4,152	33,162
	(11.1%)	(88.9%)
Region of residence	(,)	()
	58.322	99.114
London	(37.0%)	(63.0%)
	18 518	29 578
East Midlands	(38.5%)	(61 5%)
	17 513	52 085
East of England	(25.2%)	(74,8%)
	(23.276)	(74.076)
North–East	5,240	22,510
	(10.9%)	(81.1%)
North–West	10,012	58,466
	(14.6%)	(85.4%)
South–East	21,454	61,869
	(25.7%)	(74.2)
South–West	22,052	41,587
	(34.6%)	(65.3%)
West Midlands	14,520	37,618
	(27.8%)	(72.1%)
Yorkshire and Humber	22,855	39,455
	(36.7%)	(63.3%)
Year of test		
2019	81,880	298,767
	(21.5%)	(78.5%)
2020	108,606	143,515
	(43.1%)	(56.9%)
Residential area-level deprivation (Deprivation Quintile)*		
1 (most doprived)	34,679	95,436
r (most deprived)	(26.6%)	(73.3%)
0	45,911	96,871
2	(32.1%)	(67.8)
0	41,352	81,030
3	(33.8%)	(66.2%)
	35.331	71,259
4	(33.0%)	(66.8%)
	29.061	64.524
5 (least deprived)	(31.3%)	(68.7%)
	4 152	33 162
Unknown deprivation quintile	(11 1%)	(88.9%)
Total	100 /26	AAD 280

4

*Deprivation quintile is an area-level measure of deprivation and socioeconomic status based on the 2019 Index of Multiple Deprivation (IMD) quintiles

1 Table A2) Demographic characteristics of 15–24 year old females tested for chlamydia in England:

2 2019-2020

	Number and proportion tested	Number and proportion tested
	in online services	in physical services
Age group (years)		
15–19	102,232	404,106
	(20.2%)	(79.8%)
20–24	337,692	765,103
	(30.6%)	(69.4%)
Area of residence		
Rural	44,744	135,043
	(24.9%)	(75.1%)
Urban	383,826	957,935
	(28.6%)	(71.4%)
Unknown area of residence	11,354	76,231 (87,0%)
Device of regidence	(13.0%)	(87.0%)
Region of residence	400 707	242.252
London	(26.9%)	212,252
	(30.0%)	(03.2%)
East Midlands	43,623	09,923
	(32.0%)	(07.2%)
East of England	30,030	130,417
	(22.9%)	(77.0%)
North–East	(15.7%)	
	(13.7%)	(64.3%)
North–West	(15.4%)	(84.6%)
	52 636	166 789
South–East	(24.0%)	(76.0%)
	50 939	117 440
South–West	(30,2%)	(69.7%)
	33 625	101 346
West Midlands	(24.9%)	(75.1%)
	54,880	125.267
Yorkshire and Humber	(30.5%)	(30.5%)
Year of test	()	()
2019	188,169	751,914
	(20.0%)	(80.0%)
2020	251,755	417,295
	(37.6%)	(62.4%)
Residential area-level deprivation	х <i>У</i>	, , , , , , , , , , , , , , , , , , ,
(Deprivation Quintile)*		
1 (most deprived)	84,665	273,626
r (most deprived)	(23.6%)	(76.4)
3	105,764	255,330
2	(29.3%)	(70.7)
2	94,142	216,725
3	(30.3%)	(69.7%)
4	79,044	183,472
4	(30.1%)	(69.9%)
5 (least deprived)	64,955	163,825
o (least deprived)	(28.4%)	(71.6%)
Linknown deprivation quintilo	11,354	76,231
	(13.0%)	(87.0%)
Total	439,924	1,169,209

*Deprivation quintile is an area-level measure of deprivation and socioeconomic status based on the 2019 Index of Multiple Deprivation (IMD) quintiles

1 Appendix B

2 **Table B1)** Adjusted logistic regression analysis of the association between deprivation quintile^{*} and

3 chlamydia testing via an online service among 15–24 year old males in England: 2019–2020

	Adjusted Odds Ratio (95% CI)
Age group (years)	
15–19	1
20–24	1.47 (1.45–1.49)
Area	
Rural	1
Urban	1.17 (1.15–1.20)
Region of residence	
London	1
East Midlands	1.03 (1.01–1.05)
East of England	0.49 (0.48–0.50)
North-East	0.37 (0.35–0.38)
North–West	0.26 (0.25–0.26)
South–East	0.56 (0.55–0.58)
South–West	0.84 (0.82–0.85)
West Midlands	0.62 (0.60–0.63)
Yorkshire and Humber	0.93 (0.91–0.95)
Year of test	
2019	1
2020	2.81 (2.77–2.84)
Residential area-level deprivation (Deprivation Quintile)*	
1 (most deprived)	1
2	1.18 (1.16 –1.20)
3	1.35 (1.32–1.37)
4	1.37 (1.35–1.40)
5 (least deprived)	1.29 (1.27–1.32)

⁴

5 *Deprivation quintile is an area-level measure of deprivation and socioeconomic status based on the 2019 Index of Multiple Deprivation (IMD) quintiles

6 ** All p-values <0.05

1 Table B2) Adjusted logistic regression analysis of the association between deprivation quintile^{*} and

chlamydia testing via an online service among 15-24 year old females in England: 2019-2020 2

	Adjusted Odds Ratio (95% CI)
Age group (years)	
15–19	1
20–24	1.63 (1.61–1.64)
Area	
Rural	1
Urban	1.16 (1.15–1.17)
Region of residence	
London	1
East Midlands	0.80 (0.79–0.81)
East of England	0.44 (0.43–0.44)
North–East	0.31 (0.30–0.31)
North–West	0.28 (0.27–0.28)
South–East	0.51 (0.50–0.51)
South-West	0.68 (0.67–0.69)
West Midlands	0.53 (0.52–0.54)
Yorkshire and Humber	0.71 (0.70–0.72)
Year of test	
2019	1
2020	2.45 (2.44–2.47)
Residential area-level deprivation (Deprivation Quintile)	
1 (most deprived)	1
2	1.20 (1.19 –1.21)
3	1.33 (1.32–1.35)
4	1.38 (1.36–1.39)
5 (least deprived)	1.32 (1.30–1.34)

3

*Deprivation quintile is an area-level measure of deprivation and socioeconomic status based on the 2019 Index of Multiple Deprivation (IMD) quintiles 4

** All p-values <0.05 5

1 Appendix C

2 **Table C1**) Number and proportion of chlamydia tests by testing service and deprivation quintile^{*} amongst 15–24 year old males in England: 2019–2020

			De	privation Quintile	*	
	Testing service	1 (most deprived)	2	3	4	5 (least deprived)
	Specialist sexual health service	69,860	71,130	57,781	49,387	43,850
		(53.7%)	(49.8%)	(47.2%)	(46.3%)	(46.9%)
	Non-specialist sexual health service	5,721	3,591	2,276	1,612	1,462
		(4.4%)	(2.5%)	(1.9%)	(1.5%)	(1.6%)
	GP	9,207	10,366	9,123	7,615	6,358
Physical		(7.1%)	(7.3%)	(7.5%)	(7.1%)	(6.8%)
services	Pharmacy	647	978	1,046	915	836
		(0.5%)	(0.7%)	(0.9%)	(0.9%)	(0.9%)
	Termination of Pregnancy centres	5	6	5	6	3
		(0.0%)	(0.0%)	(0.0%)	(0.0%)	(0.0%)
	Other	8,687	9,847	9,968	10,364	11,234
		(6.7%)	(6.9%)	(8.1%)	(9.7%)	(12.0%)
Online	Online	34,679	45,911	41,352	35,331	29,061
services		(26.7%)	(32.2%)	(33.8%)	(33.2%)	(31.1%)
	Total**	130,115	142,782	122,382	106,590	93,585

³

4 *Deprivation quintile is an area-level measure of deprivation and socioeconomic status based on the 2019 Index of Multiple Deprivation (IMD) quintiles

5 ** Totals include tests from unknown testing services

	Deprivation Quintile*					
	Testing service	1 (most deprived)	2	3	4	5 (least deprived)
Physical services Online services	Specialist sexual health service	138,082	133,456	107,714	89,520	78,302
		(38.5%)	(37.0%)	(34.6%)	(34.1%)	(34.2%)
	Non-specialist sexual health service	16,847	11,184	6,816	4,893	3,792
		(4.7%)	(3.1%)	(2.2%)	(1.9%)	(1.7%)
	GP	67,115	64,873	60,091	53,046	49,186
		(18.7%)	(18.0%)	(19.3%)	(20.2%)	(21.5%)
	Pharmacy	1,834	2,653	3,063	2,598	2,561
		(0.5%)	(0.7%)	(1.0%)	(1.0%)	(1.1%)
	Termination of Pregnancy centres	8,325	7,248	5,142	3,708	2,971
		(2.3%)	(2.0%)	(1.7%)	(1.4%)	(1.3%)
	Other	35,464	31,428	29,881	24,528	23,889
		(9.9%)	(8.7%)	(9.6%)	(9.3%)	(10.4%)
	Online	84,665	105,764	94,142	79,044	64,955
		(23.6%)	(29.3%)	(30.3%)	(30.1%)	(28.4%)
	Total ^{**}	358,291	361,094	310,867	262,516	228,780

1 **Table C2)** Number and proportion of chlamydia tests by testing service and deprivation quintile^{*} amongst 15–24 year old females in England: 2019–2020

3 *Deprivation quintile is an area-level measure of deprivation and socioeconomic status based on the 2019 Index of Multiple Deprivation (IMD) quintiles

4 ** Totals include tests from unknown testing services

²