

## **Proteomic predictors of physical, cognitive and imaging outcomes in multiple sclerosis: 5-year follow-up study**

Kian Jalaeddini, PhD<sup>1\*</sup>, Dejan Jakimovski MD, PhD<sup>2\*</sup>, Anisha Keshavan, PhD<sup>1</sup>, Shannon McCurdy, PhD<sup>1</sup>, Kelly Leyden, MRes<sup>1</sup>, Ferhan Qureshi<sup>1</sup>, Atiyeh Ghoreyshi, PhD<sup>1</sup>, Niels Bergsland, PhD<sup>2</sup>, Michael G. Dwyer, PhD<sup>2</sup>, Murali Ramanathan, PhD<sup>3</sup>, Bianca Weinstock-Guttman, MD<sup>4</sup>, Ralph HB Benedict, PhD<sup>2</sup>, Robert Zivadinov, MD, PhD<sup>2,5</sup>

\* - Equally contributing

<sup>1</sup>Octave Biosciences, Menlo Park, CA, USA; <sup>2</sup>Buffalo Neuroimaging Analysis Center, Department of Neurology, Jacobs School of Biomedical Sciences, University at Buffalo, State University of New York, Buffalo, NY, USA; <sup>3</sup>Department of Pharmaceutical Sciences, State University of New York, Buffalo, Buffalo, NY, <sup>4</sup>Jacobs MS Center, Department of Neurology, Jacobs School of Biomedical Sciences, University at Buffalo, State University of New York, Buffalo, NY, USA; <sup>5</sup>Center for Biomedical Imaging at the Clinical Translational Science Institute, University at Buffalo, State University of New York, Buffalo, NY, USA

### **Corresponding Author:**

Dejan Jakimovski M.D., Ph.D.  
Research Assistant Professor  
Buffalo Neuroimaging Analysis Center  
Department of Neurology  
Jacobs School of Medicine and Biomedical Sciences  
University at Buffalo, State University of New York  
100 High Street  
Buffalo, NY 14203  
Phone: 716-859-7040  
Fax: 716-859-7066

Email: [djakimovski@bnac.ne](mailto:djakimovski@bnac.ne)

**Keywords:** proteomics, multiple sclerosis, disability, cognitive performance, atrophy

**Running title:** Proteomic Analyses in a Longitudinal MS Study

**Abstract word count:** 250, **Title characters:** 97/110, **Word count:** 3,755 (until the end of discussion) and 4002 (until the beginning of references), **Number of Tables:** 4, **Number of Figures:** 4, **Supplement Tables:** 6, **Supplement Figures:** 4; **Number of references:** 44.

## **Abstract**

**Background:** A quantitative measurement of serum proteome biomarkers that would associate with disease progression endpoints can provide risk stratification for persons with multiple sclerosis and supplement the clinical decision-making process.

**Materials and Methods:** 202 persons with multiple sclerosis were enrolled in a longitudinal study with measurements at two time points with an average follow-up time of 5.4 years. Clinical measures included the Expanded Disability Status Scale, Timed 25-foot Walk, 9-Hole Peg and Symbol Digit Modalities Tests. Subjects underwent magnetic resonance imaging to determine the volumetric measures of the whole brain, gray matter, deep gray matter and lateral ventricles. Serum samples were analyzed using a custom immunoassay panel on the Olink™ platform and concentrations of 18 protein biomarkers were measured. Linear mixed-effects models and adjustment for multiple comparisons were performed.

**Results:** Subjects had a significant 55.6% increase in Chemokine Ligand 20 (9.7pg/mL vs. 15.1pg/mL,  $p<0.001$ ) and Neurofilament light polypeptide (10.5 pg/ml vs. 11.5 pg/ml,  $p=0.003$ ) at the follow-up time point. Additional changes in CUB domain-containing protein 1, Contactin 2, Glial fibrillary acidic protein, Myelin oligodendrocyte glycoprotein, and Osteopontin were noted but did not survive multiple comparisons correction. Worse clinical performance in the 9-HPT was associated with Neurofilament light polypeptide ( $p=0.001$ ). Increases in several biomarker candidates were correlated with greater neurodegenerative changes as measured by different brain volumes.

**Conclusion:** Multiple proteins representing diverse biological pathways (neuroinflammation, immune modulation, and neuroaxonal integrity) associate with physical, cognitive and

radiographic outcomes. Future studies should determine the utility of multiple protein assays in routine clinical care.

## Introduction

Multiple sclerosis (MS) is a chronic, inflammatory, demyelinating, and degenerative disease of the central nervous system (CNS) resulting in progressive accrual of physical and cognitive disability.<sup>1</sup> People with multiple sclerosis (pwMS) have heterogeneous clinical presentation and multiple demographics, clinical, and paraclinical risk factors have been associated with poorer long-term outcomes.<sup>1</sup> Over the last three decades, a plethora of highly effective disease modifying therapies (DMTs) have been developed that can significantly lessen the long-term disability.<sup>1</sup> They generally range from highly effective but immunosuppressive therapies to moderately effective immunomodulators with lower rate of adverse events.<sup>2</sup> Therefore, development of attainable and cost-effective biomarkers can provide risk stratification for pwMS and supplement the clinical decision-making process.

In addition to validated imaging biomarkers such as lesion pathology and whole brain atrophy, recent developments in proteomic assay technology have allowed detection and measurement of picomolar concentrations of blood biomarkers.<sup>3</sup> These biomarkers can provide proxy measures regarding the occurrence and extent of pathological changes in the CNS.<sup>4</sup> For example, higher blood levels of neurofilament light chain (NfL), an intermediate filament present in neurons, can indicate greater neuroaxonal destruction, and higher levels of glial fibrillary acidic protein (GFAP) can indicate glial cell activation both indicative of presence of disease activity.<sup>5,6</sup> Although these individual biomarkers have already emerged as candidate outcomes measures in MS, a single protein biomarker can have limited ability in capturing changes within multiple parallel pathophysiological MS pathways.<sup>7</sup> When compared to MRI measures, blood-derived analyses are less costly, more accessible, and can be bundled together with routine clinical blood work.<sup>8</sup>

The Multiple Sclerosis Disease Activity (MSDA) is a recently developed and analytically validated<sup>9</sup> panel of 18 biomarkers that represent changes within four main pathophysiological pathways of neuroinflammation, immunomodulation, myelin biology and neuroaxonal integrity.<sup>9</sup> In the first clinical validation study, the MSDA platform was trained and tested as a predictor for presence of gadolinium-enhancing lesions or new/newly enlarging T2 lesions in a cohort of 614 samples.<sup>10</sup> The multi-protein scores outperformed the best individual protein (NfL) with area under curve change from 0.726 to 0.781.<sup>10</sup> Determining the relation of the MSDA panel to long-term disability outcomes and examining the longitudinal predictive properties of such an assay are essential for clinical adoption and wide-spread clinical utility.

The aims of this study were to determine the relationship between the multiple proteomic biomarkers and cross-sectional and longitudinal MS outcomes, including physical disability, cognitive performance and conventional MRI outcomes. We hypothesize that proteomic analysis could improve the clinical and MRI-based risk stratification when utilized in a heterogenous group of pwMS.

## **Methods**

### *Study population:*

A total of 202 patients were assessed in these analyses and derived from a larger longitudinal, case-controlled study to explore the role of cardiovascular, environmental and genetic risk factors in multiple sclerosis patients (CEG-MS).<sup>11</sup> In particular, for this study, the pwMS were enrolled at baseline between 2009-2012 and returned for a follow-up visit in years 2014-2017. The inclusion criteria were: 1) baseline age of 18-75 years old; 2) diagnosed with either MS or clinically isolated syndrome (CIS), defined by the 2010-revised McDonald criteria<sup>12</sup>; 3) availability of either baseline or follow-up serum sample, MRI, clinical and neuropsychological

assessments within 30 days of each other. The exclusion criteria were: 1) having clinical relapse or receiving intravenous corticosteroid therapy within 30 days before the MRI and serum sampling, 2) not able to undergo any of the aforementioned study procedures, and 3) pregnant or nursing mothers.

The CEG-MS study and the retrospective proteomic analyses were approved by the University at Buffalo Institutional Review Board (IRB) and all subjects provided a signed consent form.

*Physical and cognitive disability measures:*

A board-certified neurologist evaluated patients for global disability using the Expanded Disability Status Scale (EDSS) score<sup>13</sup>, a board certified neuropsychologist oversaw a clinical assessment that included assessment of quantitative mobility and leg function, using the Timed 25 Foot Walk Test (T25FWT)<sup>14</sup>, quantitative finger dexterity using the 9-Hole Peg Test (9HPT)<sup>15</sup>, and cognitive efficiency and speed performance using the Symbol Digit Modalities Test (SDMT) and the Paced Auditory Serial Addition Test (PASAT)<sup>16</sup>. A structured questionnaire was also used to collect demographic and clinical information. According to the clinical presentation and disease history, pwMS were categorized as CIS, relapsing-remitting MS (RRMS), progressive MS (PMS) (further categorized to primary (PPMS), or secondary progressive (SPMS)).<sup>17</sup>

Presence of disability progression (DP) over the follow-up was defined using standard criteria of changes in EDSS scores: 1) An increase of 2 or more points if the baseline EDSS is zero; 2) an increase of 1.5 or more points if the baseline EDSS is 0.5, 3) an increase of  $\geq 1$  point if the baseline EDSS is between 1.0 and 5.0, and 4) an increase of equal or greater than 0.5 point if the baseline EDSS is  $\geq 5.5$ .<sup>18</sup> Worsening in T25FWT and 9HPT was defined as an increase of

greater than or equal to 20% from baseline to follow-up.<sup>14, 15</sup> Worsening in SDMT performance was defined using several previously used criteria: 1) decrease of 4 or more points from baseline to follow-up<sup>16</sup>; 2) decrease of 8 or more points from baseline to follow-up; 3) being classified as cognitively impaired based with  $|z\text{-scores}| > 1.5$  derived from a healthy control population published in the literature with mean and standard deviation of 55.49, 13.06 respectively<sup>5, 19</sup>.

#### *Proteomics analyses:*

All blood samples were processed within 24-hours from acquisition and stored at  $-80^{\circ}\text{C}$  until analyzed. No freeze and thaw cycles were performed in the interim period. All samples were sent to Octave Bioscience (Menlo Park, CA, USA) for proteomic analysis using the MSDA assay panel.<sup>9</sup> Proteomic analysis was performed blinded to the demographic, clinical and MRI data. The MSDA assay uses Proximity Extension Assay (PEA) methodology and is performed on the Olink<sup>TM</sup> platform. Twenty-one proteins that are associated with key biological pathways of MS pathophysiology were selected for inclusion on the panel based on results from discovery analyses investigating relative expression of 1196 proteins in previously characterized MS cohorts. The MSDA assay utilizes a stacked classifier logistic regression model of 18 age- and sex adjusted protein concentrations as previously described to determine four disease pathway scores (immunomodulation, neuroinflammation, myelin biology, and neuroaxonal integrity) as well as an overall disease activity score.<sup>9</sup> The complete list of proteins (with commonly used aliases and their abbreviations) are shown in Figure 1.

#### *MRI acquisition and analyses:*

At baseline and follow-up visits, pwMS underwent an MRI examination using the same 3T Signa Excite 12 Twin-Speed scanner (GE Healthcare, Milwaukee, WI, USA) and eight channel

head and neck coil. The standard sequences utilized in these analyses were two-dimensional (2D) fluid attenuated inversion recovery (FLAIR), 2D T1-weighted spin echo with and without use of 0.2mL/kg gadolinium (Gd) contrast acquired 5 min post-injection, and high-resolution 3D T1-weighted imaging. The sequence parameters are explained in details elsewhere.<sup>20</sup>

Lesion analysis was performed in a blinded manner with respect to the patient clinical and proteomics status. T2 lesion volume (LV), T1-LV and Gd-LV were obtained using a semi-automated contouring/thresholding technique using Java Image Manipulation (JIM) version 6.0 (Xinapse Systems Ltd, <http://www.xinapse.com/>, Essex, UK). The cross-sectional and longitudinal changes in volumes of brain regions of interest (ROIs) of whole brain (WB), white matter (WM), gray matter (GM), all normalized for head size, were measured using the SIENAX and SIENA algorithms (FMRIB Software Library, <http://www.fmrib.ox.ac.uk/fsl>).<sup>21</sup> A lesion inpainting technique was used to avoid tissue misclassification.<sup>22</sup> Total deep GM (DGM) volume and specific volume of the thalamus were obtained with FMRIB's Integrated Registration and Segmentation Tool (FIRST, <https://fsl.fmrib.ox.ac.uk/fsl>, version 2.6). The number of patients who had baseline and follow-up samples are described in Supplementary Table 1. Lastly, pathological change in whole brain volume was determined as an annualized percent brain volume reduction of greater than or equal to 0.4%<sup>23</sup> and pathological lateral ventricle volume change if an annualized percent volume expansion of greater than or equal to 3.5%.<sup>24</sup>

#### *Statistical analyses:*

Data and statistical analyses were performed using Python version 3.8.10, SciPy 1.9.3, pandas 1.5.2, pingouin 0.5.3, and NumPy 1.24.1. Ordinary and mixed-effects models were estimated using R version 4.1.3. Logistic regression models were estimated with Python statsmodels



package version 0.13.2.

Student's t-test and Analysis of Covariance (ANCOVA) were used for statistical analysis of parametric continuous variables and longitudinal analysis was performed using the paired non-parametric Wilcoxon test. Ordinary least-squares were used to estimate cross-sectional univariable models; linear mixed-effects regression models were used to estimate models on longitudinal data; linear logistic regression models were used to estimate dichotomous outcomes (e.g., pwMS disability progression yes/no).

MRI-based brain volumes, EDSS, and neuropsychological test outcomes were used as dependent variables, and age, sex, BMI, and all proteomic measures as independent predictors ( $outcome\ score = age + sex + body\ mass\ index\ (BMI) + biomarker\ concentration$ ) and for the linear mixed-effects model, subject ID was set as random effect ( $outcome\ score = age + sex + BMI + biomarker\ concentration + (1|patient\ ID)$ ). For entry into the regression models, the proteomic data, MRI-based brain volumes, EDSS and neuropsychological test scores were transformed using  $\log(10)$  and all the statistical tests were applied to the log-transformed data. Logistic regression models were similarly used if the dependent variable was of categorical nature. Outcomes such as  $R^2$  for ordinary and mixed-effects regression, McFadden's pseudo- $R^2$  for logistic regression, standardized  $\beta$  and p-values were reported. Adjusted p-values lower than 0.05 were considered statistically significant. The regression and correlation p-values underwent false discovery rate (FDR) correction using the Benjamini-Hochberg procedure.

Data were visualized using Python matplotlib 3.4.2, seaborn 0.12.1, and plotly 5.12.0 packages. The data distribution was determined using visual inspection of histograms and Q-Q plots. Volcano plots were used to visualize the significance (p-value) versus effect size.

## Results

### *Demographic and clinical characteristics:*

Table 1 describes the demographics and clinical characteristics of the pwMS. As expected, the pwPMS were significantly older, had longer disease duration and higher EDSS scores both at baseline and follow-up visits ( $p < 0.001$  for all). We failed to reject the null hypothesis that the rate of DP between pwCIS/RRMS and pwPMS were similar (28.6% vs. 37.5%,  $p = 0.251$ ). There were no significant differences in terms of baseline and follow-up DMT use. As expected, the pwPMS had significantly greater pathology measured by conventional MRI measures of T2-LV and T1-LV ( $p < 0.001$  and  $p = 0.015$ ) and global measures of WBV, WMV and GMV ( $p < 0.001$ ). The pwCIS/RRMS had on average significantly more Gd lesions when compared to the PMS group ( $p < 0.001$ ). Supplement Figure 1 depicts the distribution of disease phenotypes at baseline and transition over the follow-up.

Out of the pwMS with available longitudinal disability data, 55 out of 181 (30.4%) worsened in EDSS scores, 46 out of 186 (24.7%) had a pathological rate of whole brain atrophy and only 19 out of 186 (10.2%) has a pathological rate of ventricle enlargement. The paired statistical test revealed that all outcomes worsened significantly at the follow-up time-point except for the WMV and PASAT performance (Supplement Table 2). Supplement Figure 2 visualizes shifts in MRI-based volumes, EDSS and neuropsychological scores between the baseline and follow-up time-points. Supplement Figure 3, 4 demonstrate differences in MRI-based volumes, EDSS, neuropsychological scores and biomarker concentrations between pwCIS/RRMS and pwPMS subgroups.

### *Proteomic characteristics of the study population:*

In total, 202 pwMS had serum samples at the baseline visit and 143 pwMS had serum samples at both the baseline and follow-up visits. The baseline, follow-up and longitudinal change in each of the proteomic biomarkers (shown as median and interquartile range (IQR)) are shown in Figure 2 for biomarkers with significant shift between the time points and detailed analysis is shown in Supplementary Table 4. Over the follow-up, pwMS had a significant 56% increase in CCL20 (9.74pg/mL vs. 15.1pg/mL,  $p=0.001$ ) and 9.4% increase in NfL (10.5 pg/mL vs 11.5 pg/mL,  $p=0.003$ ). There were also significant shifts in CDCP1, CNTN2, GFAP, MOG, and OPN but they did not survive multiple comparisons correction. Supplement Figure 4 demonstrates significant shifts in the serum biomarkers between pwCIS/RRMS and pwPMS at baseline and follow-up time-points.

In the pwMS who progressed over the follow-up, there were several proteins whose levels differed between the visits. For the patients who progressed in the 9HPT test, CDCP1 increased by 31% (110 pg/ml vs. 144 pg/ml,  $p=0.003$ ), TNFRSF10A increased by 37% (5.61 pg/ml vs 7.71 pg/ml,  $p=0.005$ ) and VCAN increased by 17% (428 pg/ml vs 500 pg/ml,  $p=0.007$ ). The other proteins in Table 2 did not survive the multiple comparisons correction. Although age adjusted NfL was increased and PRTG was decreased in patients with progressive type at follow-up, these findings did not pass the multiple comparisons correction.

### *Relationship between proteomic data and outcomes in pwMS:*

#### *Longitudinal models:*

Figure 3 shows the significant biomarkers of the mixed-effects models estimated for both time-points and all pwMS. We found six biomarker candidates, namely GFAP, FLRT2, CDCP1,

TNFRSF10A, CXCL9 and CCL20, whose increases were correlated with reduction in brain volumes and one protein (APLP1) whose decrease was correlated with reduction in brain volume. In particular, GFAP ( $p=0.003$ ), FLRT2 ( $p=0.001$ ), CDCP1 ( $p=0.004$ ) and TNFRSF10A ( $p=0.003$ ) were associated with WBV decrease and survived the multiple comparisons correction. For the WMV, GFAP was the only protein that survived the statistical test ( $p<0.001$ ). For the DGM, significant associations with CCL20 ( $p=0.004$ ), CXCL9 ( $p=0.001$ ), CDCP1 ( $p=0.002$ ), FLRT2 ( $p=0.008$ ), TNFRSF10A ( $p=0.009$ ) were noted.

In terms of clinical (EDSS scores) and cognitive assessment, we found that worsening in the 9-HPT score was associated with an increase in NfL ( $p=0.001$ ) surviving the multiple comparisons correction. Table 3 details the estimated coefficients, p-values and the qualities of the fit (R-squared of the model with standardized features as biomarker, sex, age, BMI) corresponding to each biomarker and outcome measure. Note that only significant biomarkers ( $p<0.05$ ) are included in Table 3.

#### *Cross-sectional models:*

Figure 4 shows the significant biomarkers of linear single-protein models estimated individually for the baseline and follow-up outcomes. At the baseline visit, higher GFAP was correlated with lower WBV ( $p<0.001$ ), GMV ( $p<0.001$ ), thalamic volume ( $p<0.001$ ), and DGMV ( $p<0.001$ ), and higher LVV ( $p<0.001$ ). Moreover, higher GFAP was associated with higher EDSS scores ( $p=0.002$ ). Detailed analyses are shown in Table 4.

At the follow-up GFAP, was correlated with lower WBV ( $p=0.001$ ), GMV ( $p=0.001$ ), and DGMV ( $p=0.002$ ), and higher LVV ( $p<0.001$ ). Both NfL ( $p=0.001$ ) and GFAP ( $p=0.003$ ) were correlated with worse 9HPT scores. NfL was also correlated with T25FWT ( $p=0.002$ ) and EDSS scores ( $p=0.002$ ). Detailed analyses are shown in Supplement Table 5.

### *Shifts in biomarker levels as predictors of clinical change:*

It is of importance to assess whether shifts in the biomarker concentration is correlated with shifts in outcome scores. Therefore, we fitted a model with shifts in biomarker between baseline and follow-up, age, sex, BMI as independent predictors and shifts in outcome scores as dependent variables. CDCP1 ( $p=0.001$ ), FLRT2 ( $p=0.001$ ), PRTG ( $p=0.008$ ), TNFRSF10A ( $p<0.001$ ), TNFSF13B ( $p<0.001$ ) were predicted shift in the WBV. CD6 ( $p=0.001$ ) was predicted shift in the DGMV. Figure 5 and Supplement Table 6 describe the findings of these analyses.

## **Discussion**

The findings of this longitudinal proteomics study are multifold. Firstly, multiple proteomic biomarkers representing different pathophysiological MS pathways are differentially associated with phenotypical and macroscopic pathological changes. Secondly, worse physical and cognitive outcomes in pwMS were associated with blood-based measures of NfL. Lastly, worse neurodegenerative MRI outcomes were associated with a greater number of biomarkers including GFAP, CDCP1, CXCL9, CCL20, APLP1, FLRT2, and TNFRSF10A.

The NfL/GFAP relationship with clinical outcomes was recently demonstrated in a similar longitudinal Swiss study.<sup>25</sup> Over an average follow-up of 7 years, serum GFAP levels were prognostic of progression independent of relapse activity (PIRA) and complementary to the serum NfL data.<sup>25</sup> Moreover, NfL levels were prognostic of atrophy in the WMV, whereas GFAP specifically prognosticate GM atrophy.<sup>25</sup> The multi-protein panel employed in our study was also utilized in a study of 431 unique pwMS and successfully predicted the real-world disability status (patient-reported disability score and patient-reported outcomes).<sup>26</sup> The

proteomic profiles consistently outperformed individual top-ranking markers such as NfL and GFAP.<sup>26</sup> The fact that the same protein biomarkers implicated in their stacking classification algorithm (CDCP1, IL-12B and PRTG) were also seen in our objective disability findings further validates the utility and need of multi-protein proteomic assay.<sup>26</sup>

We further expand on the literature by demonstrating that the similar set of biomarkers (APLP1, CDCP1, FLRT2, TNFRSF10A and CCL20) are also relevant to MRI-based volumetric measures. Of note, the directionality of APLP1-DGMV relationship (decrease in the biomarker concentration was associated with decreased DGMV) was opposite when compared to the remaining ones. Currently, there are no comprehensive proteomic studies that investigate associations with MRI measures in pwMS. The literature most commonly describes individual associations with one or two proteomic measures (NfL and GFAP).<sup>20, 27, 28</sup> Despite the high collinearity between serum NfL and GFAP levels, a cross-sectional study of 129 pwMS showed that the amount of lesion pathology (T2-LV) and WM/GM volumes were associated only with GFAP levels and not with NfL.<sup>29</sup> We corroborate these findings with GFAP remaining a strong predictor all MRI measures acquired in our study (WBV, DGM, LVV and thalamic volume). Serum GFAP levels were also recently associated with greater microstructural pathology in 62 pwMS assessed by diffusion tensor imaging.<sup>30</sup> Our results showed that CCL20 has strong association with the DGM volume and it was increased at the follow-up time point. This protein was previously shown to be increased in PwMS and specifically with a progression index and was higher during remission than in relapse periods.<sup>31, 32</sup>

The multiplex assays could broaden our understanding of key mechanisms underlying progression by taking a biological-based approach to objectively quantify disease progression.<sup>33</sup> They have been demonstrated in other neurological disorders as well.<sup>34, 35</sup> For example,

proteomic data from only 4-Plex assay (NfL, GFAP, tau protein and ubiquitin c-terminal hydrolase L1; UCH-L1) better classified people with traumatic brain injury when compared to only single proteomic measure.<sup>34</sup> Similarly, cognitive performance in Alzheimer's Disease and amyloid PET status can be predicted and classified by a combination of GFAP, amyloid beta and neurofilament light chain.<sup>35</sup>

While cut-offs of normal versus pathological levels of NfL in pwMS have been previously published<sup>36,37</sup>, this information is not available for the majority of proteomic biomarkers utilized in this multi-protein assay. A limited number of studies report the reference intervals and preanalytical GFAP levels.<sup>38,39</sup> For example, a Danish-based analysis of 371 apparently healthy subjects reported fairly large ranges with GFAP levels of 25-136 ng/L (20-39 years old), 34-242 ng/L (40-64 years old) and 4-438 ng/L (for 65-90 years old).<sup>38</sup> Moreover, there was ~10% variability after three freeze-thaw cycles or storing serum samples at -20 °C for an average of 133 days.<sup>38</sup> Significant semidiurnal variations in GFAP have been reported (9 AM vs. 12 PM vs 9 PM blood draw).<sup>40</sup> Based on these references, none of the median pwMS values would be considered “pathological”. Other biomarkers such as contactin-1 may be more susceptible to pre-analytical factors and have even greater variability.<sup>41</sup> After the selection of best performing biomarkers and creation of multi-protein scores, future studies should aim at determining appropriate cut-offs for best differentiation between normal and pathological states.

The linear modeling approach taken in this study can be considered as a study limitation. Since there is no guarantee that the underlying interaction is inherently linear, an area of future work is to consider nonlinear regression techniques to capture more complex interactions of serum biomarker concentrations and metrics of disease progression. In addition, multi-protein regression approaches in cross-validation studies can be considered to integrate the predictive

power of individual biomarkers in one model.<sup>42</sup> The shifts in DP outcome measures were not uniform. Such imbalance in the training data can deteriorate generalizability of the model.<sup>43, 44</sup> In the future, it will be of interest to balance the data using up, down techniques or using weighted learners. The significantly lower number of available cognitive measures (only 25% of the pwMS) for the baseline time point presents as another study limitation. Moreover, the initial determination of biomarkers within the assay were based on ability to predict presence of contrast-enhancing and new/newly enlarging lesions.<sup>9</sup> In comparison to younger more active pwMS from the literature, our population was relatively older and had very limited neuroinflammatory activity. Future development of a more comprehensive assay that contains proteins specific to neurodegenerative changes (vs. neuroinflammation) could better predict the occurrence of long-term disability worsening. Moreover, the use and change in DMT should be incorporated in future statistical analyses. Additional limitation of our analysis is the lack of a third clinical visit that would allow confirmation of the disease progression. Lastly, the lack of healthy control data does not allow us to determine pathological protein cut-offs and risk stratify the pwMS based on healthy condition.

In conclusion, the clinical, cognitive, and MRI-based outcomes in pwMS are associated with more than one proteomic biomarker. While sNfL had the strongest associations with physical disability such as EDSS scores and hand dexterity, additional proteomic biomarkers related to neuroaxonal integrity were associated with cross-sectional and longitudinal MRI measures of brain atrophy. Multi-protein assays may be essential in capturing the complex MS pathophysiology as part of the disease stratification and monitoring. Before implementation into routine clinical practice, future studies should determine the treatment responsiveness of such proteomic biomarkers.



## **Acknowledgements:**

This research has been supported by Octave Bioscience Inc.

## **Author Contributions:**

KJ, AK, SM, FQ, AG contributed to the analysis of data and drafting a significant portion of the manuscript or figures

## **Potential Conflict of Interest:**

Kian Jalaledini, Ferhan Qureshi, Anisha Keshavan Shannon McCurdy, Kelly Leyden, and Ati Ghoreyshi are employees of and either hold stock or stock options at Octave Bioscience.

Dejan Jakimovski received honoraria for serving on the advisory board of AstraZeneca. He also serves as an Associate Editor for Clinical Neurology and Neurosurgery and compensated by Elsevier B.V.

Niels Bergsland has nothing to disclose.

Murali Ramanathan received research funding from the National Multiple Sclerosis Society, Department of Defense and National Institute of Neurological Diseases and Stroke.

Michael G. Dwyer received compensation from Keystone Heart for consultant fees. He received financial support for research activities from Bristol Myers Squibb, Mapi Pharma, Keystone Heart, Protembis and V-WAVE Medical.

Bianca Weinstock-Guttman received honoraria for serving in advisory boards and educational programs from Biogen Idec, Novartis, Genentech, Genzyme and Sanofi, Janssen, Abbvie and Bayer. She also received support for research activities from the National Institutes of Health, National Multiple Sclerosis Society, Department of Defense, and Biogen Idec, Novartis, Genentech, Genzyme and Sanofi.

Dr. Robert Zivadinov has received personal compensation for speaking and consulting activities with Bristol Myers Squibb, Sanofi, Novartis, EMD Serono, Janssen, Filterlex, 415 Capital, and Sana Biotechnologies. Dr. Zivadinov has received research support from the U.S. National Institutes of Health, the U.S. National Science Foundation, the U.S. Department of Defense, the U.S. National Multiple Sclerosis Society, EMD Serono, Novartis, Mapi Pharma, ProteMBis, Octave Biosciences, CorEvitas, and V-WAVE Medical. Dr. Zivadinov serves on the editorial board of *J Alzheimers Dis*, *BMC Med*, *BMC Neurol*, *Veins and Lymphatics*, *Front, Neurol*, *Trans Neurosci*, and *Clinical CNS Drugs*.

## References:

1. Thompson AJ, Baranzini SE, Geurts J, Hemmer B, Ciccarelli O. Multiple sclerosis. *Lancet*. 2018 Apr 21;391(10130):1622-36.
2. Jakimovski D, Vaughn CB, Eckert S, Zivadinov R, Weinstock-Guttman B. Long-term drug treatment in multiple sclerosis: safety success and concerns. *Expert Opin Drug Saf*. 2020 Sep;19(9):1121-42.
3. Kuhle J, Barro C, Andreasson U, et al. Comparison of three analytical platforms for quantification of the neurofilament light chain in blood samples: ELISA, electrochemiluminescence immunoassay and Simoa. *Clin Chem Lab Med*. 2016 Oct 1;54(10):1655-61.
4. Barro C, Chitnis T, Weiner HL. Blood neurofilament light: a critical review of its application to neurologic disease. *Ann Clin Transl Neurol*. 2020 Dec;7(12):2508-23.
5. Jakimovski D, Zivadinov R, Ramanathan M, et al. Serum neurofilament light chain level associations with clinical and cognitive performance in multiple sclerosis: A longitudinal retrospective 5-year study. *Mult Scler*. 2020 Nov;26(13):1670-81.
6. Barro C, Healy BC, Liu Y, et al. Serum GFAP and NfL Levels Differentiate Subsequent Progression and Disease Activity in Patients With Progressive Multiple Sclerosis. *Neurol Neuroimmunol Neuroinflamm*. 2023 Jan;10(1).
7. Kaufmann M, Schaupp AL, Sun R, et al. Identification of early neurodegenerative pathways in progressive multiple sclerosis. *Nat Neurosci*. 2022 Jul;25(7):944-55.
8. Bittner S, Oh J, Havrdova EK, Tintore M, Zipp F. The potential of serum neurofilament as biomarker for multiple sclerosis. *Brain*. 2021 Nov 29;144(10):2954-63.
9. Qureshi F, Hu W, Loh L, et al. Analytical validation of a multi-protein, serum-based assay for disease activity assessments in multiple sclerosis. *Proteomics Clin Appl*. 2023 Feb 26;n/a(n/a):e2200018.

10. Chitnis T, Foley J, Ionete C, et al. Clinical Validation of a Multi-protein, Serum-based Assay for Disease Activity Assessments in Multiple Sclerosis. medRxiv. 2023:2023.02.08.23285438.
11. Jakimovski D, Gandhi S, Paunkoski I, et al. Hypertension and heart disease are associated with development of brain atrophy in multiple sclerosis: a 5-year longitudinal study. *Eur J Neurol*. 2019 Jan;26(1):87-e8.
12. Polman CH, Reingold SC, Banwell B, et al. Diagnostic criteria for multiple sclerosis: 2010 revisions to the McDonald criteria. *Ann Neurol*. 2011 Feb;69(2):292-302.
13. Kurtzke JF. Rating neurologic impairment in multiple sclerosis: an expanded disability status scale (EDSS). *Neurology*. 1983 Nov;33(11):1444-52.
14. Motl RW, Cohen JA, Benedict R, et al. Validity of the timed 25-foot walk as an ambulatory performance outcome measure for multiple sclerosis. *Mult Scler*. 2017 Apr;23(5):704-10.
15. Feys P, Lamers I, Francis G, et al. The Nine-Hole Peg Test as a manual dexterity performance measure for multiple sclerosis. *Mult Scler*. 2017 Apr;23(5):711-20.
16. Benedict RH, DeLuca J, Phillips G, et al. Validity of the Symbol Digit Modalities Test as a cognition performance outcome measure for multiple sclerosis. *Mult Scler*. 2017 Apr;23(5):721-33.
17. Lublin FD, Reingold SC, Cohen JA, et al. Defining the clinical course of multiple sclerosis: the 2013 revisions. *Neurology*. 2014 Jul 15;83(3):278-86.
18. Hauser SL, Bar-Or A, Comi G, et al. Ocrelizumab versus Interferon Beta-1a in Relapsing Multiple Sclerosis. *N Engl J Med*. 2017 Jan 19;376(3):221-34.
19. Jakimovski D, Benedict RH, Marr K, et al. Lower total cerebral arterial flow contributes to cognitive performance in multiple sclerosis patients. *Mult Scler*. 2020 Feb;26(2):201-9.
20. Jakimovski D, Kuhle J, Ramanathan M, et al. Serum neurofilament light chain levels associations with gray matter pathology: a 5-year longitudinal study. *Ann Clin Transl Neurol*. 2019 Sep;6(9):1757-70.
21. Smith SM, Zhang Y, Jenkinson M, et al. Accurate, robust, and automated longitudinal and cross-sectional brain change analysis. *Neuroimage*. 2002 Sep;17(1):479-89.
22. Gelineau-Morel R, Tomassini V, Jenkinson M, Johansen-Berg H, Matthews PM, Palace J. The effect of hypointense white matter lesions on automated gray matter segmentation in multiple sclerosis. *Hum Brain Mapp*. 2012 Dec;33(12):2802-14.
23. De Stefano N, Stromillo ML, Giorgio A, et al. Establishing pathological cut-offs of brain atrophy rates in multiple sclerosis. *J Neurol Neurosurg Psychiatry*. 2016 Jan;87(1):93-9.
24. Dwyer MG, Hagemeyer J, Bergsland N, et al. Establishing pathological cut-offs for lateral ventricular volume expansion rates. *Neuroimage Clin*. 2018;18:494-501.
25. Meier S, Willemse EAJ, Schaedelin S, et al. Serum Glial Fibrillary Acidic Protein Compared With Neurofilament Light Chain as a Biomarker for Disease Progression in Multiple Sclerosis. *JAMA Neurol*. 2023 Feb 6;80(3):287-97.
26. Zhu W, Chen C, Zhang L, et al. Association between Serum Biomarker Profile and Real-World Evidence of Disability in Multiple Sclerosis. 2022.
27. Barro C, Benkert P, Disanto G, et al. Serum neurofilament as a predictor of disease worsening and brain and spinal cord atrophy in multiple sclerosis. *Brain*. 2018;141(8):2382-91.
28. Cantó E, Barro C, Zhao C, et al. Association Between Serum Neurofilament Light Chain Levels and Long-term Disease Course Among Patients With Multiple Sclerosis Followed up for 12 Years. *JAMA Neurol*. 2019 Nov 1;76(11):1359-66.

29. Ayrignac X, Le Bars E, Duflos C, et al. Serum GFAP in multiple sclerosis: correlation with disease type and MRI markers of disease severity. *Sci Rep*. 2020 Jul 2;10(1):10923.
30. Saraste M, Bezukladova S, Matilainen M, et al. Increased serum glial fibrillary acidic protein associates with microstructural white matter damage in multiple sclerosis: GFAP and DTI. *Mult Scler Relat Disord*. 2021 May;50:102810.
31. Cui LY, Chu SF, Chen NH. The role of chemokines and chemokine receptors in multiple sclerosis. *Int Immunopharmacol*. 2020 Jun;83:106314.
32. El Sharkawi FZ, Ali SA, Hegazy MI, Atya HB. The combined effect of IL-17F and CCL20 gene polymorphism in susceptibility to multiple sclerosis in Egypt. *Gene*. 2019 Feb 15;685:164-9.
33. Kuhlmann T, Moccia M, Coetzee T, et al. Multiple sclerosis progression: time for a new mechanism-driven framework. *Lancet Neurol*. 2023 Jan;22(1):78-88.
34. Korley FK, Yue JK, Wilson DH, et al. Performance Evaluation of a Multiplex Assay for Simultaneous Detection of Four Clinically Relevant Traumatic Brain Injury Biomarkers. *J Neurotrauma*. 2018 Jul 23;36(1):182-7.
35. Verberk IMW, Thijssen E, Koelewijn J, et al. Combination of plasma amyloid beta((1-42/1-40)) and glial fibrillary acidic protein strongly associates with cerebral amyloid pathology. *Alzheimers Res Ther*. 2020 Sep 28;12(1):118.
36. Disanto G, Barro C, Benkert P, et al. Serum Neurofilament light: A biomarker of neuronal damage in multiple sclerosis. *Ann Neurol*. 2017 Jun;81(6):857-70.
37. Benkert P, Meier S, Schaedelin S, et al. Serum neurofilament light chain for individual prognostication of disease activity in people with multiple sclerosis: a retrospective modelling and validation study. *Lancet Neurol*. 2022 Mar;21(3):246-57.
38. Tybirk L, Hviid CVB, Knudsen CS, Parkner T. Serum GFAP - reference interval and preanalytical properties in Danish adults. *Clin Chem Lab Med*. 2022 Oct 26;60(11):1830-8.
39. Ashton NJ, Suarez-Calvet M, Karikari TK, et al. Effects of pre-analytical procedures on blood biomarkers for Alzheimer's pathophysiology, glial activation, and neurodegeneration. *Alzheimers Dement (Amst)*. 2021;13(1):e12168.
40. Christensen SH, Hviid CVB, Madsen AT, Parkner T, Winther-Larsen A. Short-term biological variation of serum glial fibrillary acidic protein. *Clin Chem Lab Med*. 2022 Oct 26;60(11):1813-9.
41. van Lierop Z, Verberk IMW, van Uffelen KWJ, et al. Pre-analytical stability of serum biomarkers for neurological disease: neurofilament-light, glial fibrillary acidic protein and contactin-1. *Clin Chem Lab Med*. 2022 May 25;60(6):842-50.
42. Pedregosa F, Varoquaux G, Gramfort A, et al. Scikit-learn: Machine Learning in Python. *J Mach Learn Res*. 2011;12(null):2825–30.
43. Gado JE, Beckham GT, Payne CM. Improving Enzyme Optimum Temperature Prediction with Resampling Strategies and Ensemble Learning. *J Chem Inf Model*. 2020 Aug 24;60(8):4098-107.
44. Branco P, Torgo L, Ribeiro RP. Pre-processing approaches for imbalanced distributions in regression. *Neurocomputing*. 2019 2019/05/28/;343:76-99.

## Figure Legends:

**Figure 1.** The Octave Bioscience Multiple Sclerosis Disease Activity (MSDA) test was developed using Proximity Extension Assay (PEA) methodology on the Olink™. It measures the concentrations of 18 proteins, and utilizes an algorithm to determine 4 disease pathways scores (immunomodulation, neuroinflammation, myelin biology, and neuroaxonal integrity) and an overall disease activity score.

**Figure 2.** Changes in blood serum biomarker concentration between the baseline and follow-up time points for those with p-value < 0.05. **Legend:** Paired Wilcoxon signed-rank test was used to compare between baseline and follow-up time points. P-value annotation legend: ns:  $5.00e-02 < p \leq 1.00e+00$ , \*:  $1.00e-02 < p \leq 5.00e-02$ , \*\*:  $1.00e-03 < p \leq 1.00e-02$ , \*\*\*:  $1.00e-04 < p \leq 1.00e-03$ , \*\*\*\*:  $p \leq 1.00e-04$ .

**Figure 3.** Longitudinal single-protein model parameters with adjustment for age, sex and BMI.

**Legend:** The radius of each circle is proportional to the estimated standardized coefficient of the corresponding protein, red (blue) circles represent proteins with positive (negative) effects in estimating the second-class label. The opacity of each circle represents the p-value; a p-value of  $< 0.001$  corresponds to full opacity and a p-value of 0.05 corresponds to the least opacity. Biomarkers that survived the multiple comparisons correction are marked with a gold star (\*).

**Figure 4** Cross-sectional single-protein model parameters with adjustment for age, sex and BMI for baseline (left) and follow-up (right). **Legend:** The radius of each circle is proportional to the estimated standardized coefficient of the corresponding protein, red (blue) circles represent proteins with positive (negative) effects in estimating the second-class label. The opacity of each circle represents the p-value; a p-value < 0.001 corresponds to full opacity and a p-value of 0.05 corresponds to the least opacity. Biomarkers that survived the multiple comparisons correction are marked with a gold star (\*).



**Figure 5.** Single-protein model parameters predicting shifts in outcome score using shifts in biomarker concentration with adjustment for age, sex and BMI. **Legend:** The radius of each circle is proportional to the estimated standardized coefficient of the corresponding protein, red (blue) circles represent proteins with positive (negative) effects in estimating the second-class label. The opacity of each circle represents the p-value; a p-value < 0.001 corresponds to full opacity and a p-value of 0.05 corresponds to the least opacity. Biomarkers that survived the multiple comparisons correction are marked with a gold star (\*).

## **Tables:**

**Table 1.** Demographic, clinical and conventional MRI characteristics of the study population.

Demographic and clinical characteristics	pwMS (n=202)	CIS/RRMS (n=148)	PMS (n=54)	p-value
Female, n (%)	151 (74.8)	106 (71.6)	45 (83.3)	0.09 <sup>a</sup>
Age at baseline, mean (SD)	47.1 (11.1)	44.1 (10.6)	55.3 (7.9)	<b>&lt;0.001<sup>b</sup></b>
Time of follow-up, mean (SD)	5.4 (0.6)	5.4 (0.6)	5.5 (0.6)	0.732 <sup>b</sup>
BMI at baseline, mean (SD)	27.5 (5.8)	27.9 (6.2)	26.5 (4.5)	0.1 <sup>b</sup>
Age of disease onset, mean (SD)	32.9 (9.8)	32.6 (9.0)	33.6 (11.8)	0.6 <sup>b</sup>
Disease duration at baseline, mean (SD)	13.4 (10.2)	11.1 (8.5)	21.7 (10.5)	<b>&lt;0.001<sup>b</sup></b>
EDSS at baseline, median (IQR)	2.5 (1.5-5.0)	1.5 (1.5-2.5)	6.0 (4.0-6.5)	<b>&lt;0.001<sup>c</sup></b>
EDSS at follow-up, median (IQR)	3.0 (1.6-6.0)	2.0 (1.5-3.5)	6.5 (4.0-6.5)	<b>&lt;0.001<sup>c</sup></b>
EDSS absolute change, mean (SD)	0.4 (0.9)	0.4 (0.9)	0.4 (0.7)	<b>&lt;0.001<sup>b</sup></b>
Disability progression, n (%) <sup>*</sup>	56 (30.9)	38 (28.6)	18 (37.5)	0.251 <sup>a</sup>
Relapse rate over the follow-up, mean (SD)	0.172 (0.369)	0.204 (0.4)	0.09 (0.24)	<b>&lt;0.001<sup>d</sup></b>
DMT at baseline, n (%)				
IFN-β	85 (42.1)	60 (40.5)	25 (46.3)	0.271 <sup>a</sup>
Glatiramer acetate	37 (18.3)	24 (16.2)	13 (24.1)	
Natalizumab	29 (14.4)	25 (16.9)	4 (7.4)	
Off-label DMT	5 (2.5)	3 (2.0)	2 (3.7)	
No DMT	46 (22.8)	36 (24.3)	10 (18.5)	
DMT at follow-up, n (%)				
IFN-β	68 (33.7)	52 (35.1)	16 (29.6)	0.797 <sup>a</sup>
Glatiramer acetate	45 (22.3)	31 (20.9)	14 (25.9)	
Natalizumab	15 (7.4)	12 (8.1)	3 (5.6)	
Oral DMT	28 (13.9)	22 (14.9)	6 (11.1)	
Off-label DMT	12 (5.9)	8 (5.4)	4 (7.4)	
No DMT	34 (16.8)	23 (15.5)	11 (20.4)	
T2-LV, mean (SD)	13.5 (16.7)	10.3 (14.09)	22.2 (20.1)	<b>&lt;0.001<sup>b</sup></b>
T1-LV, mean (SD)	3.0 (7.2)	2.2 (6.44)	5.5 (8.6)	<b>0.015<sup>b</sup></b>
Gd-LN, mean (SD)	0.05 (3.2)	0.7 (3.7)	0.04 (0.2)	<b>&lt;0.001<sup>c</sup></b>
Gd-LV, mean (SD)	0.07 (0.4)	0.1 (0.48)	0.01 (0.03)	0.194 <sup>b</sup>
WBV, mean (SD)	1466.4 (94.3)	1490.2 (87.7)	1401.7 (80.8)	<b>&lt;0.001<sup>b</sup></b>
WMV, mean (SD)	725.9 (62.2)	738.7 (62.1)	689.8 (46.7)	<b>&lt;0.001<sup>b</sup></b>
GMV, mean (SD)	740.8 (63.9)	751.4 (65.9)	711.9 (48.1)	<b>&lt;0.001<sup>b</sup></b>

**Legend:** MS – multiple sclerosis, CIS – clinically isolated syndrome, RRMS – relapsing remitting multiple sclerosis, PMS – progressive multiple sclerosis, BMI – body mass index, EDSS – Expanded Disability Status Scale, DMT – disease modifying therapy, IFN – interferon, SD – standard deviation, IQR – interquartile range. LV – lesion volume, LN – lesion number, WBV – whole brain volume, WMV – white matter volume, GMV – gray matter volume. Thirteen (13) CIS/RRMS patients transitioned into PMS over the follow-up. Parametric data is shown as mean (standard deviation), whereas non-parametric data is shown as median (interquartile range). The specific comparisons were performed using; a –  $\chi^2$  test, b – Student’s t-test, c – Mann Whitney U test, c – Negative binomial regression. P-values lower than 0.05 were considered statistically significant and shown in bold. \* - Disability progression was available for 181 out of 202 pwMS due to missing EDSS values at either baseline or follow-up visit.

**Table 2.** Changes in blood serum protein concentrations between baseline and the follow-up for pwMS with worsening in disability measures.

<b>Endpoint</b>	<b>Biomarker</b>	<b>Baseline median (IQR)</b>	<b>Follow-up median (IQR)</b>	<b>Percentage Change (%)</b>	<b>p-value</b>
EDSS worsening	CCL20	11.5 (6.12, 19.3)	15.9 (10.3, 41.5)	38	<b>0.041</b>
	CDCP1	108 (81.9, 126)	135 (86.7, 180)	25	<b>0.005</b>
	TNFSF13B	4.82 (4.0, 5.95)	5.19 (4.51, 6.27)	7.6	<b>0.008</b>
20% 9HPT worsening	CCL20	12.4 (7.68, 19.1)	28.1 (13.9, 42.0)	130	<b>0.027</b>
	CDCP1	110 (105, 125)	144 (112, 193)	31	<b>0.003*</b>
	CXCL13	51.7 (37.8, 78.1)	61.9 (51.0, 103)	20	<b>0.012</b>
	CXCL9	55.7 (39.1, 77.8)	69.1 (43.9, 101)	24	<b>0.042</b>
	OPN	20.9 (16.6, 30.6)	25.2 (19.4, 33.5)	20	<b>0.042</b>
	TNFRSF10A	5.61 (4.66, 7.65)	7.71 (6.51, 8.5)	37	<b>0.005*</b>
VCAN	428 (393, 473)	500 (460, 571)	17	<b>0.007*</b>	
	4 Points SDMT Worsening	MOG	29.9 (23.0, 38.5)	32.0 (25.2, 43.4)	6.9
8 Points SDMT Worsening	SERPINA9	61.3 (34.3, 77.4)	32.1 (16.1, 59.3)	-48	<b>0.01</b>
%0.4 loss in WBV	CDCP1	105 (66.3, 132)	107 (83.7, 147)	1.5	<b>0.014</b>
3.5% increase in LV	SERPINA9	73.0 (57.1, 89.5)	64.4 (38.2, 76.6)	-12	<b>0.049</b>

**Legend:** EDSS – Expanded Disability Status Scale, 9HPT – 9-Hole Peg Test, WBV – whole brain volume. Wilcoxon signed-rank tested the significance in shifts. P-values smaller than 0.05 were considered significant and are highlighted with bold fonts, and those with asterisks (\*) survived the Benjamini-Hochberg correction for false discovery rate (FDR). All measures are shown as pg/mL except for TNFSF13B and OPN that are shown as ng/mL.

**Table 3.** Parameters of the linear mixed-effects model predicting outcome score using single-protein models consisting of a biomarker protein concentration, age, sex, BMI and time-point that passed the significant threshold of  $p < 0.05$ .

Endpoint	Biomarker	Estimate	R-squared	P-value
<b>DGMV</b>	CD6	-0.003	0.15	0.035
	TNFRSF10A	-0.004	0.17	0.009*
	MOG	0.003	0.18	0.045
	FLRT2	-0.004	0.19	0.008*
	CXCL9	-0.005	0.16	0.001*
	CDCP1	-0.005	0.18	0.002*
	CCL20	-0.003	0.16	0.004*
	APLP1	0.004	0.17	0.011*
<b>GMV</b>	FLRT2	-0.004	0.27	0.017
	CXCL9	-0.003	0.25	0.04
	CCL20	-0.003	0.26	0.006
<b>LVV</b>	MOG	-0.008	0.14	0.031
	OPN	-0.007	0.14	0.041
<b>Thalamus</b>	CCL20	-0.003	0.16	0.006
	CXCL9	-0.003	0.15	0.031
	FLRT2	-0.003	0.18	0.037
	CDCP1	-0.005	0.17	0.007
<b>WMV</b>	GFAP	-0.008	0.12	<0.001*
	TNFSF13B	-0.003	0.11	0.046
<b>WBV</b>	CDCP1	-0.003	0.24	0.004*
	CXCL9	-0.002	0.23	0.014
	FLRT2	-0.003	0.25	0.001*
	GFAP	-0.004	0.28	<0.001*
	OPG	-0.002	0.23	0.05
	TNFSF13B	-0.002	0.24	0.028
	TNFRSF10A	-0.003	0.24	0.003*

<b>EDSS</b>	NfL	0.023	0.24	0.021
<b>9HPT</b>	NfL	0.031	0.21	0.001*
	GFAP	0.022	0.2	0.037
	CDCP1	0.02	0.16	0.028
<b>PASAT</b>	GFAP	-0.038	0.07	0.012
	CCL20	-0.018	0.02	0.042
<b>SDMT</b>	CDCP1	-0.021	0.11	0.042
	CCL20	-0.017	0.09	0.008
<b>T25FWT</b>	CXCL9	0.041	0.08	0.014

**Legend:** pwMS – people with multiple sclerosis, DGMV – deep gray matter volume, GMV – gray matter volume, LVV – lateral ventricular volume, WMV – white matter volume, WBV – whole brain volume, EDSS – Expanded Disability Status Scale, 9HPT – 9-Hole Peg Test, PASAT – Paced Auditory Serial Addition Test, SDMT – Symbol Digit Modalities Test, T25FWT – Timed 25-Foot Walk Test. P-values smaller than 0.05 were considered significant and are highlighted with bold fonts, and those with asterisks (\*) survived the Benjamini-Hochberg correction for false discovery rate (FDR).

**Table 4.** Parameters of the linear model predicting baseline outcome score using single-protein models consisting of baseline biomarker protein concentration, age, sex, BMI that passed the p=0.05 significance threshold.


<b>Endpoint</b>	<b>Biomarker</b>	<b>Estimate</b>	<b>R-squared</b>	<b>p-value</b>
<b>DGMV</b>	FLRT2	-0.008	0.16	0.035
	GFAP	-0.015	0.19	<0.001*
	MOG	0.008	0.15	0.037
<b>GMV</b>	CCL20	-0.006	0.25	0.016
	GFAP	-0.009	0.33	<0.001*
	FLRT2	-0.005	0.25	0.036
<b>LVV</b>	MOG	-0.028	0.19	0.024
	GFAP	0.060	0.26	<0.001*
<b>Thalamus</b>	MOG	0.011	0.17	0.008
	GFAP	-0.016	0.18	0.001*
	FLRT2	-0.008	0.16	0.043
<b>WMV</b>	GFAP	-0.008	0.1	0.006
	CD6	0.005	0.1	0.048
<b>WBV</b>	GFAP	-0.009	0.29	<0.001*
	FLRT2	-0.004	0.25	0.021
<b>EDSS</b>	PRTG	-0.034	0.23	0.023
	GFAP	0.050	0.24	0.002*
	CD6	-0.034	0.23	0.018
<b>9HPT</b>	PRTG	-0.037	0.25	0.043
<b>PASAT</b>	GFAP	-0.092	0.11	0.022
	APLP1	-0.079	0.09	0.038
<b>SDMT</b>	GFAP	-0.055	0.18	0.013
<b>T25FWT</b>	CXCL13	0.074	0.17	0.017

**Legend:** pwMS – people with multiple sclerosis, DGMV – deep gray matter volume, GMV – gray matter volume, LVV – lateral ventricular volume, WMV – white matter volume, WBV – whole brain volume, EDSS – Expanded Disability Status Scale, 9HPT – 9-Hole Peg Test, PASAT – Paced Auditory Serial Addition Test, SDMT – Symbol Digit Modalities Test, T25FWT – Timed 25-Foot Walk Test. P-values smaller than 0.05 were considered significant and are highlighted with bold fonts, and those with asterisks (\*) survived the Benjamini-Hochberg correction for false discovery rate (FDR).




IM Immuno-modulation


N Neuroinflammation



M Myelin Biology

NI Neuroaxonal Integrity







# Shifts in biomarker to predict shifts in outcome score

9HPT

SDMT

Gray Matter

White Matter

EDSS

PASAT

Deep Gray Matter

Thalamus

Whole Brain

CNTN2

TNFRSF10A

GFAP

FLRT2

CXCL9

CD6

APLP1

PRTG

VCAN

OPN

NfL

CDCP1

TNFSF13B

