

1 **Association between physical multimorbidity and common mental health disorders in rural and**
2 **urban Malawian settings: preliminary findings from Healthy Lives Malawi long-term conditions**
3 **survey**

4 Owen Nkoka^{1,2}, Shekinah Munthali-Mkandawire¹, Kondwani Mwandira¹, Providence Nindi¹, Albert
5 Dube¹, Innocent Nyanjagha¹, Angella Mainjeni¹, Jullita Malava¹, Abena S Amoah^{1,3,4}, Estelle
6 McLean^{1,3}, Robert Stewart^{1,5}, Amelia C Crampin^{1,2,3}, Alison J Price^{1,3}

7 ¹Malawi Epidemiology and Intervention Research Unit, Lilongwe, Malawi; ²School of Health and
8 Wellbeing, University of Glasgow, Glasgow, UK; ³London School of Hygiene and Tropical Medicine,
9 Department of Population Health, London, UK; Department of Parasitology, ⁴Leiden University
10 Medical Centre, The Netherlands, ⁵Division of Psychiatry in the College of Medicine and Veterinary
11 Medicine at the University of Edinburgh

12

13

14

15 Target Journal: PloS Global Public Health (Multimorbidity in Africa Collection)

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

17 **Abstract**

18 In low-income Africa, the epidemiology of physical multimorbidity and associated mental health
19 conditions is not well described. We investigated the multimorbidity burden, disease combinations,
20 and relationship between physical multimorbidity and common mental health disorders in rural and
21 urban Malawi using early data from 9,849 adults recruited to an on-going large cross-sectional study
22 on long-term conditions, initiated in 2021. Multimorbidity was defined as having two or more
23 measured (diabetes, hypertension) or self-reported (diabetes, hypertension, disability, chronic pain,
24 HIV, asthma, stroke, heart disease, and epilepsy). Depression and anxiety symptoms were scored using
25 the 9-item Patient Health Questionnaire and 7-item General Anxiety Disorder scale and defined by the
26 total score (range 0–27 and 0–21, respectively). We determined age-standardized multimorbidity
27 prevalence and condition combinations. Additionally, we used multiple linear regression models to
28 examine the association between physical multimorbidity and depression and anxiety symptom
29 scores. Of participants, 81% were rural dwelling, 56% were female, and the median age was 30 years
30 (IQR 21–43). The age-standardized urban and rural prevalence of multimorbidity was 14.1% (95% CI,
31 12.5–15.8%) and 12.2% (95% CI, 11.6–12.9%), respectively. In adults with two conditions,
32 hypertension, and disability (18%) co-occurred most frequently, and in those with three conditions,
33 hypertension, disability, and chronic pain (23%) were the most common combination. Compared to
34 adults without physical conditions, having one (*B*-Coefficient (*B*) 0.78; 95% CI 0.63–0.95%), two- (*B*
35 1.36; 95% CI 1.14–1.58%), and three- or more- physical conditions (*B* 2.23; 95% CI 1.87–2.59%) were
36 associated with increasing depression score, *p*-trend <0.001. A comparable ‘dose-response’
37 relationship was observed between physical multimorbidity and anxiety symptom scores. While the
38 direction of observed associations cannot be determined with these cross-sectional data, our findings
39 highlight the burden of multimorbidity and need for integrating mental and physical health service
40 delivery in Malawi.

41 **Keywords:** Multimorbidity, mental health, depression, anxiety, Malawi

42 Introduction

43 In low- and middle-income countries (LMIC), population life expectancy has improved in recent
44 decades, with a 12 year increase in life expectancy at birth observed between 1990 and 2015 [1, 2] .
45 Improved access to healthcare and other improvements in living standards have contributed to an
46 increase in life expectancy in some LMIC [3, 4]. In LMIC of Africa with generalised HIV-infection
47 epidemics, the widespread availability of free, at point of care, highly effective antiretroviral
48 medication for HIV infection has played a crucial role [5]. This medication has significantly prolonged
49 the lives of individuals living with HIV, with adults surviving to experience other health conditions that
50 may also be exacerbated by long-term treated HIV, such as dyslipidaemia, hypertension and diabetes
51 [6]. As such, the burden of people living with multiple long-term health conditions is increasing, with
52 implications for individual wellbeing and wider societal and economic impacts [7].

53 Multimorbidity (MM), defined as the presence of two or more long-term conditions, is a concern for
54 healthcare systems worldwide [8, 9]. Recent multimorbidity data from LMIC show prevalence
55 estimates as high as 60% [10] with burden evident in young adults (<25 years)[11-13], which contrasts
56 to earlier findings that observed the bulk of burden in older adults (≥65 years) [14]. Observed variation
57 in multimorbidity prevalence estimates in different populations [15] highlights the need for country-
58 specific studies to determine the distinct socioeconomic and lifestyle factors that influence burden
59 [16] and to identify high-risk and vulnerable groups.

60 Mental health disorders are an emerging health burden in LMICs, accounting for 80% of people with
61 mental health conditions worldwide [17]. However, 90% of individuals with mental health disorders
62 in LMIC do not have access to mental health services [18-20]. This treatment gap is further
63 exacerbated by inequalities in accessing mental health services between rural and urban residents
64 [21]. Previous studies in LMIC have reported links between mental health disorders and
65 multimorbidity. In rural and urban India, compared to those with one or no health conditions,

66 multimorbid individuals exhibited a higher risk of depressive symptoms measured using the Centre
67 for Epidemiologic Studies Depression Scale (CES-D) [22].

68 In Malawi, a low-income, high HIV prevalence (8.9%)[23] African country, the primary healthcare
69 system is under-resourced and based largely on a single-disease delivery paradigm [24] despite a rising
70 burden of non-communicable diseases (NCDs) – including hypertension, diabetes and chronic
71 respiratory conditions [25, 26]. Furthermore, several facility-based studies in Malawi have reported a
72 high burden of mental health disorders [27, 28]. Nonetheless, most mental health conditions remain
73 under diagnosed in the community due to limited availability of mental health services in primary
74 care, and a lack of mental health funding and trained mental health professionals [29].

75 The relationship between multimorbidity and mental health disorders may be biological as well as
76 social. For example, elevated inflammatory markers and stress, have been found to be associated with
77 an increased risk of developing depression [30, 31] as well as with physical long-term conditions [32].
78 Epidemiological studies have also established links between depression, anxiety, and cardiovascular
79 diseases [33-35].

80 In LMIC, there is some evidence of the impact of individual physical health conditions on mental health
81 [36-38] yet, few studies have investigated the relationship between multiple physical health
82 conditions and depression and anxiety. A population-based study in South Africa reported no
83 significant association of hypertension alone with depression and anxiety, but revealed a significant
84 association between the co-morbidity of hypertension and another chronic physical condition and
85 anxiety [39]. A better understanding of this complex relationship is needed to provide holistic patient
86 care and to develop effective interventions for the prevention of these conditions.

87 In this study, we investigated the burden and common combinations of multiple health conditions,
88 and the association between physical multimorbidity and common mental health disorders using early
89 data from an ongoing population based cross-sectional study of long-term conditions in adolescents
90 and adults being conducted in rural and urban Malawi between 2021 and 2024.

91 **Methods**

92 Study design

93 Our long-term conditions study in rural and urban Malawi is an ongoing cross-sectional survey
94 gathering information on burden of conditions and their risk factors, along with anthropometric and
95 biological measurements. We aim to recruit 50,000 adolescents (15+ years) and adults who are
96 resident in rural and urban health and demographic surveillance systems (HDSS), which operate as
97 open-cohorts. The survey is one component of a wider “Healthy Lives Malawi” research platform,
98 investigating long-term health conditions using population-level data from the rural and urban HDSS.

99 Study setting

100 The rural Karonga HDSS was started in 2002 and currently captures information on the births, deaths
101 and movements of approximately 53,000 people covering an area of 140km² [40]. It is situated in the
102 northern part of Malawi, in Karonga district, with a predominantly subsistence economy [40]. The
103 Lilongwe HDSS (urban) covers an area of 6.68 km² and is nested within Area-25, a high-density area
104 located in the west of Lilongwe city that borders ‘Kanengo’, an industrial location including tobacco
105 auction floors. Baseline recruitment for the Lilongwe HDSS started in August 2022. At the time of
106 analysis, over 7,000 of the 65,000 people targeted by the Lilongwe HDSS have been recruited and are
107 being followed. The population includes people working in public and private sectors, including many
108 seasonal tobacco workers due to the location’s proximity to the Kanengo industrial area [41].

109 Study population

110 Individuals who are aged 15 years and above and who are usual members/residents of households (in
111 the rural or urban HDSS) are eligible for recruitment.

112 Data collection

113 Our ongoing data collection is interviewer-led using electronic data capture with Open Data Kit
114 software [42]. Questions are read to each participant in the language of their choice. Most interviews

115 in Karonga are conducted in Chitumbuka, those in Lilongwe mostly in Chichewa. All field workers were
116 trained using standard operating procedures, including obtaining informed, written (or thumbprint)
117 consent from participants, biomeasures and venepuncture. A pre-test of the survey tool and study
118 procedures was carried out from July 26 to August 06, 2021 (in rural site Karonga) and from July 18 to
119 July 29, 2022 (in urban site Lilongwe). Data collection in Karonga commenced in October 2021, and
120 Lilongwe in September 2022.

121 Measures

122 Main outcomes

123 *Depression:* To evaluate symptoms of depression, the Patient Health Questionnaire (PHQ)-9 was
124 employed to evaluate symptoms of depression [43]. The PHQ9 is a survey measure that includes items
125 that enquire about the presence symptoms of Diagnostic and Statistical Manual of Mental Disorders
126 (DSM) depressive disorder in the previous two weeks. Each item is scored from '0' (not at all), "1"
127 (sometimes), "2" (usually), and "3" (almost always). The PHQ-9 scores range from 0 to 27, with higher
128 scores corresponding to more symptoms and increasing severity of depression [44].

129 *Anxiety:* Data on anxiety symptoms were gathered using the General Anxiety Disorder (GAD)-7 scale
130 [45]. The GAD-7 is a seven-item survey measure that assesses for symptoms of DSM generalized
131 anxiety disorder in previous two weeks. Each item receives a score of "0" (not at all), "1" (sometimes),
132 "2" (usually), and "3" (almost always). The GAD-7 scale has a score range of 0 to 21, with higher values
133 indicating more symptoms and more severe anxiety [45].

134 Chichewa and Chitumbuka versions of the PHQ9 and GAD7 were developed through a rigorous
135 process of translation, adaptation, and piloting, and validated against clinical interviews.

136 Main independent variable

137 Physical health multimorbidity was defined as the presence of two or more long-term conditions.
138 These included a self-report of previously clinician-diagnosed conditions such as heart disease,

139 asthma, epilepsy, stroke, and HIV. A question ‘have you ever been diagnosed with [*name of condition*]
140 by a doctor or other health professionals’ was asked for each of the conditions to obtain self-reported
141 medical history of the participants. The Washington Group short set of questions on disability's six
142 functional domains—difficulty seeing even with glasses, difficulty hearing even with a hearing aid,
143 difficulty walking or climbing stairs, difficulty remembering or concentrating, difficulty washing oneself
144 thoroughly or dressing—were used to capture disability [46]. Disability was defined as the inability to
145 perform any of the six disability domains indicated by responses "a lot of difficulty" or "can't do at all."
146 Participants who reported having any pain for more than three months were categorised as having
147 chronic pain [47] with information on location also collected. Diabetes was defined as fasting blood
148 glucose of ≥ 7.0 mmol/L or self-report of a prior diagnosis of diabetes by a health professional. A
149 systolic blood pressure of ≥ 140 mm Hg, a diastolic blood pressure of ≥ 90 mm Hg, or self-reporting of
150 current antihypertensive drug use or prior diagnosis were all used to define hypertension. Individuals
151 with incomplete data for whom it was not possible to determine if they had at least 2 conditions, only
152 1 or no conditions, were excluded from the multimorbidity definition.

153 Covariates

154 Covariates included: site (Lilongwe (urban), Karonga (rural)); age in years (15-17, 18-29, 30-39, 40-49,
155 50-59, and ≥ 60); sex (male vs female); education attainment (not completed any level, primary, junior
156 secondary, senior secondary, and postsecondary education); employment (not employed, full-time
157 student, regular employment, and self/irregular employment); and marital status (never married,
158 married, divorced/separated, and widowed). We also considered several lifestyle and other health-
159 related factors such as smoking status (never smoked, current [smoked in the last 6 months], and
160 former [stopped more than 6 months ago]); alcohol consumption in the past one year (yes/no); being
161 physically active as defined by report of activities that strengthen muscle at least 3 days per week (for
162 those aged 15-17 years) and at least 2 days per week (among those aged ≥ 18 years) [48]; and body
163 mass index (BMI) measured as weight in kilograms by height in meters squared (kg/m^2) based on WHO

164 conventional classification and categorized into four groups (underweight < 18.5, normal 18.5–24.9,
165 overweight 25.0–29.9 kg/m² and obesity ≥30.0 kg/m²) [49].

166 Data analysis

167 We performed univariate and bivariate analyses (using one-way analysis of variance (ANOVA) or t-
168 tests, as appropriate) to describe the main variables and the relationship between selected factors
169 and depressive and anxiety symptoms as well as multimorbidity status. To account for multiple testing,
170 we also performed a Bonferroni test on associations for which the ANOVA p value was < 0.05. Age-
171 standardized prevalence of the physical and mental conditions considered in the current analysis and
172 the mental health measures were calculated using WHO standard population[50] to allow meaningful
173 comparison with external populations.

174 To investigate the association between physical multimorbidity and depressive and anxiety symptoms,
175 we applied multiple linear regression models without accounting for clustering of individuals from the
176 same households (as interclass correlation coefficients (ICC) of responses for participants from the
177 same household in relation to depressive and anxiety symptoms were low (ICC 0.14 and 0.09,
178 respectively)). We tested for trend to determine whether there was a ‘dose-response’ relationship
179 between increasing number of physical conditions and depression and anxiety scores [51, 52].

180 We investigated the extent to which the association between physical health and multimorbidity
181 varied by key sociodemographic factors (sex, site, and age), by adding an interaction term to the
182 regression model one at a time. We further conducted stratified analyses for variables with significant
183 interactions ($p < 0.05$) as this signalled potential moderation effects.

184 We used χ^2 , t-tests and ANOVA to compare the sociodemographic characteristics (sex, age,
185 educational level, employment) and the outcome measures in the individuals that were omitted from
186 the analysis due to incomplete data on multimorbidity status to those with complete data to check
187 whether there were any significant differences.

188 All analyses were performed using Stata version 17.0 (Stata Corp LP, College Station, TX, USA), and a
189 2-tailed p -value < 0.05 was considered significant.

190 Ethical Considerations

191 The Healthy Lives Malawi long-term conditions survey protocol was reviewed and approved by the
192 Malawi National Health Sciences Research Committee (approval #2642) and the University of
193 Glasgow's College of Medical, Veterinary and Life Sciences Ethics Committee (project #200200057).
194 Written informed consent for the survey was obtained from each respondent at the start of each
195 interview. For participants aged between under 18 years, written informed consent from their legally
196 acceptable representative (parent/guardian) was obtained in addition to their assent to take part in
197 the study, unless they were regarded as emancipated minors (i.e., living independently from their legal
198 guardians).

199 Results

200 *Sample Characteristics*

201 Of the 10,011 individuals approached between 19 Oct 2021 and 28 April 2023 (8,011 and 2,000 in
202 Karonga and Lilongwe, respectively), 9,849 individuals were recruited representing a 98% response
203 rate (7,930 and 1,919 from Karonga and Lilongwe, respectively). Table 1 lists the demographic and
204 health-related characteristics of the study population by sex and site. Overall, 56% of the participants
205 were female. The overall median age of the total population was 30.0 years (IQR, 21.0 – 43.0). The
206 mean age was higher among women than men in the total population 34.9 ± 16.4 vs 33.2 ± 16.2 years,
207 $p < 0.001$, and in the rural population (35.6 ± 16.9 vs 33.6 ± 16.8) $p < 0.001$. Of the participants, 51% had
208 not completed any formal education, and 50% were self-employed or had irregular employment. In
209 terms of lifestyle factors, a majority (92%) reported to have never smoked, and 19% had not consumed
210 alcohol in the past 12 months. Among the participants, 17% met criterion for being physically active,
211 most (67%) had normal weight while 25% were overweight/obese with women having the highest

212 prevalence, at 33% (Table 1). We excluded 11% (1,095 participants) from the main analysis because
213 they had missing data on multimorbidity.

214 Table 1 Characteristics of participants in rural and urban sites

Variable	Total (n=9849)			Karonga (rural, n=7930)		Lilongwe (urban, n=1919)	
	All (n=9849)	Men (n=4287)	Women (n=5562)	Men (n=3501)	Women (4429)	Men (n=786)	Women (n=1133)
Sociodemographic factors							
Median age, IQR	30.0 (21.0 – 43.0)	28.0 (20.0 – 43.0)	31 (22.0 – 44.0)	28.0 (20.0 – 44.0)	32.0 (22.0 – 45.0)	27.0 (21.0 – 39.0)	29.0 (22.0 – 39.0)
Mean age, SD (in years) ^a	34.1 (16.3)	33.2 (16.4)	34.9 (16.2)	33.6 (16.8)	35.6 (16.9)	31.3 (14.1)	32.0 (12.8)
Age group (years)							
15 – 17	1142 (12%)	590 (14%)	552 (10%)	505 (14%)	459 (10%)	85 (11%)	93 (8%)
18 – 29	3743 (38%)	1708 (40%)	2035 (37%)	1328 (38%)	1549 (35%)	380 (48%)	486 (43%)
30 – 39	1906 (19%)	720 (17%)	1186 (21%)	581 (17%)	910 (21%)	139 (18%)	276 (24%)
40 – 49	1383 (14%)	569 (13%)	814 (15%)	484 (14%)	651 (15%)	85 (11%)	163 (14%)
50 – 59	782 (8%)	345 (8%)	437 (8%)	288 (8%)	363 (8%)	57 (7%)	74 (7%)
≥60	893 (9%)	355 (8%)	538 (10%)	315 (9%)	497 (11%)	40 (5%)	41 (4%)
Education attainment							
Not completed any level	5024 (51%)	1887 (44%)	3137 (56%)	1722 (49%)	2777 (63%)	165 (21%)	360 (32%)
Primary	1604 (16%)	745 (17%)	859 (15%)	601 (17%)	645 (15%)	144 (18%)	214 (19%)
Junior secondary	1342 (14%)	616 (14%)	726 (13%)	494 (14%)	500 (11%)	122 (16%)	226 (20%)
Senior secondary	1175 (12%)	693 (16%)	482 (9%)	484 (13%)	295 (7%)	209 (27%)	187 (17%)
Post secondary	539 (5%)	303 (7%)	233 (4%)	172 (5%)	117 (7%)	134 (17%)	116 (10%)
Missing	165 (2%)	40 (1%)	125 (2%)	28 (1%)	95 (2%)	12 (2%)	30 (3%)
Marital status							
Never married	2942 (30%)	1790 (42%)	1152 (21%)	1373 (39%)	851 (19%)	417 (53%)	301 (27%)
Married	5431 (55%)	2246 (52%)	3185 (57%)	1914 (55%)	2527 (57%)	332 (42%)	658 (58%)
Divorced/separated	902 (9%)	207 (5%)	695 (13%)	179 (5%)	584 (13%)	28 (4%)	111 (10%)
Widowed	574 (6%)	44 (1%)	530 (10%)	35 (1%)	467 (11%)	9 (1%)	63 (6%)
Employment							
Not employed	2696 (27%)	741 (17%)	1955 (35%)	575 (16%)	1470 (33%)	166 (21%)	485 (43%)
Full-time student	1486 (15%)	817 (19%)	669 (12%)	661 (19%)	506 (11%)	156 (20%)	163 (14%)
Employed/regular	704 (7%)	448 (11%)	256 (5%)	263 (8%)	146 (3%)	185 (24%)	110 (10%)
Self/irregular	4963 (50%)	2281 (53%)	2682 (48%)	2002 (57%)	2307 (52%)	279 (36%)	375 (33%)
Health-related factors							
Smoking status							
Never	9097 (92%)	3561 (81%)	5536 (99%)	2907 (83%)	4416 (99%)	654 (83%)	1120 (99%)

	Former	314 (3%)	297 (7%)	17 (<1%)	228 (7%)	7 (<1%)	69 (9%)	10 (<1%)
	Current	438 (5%)	429 (10%)	9 (<1%)	366 (10%)	6 (<1%)	63 (8%)	2 (<1%)
Alcohol consumption								
	Not in last year	8018 (81%)	2680 (63%)	5338 (96%)	2164 (62%)	4265 (96%)	516 (66%)	1073 (95%)
	In last year	1827 (19%)	1604 (37%)	223 (4%)	1337 (38%)	164 (4%)	267 (34%)	59 (5%)
	Missing	4 (<1%)	3 (<1%)	1 (<1%)	-	-	3 (<1%)	1(<15)
Physical activity								
	Not met strength guidelines	8052 (82%)	3005 (70%)	5047 (91%)	2486 (71%)	4026 (91%)	519 (66%)	1021 (90%)
	Met strength guidelines	1660 (17%)	1196 (28%)	464 (8%)	936 (27%)	355 (8%)	260 (33%)	109 (10%)
	Missing	137 (1%)	86 (2%)	51 (1%)	79 (2%)	48 (1%)	7 (<1%)	3 (<1%)
Body mass index (kg/m ²)								
	Underweight	692 (7%)	426 (10%)	226 (5%)	358 (10%)	213 (5%)	68 (9%)	53 (5%)
	Normal	6614 (67%)	3338 (78%)	3276 (59%)	2752 (79%)	2680 (61%)	586 (75%)	596 (53%)
	Overweight	1646 (17%)	396 (9%)	1250 (22%)	299 (9%)	995 (22%)	97 (12%)	255 (23%)
	Obese	820 (8%)	98 (9%)	722 (13%)	68 (2%)	505 (11%)	30 (4%)	217 (19%)
	Missing	77 (<1%)	29 (<1%)	48 (<1%)	24 (<1%)	36 (<1%)	5 (<1%)	12 (1%)
Number of physical health conditions ^b								
	None	5657 (57%)	2637 (62%)	3020 (54%)	2169 (62%)	2423 (55%)	468 (60%)	597 (53%)
	One	1865 (19%)	679 (16%)	1186 (21%)	579 (17%)	979 (22%)	100 (13%)	207 (18%)
	Two	912 (9%)	312 (7%)	600 (11%)	244 (7%)	481 (11%)	68 (9%)	119 (11%)
	Three or more	320 (3%)	94 (2%)	226 (4%)	83 (2%)	194 (4%)	11 (1%)	32 (3%)
	Incomplete data	1095 (11%)	565 (13%)	530 (10%)	426 (12%)	352 (8%)	139 (18%)	178 (16%)

215 IQR= Interquartile Range, SD = standard deviation, kg= kilograms, m=meters

216 ^a *p*-values for difference in mean age between men and women, total population <0.001, rural site <0.001, and urban site = 0.203, results from t-test

217 Body mass index defined as underweight (<18.0 kg/m²), normal weight (18.0 – 24.9 kg/m²), overweight (25.0 – 29.9 kg/m²) and obese (≥30 kg/m²)

218 ^b Physical health conditions included diabetes, hypertension, heart disease, stroke, chronic pain, disability, epilepsy, HIV, and asthma. Unadjusted prevalence of multimorbidity

219 We observed significant differences in mean depressive and anxiety symptom scores in the considered
220 sociodemographic and health-related factors as displayed in S1 Appendix. Specifically, we observed
221 an increase in the mean depressive symptom scores as the number of physical conditions increased,
222 with a higher mean score among individuals with three or more physical health conditions.
223 Additionally, there was a strong correlation between depression and anxiety scores (Spearman
224 correlation coefficient (ρ)= 0.68).

225 *Prevalence of long-term conditions and multimorbidity*

226 Figure 1 presents the age-standardized prevalence of physical health conditions and multimorbidity
227 of these conditions. The age-standardized prevalence of hypertension was significantly higher among
228 urban (24.3%, 95%CI = 22.6 – 26.0%) compared to rural (15.1%, 95%CI= 14.4 – 15.8%) participants.
229 The age-standardized prevalence of chronic pain was significantly higher in rural participants (9.3%,
230 95%CI = 8.7 – 9.9%) as compared to urban participants (6.8%, 95%CI = 5.6 – 8.1%). There were no
231 significant differences in the age-standardized prevalence of multimorbidity in urban participants
232 (14.1%,95% CI, 12.5 - 15.8%) and rural individuals (12.2%, 95% CI, 11.6 - 12.9%). Multimorbidity
233 prevalence increased with age, but no significant differences were observed between males from
234 urban and rural areas, and females from urban and rural areas (Figure 2).

235 <Insert Figure 1 >

236 Figure 1 Age-standardized prevalence of physical health conditions by site.

237 *Diabetes and hypertension included objective measures plus self-reports while the rest of*
238 *the conditions were self-reported.*

239

240 <Insert Figure 2>

241 Figure 2 Age- and sex-specific prevalence of multimorbidity by site

242 *Common physical health conditions combinations*

243 The most common disease combinations among individuals with two conditions were hypertension
244 and disability (18%), and hypertension and chronic pain (14%). Among those with three conditions,
245 the most prevalent combinations were hypertension, disability, and chronic pain (24%), and
246 hypertension, disability, and HIV (8%). Diabetes, hypertension, disability, and chronic pain were the
247 most common combination (23%) among those with four conditions (S2 Appendix).

248 *Association between physical health multimorbidity and common mental health disorders*

249 Table 2 presents the association between physical health multimorbidity and depression and anxiety
250 scores. After adjusting for covariates, depressive symptom scores significantly increased with an
251 increasing number of physical health conditions. Compared to those who had no physical health
252 condition, depressive symptom score (y) increased for each increment increase in physical health
253 conditions (x): one condition (Coefficient (B) 0.78; 95% CI 0.63–0.95), two conditions (B 1.36; 95% CI
254 1.14–1.58), and three or more conditions (B 2.23; 95% CI 1.87–2.59), p -trend <0.001. Similarly,
255 compared to those without physical health conditions, an increase in anxiety symptom score was
256 observed for each increment increase in physical health conditions; one condition (B 0.67; 95% CI
257 0.53–0.81), two conditions (B 1.15; 95%CI 0.96–1.25), and three or more conditions (B 1.58; 95% CI
258 1.27–1.90), p -trend <0.001, with adjustment for covariates. While the interaction of sex or site with
259 the association between physical multimorbidity and mental health conditions were not significant,
260 the interaction terms for age and physical multimorbidity were significant but the stratified analysis
261 by age yielded no different results (S3 Appendix).

262

263 Table 2 Association between number of physical health conditions and depression and anxiety
 264 scores

		Depression score ^a					
		Unadjusted Model		Model I		Model II	
Number of physical conditions		B coefficient (95% CI)	P-value	B coefficient (95% CI)	P-value	B coefficient (95% CI)	P-value
	None	Ref.		Ref.		Ref.	
	One	0.64 (0.48 – 0.79)	<0.001	0.78 (0.63 – 0.94)	<0.001	0.79 (0.63 – 0.94)	<0.001
	Two	1.04 (0.84 – 1.25)	<0.001	1.34 (1.12 – 1.56)	<0.001	1.36 (1.14 – 1.58)	<0.001
	Three or more	1.57 (1.23 – 1.91)	<0.001	2.16 (1.81 – 2.51)	<0.001	2.23 (1.86 – 2.59)	<0.001
	<i>p</i> value for trend	<0.001		<0.001		<0.001	
		Anxiety score ^b					
Number of physical conditions							
	None	Ref.		Ref.		Ref.	
	One	0.47 (0.32 – 0.63)	<0.001	0.67 (0.53 – 0.81)	<0.001	0.67 (0.53 – 0.81)	<0.001
	Two	0.85 (0.63 – 1.07)	<0.001	1.14 (0.94 – 1.33)	<0.001	1.16 (0.97 – 1.36)	<0.001
	Three or more	1.01 (0.76 – 1.45)	<0.001	1.52 (1.20 – 1.83)	<0.001	1.59 (1.27 – 1.91)	<0.001
	<i>p</i> value for trend	<0.001		<0.001		<0.001	

265 ^a Patient Health Questionnaire-9 depression score, measured on continuous scale from 0 – 27.

266 ^b General Anxiety Disorder-7 anxiety score, measured on a continuous scale from 0 – 21.

267 Model I: Adjusted for sex, age, site education, and employment status, marital status.

268 Model II: Adjusted for the variables in Model I plus physical activity, BMI, alcohol status, and smoking status

269

270

271 *Sensitivity analyses*

272 When we compared the sociodemographic factors and the outcomes between those that had
273 multimorbidity data and those with incomplete multimorbidity data, there were significant
274 differences in the sociodemographic factors (sex, age, site, education attainment and employment
275 status), but we detected no significant differences in the mean depressive and anxiety scores between
276 the two groups (S4 Appendix). A sensitivity analysis to exclude multimorbid individuals with missing
277 data on other conditions was conducted, and results were fairly consistent (S5 Appendix). Therefore,
278 results from a sample that included them were reported.

279 **Discussion**

280 Multimorbidity due to physical health conditions was linked to an increase in depression and anxiety
281 symptom scores in these analyses of early data from an on-going, large, cross-sectional study in urban
282 and rural Malawian settings. Notably, the greater the number of co-morbid physical health conditions,
283 the greater the depression and anxiety symptom scores. Furthermore, we observed multimorbidity in
284 16% of young adults (< 30 years) in rural and urban areas, which is comparable to the 13% reported
285 in South Africa within the same age group [53]. These findings suggest that in the future, a sizeable
286 proportion of the Malawian adult population will be living with multiple long-term health conditions.

287 In these early data, there is a suggestion of differences in prevalence of multimorbidity between urban
288 and rural dwellers, consistent with the pattern observed in rural and urban India (5.8%; 95% CI 5.7 –
289 6.0% for rural and 9.7%; 95% CI 9.4 – 19.1%, respectively) [54], albeit with materially higher
290 multimorbidity estimates in Malawi. The observed multimorbidity differences were in large part
291 driven by the higher prevalence of hypertension in urban Malawi as previously described [55]. With
292 nearly 80% of the Malawian population living in rural locations [56] where health services (including
293 diagnostics and treatments) and clinical expertise is limited, the almost comparable level of
294 multimorbidity in rural areas, poses a significant challenge for the design and delivery of health care
295 needed to sustain the health of the population.

296 We observed a strong correlation ($\rho = 0.68$) between PHQ-9 and GAD-7 scores, similar to previous
297 findings that reported a correlation of 0.65 between these scales [57, 58]. The strong correlation
298 between scores reflects the common co-existence of these conditions [59, 60], overlapping symptoms
299 [61] and shared underlying neurobiological and psychological mechanisms [62, 63].

300 We noted a “dose response” increase in depression and anxiety symptoms with an increasing number
301 of physical conditions, that did not differ between urban and rural populations, and which is consistent
302 with previous research in other settings. Findings from a nationally representative survey in India have
303 shown increased odds of having depressive symptoms with an increasing number of chronic morbidity
304 [22]. A systematic review reported a 45% increase in the likelihood of having a depressive disorder
305 with each additional chronic physical condition, compared to individuals without any chronic physical
306 conditions [64]. Similarly, a study encompassing 42 countries [65] showed that compared to
307 individuals with no physical health condition, the odds of anxiety symptoms were two-fold and five-
308 fold higher in individuals with one condition and five or more conditions, respectively.

309 In the current study we cannot elucidate the biological mechanisms underpinning the observed
310 associations between increasing number of long-term physical health conditions and higher self-
311 reported depression and anxiety symptoms. However, future studies will investigate the role of
312 inflammatory biomarkers such as C-reactive protein which have been shown to predict long-term
313 conditions such as heart disease, hypertension and diabetes [66] and may also increase susceptibility
314 to genetic and environmental stressors, which may subsequently contribute to the onset of depressive
315 symptoms [67, 68]. Polypharmacy, which is common in multimorbid individuals [64], has also been
316 linked with depression [69, 70]. Furthermore, the need for ongoing adherence to treatment programs
317 for multiple conditions [71], the detrimental impacts on daily activities [72] and increased social
318 isolation [73-75] experienced with multimorbidity, are recognised risk factors for depression. Fear and
319 worry associated with managing various diseases and possible lack of treatment or treatment
320 inefficacy has also been linked with increased anxiety [76]. We have previously shown that the burden

321 of “lack of treatment” in underserved populations in Malawi is a key factor in wellbeing in those
322 suffering from long-term conditions [77], and out-of-pocket costs of care will exacerbate that anxiety,
323 hence it is plausible that those with multiple long term conditions who, due the fragmented nature of
324 primary care services and need to attend multiple clinics to access appropriate care, may be
325 disproportionately affected compared to those with single diagnosed conditions.

326 In terms of patterns of multimorbidity, hypertension, disability, and chronic pain were the most
327 common combination in our setting. People living with disability and chronic pain may experience
328 challenges in daily living [78], including their ability to work and socialize [79], which may contribute
329 to poor mental health and which may be exacerbated by limited access to appropriate treatment and
330 support [80]. Chronic pain has been shown to negatively impact sleep quality and to increase stress
331 levels, which may also contribute to the development of depression [81]. Specifically, an earlier study
332 in rural Malawi reported a significant correlation between chronic pain and poor mental health, as
333 measured using PHQ-9 and GAD-7 [82]. To better understand the extent to which modifiable lifestyle
334 determinates differentially impact the risk for long-term health conditions in rural and urban African
335 populations, it is necessary to have detailed and accurate physical activity, sleep, and diet data from
336 large population-based studies. Some of these data will become available from the current study once
337 data collection is complete.

338 The strength of this research includes the availability of standardised, detailed, and comparable data
339 from urban and rural settings across the adult population aged 15 years and older.

340 Limitations include the cross-sectional design and potential for reverse causation in the observed
341 associations. To determine causal relationships and the extent to which the relationship between
342 physical multimorbidity and common mental health disorders is bidirectional, will require long term
343 follow-up (which will become available in future), and scheduled ascertainment of time changing data,
344 including lifestyle and physical and mental health conditions. The definition of multimorbidity was

345 limited to the conditions available in our datasets and did not include all long-term health conditions,
346 affecting adults in Malawi, such as chronic obstructive pulmonary disease [83].

347 We observed some differences in the sociodemographic factors of our participants and those of the
348 wider HDSS adult populations which may affect the generalisability of our findings and highlights the
349 importance of using alternative strategies to improve recruitment in these groups for the ongoing
350 study. We excluded individuals with incomplete data regarding their health conditions, except for
351 those cases where we could categorize them based on available data as having either two or three or
352 more physical health conditions. This approach might have resulted in some misclassification and an
353 underestimation of the strength of the observed associations. Nevertheless, observed associations did
354 not change materially following exclusion of these individuals, suggesting that the effect of
355 misclassification was minimal.

356 Lastly, it is important to note that these analyses were conducted on a subset of the study. In future
357 analyses, when larger, more population-representative data become available, we will be adequately
358 powered to investigate associations stratified by age-, sex-, and urban-rural location and to investigate
359 the impact of knowledge of multimorbidity on mental health by disaggregating those with a prior
360 diagnosis of conditions (such as diabetes or hypertension) from those who are newly diagnosed during
361 the survey. This will allow us to gain deeper insights into the unique challenges faced by each group
362 and their respective impacts on well-being.

363 Our early findings highlight the need for a holistic approach to patient care in rural and urban Malawi
364 to address the substantive proportion of adolescents and adults living with multiple physical health
365 conditions and poor mental health. Constrained by limited resources, this poses a significant
366 challenge for health policy makers to design and deliver an effective and sustainable health system.
367 Possible approaches include; clinical and nursing training curricula that address multimorbidity
368 scenarios [84], to ensure healthcare professionals are equipped to support management of a variety
369 of long-term conditions (including common mental disorders) [85] and streamlining healthcare

370 processes toward holistic care [86], through provision of integrated services for the treatment of
371 commonly co-occurring condition, including support of self-management programmes. In Malawi, this
372 necessitates the addition of mental health services within primary care settings [87], training health-
373 care workers in mental health screening and management, establishing referral pathways to mental
374 health lay counsellors or clinical specialists, and availability of psychosocial support services.
375 Interventions that are scalable and sustainable in low resource environments are essential if we are
376 to address the current challenges associated with accessing multiple treatments and services in
377 Malawi.

378 Furthermore, interventions dedicated to improving the coordination of care for the elderly
379 population, who face the highest burden of multimorbidity, are not widespread in Malawi [83]. There
380 is a pressing need for such interventions, along with effective policies for disease prevention and the
381 promotion of healthy living. These interventions might include tobacco and sugar taxation, regulated
382 advertising, salt reduction strategies, improved public infrastructure to facilitate physical activity,
383 school- and workplace-based programs promoting healthy lifestyles, as well as mental health and well-
384 being, and early identification and management of health conditions [84].

385 **Conclusion**

386 Our findings highlight the burden of common mental health disorders in individuals living with
387 multiple long term physical health conditions. Efforts are urgently needed to design and deliver
388 effective, holistic health care for people living with multiple physical and mental health conditions in
389 rural and urban Malawi.

390 **Data availability statement**

391 The anonymized data relevant to this study can be obtained by submitting a reasonable request to
392 the Malawi Epidemiology and Intervention Research Unit's Data Documentalist. For data inquiries,
393 please feel free to contact Paul Kambiya at paul.kambiya@meiru.mw.

394 **Funding**

395 This work is supported by the Wellcome Trust, awarded to ACC (217073/Z/19/Z). The funder has no
396 involvement in the study design, data collection, analysis, decision to publish, or the preparation of
397 the manuscript. All aspects of the research have been conducted independently and are solely the
398 responsibility of the authors.

399 **Competing interests**

400 The authors have no competing interests to declare.

401 **Acknowledgements**

402 The authors express their sincere gratitude to the participants for generously giving their time and
403 cooperation. We would also like to extend our appreciation to the Healthy Lives field interviewers and
404 MEIRU laboratory staff for their unwavering dedication and invaluable contributions to this ongoing
405 work.

406 For the purpose of open access, the author(s) has applied a Creative Commons Attribution (CC BY)
407 licence to any Author Accepted Manuscript version arising from this submission.

408 **Author Contributions**

409 **Conceptualization:** Owen Nkoka, Amelia C. Crampin, Alison J. Price

410 **Data curation:** Owen Nkoka, Estelle Mclean, Amelia C. Crampin, Alison J. Price

411 **Formal analysis:** Owen Nkoka, Amelia C. Crampin, Alison J. Price

412 **Funding acquisition:** Amelia C. Crampin Alison J. Price

413 **Methodology:** Owen Nkoka, Shekinah Munthali, Kondwani Mwandira, Albert Dube

414 Amelia C. Crampin, Alison J. Price

415 **Project Administration:** Owen Nkoka, Shekinah Munthali-Mkandawire, Kondwani Mwandira, Albert
416 Dube, Innocent Nyanjagha, Angella Mainjeni, Providence Nindi, Jullita Malava, Robert Stewart, Abena
417 S. Amoah, Amelia C. Crampin, Alison J. Price

418 **Resources:** Amelia C. Crampin

419 **Writing original draft:** Owen Nkoka, Amelia C. Crampin, Alison J. Price

420 **Writing – review & editing:** Owen Nkoka, Shekinah Munthali-Mkandawire, Kondwani Mwandira,
421 Providence Nindi, Albert Dube, Innocent Nyanjagha, Angella Mainjeni, Jullita Malava, Abena S. Amoah,
422 Robert Stewart, Amelia C. Crampin, Alison J. Price

423

424 **References**

- 425 1. Tabutin, D. and B. Masquelier, Tendances et inégalités de mortalité de 1990 à 2015 dans les
426 pays à revenu faible et intermédiaire. 2017. 72(2): p. 227-307.
- 427 2. Sudharsanan, N., D.E. Bloom, and N. Sudharsanan. The demography of aging in low-and
428 middle-income countries: chronological versus functional perspectives. in Future directions
429 for the demography of aging: Proceedings of a workshop. 2018.
- 430 3. Salami, D., A.N. Shaaban, and M.D.R. Oliveira Martins, Africa Rising, a Narrative for Life
431 Expectancy Gains? Evidence from a Health Production Function. *Ann Glob Health*, 2019. 85(1).
- 432 4. Chewe, M. and P. Hangoma, Drivers of Health in sub-Saharan Africa: A Dynamic Panel Analysis.
433 *Health Policy OPEN*, 2020. 1: p. 100013.
- 434 5. Trickey, A., et al., Life expectancy after 2015 of adults with HIV on long-term antiretroviral
435 therapy in Europe and North America: a collaborative analysis of cohort studies. 2023. 10(5):
436 p. e295-e307.
- 437 6. Gona, P.N., et al., Burden and changes in HIV/AIDS morbidity and mortality in Southern Africa
438 Development Community Countries, 1990–2017. *BMC Public Health*, 2020. 20(1): p. 867.
- 439 7. Callahan, D., Medical Progress and Global Chronic Disease: The Need for a New Model. *The*
440 *Brown Journal of World Affairs*, 2013. 20(1): p. 35-46.
- 441 8. Basto-Abreu, A., et al., Multimorbidity matters in low and middle-income countries. *J*
442 *Multimorb Comorb*, 2022. 12: p. 26335565221106074.
- 443 9. Chowdhury, S.R., et al., Global and regional prevalence of multimorbidity in the adult
444 population in community settings: a systematic review and meta-analysis. 2023. 57.
- 445 10. Abebe, F., et al., Multimorbidity of chronic non-communicable diseases in low-and middle-
446 income countries: a scoping review. 2020. 10: p. 2235042X20961919.
- 447 11. van den Akker, M., et al., Children and adolescents are not small adults: toward a better
448 understanding of multimorbidity in younger populations. *Journal of Clinical Epidemiology*,
449 2022. 149: p. 165-171.
- 450 12. Roche, S. and E. De Vries, Multimorbidity in a large district hospital: A descriptive cross-
451 sectional study. *S Afr Med J*, 2017. 107(12): p. 1110-1115.

- 452 13. Tegethoff, M., et al., Association between mental disorders and physical diseases in
453 adolescents from a nationally representative cohort. *Psychosom Med*, 2015. 77(3): p. 319-32.
- 454 14. Ioakeim-Skoufa, I., et al., Multimorbidity Patterns in the General Population: Results from the
455 EpiChron Cohort Study. *Int J Environ Res Public Health*, 2020. 17(12).
- 456 15. Asogwa, O.A., et al., Multimorbidity of non-communicable diseases in low-income and middle-
457 income countries: a systematic review and meta-analysis. *BMJ Open*, 2022. 12(1): p. e049133.
- 458 16. Chowdhury, S.R., et al., Global and regional prevalence of multimorbidity in the adult
459 population in community settings: a systematic review and meta-analysis. *EClinicalMedicine*,
460 2023. 57: p. 101860.
- 461 17. Rathod, S., et al., Mental Health Service Provision in Low- and Middle-Income Countries.
462 *Health Serv Insights*, 2017. 10: p. 1178632917694350.
- 463 18. Naslund, J.A., et al., Exploring opportunities to support mental health care using social media:
464 A survey of social media users with mental illness. 2019. 13(3): p. 405-413.
- 465 19. Alloh, F.T., et al., Mental Health in low-and middle income countries (LMICs): Going beyond
466 the need for funding. 2018. 17(1): p. 12-17.
- 467 20. Blasco, B.V., et al., Obesity and Depression: Its Prevalence and Influence as a Prognostic
468 Factor: A Systematic Review. *Psychiatry Investig*, 2020. 17(8): p. 715-724.
- 469 21. Morales, D.A., et al., A call to action to address rural mental health disparities. 2020. 4(5): p.
470 463-467.
- 471 22. Singh, S., N. Shri, and L.K. Dwivedi, An association between multi-morbidity and depressive
472 symptoms among Indian adults based on propensity score matching. *Scientific Reports*, 2022.
473 12(1): p. 15518.
- 474 23. Ministry of Health (MOH), Malawi. Malawi Population-based HIV Impact Assessment 2020-
475 2021 (MPHIA 2020-2021): Final Report. Lilongwe: MOH, Malawi; November, 2022.
- 476 24. Albrecht, T., et al., Many diseases, one model of care? *J Comorb*, 2016. 6(1): p. 12-20.
- 477 25. Olds, P., et al., Non-communicable disease burden among inpatients at a rural district hospital
478 in Malawi. *Global Health Research and Policy*, 2023. 8(1): p. 4.
- 479 26. Gowshall, M. and S.D. Taylor-Robinson, The increasing prevalence of non-communicable
480 diseases in low-middle income countries: the view from Malawi. *Int J Gen Med*, 2018. 11: p.
481 255-264.
- 482 27. Kim, M.H., et al., Prevalence of depression and validation of the Beck Depression Inventory-II
483 and the Children's Depression Inventory-Short amongst HIV-positive adolescents in Malawi. *J*
484 *Int AIDS Soc*, 2014. 17(1): p. 18965.
- 485 28. Udedi, M., The prevalence of depression among patients and its detection by primary health
486 care workers at Matawale Health Centre (Zomba). *Malawi Med J*, 2014. 26(2): p. 34-7.
- 487 29. Lilford, P., Mental health in Malawi. *BJPsych Int*, 2020. 17(2): p. 32-33.
- 488 30. Dregan, A., et al., Common mental disorders within chronic inflammatory disorders: a primary
489 care database prospective investigation. 2019. 78(5): p. 688-695.
- 490 31. Repousi, N., et al., Depression and metabolic syndrome in the older population: A review of
491 evidence. 2018. 237: p. 56-64.
- 492 32. Liu, Y.Z., Y.X. Wang, and C.L. Jiang, Inflammation: The Common Pathway of Stress-Related
493 Diseases. *Front Hum Neurosci*, 2017. 11: p. 316.
- 494 33. Vancampfort, D., et al., *World Psychiatry*. 2016.
- 495 34. Cohen, B.E., D. Edmondson, and I.M.J.A.j.o.h. Kronish, State of the art review: depression,
496 stress, anxiety, and cardiovascular disease. 2015. 28(11): p. 1295-1302.
- 497 35. Batelaan, N.M., et al., Anxiety and new onset of cardiovascular disease: critical review and
498 meta-analysis. 2016. 208(3): p. 223-231.
- 499 36. Daré, L.O., et al., Co-morbidities of mental disorders and chronic physical diseases in
500 developing and emerging countries: a meta-analysis. *BMC Public Health*, 2019. 19(1): p. 304.

- 501 37. Asmare, Y., A. Ali, and A. Belachew, Magnitude and associated factors of depression among
502 people with hypertension in Addis Ababa, Ethiopia: a hospital based cross-sectional study.
503 BMC Psychiatry, 2022. 22(1): p. 327.
- 504 38. Ogunsakin, R.E., et al., Meta-analysis of studies on depression prevalence among diabetes
505 mellitus patients in Africa. 2021. 7(5): p. e07085.
- 506 39. Grimsrud, A., et al., The association between hypertension and depression and anxiety
507 disorders: results from a nationally-representative sample of South African adults. PLoS One,
508 2009. 4(5): p. e5552.
- 509 40. Crampin, A.C., et al., Profile: The Karonga Health and Demographic Surveillance System.
510 International Journal of Epidemiology, 2012. 41(3): p. 676-685.
- 511 41. Munthali, K.G.J.U.D.i.A. and A.G.A.o. Metropolises, Lilongwe metropolitan area. 2017: p. 319-
512 345.
- 513 42. Hartung, C., et al. Open data kit: tools to build information services for developing regions. in
514 Proceedings of the 4th ACM/IEEE international conference on information and
515 communication technologies and development. 2010.
- 516 43. Manea, L., S. Gilbody, and D.J.G.h.p. McMillan, A diagnostic meta-analysis of the Patient
517 Health Questionnaire-9 (PHQ-9) algorithm scoring method as a screen for depression. 2015.
518 37(1): p. 67-75.
- 519 44. Kroenke, K., R.L. Spitzer, and J.B.J.J.o.g.i.m. Williams, The PHQ-9: validity of a brief depression
520 severity measure. 2001. 16(9): p. 606-613.
- 521 45. Spitzer, R.L., et al., A brief measure for assessing generalized anxiety disorder: the GAD-7.
522 2006. 166(10): p. 1092-1097.
- 523 46. Madans, J.H., M.E. Loeb, and B.M. Altman. Measuring disability and monitoring the UN
524 Convention on the Rights of Persons with Disabilities: the work of the Washington Group on
525 Disability Statistics. in BMC public health. 2011. BioMed Central.
- 526 47. Treede, R.-D., et al., A classification of chronic pain for ICD-11. 2015. 156(6): p. 1003.
- 527 48. Santé, O.M.d.l., Global Recommendations on Physical Activity for Health. 2010: World Health
528 Organization.
- 529 49. WHO. Obesity: Preventing and Managing The Global Epidemic. [Internet]. 2000. Available
530 from: https://whqlibdoc.who.int/trs/WHO_TRS_894.pdf.
- 531 50. Ahmad, O.B., et al., Age standardization of rates: a new WHO standard. 2001. 9(10): p. 1-14.
- 532 51. Patino, C.M. and J.C. Ferreira, Test for trend: evaluating dose-response effects in association
533 studies. J Bras Pneumol, 2016. 42(4): p. 240.
- 534 52. Vittinghoff, E., et al., Regression methods in biostatistics: linear, logistic, survival, and
535 repeated measures models. 2006.
- 536 53. Roomaney, R.A., et al., One in five South Africans are multimorbid: An analysis of the 2016
537 demographic and health survey. PLOS ONE, 2022. 17(5): p. e0269081.
- 538 54. Prenissl, J., et al., Patterns of multimorbidity in India: A nationally representative cross-
539 sectional study of individuals aged 15 to 49 years. PLOS Global Public Health, 2022. 2(8): p.
540 e0000587.
- 541 55. Price, A.J., et al., Prevalence of obesity, hypertension, and diabetes, and cascade of care in
542 sub-Saharan Africa: a cross-sectional, population-based study in rural and urban Malawi.
543 Lancet Diabetes Endocrinol, 2018. 6(3): p. 208-222.
- 544 56. National Statistical Office. Malawi Population and Housing Census 2018. Zomba, Malawi.
- 545 57. Smith, M., et al., Clinical judgement, case complexity and symptom scores as predictors of
546 outcome in depression: an exploratory analysis. 2020. 20: p. 1-11.
- 547 58. Boothroyd, L., D. Dagnan, and S.J.H.P.C. Muncer, Psychometric analysis of the Generalized
548 Anxiety Disorder Scale and the Patient Health Questionnaire using Mokken scaling and
549 confirmatory factor analysis. 2018. 2: p. 1-4.
- 550 59. Mathew, A., et al., Co-morbidity between major depressive disorder and anxiety disorders:
551 shared etiology or direct causation? 2011. 41(10): p. 2023-2034.

- 552 60. Kalin, N.H.J.A.J.o.P., The critical relationship between anxiety and depression. 2020, Am
553 Psychiatric Assoc. p. 365-367.
- 554 61. Zbozinek, T.D., et al., Diagnostic overlap of generalized anxiety disorder and major depressive
555 disorder in a primary care sample. *Depress Anxiety*, 2012. 29(12): p. 1065-71.
- 556 62. Kovner, R., J.A. Oler, and N.H.J.A.J.o.P. Kalin, Cortico-limbic interactions mediate adaptive and
557 maladaptive responses relevant to psychopathology. 2019. 176(12): p. 987-999.
- 558 63. Etkin, A. and A.F.J.A.J.o.P. Schatzberg, Common abnormalities and disorder-specific
559 compensation during implicit regulation of emotional processing in generalized anxiety and
560 major depressive disorders. 2011. 168(9): p. 968-978.
- 561 64. Read, J.R., et al., Multimorbidity and depression: a systematic review and meta-analysis. 2017.
562 221: p. 36-46.
- 563 65. Vancampfort, D., et al., The relationship between chronic physical conditions, multimorbidity
564 and anxiety in the general population: A global perspective across 42 countries. *Gen Hosp
565 Psychiatry*, 2017. 45: p. 1-6.
- 566 66. Juber, N., C.-C. Lee, and J.J.J.B. Liu, Associations between high-sensitivity C-reactive protein
567 and non-communicable diseases in an Asian population: findings from the IFLS study. 2021.
568 26(6): p. 548-556.
- 569 67. Miller, A.H. and C.L. Raison, The role of inflammation in depression: from evolutionary
570 imperative to modern treatment target. *Nat Rev Immunol*, 2016. 16(1): p. 22-34.
- 571 68. van den Bosch, M. and A.J.A.r.o.p.h. Meyer-Lindenberg, Environmental exposures and
572 depression: biological mechanisms and epidemiological evidence. 2019. 40: p. 239-259.
- 573 69. Palapinyo, S., et al., Association between polypharmacy and depression: a systematic review
574 and meta-analysis. 2021. 51(4): p. 280-299.
- 575 70. Wiersema, C., et al., Determinants and consequences of polypharmacy in patients with a
576 depressive disorder in later life. *Acta Psychiatr Scand*, 2022. 146(1): p. 85-97.
- 577 71. Heckman, B.W., A.R. Mathew, and M.J. Carpenter, Treatment Burden and Treatment Fatigue
578 as Barriers to Health. *Curr Opin Psychol*, 2015. 5: p. 31-36.
- 579 72. Ralph, N.L., et al., Peer Reviewed: Multiple chronic conditions and limitations in activities of
580 daily living in a community-based sample of older adults in New York City, 2009. 2013. 10.
- 581 73. Leigh-Hunt, N., et al., An overview of systematic reviews on the public health consequences
582 of social isolation and loneliness. 2017. 152: p. 157-171.
- 583 74. Santini, Z.I., et al., The association between social relationships and depression: A systematic
584 review. 2015. 175: p. 53-65.
- 585 75. Ma, R., et al., Physical Multimorbidity and Social Participation in Adult Aged 65 Years and Older
586 From Six Low- and Middle-Income Countries. *The Journals of Gerontology: Series B*, 2021.
587 76(7): p. 1452-1462.
- 588 76. Brown, A., et al., Managing uncertainty in chronic illness from patient perspectives. 2020. 2(1):
589 p. 1.
- 590 77. Chikumbu, E.F., et al., Experiences of multimorbidity in urban and rural Malawi: An interview
591 study of burdens of treatment and lack of treatment. *PLOS Global Public Health*, 2022. 2(3):
592 p. e0000139.
- 593 78. Grant, M., et al., Obstacles to returning to work with chronic pain: in-depth interviews with
594 people who are off work due to chronic pain and employers. *BMC Musculoskeletal Disorders*,
595 2019. 20(1): p. 486.
- 596 79. Jung, Y.H., et al., Impact of the Acceptance of Disability on Self-Esteem among Adults with
597 Disabilities: A Four-Year Follow-Up Study. *Int J Environ Res Public Health*, 2022. 19(7).
- 598 80. Harrison, J.A.K., et al., Access to health care for people with disabilities in rural Malawi: what
599 are the barriers? *BMC Public Health*, 2020. 20(1): p. 833.
- 600 81. Alhalal, E.A., et al., Effects of chronic pain on sleep quality and depression: A cross-sectional
601 study. *Saudi Med J*, 2021. 42(3): p. 315-323.

- 602 82. Kohler, I.V., et al., Pain Is Widespread and Predicts Poor Mental Health Among Older Adults in
603 Rural Malawi. *Innov Aging*, 2022. 6(3): p. igac008.
- 604 83. Njoroge, M.W., et al., Changing lung function and associated health-related quality-of-life: A
605 five-year cohort study of Malawian adults. *eClinicalMedicine*, 2021. 41: p. 101166.
- 606 84. Whitty, C.J., et al., Rising to the challenge of multimorbidity. 2020, British Medical Journal
607 Publishing Group.
- 608 85. Leiva-Fernández, F., et al., Training primary care professionals in multimorbidity management:
609 educational assessment of the eMULTIPAP course. 2020. 192: p. 111354.
- 610 86. The, P.M.E., Multimorbidity: Addressing the next global pandemic. *PLOS Medicine*, 2023.
611 20(4): p. e1004229.
- 612 87. Stein, D.J., et al., Integrating mental health with other non-communicable diseases. 2019. 364.
- 613
- 614

S1 Appendix It is made available under a CC-BY 4.0 International license .			
S1 Appendix Association between sociodemographic and lifestyle factors and depression and anxiety scores			
Variable		PHQ-9, mean (SD)	GAD7, mean (SD)
Site			
	Karonga (rural)	1.6 (2.8)	1.3 (2.5)
	Lilongwe (Urban)	2.8 (3.5)	1.9 (2.9)
	t, p-value	- 15.5, <0.001	-8.8. <0.001
Age group (years) ^a			
	15 – 17	1.1 (2.0)	0.8 (1.7)
	18 – 29	2.0 (3.1)	1.5 (2.6)
	30 – 39	2.1 (3.3)	1.6 (2.9)
	40 – 49	1.8 (3.0)	1.4 (2.5)
	50 – 59	1.7 (2.9)	1.4 (2.5)
	≥60	1.5 (2.8)	1.1 (2.5)
	F, p-value	21.1, <0.001	18.1, <0.001
Sex			
	Male	1.7 (2.9)	1.3 (2.4)
	Female	1.9 (3.0)	1.5 (2.7)
	t, p-value	- 2.2, 0.031	-4.4. <0.001
Education attainment ^a			
	Not completed any level	1.7 (2.8)	1.4 (2.5)
	Primary	1.8 (3.1)	1.3 (2.7)
	Junior secondary	1.9 (2.8)	1.4 (2.5)
	Senior secondary & post-secondary	2.2 (3.2)	1.6 (2.6)
	Missing	1.6 (2.5)	1.1 (2.1)
	F, p-value	10.0, <0.001	2.9, 0.023
Marital status ^a			
	Never married	1.6 (2.7)	1.2 (2.2)
	Married	1.9 (3.1)	1.5 (2.7)
	Divorced/separated/Widowed	2.0 (3.1)	1.6 (2.8)
	F, p-value	9.0, <0.001	14.3, <0.001
Employment ^a			
	Not employed	2.1 (3.2)	1.5 (2.7)
	Full-time student	1.4 (2.4)	1.0 (2.0)
	Self/irregular	1.8 (2.9)	1.4 (2.6)
	Employed/regular	2.2 (3.1)	1.5 (2.6)
	F, p-value	19.4, <0.001	13.9, <0.001
Smoking status			
	Never	1.8 (2.9)	1.4 (2.6)
	Former/Current	2.5 (3.3)	1.6 (2.9)
	t, p-value	-3.5, <0.001	-2.5, 0.013
Alcohol consumption			
	Not in last year	1.8 (2.9)	1.3 (2.5)
	In last year	2.1 (3.1)	1.6 (2.7)
	Missing	5.8 (5.5)	1.7 (2.1)
	F, p-value	16.7, <0.001	9.2, <0.001
Physical activity ^a			
	Not met strength guidelines	1.8 (3.0)	1.4 (2.6)
	Met strength guidelines	1.9 (2.9)	1.4 (2.4)
	Missing	0.8 (1.6)	0.5 (1.3)
	F, p-value	9.9, <0.001	8.4, <0.001
Body mass index (kg/m²)			
	Underweight	1.6 (2.8)	1.2 (2.5)
	Normal	1.8 (3.0)	1.4 (2.6)
	Overweight	1.9 (2.9)	1.4 (2.6)
	Obese	2.0 (2.9)	1.5 (2.6)
	Missing	2.0 (3.5)	1.2 (2.5)

615
 616
 617
 618
 619
 620

	F, <i>p</i> -value	2.3 , 0.059	1.9, 0.107
	Number of physical health conditions		
	None	1.5 (2.6)	1.2 (2.2)
	One	2.2 (3.2)	1.7 (3.0)
	Two	2.6 (3.7)	2.0 (3.3)
	Three or more	3.1 (3.9)	2.2 (3.4)
	Incomplete data	1.7 (2.7)	1.3 (2.5)
	F, <i>p</i> -value	50.2 , <0.001	41.8, <0.001
F means F-value from ANOVA t means t-value from t-test ^a Bonferroni post-hoc test was done for significant ANOVA			

621 **S2 Appendix**

622

623

624

625

626

627

628

629

630

631

632

633

634

635

636

637

638

639

S2 Appendix Common combinations of multiple health conditions	
Health condition combination ^a	n (%)
<i>Two conditions (n = 912)</i>	
Hypertension + disability	164 (18%)
Hypertension + chronic pain	127 (14%)
Disability + chronic pain	87 (10%)
Hypertension + HIV	78 (9%)
Hypertension + asthma	66 (7%)
Hypertension + diabetes	60 (6%)
<i>Three conditions (n = 263)</i>	
Hypertension + disability + chronic pain	64 (24%)
Hypertension + disability + HIV	22 (8%)
Diabetes + hypertension + disability	21 (8%)
Hypertension + disability + asthma	14 (5%)
Hypertension+ disability + stroke	13 (5%)
Disability + asthma + chronic pain	13 (5%)
<i>Four conditions (n = 47)</i>	
Diabetes + hypertension + disability + chronic pain	11 (23%)
Hypertension + disability + asthma + chronic pain	6 (13%)
Hypertension + disability + heart disease + chronic pain	3 (6%)
<i>Five conditions (n = 10)</i>	
Diabetes, hypertension, disability, heart disease, chronic pain	2 (20%)
Diabetes, hypertension, disability, asthma, chronic pain	2 (20%)
^a Only combinations of conditions with prevalence ≥ 5% are presented in table S1	

S3 Appendix

S3 Appendix Sensitivity analysis of the association between number of physical health conditions and depression and anxiety scores, stratified by age											
		Depression score ^a									
		15 – 17 years		18 – 29 years		30 – 39 years		40 – 49 years		≥50 years	
Number of physical conditions		B coefficient (95% CI)	P-value	B coefficient (95% CI)	P-value	B coefficient (95% CI)	P-value	B coefficient (95% CI)	P-value	B coefficient (95% CI)	P-value
None		Ref.		Ref.		Ref.		Ref.		Ref.	
One		0.65 (0.07 – 1.23)	0.028	1.02 (0.74 – 1.29)	<0.001	0.91 (0.56 – 1.26)	<0.001	0.30 (-0.07 – 0.69)	0.113	0.45 (0.07 – 0.82)	0.019
Two		1.28 (0.34 – 2.22)	0.008	1.51 (1.02 – 2.01)	<0.001	1.57 (1.03 – 2.11)	<0.001	1.57 (1.09 – 2.05)	<0.001	0.56 (0.16 – 0.96)	0.007
Three or more		-	-	3.80 (2.46 – 5.13)	<0.001	2.56 (1.29 – 3.82)	<0.001	1.96 (1.20 – 2.71)	<0.001	1.52 (1.01 – 2.03)	<0.001
		Anxiety score ^b									
Number of physical conditions											
None		Ref.		Ref.		Ref.		Ref.		Ref.	
One		0.82 (0.32 – 1.32)	0.001	0.74 (0.49 – 0.97)	<0.001	0.84 (0.52 – 1.67)	<0.001	0.39 (0.08 – 0.70)	0.013	0.30 (-0.04 – 0.64)	0.079
Two		1.57 (0.76 – 2.38)	<0.001	1.59 (1.15 – 2.01)	<0.001	1.18 (0.68 – 1.67)	<0.001	1.18 (0.78 – 1.58)	<0.001	0.51 (0.15 – 0.87)	0.006
Three or more		-	-	3.15 (1.98 – 4.31)	<0.001	0.65 (-0.50 – 1.81)	0.267	1.78 (1.15 – 2.41)	<0.001	0.97 (0.52 – 1.43)	<0.001
^a Patient Health Questionnaire-9 depression score , measured on continuous scale from 0 – 27. ^b General Anxiety Disorder-7 anxiety score, measured on a continuous scale from 0 – 21. Model I: Adjust for sex, age, site education, and employment status, marital status. Model II: Adjust for the variables in Model I plus physical activity, BMI, alcohol status, and smoking status											

S4 Appendix

S4 Appendix Distribution of sociodemographic characteristics by availability of multimorbidity data			
Variable	Sufficient ⁺ data (n=8754)	Incomplete multimorbidity data (n=1094)	<i>p</i> -value ^a
	n (%)	n (%)	
Sex			<0.001
Male	3722 (43%)	565 (52%)	
Female	5032 (57%)	530 (48%)	
Age group (years)			<0.001
15 – 17	857 (10%)	285 (26%)	
18 – 29	3332 (38%)	411 (38%)	
30 – 39	1782 (20%)	124 (11%)	
40 – 49	1293 (15%)	90 (8%)	
50 – 59	725 (8%)	57 (5%)	
≥60	765 (9%)	128 (12%)	
Site			<0.001
Lilongwe	1602 (18%)	317 (29%)	
Karonga	7152 (82%)	778 (71%)	
Education attainment			<0.001
Not completed any level	4394 (50%)	630 (58%)	
Primary	1441 (17%)	163 (15%)	
Junior secondary	1232 (14%)	110 (10%)	
Senior secondary	1057 (12%)	118 (11%)	
Post secondary	491 (6%)	48 (4%)	
Missing	139 (2%)	26 (2%)	
Employment status			<0.001
Not employed	2385 (27%)	311 (28%)	
Full-time student	1161 (13%)	325 (30%)	
Self or irregular	4565 (52)	398 (36%)	
Employed or regular	643 (7%)	61 (6%)	
Depressive symptoms score Mean (SD) ^b	1.8 (3.0)	1.7 (2.7)	0.281
Anxiety symptoms score Mean (SD) ^b	1.4 (2.6)	1.3 (2.5)	0.467
SD = standard deviation ^a <i>p</i> -value from chi-square test unless specified. ^b <i>p</i> -value from t-test ⁺ Sufficient data if self-reported or measured with two or more conditions, irrespective of whether missing data on one or more conditions.			

S5 Appendix

S5 Appendix Sensitivity analysis of the association between number of physical health conditions and depression and anxiety scores, excluding those with missing data on one condition despite having data on the other conditions							
		Depression score ^a					
		Unadjusted Model		Model I		Model II	
Number of physical conditions		B coefficient (95% CI)	P-value	B coefficient (95% CI)	P-value	B coefficient (95% CI)	P-value
	None	Ref.		Ref.		Ref.	
	One	0.57 (0.39 – 0.74)	<0.001	0.73 (0.55 – 0.90)	<0.001	0.74 (0.56 – 0.91)	<0.001
	Two	1.13 (0.88 – 1.38)	<0.001	1.44 (1.18 – 1.71)	<0.001	1.47 (1.21 – 1.74)	<0.001
	Three or more	1.53 (1.13 – 1.92)	<0.001	2.08 (1.67 – 2.50)	<0.001	2.19 (1.76 – 2.61)	<0.001
	<i>p</i> value for trend	<0.001		<0.001		<0.001	
		Anxiety score ^b					
Number of physical conditions							
	None	Ref.		Ref.		Ref.	
	One	0.57 (0.43 – 0.70)	<0.001	0.61 (0.46 – 0.77)	<0.001	0.62 (0.46 – 0.77)	<0.001
	Two	0.89 (0.71 – 1.07)	<0.001	1.14 (0.91 – 1.37)	<0.001	1.17 (0.93 – 1.40)	<0.001
	Three or more	1.04 (0.74 – 1.33)	<0.001	1.58 (1.22 – 1.95)	<0.001	1.65 (1.28 – 2.03)	<0.001
	<i>p</i> value for trend	<0.001		<0.001		<0.001	

^a Patient Health Questionnaire-9 depression score , measured on continuous scale from 0 – 27.
^b General Anxiety Disorder-7 anxiety score, measured on a continuous scale from 0 – 21.
 Model I: Adjust for sex, age, site education, and employment status, marital status.
 Model II: Adjust for the variables in Model I plus physical activity, BMI, alcohol status, and smoking status




Figure 1





Figure 2