- 1 Social contact patterns following the COVID-19 pandemic: a snapshot of post-pandemic
- 2 behaviour from the CoMix study
- 3 Authors: Christopher I Jarvis^{*1}, Pietro Coletti^{*2}, Jantien A Backer³, James D
- 4 Munday^{1,4}, Christel Faes², Philippe Beutels^{2,5}, Christian L. Althaus⁶, Nicola Low⁶,
- 5 Jacco Wallinga^{3,7}, Niel Hens^{2,5}, W John Edmunds¹
- 6 * Contributed equally
- 7 Corresponding author: pietro.coletti@uhasselt.be
- 8 1 Centre for Mathematical Modelling of Infectious Diseases, Department of Infectious Disease
- 9 Epidemiology, London School of Hygiene and Tropical Medicine, Keppel Street, WC1E 7HT
- 10 London, UK
- 11 2. Data Science Institute, I-Biostat, Hasselt University, Agoralaan Gebouw D, Diepenbeek,
- 12 3590, Belgium
- 13 3 Centre for Infectious Disease Control, National Institute for Public Health and the
- 14 Environment, Bilthoven, the Netherlands
- 15 4 Department of Biosystems Science and Engineering, ETH Zürich, Switzerland
- 16 5. Centre for Health Economics Research and Modelling Infectious Diseases, Vaccine and
- 17 Infectious Disease Institute, University of Antwerp, Universiteitsplein 1, Wilrijk, 2610, Belgium
- 18 6 Institute of Social and Preventive Medicine, University of Bern, Bern, Switzerland
- 19 7 Department of Biomedical Data Sciences, Leiden University Medical Center, Leiden, the
- 20 Netherlands

21 Abstract

22 Background

The COVID-19 pandemic led to unprecedented changes in behaviour. To estimate if these persisted a final new round of the CoMix survey was conducted in four countries at a time when all societal restrictions had been lifted for several months.

26 Methods

We conducted a survey on a nationally representative sample in the UK, Netherlands (NL), Belgium (BE), and Switzerland (CH). Participants were asked about their contacts and behaviours on the previous day. We calculated contact matrices and compared the contact levels to a pre-pandemic baseline to estimate R₀.

31 **Results**

32 Data collection occurred from 17 November to 7 December 2022. 7,477 participants were 33 recruited. Some were asked to undertake the survey on behalf of their children. Only 14.4% of 34 all participants reported wearing a facemask on the previous day, varying between 6.7% in NL 35 to 17.8% in CH. Self-reported vaccination rates in adults were similar for each country at around 36 86%. Trimmed mean recorded contacts were highest in NL with 9.9 (95% confidence interval 37 [CI] 9.0 to 10.8) contacts per person per day and lowest in CH at 6.0 (95% CI 5.4 to 6.6). The 38 number of contacts at home were similar between the countries. Contacts at work were lowest 39 in the UK (1.4 contacts per person per day) and highest in NL at 2.8 contacts per person per 40 day. Other contacts were also lower in the UK at 1.6 per person per day (95% CI 1.4 to 1.9) and

3

highest in NL at 3.4 recorded per person per day (95% CI 4.0 to 4.0). Using the next-generation
approach suggests that R₀ for a close-contact disease would be roughly half pre-pandemic
levels in the UK, 80% in NL and intermediate in the other two countries.

44 Conclusions

- 45 The pandemic appears to have resulted in lasting changes in contact patterns that would be
- 46 expected to have an impact on the epidemiology of many different pathogens. Further post-
- 47 pandemic surveys are necessary to confirm this finding.

48 Key words

- 49 COVID-19 pandemic, post-pandemic, contact survey, social contacts, social distance, physical
- 50 distancing, Europe

52 Pandemics do not end with a bang [1, 2] but if you've seen one pandemic, then you've seen one 53 pandemic [3]! The much-desired return to normality following the COVID-19 pandemic was 54 always going to be difficult to determine both in what it means and when, if ever, it might 55 happen. The expectation that things will be the same as before is also complicated by the 56 pandemic leaving an indelible mark on society. The demonstration for the capacity of remote 57 working, where possible, may mean the number of people in offices will be lower. Socialising 58 when ill could become taboo. Facemasks may become routine for some. Sentiments towards 59 vaccines, perhaps more complex. One way we can assess the return is by conducting contacts 60 surveys [4] to measure who mixes with whom.

61 During the pandemic, the CoMix study recorded epidemiologically relevant (i.e. face-to-face) 62 social interactions in representative samples of individuals from a number of European countries 63 (21 countries in total collected data as part of the project) [5–11]. Different countries collected 64 data at different points during the pandemic[12–15]. However, the UK, Netherlands and Belgium 65 initiated their surveys during the first lockdowns in Spring 2020 and collected data more or less 66 continually for about two years. Switzerland collected data between January 2021 and May 67 2022. The surveys were used to provide rapid insights on how social contact behaviour adapted 68 as a result of the pandemic and the restrictions that governments put in place. Data collection 69 was wound up at different times, the Netherlands stopped in September 2021 and the rest of 70 the countries in this study all stopped their CoMix surveys by Spring or early Summer 2022, as 71 pandemic-specific restrictions were being lifted across Europe.

72

4

5

73 In this study, we return to measure epidemiologically relevant social contacts during late 74 November and early December 2022 in the UK, Netherlands, Belgium, and Switzerland, using 75 identical methods as for the main CoMix study. Moreover, we provide quantitative estimates of 76 contact patterns some months after all restrictions were lifted. We compare estimates of contact 77 patterns in this post-pandemic period (in which high rates of infection with Omicron subvariants 78 as well as other respiratory infections was relatively common) with those measured prior to and 79 during the pandemic. We compare the levels of mixing across the four countries and in different 80 settings. We may not yet be at a stable post-pandemic period of behaviour, with adaptations still 81 to come, but this study provides a bridge between how we behaved during 2020, the acute 82 phase of the COVID-19 pandemic, and the evolving picture of where we might be heading in the 83 years to come.

84 Methods

85 Ethics Statement

86 Participation in this opt-in study was voluntary, and all analyses were carried out on anonymised 87 data. The study was approved in the UK by the ethics committee of the London School of 88 Hygiene & Tropical Medicine Reference number 21795. The study to collect CoMix data in 89 Belgium was approved by the Ethics Committee of UZA with reference 3236 - BUN 90 B3002020000054. The Medical Research Ethics Committee (MREC) NedMec confirmed that 91 the Medical Research Involving Human Subjects Act (WMO) does not apply to the CoMix study 92 in the Netherlands (research protocol number 22/917). Therefore an official approval of this 93 study by the MREC NedMec is not required under the WMO. The study to collect CoMix data in

6

94 Switzerland was approved by the ethics committee of the Canton of Bern (project number 2020-95 02926).

96 Study design

97 We conducted an online behavioural survey called CoMix where individuals recorded details of 98 direct contacts in the day prior to the survey. We defined a direct contact as anyone who was 99 met in person and with whom at least one word was exchanged, or anyone with whom the 100 participants had any sort of skin-to-skin contact. Contacts of individuals under the age of 18 101 were collected by asking parents to answer on behalf of their child.

The design of the CoMix survey is based on the POLYMOD contact survey. The POLYMOD
survey was a self-administered paper survey in the form of a daily diary recording participants'
social contacts [16]. In the CoMix study, participants consented to self-report their social
contacts made on the day prior to survey participation. Other survey questions in CoMix
included participants' work attendance, self-reported risk status, use of facemasks, presence of
recent symptoms, and vaccination history. Details of the CoMix study including the protocol,
methodology, and survey instrument have been published previously [5, 8, 17].

CoMix was conducted in 21 European countries between March 2020 and July 2022. In this paper, we present the final additional round of data collected between Nov 2022 and Dec 2022 in the UK, the Netherlands, Belgium, and Switzerland. In each study country, a nationally representative sample was recruited using quota sampling based on age, gender, geographic region, and where possible, socioeconomic status to reflect the distribution within the national population. The market research company Ipsos recruited participants through a combination of social media, web advertising, and email campaigns to meet quotas.

7

116 Study participants

The final round of CoMix ran from 17 November 2022 to 7 December 2022. Data was collected
at similar times for all countries; starting first in the UK (17 Nov to 29 Nov), then the
Netherlands (21 Nov to 3 Dec), Switzerland (22 Nov to 7 Dec), and finally Belgium (23 Nov to 5
Dec). As per prior rounds of CoMix and due to differing funding levels, the UK panel was double
the size of the other countries with 2,991 participants (Netherlands 1,491, Switzerland 1,495,
Belgium 1,500). Most of the data was collected in adults, with a proportion of parents reporting
on behalf of their children.

124 Data

125 Reporting of contacts

126 The participants reported their contacts from the day prior to the survey in two ways: individual 127 contacts and group contacts. Individual contacts were recorded by asking the participant to list 128 each contact and their characteristics separately. Following this, we asked whether they had 129 recorded all their contacts. If they had not, then they provided details of the total number of 130 contacts they had at work, school, or other settings for the age groups 0 to 17, 18 to 59, and 131 60+, both overall and for physical contacts only ('group contacts'). They were also asked how 132 often they met each contacted person, how much time was spent with them, and their 133 relationship with the contacted person. Further details of the CoMix survey have been reported 134 extensively previously [5, 6, 8, 9].

135 Demographic information

The survey captures information about participants' demographics. Participants' ages were
grouped into categories of 0-4, 5-11, 12-17, 18-29, 30-39, 40-49, 50-59, 60-69, and 70 years
and above. Participants were asked to report how they describe their gender, with the options of

8

139 "Female", "Male", "In another way", or "Prefer not to answer". Participants were also asked140 about their household size.

141 Risk perception, status, and mitigation

Participants reported about their uptake of risk mitigating activities and responded to statements regarding their perception of risk. Participants were asked to rate the following statements: (i) "I am likely to catch coronavirus"; (ii) "I am worried that I might spread coronavirus to someone who is vulnerable"; and (iii) "Coronavirus would be a serious illness for me" with the Likert scale of "Strongly agree," "Tend to agree", "Neutral", "Tend to disagree" and "Strongly disagree". Participants self-reported whether they considered themselves to be high risk, whether they wore a face covering at least once on the prior day, and their COVID-19 vaccination status.

149 Presentation of COVID-like symptoms

Participants reported COVID-19-compatible symptoms in the 7 days prior to survey participation. These symptoms included: fever or chills, cough, shortness of breath (or difficulty breathing), fatigue (or extreme tiredness), muscle or body aches or headache, congestion (or runny nose), and sore throat.

154 Work status and attendance

Participants were asked to report if they were employed, and if so, whether they were full time, part time, or self-employed. They reported whether their work place was open and whether they attended work on the day prior to responding to the survey (the day on which they reported contacts for).

159 Statistical analysis

9

R version 4.1.1 [18] was used for all analyses, and the code and data are available online (see
Data Availability Statement). The code for the analyses conducted in this study are available on
https://github.com/jarvisc1/cmix_post_pandemic.

163 **Descriptive**

164 We calculated the counts and percentages for contacts, risk perceptions, mitigations,

symptoms, and employment related questions stratified by age, gender, household size, day of the week. While parents answer as proxies for children in the study, we describe the designated child as the "participant" where applicable. We restricted the analysis to adults only for risk perception, mitigation, symptoms, and employment questions, as we consider the data to be more reliable than those reported for children by their parents. For risk perception, we present the number and percentage of adults who strongly agreed with the statements asked.

171 Mean number of contacts

172 We calculated the mean number of contacts for each of the characteristics presented in the 173 descriptive analysis. We used a cut-off value of 100 as the maximum for contacts. This means 174 we counted any individual who reported more than 100 contacts as if they reported 100 contacts 175 to reduce the weight of individuals reporting high numbers of contacts on the mean. Previous 176 publications, specifically for the UK papers for CoMix have used a cut-off of 50 [8]. The value of 177 100 was chosen for two reasons, 1) Over 99.9% of participants reported contacts of less than 178 100, 2) The previous publication of CoMix comparing 21 countries [17] used a cutt-off of 100, so 179 for sake of consistency we used this threshold. For mean contacts by setting and country we 180 calculated 95% confidence intervals (95% CI) using bootstrapping, similar to the approach used 181 in a previous CoMix publication [17]. For mean contacts by characteristics we present means

10

with standard deviations, as this makes comparison easier with those presented in POLYMOD [16] and in other social contact surveys [4]. As per previous studies, the sample was also weighted by 2/7 for weekends and 5/7 for weekdays to account for differences in sampling of weekend and weekend days and the difference between weekend and weekday contacts.

186 Frequency and time spent with contacts

We explored types of behaviour with the frequency that participants met a contacted person, and with how long they spent with them. For this, we calculated the proportion of contacts that were physical, where a 2 metre distance was maintained, where a face-mask was used, and where they met outside. These were presented visually using stacked percentage bar charts. This approach was chosen as it allows for more direct comparison with the original POLYMOD paper [16] which explored duration and frequency with physical contact. We extend that analysis to include more pandemic specific behaviours.

194 Contact matrices

195 For each country, we constructed age-stratified contact matrices for nine age groups (0 to 4, 5 196 to 11, 12 to 17, 18 to 29, 30 to 39, 40 to 49, 50 to 59, 60 to 69, and 70+ years old). For child 197 participants and contacts, we did not record exact ages and therefore sampled from the 198 reported age-group with a weighting consistent with the age distribution of contacts for the 199 participants' own age group, according to the POLYMOD survey methods [16]. Observations 200 were weighted by 2/7 for weekends and 5/7 for weekdays. We fitted a negative binomial model 201 censored to 50 per matrix cell, due to dispersion of the reported number of contacts, to calculate 202 mean contacts between each participant and contact age groups. The value for censoring was 203 chosen to be consistent and to ease comparison with previously published contact matrix

11

estimates [8, 19]. To find the population normalised reciprocal contact matrix, we first multiplied the columns of the matrix by the mean-normalised proportion of the relevant country population in each age-group [16, 20]. Then we took the cross-diagonal mean of each element of the contact matrix. Finally, we divided the resulting symmetrical matrix by the population meannormalised proportion of the country's population in each age-group.

209 Comparison to pre-pandemic and pandemic contact levels

210 We estimated the potential relative change in basic reproduction number R₀ of an infection (that 211 spreads along the contacts, assuming everyone would be susceptible to that infection) due to 212 change in contact levels compared to pre-pandemic levels by calculating the ratio of the 213 dominant eigenvalues of the CoMix matrices to those from POLYMOD, using the same 214 approach as previously published [5]. Switzerland did not participate in the POLYMOD study 215 and we therefore used an average of the eight countries for which data was collected to provide 216 the pre-pandemic dominant eigenvalue for Switzerland. We also considered as an alternative 217 the projected synthetic contact matrix for Switzerland from Prem et al[21], as a sensitivity 218 analysis. Uncertainty for the ratios were provided by calculating the dominant eigenvalues from 219 1,000 bootstrap samples for the CoMix matrices for each country and the dominant eigenvalue 220 from 1,000 bootstrap samples for the POLYMOD matrices for each country.

We further compared POLYMOD to the earliest estimates of contact levels during the 1st
lockdown in the UK and BE. This estimate was not repeated for Switzerland and the
Netherlands as data from children in these countries was not collected until later (December
2020).

12

225 Comparison of post-pandemic and pandemic behaviours

We compared several of the measurements made during this final round of CoMix to those previously published from the prior rounds of the survey in order to frame the current findings in relation to those during the pandemic. We provide an exploratory but non-comprehensive comparison in order to reduce the burden for the reader to compare across multiple publications.

231 **Results**

232 Participant characteristics

Overall, we recorded observations on 7,477 participants who reported 74,534 contacts between
16 November 2022 and 6 December 2022 in the UK, Belgium, Netherlands, and Switzerland
(Table 1). Just under 20% (1,336) were proxy respondents (i.e. the survey was completed by
parents on behalf of children), and 6,141 were adults. The UK has the highest number of
participants at 2,991, almost double the number of the other countries.

238 The age distributions were broadly similar across the four countries, with Switzerland the most 239 different with slightly more over 70s and fewer 60-69, and more 5-11s and fewer 12-17 year 240 olds. There were 3,781 (50.8%) females and 3,667 (49.2%) males, with a similar roughly equal 241 split in all countries. The majority of households consisted of 3-5 people in total with less than 242 2.5% of participants in any country being in a household size of six or more. Contact data was 243 collected on every day of the week for all countries, though some days had lower participation 244 such as 24 (0.8%) and 32 (1.1%) responses in the UK on Friday and Saturday, and 41 (2.7%) in 245 Belgium on Monday, and 26 (1.7%) in the Netherlands on Tuesdays.

246 **Risk Perception**

Overall, 7.6% of the sample (ranging from 6.2% in the UK to 10.3% in the Netherlands) strongly agreed that they were at risk of catching coronavirus and 9.5% strongly agreed that they were at high risk of severe disease if they did catch coronavirus (ranging from 6.3% in Switzerland to 13.4% in the Netherlands). A slightly higher fraction (12.4%) strongly agreed that they were likely to spread the virus to someone vulnerable, varying from 7.7% in Belgium to 15.3% in the UK.

253 Risk Mitigation

Only 14.4% of participants reported wearing a facemask on the previous day. The Netherlands
had the lowest with 6.7% participants wearing a facemask and Switzerland the highest with
17.8% (Table 1). Self-reported vaccination in adults was similar for each country at around 85%
vaccinated. The UK had the lowest percentage of people self-reporting as being high risk at
17.2% versus 31.2% in the Netherlands.

259 Symptoms

260 Nearly 40% of participants reported at least one of the following symptoms: fever or chills,

- 261 cough, shortness of breath (or difficulty breathing), fatigue (or extreme tiredness), muscle or
- body aches or headache, congestion (or runny nose), and sore throat.

263 Employment

About 43% of adult participants were employed, though this includes individuals who may be retired as unemployed in the denominator. Of those that were employed, the majority (between 60 to 80%) in each country were in full time employment, and around 5% were self employed.

14

For those in employment the vast majority (~90%) reported their workplaces were open and around two thirds attended work in person on the day they made their contacts (Table 1).

²⁶⁹ Mean Contacts by country and setting

Participants from the Netherlands recorded considerably more contacts than the other three 270 271 countries with 9.9 (95% CI 9.0 to 10.8) contacts per person per day, as compared to 6.5 (95% CI 272 6.0 to 7.0) contacts in UK, 6.7 (95% CI 6.0 to 7.3) in Belgium and 6.0 (95% CI 5.4 to 6.6) in 273 Switzerland. (Table 2). This pattern was also seen for adults and children (8.8, 95% CI 7.9 to 9.8 274 for adults; 14.8, 95% CI 12.6 to 16.8 for children in the Netherlands). As well as overall contacts, 275 we measured contacts for the four settings of home, work, school, and other. Contacts at home 276 were very similar between the countries, with an average of about 1.5 contacts per person per 277 day recorded, which is consistent with the household sizes seen in Table 1 (a mean of 2.6 278 overall for the study). Contacts at work for adults were lowest in the UK (a mean of 1.5 contacts 279 recorded per person per day, 95% CI 1.2 to 1.9) and highest in the Netherlands at 3.3 contacts 280 per person per day (95% CI 2.7 to 4.0). Other contacts (mostly in social settings) were also 281 lowest in the UK at 1.6 per person per day (95% CI 1.4 to 1.9) and highest in the Netherlands at 282 3.3 recorded per person per day (95% Cl 2.7 to 4.0).

²⁸³ Frequency and time spent with contacts

Higher frequency contacts (1-2 days) were more likely to include physical touch (> 50%)
compared to less frequent contacts (e.g. never met <25%) (Figure 1). Similarly, physical contact

- 286 was more likely for those spending 4 hours or more with a contact, with the proportion of
- 287 physical contacts observed in the data reducing as the duration of contact reduced (Figure 2).

15

The percentage of contacts met every 1-2 days that were physical was similar to those seen in POLYMOD for the UK, Netherlands, and Belgium (Figure 1). Though for those meeting less often, the percentage of physical contacts appears lower than POLYMOD for the UK and Belgium but similar for the Netherlands (Figure 1). The patterns were somewhat consistent for time spent with contacts and percentage of physical contact, the Netherlands had near identical percentages, Belgium had slightly lower for all but the shortest durations of contacts, and the UK had slightly lower for all but the longest duration of contacts (Figure 2).

The percentage of participants staying at least 2 metres away was slightly higher in the Netherlands though still less than 25% for all countries, with only those who were met every 1-2 days being less likely to wear a mask compared to when meeting a less frequent contact (Figure 1). Maintenance of a two metre distance appears to be more common for shorter interactions (Figure 2),

Mask wearing was infrequent (<15%) in all countries and for all types of contact, with
participants less likely to wear a mask when meeting someone often and for shorter periods
(<5m) or longer (4h+) periods of time (Figures 1 and 2).

The fraction of contacts who met outside were similar for all frequency of contact and across the four countries (Figure 1). There was a slight trend (in each country) for longer-duration contacts to have occurred outside (Figure 2).

306 Mean contacts by characteristics

307 Age, Gender, Households size

The reported mean contacts for school-aged (5-11 and 12-17 years of age) in the UK and
Netherlands were similar at around 14 contacts per person per day, whereas Belgium and

16

Switzerland were lower with both at around 10 contacts (Table 3). This pattern was different
amongst adults, with the UK reporting the lowest levels of contacts in most adult age groups.
Young adults (18-29 years old) in Belgium and the Netherlands reported the highest mean
contact rates (7.6 and 10.4 per person per day, respectively, as compared to 4.8 in the UK and
5.9 in Switzerland).

315 Females generally reported more contacts than males, though this pattern was not consistent in

and each country. As expected, household size was positively correlated with the number of

317 reported contacts with some slight departures from this pattern in Belgium and the Netherlands.

318 Day of the week

Contacts by day suggest a strong weekend effect for all countries, with far lower contacts on the
weekend and also on a day either side of the weekend for the UK (Friday) and Belgium and the
Netherlands (Monday) (Table 3).

322 Risk mitigation

Those who reported wearing a facemask tended to report fewer contacts in all countries other than Belgium. Those self reporting as high risk reported lower contacts across all four countries. Those who were vaccinated tended to report fewer contacts than those who said they had not been vaccinated (except for in Belgium), though it should be stressed that this is a univariate analysis and the unvaccinated tended to be younger in age.

328 Employment

Number of contacts were highest for employed people in the Netherlands, with self employed
people in Belgium and the Netherlands reporting about 20 contacts per person per day. With the
vast majority of workplaces being open now, contacts still tended to be higher for people whose

17

workplace was open. As expected, there was still a considerable difference in the meancontacts for those who attended work versus those who did not.

334 Contact matrices and changes in pre-pandemic and post-

335 pandemic R₀

Contact matrices were similar across the four nations, with high rates of recorded contacts along the leading diagonal (suggesting that contact is age-assortative) and the highest rates of recorded contacts being for children (Figure 3A). The Netherlands had the highest levels of contacts overall. There were comparatively high levels of contact between over 70s in all countries, except Belgium.

341 Using the next-generation approach [22], these contact matrices can be used to estimate R_0 for 342 close-contact infections spread through physical or conversational contacts (as measured here), 343 assuming that everyone is susceptible to infection. The relative change in R_0 for reported 344 contacts, compared to contacts at pre-pandemic levels (as measured in the POLYMOD study) 345 is shown in Figure 3B (Table S1). The reduction in contacts, compared with POLYMOD, would 346 lead to a significant reduction in the reproduction number R_0 in each of the four countries, with 347 the UK's R₀ being roughly half of pre-pandemic levels and the Netherlands about 80% of the 348 pre-pandemic level (with the other two countries being intermediate). For context, Figure 3B 349 also shows the relative reduction in R_0 during the first lockdown in the UK, which was 25% of 350 the pre-pandemic level and Belgium which was 20% of pre-pandemic levels (Figure 3B and 351 Table S1) [6]. Using the projected contact matrix for Switzerland by Prem et al. [21] instead of 352 the POLYMOD average leads to comparable results (Figure SI1).

353 Comparison of post-pandemic and pandemic behaviours and

354 contacts

In this section comparisons are made between the results from this round and those found for
the UK, Netherlands, Belgium, and Switzerland in the analysis of the 2-year 21 country study by
Wong et al [9].

358 Risk Perception, Mitigation, and Symptoms

359 It appears that risk perceptions may have slightly changed since the pandemic with participants

360 reporting a slightly higher belief that they will catch coronavirus (6-10% versus 4-5% in Wong),

and a lower concern that it would be a serious illness for them (6 to 13% versus 14 to 25% in

Wong), or that they will pass it on to someone considered vulnerable (8 to 15% versus 19 to 26% in Wong).

The percentage of individuals wearing facemasks (7 to 18%) were considerably lower than levels measured in each individual country over the course of the pandemic (UK, 58.2%, BE 61.4%, NL 34.3%, CH 76.7%) [9].

The percentage of participants reporting at least one symptom (38 to 43%) was quite a bit higher than reported over the two years of the pandemic which was between 21 to 26% [17]. (Table 1 in this paper versus Table 1 in Wong et al [9]).

370 Contacts

The mean contacts measured in this survey were somewhat higher at between 6-10 for the mean contacts for the four countries compared to 3-4 contacts per day measured during the pandemic [9]. Apart from a general increase in the level of contacts the main change appears to be in those 70+, an age group with very few contacts made during the pandemic, especially in

19

the UK (see Figure 6A in Gimma et al [8]). In contact matrices measured during the pandemic in the UK, those aged 70 or older never had more than 1 contact on average with those also aged 70+ and less than 0.4 for contacts with other age groups. In contrast, we estimate a value of 1.7 for 70+ year olds mixing with 70+ year olds and values as high as 0.7 for mixing with other age groups.

380 Discussion

We estimate that contact levels have increased compared to those measured during the pandemic but still remain lower than those measured prior to the pandemic. These reduced levels are likely to have a big impact on transmission with a reduction of R₀ of between 20% to 50% compared to pre-pandemic levels across the four nations. The consequences of this change in behaviour extends well beyond Covid and would be expected to have an impact on a range of infections that are spread person-to-person.

The use of facemasks has dropped considerably compared to the levels measured during the pandemic. We estimated around 15% of people wore a face mask on the day of the study across the four countries which is considerably lower than the 64% average observed during the pandemic across 21 European countries [17].

Contacts amongst the individuals over the age of 70 were consistently low during the pandemic
and we observed a bounce back in the number of contacts over 70s make especially in the
social setting.

Contact patterns were broadly similar across the four countries, with the Netherlands generally
reporting a higher level of contacts. The patterns of the frequency of contacts, whether they're
physical or not, and the duration of contacts were somewhat similar to those seen prior to the
pandemic.

20

We also observed that the proportion of individuals who think they are likely to get Covid was higher than those measured during the pandemic but there has potentially been a shift that it is considered less serious for them and there is less concern about giving it to someone vulnerable. Given the relationship between perceived severity and contacts measured during the pandemic, this is one of the potential explanation for the increase in contacts that we observed [23].

The CoMix study was nearly identical in the four countries, with the same questionnaire (apart from translation issues) and a similar sampling frame, and collected by the same survey organisation at the same calendar time. The study design was also the same as those used for the previous rounds of CoMix which allows for more straightforward comparison to the estimate calculated during the pandemic. We also structured our analyses to be consistent with previous analyses conducted for POLYMOD and CoMix.

A difficulty of our study design is that it is retrospective (individuals were asked about their contacts on the previous day), so may miss contacts, particularly those that would be short lasting. Furthermore, the children's contacts are a proxy with parents reporting on behalf of those under-18. We also allow individuals to estimate mass contacts that they were unable to report individually, which results in skewed distributions of contacts and is why a maximum threshold value of 100 contacts per person is used for estimates of the mean.

This research provides a snapshot picture of contacts in four European nations during the return to post-pandemic patterns of behaviour. We have measurements that are higher than those seen during the pandemic but are still considerably lower than those prior to the pandemic. It may be that the huge changes we saw during the pandemic are not over, and it will be important to monitor changes in contacts that may occur over the coming years.

21

It appears that the pandemic, at least in terms of behaviour, is ending very slowly and we are seeing a long return to contact level prior to 2019. However, it could be that we may never return to the levels of contacts seen before the pandemic. The changes in work patterns, and behaviour may have resulted in long-lasting impacts with implications on the epidemiology of a wide range of infections, as well as on important societal and economic outcomes.

426 Conclusions

Despite the number of contacts being higher compared to pandemic levels, we are not back to the levels seen prior to the pandemic. The Netherlands and Belgium appear closer to prepandemic levels with the UK further behind. These divergences between countries may represent long-term changes and measuring the level of social interactions in the years to come will allow this to be assessed. Pandemics may not end with a bang but perhaps rather a slow and cautious trudge back to newly considered risky behaviour that was previously part of everyday life.

- 436 Figure 1: Frequency of meeting the contacts for each country by: **A**. Whether it was a physical
- 437 contact or not, **B** whether they were further than 2 metres from the contact, **C** Whether they
- 438 wore a mask when meeting the contact, **D** whether they met outside when meeting the contact



	23
23	

- Figure 2: Time spent with contacts for each country by: **A**. Whether it was a physical contact or
- not, **B** whether they were further than 2 metres from the contact, **C** Whether they wore a mask
- 446 when meeting the contact, **D** whether they met outside when meeting the contact



Figure 3: **A**: Contact matrices for each country. **B**: Points show relative change in R₀ (compared to POLYMOD) based on the dominant eigenvalues of contact matrices.



-

25

25

453

Category Value All UK BE NL СН 7,477 All 2,991 1,500 1,491 1,495 Adult N (%) 6,141 (82.1%) 2,488 (83.2%) 1,200 (80.0%) 1,215 (81.5%) 1,238 (82.8%) Child 1,336 (17.9%) 503 (16.8%) 300 (20.0%) 276 (18.5%) 257 (17.2%) Age group (Children) 0-4 176 (15.4%) 49 (14.0%) 33 (11.3%) 42 (16.3%) 52 (21.4%) 5-11 424 (37.1%) 127 (36.4%) 110 (37.7%) 81 (31.4%) 106 (43.6%) 12-17 542 (47.5%) 173 (49.6%) 149 (51.0%) 135 (52.3%) 85 (35.0%) Unknown 194 154 8 18 14 212 (17.7%) 18-29 373 (16.6%) 202 (16.9%) 992 (17.0%) 205 (17.3%) Age group (Adult) 30-39 999 (17.1%) 411 (18.3%) 196 (16.3%) 188 (15.8%) 204 (17.1%) 40-49 325 (14.5%) 189 (15.8%) 193 (16.3%) 906 (15.5%) 199 (16.7%) 50-59 988 (16.9%) 403 (17.9%) 206 (17.2%) 190 (16.0%) 189 (15.8%) 60-69 1,203 (20.6%) 474 (21.1%) 262 (21.8%) 264 (22.2%) 203 (17.0%) 70+ 743 (12.7%) 263 (11.7%) 135 (11.2%) 147 (12.4%) 198 (16.6%) Unknown 43 310 239 28 Gender Female 1,564 (52.5%) 733 (49.0%) 759 (51.1%) 3,781 (50.8%) 725 (48.7%) Male 726 (48.9%) 3,667 (49.2%) 1,414 (47.5%) 762 (51.0%) 765 (51.3%) Other 29 13 5 6 5 Household size 1 1,508 (20.2%) 538 (18.0%) 295 (19.7%) 339 (22.7%) 336 (22.5%) 2 2,520 (33.7%) 1,062 (35.5%) 473 (31.5%) 487 (32.7%) 498 (33.3%) 3-53,292 (44.0%) 1,323 (44.2%) 699 (46.6%) 638 (42.8%) 632 (42.3%) 6+ 157 (2.1%) 68 (2.3%) 33 (2.2%) 27 (1.8%) 29 (1.9%) Day of week Sun 1,796 (24.0%) 533 (17.8%) 317 (21.1%) 656 (44.0%) 290 (19.4%) Mon 856 (11.4%) 357 (11.9%) 41 (2.7%) 111 (7.4%) 347 (23.2%) Tue 1,663 (22.2%) 676 (22.6%) 570 (38.0%) 391 (26.2%) 26 (1.7%) Wed 1,704 (22.8%) 950 (31.8%) 256 (17.1%) 322 (21.6%) 176 (11.8%) Thu 848 (11.3%) 419 (14.0%) 117 (7.8%) 234 (15.7%) 78 (5.2%) Fr 366 (4.9%) 24 (0.8%) 132 (8.8%) 88 (5.9%) 122 (8.2%) Sat 244 (3.3%) 32 (1.1%) 67 (4.5%) 54 (3.6%) 91 (6.1%) Risk perception (Adults) Catching 441 (7.6%) 139 (6.2%) 89 (7.4%) 122 (10.3%) 91 (7.6%) coronavirus Strongly agree only Serious illness 554 (9.5%) 187 (8.3%) 133 (11.1%) 159 (13.4%) 75 (6.3%) from coronavirus Spreading 723 (12.4%) 344 (15.3%) 92 (7.7%) 157 (13.2%) 130 (10.9%) coronavirus to vulnerable people Risk mitigation (Adults) Face mask 886 (14.4%) 393 (15.8%) 191 (15.9%) 82 (6.7%) 220 (17.8%) Vaccinated 5,269 (85.8%) 2,196 (88.3%) 1,044 (87.0%) 1,044 (85.9%) 985 (79.6%) High risk 1,468 (24.2%) 423 (17.2%) 347 (29.3%) 372 (31.2%) 326 (26.8%) Symptoms (Adults) Fever 247 (4.2%) 84 (3.7%) 47 (3.9%) 46 (3.9%) 70 (5.9%) Cough 835 (14.3%) 326 (14.5%) 153 (12.8%) 157 (13.2%) 199 (16.7%) Shortness of 311 (5.3%) 146 (6.5%) 50 (4.2%) 67 (5.6%) 48 (4.0%) breath Congestion 892 (15.3%) 315 (14.0%) 180 (15.0%) 198 (16.7%) 199 (16.7%) Sore throat 554 (9.5%) 199 (8.8%) 118 (9.8%) 106 (8.9%) 131 (11.0%) Fatigue or 551 (9.4%) 237 (10.5%) 97 (8.1%) 115 (9.7%) 102 (8.5%) tiredness Any symptoms 2,324 (39.9%) 872 (38.8%) 462 (38.5%) 473 (39.8%) 517 (43.3%) Full time Employed (Adults) 1,772 (68.2%) 677 (69.1%) 331 (59.4%) 385 (67.3%) 379 (77.3%) Part time 251 (25.6%) 682 (26.2%) 87 (17.8%) 192 (34.5%) 152 (26.6%)

Self employed

145 (5.6%)

52 (5.3%)

24 (4.9%)

34 (6.1%)

35 (6.1%)

454	Table 1:	: Participants	characteristics i	n the	CoMix	surve	y for	each	country
-----	----------	----------------	-------------------	-------	-------	-------	-------	------	---------

Work open (Adults)	Closed	297 (9.1%)	148 (11.2%)	48 (8.0%)	51 (7.5%)	50 (7.5%)
	Open	2,980 (90.9%)	1,179 (88.8%)	552 (92.0%)	630 (92.5%)	619 (92.5%)
Attended work (Adults)	Yes	1,632 (61.6%)	598 (60.0%)	307 (61.5%)	299 (52.3%)	428 (73.8%)

Table 2: Mean contacts by country and setting

Category	Setting	UK	BE	NL	СН
All participants		Mean (95% CI*)			
	All	6.5 (6.0 to 7.0)	6.7 (6.0 to 7.3)	9.9 (9.0 to 10.8)	6.0 (5.4 to 6.6)
	Home	1.5 (1.5 to 1.6)	1.6 (1.5 to 1.6)	1.6 (1.5 to 1.7)	1.5 (1.4 to 1.5)
	Work	1.4 (1.1 to 1.7)	1.7 (1.3 to 2.1)	2.8 (2.3 to 3.3)	1.6 (1.3 to 1.9)
	School	2.2 (1.9 to 2.6)	1.7 (1.3 to 2.1)	3.0 (2.5 to 3.5)	1.1 (0.8 to 1.4)
	Other	1.6 (1.4 to 1.9)	2.2 (1.9 to 2.6)	3.4 (3.0 to 4.0)	2.2 (1.9 to 2.7)
Adults					
	All	5.4 (5.0 to 5.9)	5.5 (4.8 to 6.2)	8.8 (7.9 to 9.8)	5.3 (4.8 to 5.9)
	Home	1.4 (1.3 to 1.4)	1.3 (1.3 to 1.4)	1.4 (1.3 to 1.5)	1.3 (1.2 to 1.4)
	Work	1.5 (1.2 to 1.9)	2.1 (1.7 to 2.7)	3.3 (2.7 to 4.0)	1.9 (1.5 to 2.3)
	School	1.2 (1.0 to 1.5)	0.5 (0.3 to 0.8)	1.9 (1.4 to 2.4)	0.5 (0.3 to 0.7)
	Other	1.6 (1.4 to 1.9)	2.0 (1.6 to 2.3)	3.3 (2.7 to 4.0)	2.1 (1.7 to 2.5)
Children					
	All	11.1 (9.4 to 12.7)	10.4 (8.7 to 12.3)	14.8 (12.6 to 16.8)	9.1 (7.4 to 11.1)
	Home	2.2 (2.1 to 2.3)	2.2 (2.1 to 2.4)	2.6 (2.4 to 2.8)	2.1 (1.9 to 2.3)
	Work	0.9 (0.4 to 1.5)	0.1 (0.0 to 0.3)	0.3 (0.1 to 0.4)	0.4 (0.2 to 0.7)
	School	6.7 (5.3 to 8.1)	5.5 (4.3 to 6.9)	8.1 (6.6 to 9.6)	4.0 (2.8 to 5.2)
	Other	1.7 (1.2 to 2.3)	3.1 (2.2 to 4.0)	4.0 (3.1 to 4.9)	3.1 (2.1 to 4.2)

*Bootstrapped mean and 95% percentage confidence interval from 1,000 samples. Sample weighted by 2/7 for weekends and 5/7 for weekdays. 458

Value UK BE Category NL СН All Mean (SD) 6.1 (13.6) 6.5 (13.5) 9.2 (17.1) 5.8 (11.3) Adult 5.2 (12.1) 5.6 (12.5) 8.2 (16.8) 5.2 (10.0) Child 10.9 (18.8) 10.4 (16.2) 14.1 (17.6) 9.0 (15.9) Age group (Children) 12.4 (12.6) 0-4 10.2 (17.6) 11.8 (14.9) 6.2 (10.4) 5-11 14.3 (18.7) 11.5 (16.7) 14.3 (17.7) 10.6 (18.5) 12-17 14.2 (22.6) 9.8 (16.4) 15.4 (19.4) 9.7 (16.4) Unknown 18-29 4.8 (10.9) 7.6 (16.1) 10.4 (22.1) 5.9 (10.3) Age group (Adult) 30-39 4.8 (13.1) 6.7 (12.3) 5.8 (12.6) 7.5 (13.1) 40-49 4.4 (9.8) 6.9 (15.7) 8.5 (16.2) 5.6 (9.9) 50-59 5.8 (15.1) 5.5 (12.5) 8.4 (17.6) 5.3 (12.0) 60-69 2.7 (3.7) 3.9 (8.1) 7.0 (15.7) 3.6 (6.8) 70+ 4.0 (10.7) 3.3 (5.2) 5.5 (11.7) 3.3 (6.4) Unknown Gender Female 6.2 (12.8) 9.7 (17.8) 6.3 (12.3) 7.1 (15.5) Male 5.1 (11.3) 6.9 (14.1) 8.9 (16.3) 5.5 (10.4) Other Household size 1 3.8 (13.7) 3.7 (11.1) 4.7 (12.1) 3.6 (8.2) 2 4.4 (11.2) 5.1 (11.0) 8.0 (16.8) 5.1 (10.9) 3-5 8.7 (15.6) 12.7 (19.0) 7.1 (11.9) 8.2 (15.0) 6+ 10.4 (14.2) 6.9 (6.2) 8.8 (11.5) 17.5 (22.2) Day of week Sun 3.3 (8.9) 6.3 (14.3) 7.6 (15.2) 4.0 (9.0) Mon 10.1 (16.8) 4.8 (5.9) 5.6 (9.0) 5.8 (11.5) Tue 5.6 (13.7) 5.5 (11.9) 17.2 (29.0) 5.9 (9.9) Wed 6.4 (14.1) 10.7 (18.7) 7.5 (13.5) 10.2 (16.7) Thu 7.1 (14.5) 10.6 (18.1) 7.2 (13.1) 5.0 (7.6) Fr 2.2 (2.6) 6.1 (13.4) 15.2 (22.7) 6.8 (15.1) Sat 2.5 (2.8) 3.9 (8.8) 9.0 (14.2) 7.5 (14.6) Face mask Yes 4.0 (6.7) 5.9 (14.8) 7.7 (18.9) 3.8 (4.7) No 5.4 (12.9) 5.5 (12.0) 8.2 (16.6) 5.5 (10.8) Unknown Vaccinated Yes 4.0 (6.7) 5.9 (14.8) 7.7 (18.9) 3.8 (4.7) 5.4 (12.9) 5.5 (10.8) No 5.5 (12.0) 8.2 (16.6) Unknown High risk Yes 4.5 (11.7) 4.1 (9.0) 7.6 (17.1) 4.9 (10.3) No 5.3 (12.3) 6.0 (12.9) 8.5 (16.8) 5.3 (9.9) Unknown Employed (Adults) Full time 5.6 (13.7) 6.7 (13.1) 10.4 (18.8) 5.9 (9.6) Part time 7.6 (16.1) 7.4 (13.7) 10.4 (19.3) 7.1 (13.6) Self employed 2.9 (3.7) 19.7 (29.4) 19.9 (33.2) 4.1 (5.4) Work open (Adults) No 3.2 (8.8) 9.5 (24.7) 8.4 (19.5) 5.6 (8.2) Yes 5.4 (12.9) 7.6 (15.6) 10.6 (19.8) 6.1 (11.1) Attended work (Adults) No 3.2 (5.3) 4.7 (7.2) 9.1 (17.8) 4.4 (7.0) Yes 7.8 (17.2) 9.3 (18.2) 13.1 (22.7) 7.0 (12.5)

459 Table 3: Mean contacts by characteristics.

- 461 Abbreviations
- 462 CI confidence interval
- 463 UK United Kingdom
- 464 CH Switzerland
- 465 BE Belgium
- 466 NL Netherlands
- 467 Declarations
- 468 Authors' contributions
- 469 WJE, and CIJ designed the CoMix contact survey. CIJ conceived of and planned the analysis.
- 470 CIJ, PC, JAB, JDM, PB, NH, CLA, JW, CF, and WJE provided comments and discussions on
- 471 analytical methods. CIJ, JDM, and PC conducted the analysis. CIJ wrote the first draft of the
- 472 manuscript with feedback from all other authors.
- 473
- 474 Acknowledgements
- 475 We acknowledge support from the European Centre for Disease Prevention and Control
- 476 (ECDC) in setting up the collaborations between the Epipose consortium, and universities and
- 477 public health institutions in all other countries. We gratefully acknowledge the tremendous
- 478 efforts put in all the steps by the EpiPose consortium, its collaborators and Ipsos.

- 480 We would also like to thank the team at Ipsos who have been excellent in running the survey,
- collecting the data, and allowing for this study to, happen at a rapid speed. We acknowledge the
- 482 exceptional project management support given by Sarah Vercruysse, Bieke Vanhoutte and
- 483 Anna Carnegie.

29

484 Ethics approval and consent to participate

485 Participation in this opt-in study was voluntary, and all analyses were carried out on anonymised 486 data. The study was approved in the UK by the ethics committee of the London School of 487 Hygiene & Tropical Medicine Reference number 21795. The study to collect CoMix data in 488 Belgium was approved by the Ethics Committee of UZA with reference 3236 - BUN 489 B3002020000054. The Medical Research Ethics Committee (MREC) NedMec confirmed that 490 the Medical Research Involving Human Subjects Act (WMO) does not apply to the CoMix study 491 in the Netherlands (research protocol number 22/917). Therefore an official approval of this 492 study by the MREC NedMec is not required under the WMO. The study to collect CoMix data in 493 Switzerland was approved by the ethics committee of the Canton of Bern (project number 2020-494 02926).

495 Consent for publication

496 Not applicable. We do not report individual patient data.

- 497 Availability of data and materials
- 498 The code and data used to conduct these analyses are found at
- 499 <u>https://github.com/jarvisc1/cmix_post_pandemic</u>
- 500 Competing interests
- 501 None

502 Funding

The EU have been the primary funder for this study. Over the course of the study CoMix received funding from: EU Horizon 2020 Research and Innovations Programme - project EpiPose (Epidemic Intelligence to Minimize COVID-19's Public Health, Societal and Economical Impact, No 101003688); Medical Research Council (MC_PC_19065); the NIHR (CV220-088 -COMIX); HPRU in Modelling & Health Economics (NIHR200908); and UKHSA. This work reflects only the authors' view. The European Commission is not responsible for any use that may be made of the information it contains The following funding sources are acknowledged as providing funding for the named authors: CIJ received funding from the LSHTM COVID-19 response fund. DFID/Wellcome Trust (Epidemic Preparedness Coronavirus research programme 210758/Z/18/Z: JDM); The views expressed in this publication are those of the author(s) and not necessarily those of the NIHR or the UK Department of Health and Social Care NIHR (PR-OD-1017-20002: WJE). UK MRC (MC_PC_19065: WJE) .

532 References

- 534 1. Wilson C. "Pandemics don"t end with a bang' lessons from the "Spanish" Flu. 2022.
- 535 https://www.rte.ie/news/primetime/2022/0125/1275848-spanish-flu-lessons-covid-19/. Accessed 536 24 Aug 2023.
- 537 2. Milne I. Influenza, war and revolution in Ireland, 1918–19. Manchester University Press;538 2018.
- 539 3. Osterholm MT. Pandemic preparedness after H1N1: Remember--If you've seen one
- 540 pandemic, you've seen one pandemic. CIDRAP. 2011. https://www.cidrap.umn.edu/business-
- 541 preparedness/pandemic-preparedness-after-h1n1-remember-if-youve-seen-one-pandemic-542 vouve. Accessed 24 Aug 2023.
- 4. Hoang T, Coletti P, Melegaro A, Wallinga J, Grijalva CG, Edmunds JW, et al. A Systematic
 Review of Social Contact Surveys to Inform Transmission Models of Close-contact Infections.
 Epidemiology. 2019;30:723–36.
- 546 5. Jarvis CI, Van Zandvoort K, Gimma A, Prem K, CMMID COVID-19 working group, Klepac P, 547 et al. Quantifying the impact of physical distance measures on the transmission of COVID-19 in 548 the UK. BMC Med. 2020;18:124.
- 6. Coletti P, Wambua J, Gimma A, Willem L, Vercruysse S, Vanhoutte B, et al. CoMix:
 comparing mixing patterns in the Belgian population during and after lockdown. Sci Rep.
 2020;10:21885.
- 552 7. Steens A, Freiesleben de Blasio B, Veneti L, Gimma A, Edmunds WJ, Van Zandvoort K, et al.
 553 Poor self-reported adherence to COVID-19-related quarantine/isolation requests, Norway, April
 554 to July 2020. Euro Surveill. 2020;25.
- 8. Gimma A, Munday JD, Wong KLM, Coletti P, van Zandvoort K, Prem K, et al. Changes in
 social contacts in England during the COVID-19 pandemic between March 2020 and March
 2021 as measured by the CoMix survey: A repeated cross-sectional study. PLoS Med.
 2022;19:e1003907.
- 9. Wong KLM, Gimma A, Coletti P, CoMix Europe Working Group, Faes C, Beutels P, et al.
 Social contact patterns during the COVID-19 pandemic in 21 European countries evidence
 from a two-year study. BMC Infect Dis. 2023;23:268.
- 10. Reichmuth ML, Heron L, Riou J, Moser A, Hauser A, Low N, et al. Socio-demographic
 characteristics associated with COVID-19 vaccination uptake in Switzerland: longitudinal
 analysis of the CoMix study. BMC Public Health. 2023;23:1523.
- 11. Backer JA, Bogaardt L, Beutels P, Coletti P, Edmunds WJ, Gimma A, et al. Dynamics of
 non-household contacts during the COVID-19 pandemic in 2020 and 2021 in the Netherlands.
 Sci Rep. 2023;13:5166.
- 568 12. Latsuzbaia A, Herold M, Bertemes J-P, Mossong J. Evolving social contact patterns during

- the COVID-19 crisis in Luxembourg. PLoS One. 2020;15:e0237128.
- 570 13. Liu CY, Berlin J, Kiti MC, Del Fava E, Grow A, Zagheni E, et al. Rapid Review of Social 571 Contact Patterns During the COVID-19 Pandemic. Epidemiology. 2021;32:781–91.
- 14. Trentini F, Manna A, Balbo N, Marziano V, Guzzetta G, O'Dell S, et al. Investigating the
 relationship between interventions, contact patterns, and SARS-CoV-2 transmissibility.
 Epidemics. 2022;40:100601.
- 575 15. Dorélien AM, Venkateswaran N, Deng J, Searle K, Enns E, Alarcon Espinoza G, et al.
 576 Quantifying social contact patterns in Minnesota during stay-at-home social distancing order.
- 577 BMC Infect Dis. 2023;23:324.
- 578 16. Mossong J, Hens N, Jit M, Beutels P, Auranen K, Mikolajczyk R, et al. Social contacts and 579 mixing patterns relevant to the spread of infectious diseases. PLoS Med. 2008;5:e74.
- 580 17. Wong KLM, Gimma A, Coletti P, Faes C, Beutels P, Hens N, et al. Social contact patterns
 581 during the COVID-19 pandemic in 21 European countries evidence from a two-year study.
 582 bioRxiv. 2022.
- 583 18. R Core Team. R: A Language and Environment for Statistical Computing. 2017.
- 19. Munday JD, Jarvis CI, Gimma A, Wong KLM, van Zandvoort K, CMMID COVID-19 Working
 Group, et al. Estimating the impact of reopening schools on the reproduction number of SARS-
- 586 CoV-2 in England, using weekly contact survey data. BMC Med. 2021;19:233.
- 587 20. Klepac P, Kucharski AJ, Conlan AJK, Kissler S, Tang ML, Fry H, et al. Contacts in context:
 588 large-scale setting-specific social mixing matrices from the BBC Pandemic project. bioRxiv.
 589 2020.
- 590 21. Prem K, Cook AR, Jit M. Projecting social contact matrices in 152 countries using contact 591 surveys and demographic data. PLoS Comput Biol. 2017;13:e1005697.
- 592 22. Diekmann O, Heesterbeek JAP, Roberts MG. The construction of next-generation matrices 593 for compartmental epidemic models. J R Soc Interface. 2010;7:873–85.
- 23. Wambua J, Loedy N, Jarvis CI, Wong KLM, Faes C, Grah R, et al. The influence of COVID-
- 595 19 risk perception and vaccination status on the number of social contacts across Europe: 596 insights from the CoMix study. BMC Public Health. 2023;23:1350.

597 Additional Files

period	R₀	2.5%	97.5%
Final CoMix round	0.526	0.494	0.557
Final CoMix round	0.682	0.631	0.741
Final CoMix round	0.813	0.749	0.881
Final CoMix round	0.583	0.543	0.627
1st Lockdown	0.234	0.225	0.244
1st Lockdown	0.186	0.171	0.203
	period Final CoMix round Final CoMix round Final CoMix round Final CoMix round 1st Lockdown 1st Lockdown	periodR₀Final CoMix round0.526Final CoMix round0.682Final CoMix round0.813Final CoMix round0.5831st Lockdown0.2341st Lockdown0.186	period R₀ 2.5% Final CoMix round 0.526 0.494 Final CoMix round 0.682 0.631 Final CoMix round 0.813 0.749 Final CoMix round 0.583 0.543 1st Lockdown 0.234 0.225 1st Lockdown 0.186 0.171

598 Table S1: Estimates of R0 from Figure 3B

Figure S1: Reduction in R₀ for Switzerland when using different baseline: Average of the
Polymod data ("CH (pmod) " bottom row) vs projected contact matrix from Prem et al [21] ("CH
(prem)", bottom row).

