1 SARS-CoV-2 quarantine mandated by contact tracing: burden and infection rate among close

2 contacts in Zurich, Switzerland, 2020-2021

Hélène E. Aschmann^{1,2}, Anja Domenghino^{2,3}, Ruedi Jung², Tala Ballouz², Dominik Menges², Jan Fehr²,
Milo A. Puhan²

- 5 1. Department of Pulmonary and Critical Care Medicine & Department of Epidemiology and
- Biostatistics, University of California San Francisco (UCSF), San Francisco General Hospital, 1001
 Potrero Avenue, San Francisco, CA 94110, San Francisco, USA
- 8 2. Epidemiology, Biostatistics and Prevention Institute (EBPI), University of Zurich (UZH),
- 9 Hirschengraben 84, CH-8001 Zurich, Switzerland
- 10 3. Department of Visceral and Transplantation Surgery, University Hospital Zurich, Rämistrasse
- 11 100, CH-8091 Zurich, Switzerland
- Correspondence: Hélène E. Aschmann (<u>helene.aschmann@ucsf.edu</u>)
 Division of Pulmonary and Critical Care Medicine
 San Francisco General Hospital
 1001 Potrero Avenue
 San Francisco, CA 94110

17 Abstract

- 18 Objectives
- 19 Before vaccines and effective treatments were available, quarantine of close contacts was important
- 20 to limit the spread of SARS-CoV-2. To evaluate potential benefits and harms of quarantine, we aimed
- 21 to estimate infection rates and describe experiences and mental health among persons in mandated
- 22 quarantine during the early SARS-CoV-2 pandemic.
- 23 Methods
- 24 We invited adults in mandated quarantine after an exposure to SARS-CoV-2 identified through
- contact tracing of the Canton of Zurich, Switzerland, between August 2020 and January 2021.
- 26 Participants completed two questionnaires and received up to two SARS-CoV-2 polymerase chain
- 27 reaction tests, during and at the end of quarantine.
- 28 Results

29 Among 395 participants, quarantine duration ranged from 2 to 20 days. By day 11 since the last

30 contact, 11.1% [95% CI 8.4%–14.7%] were infected with SARS-CoV-2. The proportion of participants

31 with symptoms of depression doubled from 9.3% before quarantine to 18.9% during quarantine, and

32 12.1% reported quarantine was very or extremely difficult.

33 Conclusions

Although quarantine was only moderately burdensome for most participants, some experienced
 significant difficulties and burden. Policymakers need to balance infection control with potential
 harms placed on individuals.

37 Introduction

38 Authorities mandating quarantine of close contacts of persons infected with SARS-CoV-2 need to 39 balance quarantine's efficacy in limiting viral spread against its impact on individuals, society, and 40 economy. The balance of quarantine's benefits and harms should not only be counted with mortality 41 and morbidity, but should also consider protection from financial, social, and psychological harm on 42 quarantined persons [1]. Quarantine was especially important at the beginning of the SARS-CoV-2 43 pandemic when vaccines and effective medical treatment were not available. Currently, only few 44 countries still impose quarantine and in Switzerland, quarantine mandates were lifted in April 2022. 45 But with new variants and waning vaccine effectiveness, quarantine remains an option according to the World Health Organization [2]. 46

Quarantine is associated with poor psychological outcomes, in particular after a duration of one
week or longer [3,4]. Individuals in quarantine or isolation have 2-3 times increased odds for
depressive, anxiety or stress-related disorders [4]. Furthermore, low adherence (42% in the UK [5],
28% in Norway [6]) limited its efficacy. While at the beginning of the pandemic, some called for
quarantine as long as 21 days based on incubation time [7–9], testing soon allowed to reduce
quarantine duration [10–15]. Later some persons were exempted from quarantine due to prior
infections or vaccination [2].

To evaluate or inform policy makers, a detailed evaluation of the benefits and harms of quarantine measures is warranted. Our aims were (1) to estimate infection rates among close contacts in mandated quarantine in the Canton of Zurich, Switzerland, during the early SARS-CoV-2 pandemic (i.e., the secondary attack rate), and (2) to assess mental health, difficulties and worries during quarantine as well as adherence, and motivation to adhere to quarantine.

59 Methods

60 Study Population

This analysis is based on the Zurich SARS-CoV-2 Cohort, a prospective, observational, populationbased study of an age-stratified random sample of 1106 SARS-CoV-2 positive individuals and their close contacts based on contact tracing in the Canton of Zurich, Switzerland. All participants provided electronic consent. The study was prospectively registered (ISRCTN14990068) and approved by the ethics committee of the Canton of Zurich (BASEC 2020-01739).

More detailed information about the study enrolment procedures are reported elsewhere [16]. In
short, close contacts were identified through contact tracing and randomly sampled in clusters based
on their index case (the person with a positive SARS-CoV-2 test in the cohort with whom a potentially
infectious contact occurred). We recruited close contacts from August 7, 2020, to January 15, 2021.
Eligible persons included adults (aged ≥18), fluent in German, resident in the canton of Zurich, and
consenting to participate. Persons were not eligible if they declined to be re-contacted for study
purposes or if the index case was not recorded (e.g., the contact occurred in another country).

73 Data collection

Close contacts were invited to complete a baseline questionnaire upon enrolment. A second questionnaire was sent on the second to last day of quarantine. Moreover, they were invited to receive a polymerase chain reaction (PCR) test for SARS-CoV-2 at the beginning of quarantine (preferably on day 5 or 6) and at the end of quarantine. Close contacts testing positive were invited

to participate in a separate arm of the study instead. We reported mental health outcomes in
persons who tested positive separately [17].

The questionnaires included the German version of short form of the Depression, Anxiety, and Stress Scale (DASS-21) [18], and a subset of questions from the COVID-19 Pandemic Mental Health Questionnaire [19]. We asked how well informed and prepared participants felt, how difficult quarantine measures were in general to follow, and how difficult the specific rules published by the Swiss Federal Office of Public Health were [20]. Potential worries, reasons for motivation and positive experiences were piloted with lay persons and we added open text comments to allow participants to add others.

To compare participants with non-participants, aggregated data were analysed based on data from
contact tracing. These data were recorded in Microsoft Office Excel forms and in the Surveillance
Outbreak Response Management and Analysis System (SORMAS) [21] a web application developed
for contact tracing.

91 Sample size estimation

The sample size calculation was based on the proportion of close contacts with a positive PCR test after day 5 of quarantine. We estimated 5% would test positive until day 5, and 3% with an initial negative test would test positive by the end of quarantine. To show a difference of 1% at a significance level of 0.05, a sample size of 294 was needed. However, the sample size was increased because not all participants agreed to get tested. Recruitment was stopped when the targeted sample size was reached for SARS-CoV-2 positive individuals of our cohort study, as the study had several aims.

99 Statistical analysis

We modelled the cumulative probability over time of converting during the mandated quarantine for
 all invited close contacts. This cumulative probability was derived by modelling the proportion of
 persons who remained without a positive test using a survival analysis with the tram package

103 (version 0.6-4) in R [22]. For each positive test, we assumed the conversion to occur earliest on the 104 day after the last negative test or, in the absence of a test, on the day of last contact, and latest on 105 the day the sample for the positive test was taken (interval censoring) [23,24]. For close contacts 106 without a positive test, the observations were censored after the last negative test. Because our 107 recruitment in the close contact sub-study did not include contacts who converted before they 108 consented to participate, we also included data from non-participants in the model, to account for 109 contacts who converted early. For non-participants, we obtained the sampling day of positive tests 110 from contact tracing data. We assumed a similar frequency of negative tests as for participants, 111 except for the tests on day 10 and 11 of quarantine. Since testing at the end of quarantine was a 112 study-specific procedure, and not mandated by the Department of Health, we assumed non-113 participants had only as many negative tests on day 10 and 11 as participants on day 9. 114 Descriptive analyses were performed for all other outcomes. For DASS-21, missing data were 115 imputed for each subscale if two or less answers were missing using the mean of the remaining 116 answers. We explored associations of (1) overall motivation, (2) feeling prepared, (3) difficulty of 117 quarantine measures overall, and (4) having more time to relax during quarantine with baseline 118 characteristics based on a priori hypotheses and hypotheses generated based on participants' 119 comments. We included age and sex in all regressions. All outcomes were elicited as five-point Likert-120 type scales, and we used logistic ordinal regression to analyse associations [25]. 121 Informed consent, recruitment and survey data were collected and managed using REDCap 122 electronic data capture tools [26,27]. For participants who preferred not to complete the survey 123 online, written consent was sought via letters and questionnaires were completed via phone. All

analyses were performed using R version 4.1.1 [28].

125 Results

126 Study Population

127 Of 21 316 records of close contacts, 10 110 were considered eligible during the initial screening

128 (Figure 1). We invited 1484 persons (and 8 duplicates). Of these, 129 were not eligible, and 526 had

129 finished quarantine or converted to a positive case. Of the remaining 829 persons, 395 consented to

130 participate, corresponding to a response rate of 48%.

131 We compared baseline characteristics between 395 participants and 1018 non-participants. This

analysis did not consider 79 persons who should not have been invited (8 duplicates, 56 underage, 14

133 not close contacts, 1 already converted before we invited them). Participants and non-participants

134 were similar in sex and duration of quarantine (Table S1). Age was missing for 36% of non-

participants, so a direct comparison is difficult. However, participants were likely younger than non-

136 participants because their index cases (persons with a positive SARS-CoV-2 test) were younger (Table

137 S1), and the age of index cases and close contacts was correlated (Table S2).

138 Almost all participants (94%) performed the mandated quarantine at home (Table 1). Some

participants (22%) had children in their household. Many were household contacts, meaning they

140 were living in the same household as an individual with a SARS-CoV-2 infection (35%). Most

141 participants were employed (70%). Of those who worked, most could work at home (47%) or at least

142 partially (24%). Some self-quarantined before receiving the official mandate because they knew of a

143 contact with an infected person (38%). Among household contacts, 47% (64/135) self-quarantined,

144 while 37% (50/135) answered that their reason to initiate quarantine was the official mandate.

145 PCR test results during quarantine

At least one PCR test was performed in 358 of 395 participants during their quarantine. Our analysis of the secondary attack rate considered a total of 1404 invited adult close contacts in whom we could have noted a conversion (Figures S1 and S2). In a survival analysis that considers when persons were tested (positive and negative), the estimated cumulative probability for a conversion by day 11

- 150 was 11.1% [95% CI: 8.4–14.7%] (Figure 2A). Most conversions occurred within 5 days after the last
- 151 potentially infectious contact: the estimated cumulative probability reached 9.1% [95% CI: 6.6–
- 152 12.7%] by day 5. The mean follow-up time up to the last test result was 7.4 days.

153 Quarantine duration

In theory, household contacts should always have had a quarantine duration of at least 10 or 11 days
(quarantine duration was decreased from 11 to 10 days in November 2020). However, 24 of 135
household contacts (17.8%) reported durations of 5 to 9 days (Figure 2B). A regular duration of 10 or
11 days was reported by 89 (65.9%) and an extended duration by 22 (16.3%) household contacts.
Non-household contacts declared shorter durations of quarantine, with a median of 8 days and
ranging from 2 to 20 days. Among all contacts, quarantine was extended in 29 (7.4%) persons; this
occurred when persons spent their quarantine together and one of them converted during that time.

161 Motivation to adhere to quarantine measures

162 Most participants reported that they were motivated to adhere to quarantine measures overall: 338

163 (86.7%) answered that they were motivated, very motivated, or extremely motivated (Figure 3).

164 Participants were most motivated to get tested when symptoms appeared (360, 92.3%). However,

less participants were motivated not to leave their home for 10 days (275, 70.5%), not to meet family

and friends (269, 69.0%), to follow hygiene rules (220, 56.4% among those not living alone), and to

167 stay at home longer if necessary (198, 50.8%) (e.g., if they or others in their household developed

168 symptoms).

The most important motivations to stay in quarantine were to protect family and friends, protect the community at large, and contribute to fighting the pandemic, followed by the official mandate from authorities (Figure 3). Less participants considered others knowing they should be in quarantine, others being afraid to contract the virus, or possible punishments as important reasons.

173 Worries during quarantine

Participants worried most that their loved ones could contract the coronavirus: 289 (74.1%) were
worried, very, or extremely worried (Figure 3). Most worried about the health of the person infected
with SARS-CoV-2 (i.e., their index case) (63.8%) and becoming inactive (61.8%). Many worried about
contracting the virus themselves (43.1%). Some participants worried about becoming depressed
(22.3%).

Worries were associated with the overall motivation (Table 2). Participants who worried about
contracting the virus themselves had 2.08 [95% CI: 1.41–3.09] times higher odds of reporting higher
levels of overall motivation. By contrast, worries about becoming depressed were associated with
lower motivation (0.45 [95% CI: 0.28–0.72] times smaller odds for higher levels of motivation).

183 Difficulty of quarantine measures

184 Quarantine measures were perceived as difficult or very difficult by 84 participants (21.5%) during 185 quarantine (Figure S3) and by 65 participants (17.5%) at the end of quarantine (Figure S4). In total, 186 109 participants (27.9%) found quarantine measures difficult or very difficult at either time point. 187 Difficulty overall was lower for older adults (odds ratio 0.84 [95% CI: 0.75–0.95] for each additional 10 years), and lower for male participants (odds ratio 0.50 [95% CI: 0.34–0.72]) (Table 2). Although 188 189 several participants commented on the difficulty of taking care of children and the infected person, 190 we could not confirm with statistical significance that difficulty was higher for participants living with 191 children (odds ratio 1.49 [95% CI: 0.96–2.31]), or for household contacts (odds ratio 1.39 [95% CI: 192 0.95–2.04]). There was no association between the overall difficulty and work from home during 193 quarantine (odds ratio 1.00 [95% CI: 0.67–1.47]).

194 Symptoms of depression, anxiety, and stress

195 370 participants answered the DASS-21 questionnaire both at baseline and at the end of quarantine.

196 The proportion of participants with depressive symptoms (of any severity) doubled from 9.3% before

197 quarantine to 18.9% during quarantine (Figure 4). A sensitivity analysis with all participants showed

198 similar results (Figure S5). Overall, 65 participants (17.6%) had a relevant worsening by at least one 199 category in depression, anxiety, or stress (Figure S6). However, 27 participants (7.3%) improved by at 200 least one category in at least one subscale. Depression and stress scores increased with statistical 201 significance (depression: +1.70 [95% CI: 1.19–2.22], stress +1.06 [95% CI: 0.47–1.66]) (Figure S6). The 202 change in depression scores was higher in men than in women and higher in those with higher 203 baseline scores, i.e., more pronounced in those with pre-existing depressive symptoms (Table 2). 204 In direct comparisons, some participants reported feeling more isolated (22.0%) and increased 205 trouble sleeping (14.0%) during quarantine compared to two weeks prior (Figure S7). Feeling more 206 impatient or angry, consuming more alcohol, having more nightmares, feeling more worried, 207 nervous, or depressed was reported by 8.3% to 11.0%.

208 Positive experiences

Most participants reported more time to relax (253, 68.0%). Some commented that quarantine was stressful because they took care of children or of the SARS-CoV-2 positive person. An ordinal logistic regression confirmed that persons living with children (odds ratio 0.59 [95% CI: 0.37–0.92]) and persons working from home (odds ratio 0.66 [95% CI: 0.44–0.98]) had lower odds of reporting more time to relax.

214 Information and preparation for quarantine

Most participants felt well or very well prepared for quarantine (47.2%) or neutral (40.3%). Older adults felt better prepared (odds ratio 1.19 [95% CI: 1.05–1.34] per additional 10 years), as did participants who felt well or very well informed (odds ratio 5.27 [95% CI: 3.03–9.32]). Participants worried about becoming depressed felt less well prepared (odds ratio 0.37 [95% CI: 0.23–0.58]).

219 Adherence with quarantine

220 By the end of quarantine, 74 participants (19.9%) had left their house or met people during

- 221 quarantine. One participant decided to stop quarantine early after a negative test based on the
- 222 mistaken belief that a negative test ends quarantine.

We also asked household contacts about their adherence to the recommendations, in particular
interacting with the person infected with SARS-CoV-2 (Figure S7). The recommendation that was
least followed was wearing a mask when entering this person's room – 20% rarely or never followed
this rule.

Several participants expressed a need for physical activity outdoors and took precautions. One
person described: "I badly needed a 20-minute walk or run. I went once early in the morning and
once in the evening, so I virtually met nobody [...]. I always had a mask with me. This should be
allowed in my opinion." Others answered that quarantine rules did not consider the need to walk the
dog or helping people with disabilities.

232 Financial hardship

In total, 62 of 390 participants worried about financial consequences: they were worried about losing
their job or getting into financial difficulties or expected income loss due to quarantine (with no or
only partial compensation). Only 14% (53 of 390) expected a reduced income; among them, 20
expected a partial, 25 no compensation. Twenty-two persons (6%) were worried, very, or extremely
worried about getting into financial difficulties due to quarantine, and 39 persons (10%) were a little
worried. 26 (7%) were a little worried about losing their job due to quarantine, and 12 persons were
worried, very, or extremely worried.

240 Burden of quarantine

In total, 176 of 390 participants (45.1%) found quarantine difficult, expected negative financial
consequences or had symptoms of decreased mental health. There was little overlap between the 65
participants who worsened significantly in a DASS-21 dimension and the 62 participants who
expected negative financial consequences; 117 of 390 participants either had worsened mental
health or expected financial consequences (29.2%). Finally, 109 of 390 participants felt that
quarantine measures were difficult or very difficult.

247 Discussion

This study puts infection rates in close contacts (i.e., the secondary attack rate) into context with the negative and positive experiences of persons in quarantine. We found that about 11% of contacts were infected, and the majority (9%) had already converted by day 5 after the last (potentially infectious) contact. While some persons had positive experiences (e.g., more time to relax was reported by 68%), this stands in contrast to significant difficulties and burden experienced by almost half of the close contacts, including decreased mental health and financial consequences. Our results compare to published secondary attack rates in close contacts, although these varied

widely, partly due to different definitions of close contacts [29–34]. Even in household contacts,
estimates ranged from 5% to 48% [29–34]. Highest estimates come from disadvantaged populations
[34], reflecting that our population had comparatively good resources to protect themselves from
transmission within a household.

259 Our study found a comparable increase in the risk for depressive symptoms as studies from earlier 260 epidemics, which reported a 2.0-2.8 fold increase [4]. Prior studies also reported a higher risk among 261 those with a history of depressive symptoms or mental illness, but did not report that the increase 262 was higher in men [4]. Although others found associations of the duration of quarantine with 263 negative psychological effects [3,36], we did not find a similar effect. A study in Germany found 264 higher psychological distress among infected persons in isolation than in close contacts [37]. In 265 comparison, our cohort study found similar rates of depressive symptoms for both groups [17]. 266 However, in our study close contacts had higher rates of stress but lower rates of anxiety [17]. 267 Although close contacts reported high motivation, adherence was moderate and comparable to 268 other studies [1,5,6,38]. About 20% left their house, e.g., for groceries, or met persons from other 269 households, and only half the household contacts reported voluntary self-quarantine before

270 receiving the official mandate. In comparison, in the index case study arm of our cohort, 14% did not

comply with measures before their positive result, and 3% after their positive result [17]. This is in

line with other studies where adherence was lower in persons without symptoms [38]. We found
that contacts who worried less about contracting the virus were less motivated to adhere. Contact
tracers should therefore carefully explain why the exposure was potentially infectious.

Quarantine should be limited to the shortest possible but necessary duration. A 10-11 day quarantine was sufficient in our population, in a setting with high testing resources. Our study also adds to other evidence that supported a shortened 7-day quarantine with a negative test at the end of quarantine, as introduced soon after our study concluded [12–14,39]. Public health authorities should provide better information on testing. For example, 7% of our participants experienced long quarantine durations of 12 to 20 days due to others in their household testing positive. These

Future interventions should aim to reduce the burden of quarantine for parents and persons with depressive symptoms [3]. Participants in our survey expressed a strong desire for outdoor physical activity, walking their dog, and letting children go outside. Experiences among parents are in line with our findings in infected persons in isolation, where parents had higher difficulty with isolation measures than persons without children [17].

extensions could have been reduced if their household members had been tested earlier.

281

287 A limitation of our study is that we only included close contacts of adult index cases, although 288 sometimes children in the same household were also positive. Parents likely experienced more 289 difficult guarantines, so we may have underestimated difficulties. Furthermore, we may have 290 overestimated motivation and adherence as more motivated persons may have been more likely to 291 consent to participate in the study. Additionally, we did not measure long term effects on mental 292 health, although a prior study reported sustained negative effects 6 months after quarantine [3]. 293 Finally, infection rates among close contacts alone are not sufficient to measure quarantine's 294 effectiveness, which also depends on epidemic severity, timeliness of contact tracing, duration and 295 adherence with guarantine [40]. Delays in contact tracing and many undetected or not traced COVID-

- 19 cases during the Fall 2020 surge likely limited quarantine's effectiveness, but we did not aim toestimate averted infections.
- 298 To inform future pandemic response and evaluate past public health interventions, it is important to
- 299 compare the effectiveness with the burden placed on quarantined persons. Our analysis exemplifies
- 300 that this comparison should ideally be performed in the same population, as many factors, such as
- 301 mental health, motivation, and adherence, influence each other. Going forward, policy makers
- 302 should critically assess the need for quarantine and strategically study and reduce negative impact on
- 303 quarantined persons.

304 Acknowledgments

305 We thank Prof. Torsten Hothorn for his support with the survival analysis.

306 Potential conflicts of interest

307 The authors declare no conflicts of interest.

308 Patient consent statement

- 309 All participants provided electronic consent. The study was prospectively registered
- 310 (ISRCTN14990068) and approved by the ethics committee of the Canton of Zurich (BASEC 2020-

311 01739).

312 Funding

313 The ZSAC study was funded through the Health Directorate of the Canton of Zurich and the Pandemic 314 Fund of the University of Zurich. ZSAC was also part of the Corona Immunitas research program, 315 coordinated by the Swiss School of Public Health (SSPH+) and funded through SSPH+ fundraising, 316 including funding by the Swiss Federal Office of Public Health, the Cantons of Switzerland (Basel, 317 Vaud, and Zurich), private funders (ethical guidelines for funding stated by SSPH+ were respected) 318 and institutional funds of the participating universities. HEA was also supported by a Swiss National 319 Science Foundation Early Postdoc. Mobility Fellowship [191414] and a Postdoc. Mobility Fellowship 320 [214129]. TB received funding from the European Union's Horizon 2020 research and innovation 321 program under the Marie Skłodowska-Curie grant agreement No 801076, through the SSPH+ Global 322 PhD Fellowship Program in Public Health Sciences (GlobalP3HS) of the Swiss School of Public Health. 323 Study funders had no role in the study design, data collection, analysis, interpretation, or writing of 324 this report.

325

326 References

- 327 1. Sopory P, Novak JM, Noyes JP. Quarantine acceptance and adherence: qualitative evidence
 328 synthesis and conceptual framework. J Public Heal. Journal of Public Health; 2021;
- 329 2. World Health Organization. Contact tracing and quarantine in the context of the Omicron SARS-
- CoV-2 variant. 2022;1–9. Available from: https://www.who.int/publications/i/item/WHO-2019-nCoV Contact_tracing_and_quarantine-2022.1
- 332 3. Brooks SK, Webster RK, Smith LE, Woodland L, Wessely S, Greenberg N, et al. The psychological 333 impact of quarantine and how to reduce it: rapid review of the evidence. Lancet. 2020;395:912–20.
- 4. Henssler J, Stock F, van Bohemen J, Walter H, Heinz A, Brandt L. Mental health effects of infection
 containment strategies: quarantine and isolation—a systematic review and meta-analysis. Eur Arch
 Psychiatry Clin Neurosci. 2021;271:223–34.
- 5. Smith LE, Potts HWW, Amlôt R, Fear NT, Michie S, Rubin GJ. Adherence to the test, trace, and
 isolate system in the UK: Results from 37 nationally representative surveys. BMJ. 2021;372.
- 6. Steens A, De Blasio BF, Veneti L, Gimma A, Edmunds WJ, Van Zandvoort K, et al. Poor self-reported
 adherence to COVID-19-related quarantine/isolation requests, Norway, April to July 2020.
 Eurosurveillance. 2020;25:1–6.
- 7. Li ZY, Zhang Y, Peng LQ, Gao RR, Jing JR, Wang J Le, et al. Demand for longer quarantine period
 among common and uncommon COVID-19 infections: a scoping review. Infect Dis Poverty.
 2021;10:5–13.
- 8. Khadem Charvadeh Y, Yi GY, Bian Y, He W. Is 14-Days a Sensible Quarantine Length for COVID-19?
 Examinations of Some Associated Issues with a Case Study of COVID-19 Incubation Times. Stat Biosci.
 2022;14:175–90.
- 9. Wang R, Wang Q. Determination and estimation of optimal quarantine duration for infectious
 diseases with application to data analysis of COVID-19. Biometrics. 2021;1–10.
- 10. Wells CR, Townsend JP, Pandey A, Moghadas SM, Krieger G, Singer B, et al. Optimal COVID-19
 quarantine and testing strategies. Nat Commun. Springer US; 2021;12:1–9.
- 352 11. Reid MJA, Prado P, Brosnan H, Ernst A, Spindler H, Celentano J, et al. Assessing Testing Strategies
- and Duration of Quarantine in Contact Tracing for SARS-CoV-2: A Retrospective Study of San
- Francisco's COVID-19 Contact Tracing Program, June-August 2020. Open Forum Infect Dis. 2021;8:2–
 5.
- 12. Quilty BJ, Clifford S, Hellewell J, Russell TW, Kucharski AJ, Flasche S, et al. Quarantine and testing
 strategies in contact tracing for SARS-CoV-2: a modelling study. Lancet Public Heal. 2021;6:e175–83.
- Jones A, Fialkowski V, Prinzing L, Trites J, Kelso P, Levine M. Assessment of Day-7 Postexposure
 Testing of Asymptomatic Contacts of COVID-19 Patients to Evaluate Early Release from Quarantine —
 Vermont, May–November 2020. MMWR Morb Mortal Wkly Rep. 2021;70:12–3.
- 14. Ashcroft P, Lehtinen S, Angst DC, Low N, Bonhoeffer S. Quantifying the impact of quarantine
 duration on covid-19 transmission. Elife. 2021;10:1–33.
- 15. Peng B, Zhou W, Pettit RW, Yu P, Matos PG, Greninger AL, et al. Reducing COVID-19 quarantine
 with SARS-CoV-2 testing: A simulation study. BMJ Open. 2021;11:1–13.
- 365 16. Ballouz T, Menges D, Aschmann HE, Jung R, Domenghino A, Fehr JS, et al. Individual-Level
 366 Evaluation of the Exposure Notification Cascade in the SwissCovid Digital Proximity Tracing App:

- 367 Observational Study. JMIR Public Heal Surveill. 2022;8:1–14.
- 368 17. Domenghino A, Aschmann HE, Ballouz T, Menges D, Strebel D, Derfler S, et al. Mental health of
- 369 individuals infected with SARS-CoV-2 during mandated isolation and compliance with
- 370 recommendations A population-based cohort study. 2022;1–20.
- 18. Nilges P, Essau C. Die Depressions-Angst-Stress-Skalen: Der DASS ein Screeningverfahren nicht
 nur für Schmerzpatienten. Obere Extrem. 2015;10:649–57.
- 19. Rek S V., Bühner M, Reinhard MA, Freeman D, Keeser D, Adorjan K, et al. The COVID-19 Pandemic
- 374 Mental Health Questionnaire (CoPaQ): psychometric evaluation and compliance with
- countermeasures in psychiatric inpatients and non-clinical individuals. BMC Psychiatry. 2021;21:1–
 15.
- 20. Federal Office of Public Health FOPH. COVID-19: Instructions on quarantine. 2021.
- 21. Perscheid C, Benzler J, Hermann C, Janke M, Moyer D, Laedtke T, et al. Ebola outbreak
- 379 containment: Real-time task and resource coordination with SORMAS. Front ICT. Frontiers Media380 S.A.; 2018;5.
- 381 22. Hothorn T. Most likely transformations: The mlt package. J Stat Softw. 2020;92.
- 382 23. Therneau T, Grambsch PM. Modeling Survival Data: Extending the Cox Model. New York:383 Springer; 2000.
- 24. Therneau T. A Package for Survival Analysis in R. R Packag. version 3.2-13. 2021. Available from:
 https://cran.r-project.org/package=survival
- 25. Venables WN, Ripley BD. Modern Applied Statistics with S. Fourth edition. New York: Springer;
 2002. Available from: https://www.stats.ox.ac.uk/pub/MASS4/
- 26. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture
 (REDCap)-A metadata-driven methodology and workflow process for providing translational research
- informatics support. J Biomed Inform. 2009;42:377–81.
- 391 27. Harris PA, Taylor R, Minor BL, Elliott V, Fernandez M, O'Neal L, et al. The REDCap consortium:
- Building an international community of software platform partners. J Biomed Inform.
- 393 2019;95:103208.
- 28. R Core Team. R: a language and environment for statistical computing. R Foundation for
 Statistical Computing. Vienna, Austria; 2020. p. https://www.R-project.org.
- 29. Hu S, Wang W, Wang Y, Litvinova M, Luo K, Ren L, et al. Infectivity, susceptibility, and risk factors
 associated with SARS-CoV-2 transmission under intensive contact tracing in Hunan, China. Nat
 Commun. 2021;12:1–11.
- 30. Proesmans K, Hancart S, Braeye T, Klamer S, Robesyn E, Djiena A, et al. COVID-19 contact tracing
 in Belgium: main indicators and performance, January September 2021. Arch Public Heal. BioMed
 Central; 2022;80:1–10.
- 402 31. Cheng HY, Jian SW, Liu DP, Ng TC, Huang WT, Lin HH. Contact Tracing Assessment of COVID-19
- 403 Transmission Dynamics in Taiwan and Risk at Different Exposure Periods before and after Symptom
 404 Onset. JAMA Intern Med. 2020;180:1156–63.
- 405 32. Tsang TK, Fang LQ, Zhang A, Jiang FC, Ruan SM, Liu LZ, et al. Variability in transmission risk of
- SARS-CoV-2 in close contact settings: A contact tracing study in Shandong Province, China. Epidemics.
 2022;39:100553.
- 408 33. Ma X, Wu K, Li Y, Li S, Cao L, Xie H, et al. Contact tracing period and epidemiological

- 409 characteristics of an outbreak of the SARS-CoV-2 Delta variant in Guangzhou. Int J Infect Dis.
 410 2022;117:18–23.
- 411 34. Vallès X, Roure S, Valerio L, López-Muñoz I, Pérez-Quílez O, Soldevila L, et al. SARS-CoV-2 contact
- tracing among disadvantaged populations during epidemic intervals should be a priority strategy:
- 413 results from a pilot experiment in Barcelona. Public Health. 2021;195:132–4.
- 35. Jassim G, Jameel M, Brennan E, Yusuf M, Hasan N, Alwatani Y. Psychological Impact of COVID-19,
 Isolation and Quarantine: A Comparative Study. SSRN Electron J. 2021;1413–21.
- 416 36. Kwon H-Y, Kim Y, Lee S-Y. What Matters for Depression and Anxiety During the COVID-19
- 417 Quarantine?: Results of an Online Cross-Sectional Survey in Seoul, South Korea. Front Psychiatry.
 418 2022;13:1–9.
- 419 37. Klee L, Fabrice A, Eisenburger N, Feddern S, Gabriel C, Kossow A, et al. Coping strategies during
- 420 legally enforced quarantine and their association to psychological distress level: a cross-sectional
 421 study. Public Health. 2022;209:52–60.
- 422 38. Eraso Y, Hills S. Self-isolation and quarantine during the UK's first wave of covid-19. A mixed-423 methods study of non-adherence. Int J Environ Res Public Health. 2021;18.
- 39. Kolesar JM, Gayheart T, Poston L, Monday E, Forster D, Belcher E, et al. Preliminary Research on a
 COVID-19 Test Strategy to Guide Quarantine Interval in University Students. Covid. 2022;2:254–60.
- 426 40. Craig KJT, Rizvi R, Willis VC, Kassler WJ, Jackson GP. Effectiveness of contact tracing for viral
- 427 disease mitigation and suppression: Evidence-based review. JMIR Public Heal Surveill. 2021;7.
- 428

430 Tables

		N of 390	(%)
Age	18-39 years	196	(50)
-	40-64 years	156	(40)
	65+ years	38	(10)
Sex	Female	195	(50)
Education	No school diploma	1	(0)
	Mandatory school	12	(3)
	Vocational education	98	(25)
	Upper or post-secondary education	41	(11)
	Tertiary education other than university	110	(28)
	University bachelor or higher	126	(32)
Job situation	Employed	273	(70)
	Self-employed	24	(6)
	In education	47	(12)
	Retired	32	(8)
	Not working / unemployed / housewife	12	(3)
Knows who the index case is	Yes	364	(93)
	Strong suspicion	17	(4)
	No	7	(2)
Type of contact	Household contact	135	(35)
	Workplace contact	49	(13)
	Other	197	(51)
Quarantine place	At home	371	(94)
	At someone else's home	9	(2)
	Vacation home	9	(2)
	Hotel	1	(0)
	Hospital	1	(0)
	Other	1	(0)
Reason to initiate quarantine	Mandated through Department of Health	163	(42)
	Self-quarantined due to known contact	161	(38)
	Asked by employer	28	(7)
	Asked by health care provider	10	(2)
	Recommended by close ones	15	(4)
	SwissCovid App warning	6	(2)
	Always stayed at home	1	(0)
Household	Living alone	51	(13)
	Living with one other adult	121	(31)
	Living with several adults, no children	131	(34)
	Living with children	87	(22)
Care giving of index case	(among household contacts)	N of 135	
	Yes	91	(67)
Possibility to work from home	(among those working or in education)	N of 342	I.
	Possible	162	(47)
	Partially	82	(24)
	Not possible	76	(22)
	On sick leave	8	(2)
	Vacation or semester break	6	(2)

Table 1: Baseline characteristics of participants and information relevant to their quarantine

- 433 **Table 2:** Associations of baseline participant characteristics with motivation, feeling prepared,
- 434 difficulty with quarantine measures, time to relax (all ordinal logistic regressions where the outcome
- 435 was a five-point Likert-type scale), and change in depression score (linear regression). *Statistically
- 436 significant at 0.05 significance level, not corrected for multiple testing.

Outcome: Higher overall motivation to comply with quarantine measures	OR [95% CI]		
Age (per 10 years, reference age 18)	1.01 [0.90–1.14]		
Sex (male vs female)	0.96 [0.66–1.40]		
Worried about getting sick with COVID-19 (own health) (worried, very	2.08 [1.41–3.09]*		
worried, or extremely worried vs little or not at all worried)			
Worried about becoming depressed (worried, very worried, or	0.45 [0.28–0.72]*		
extremely worried vs little or not at all worried)			
Outcome: Feeling better prepared			
Age (per 10 years, reference age 18)	1.19 [1.05–1.34]*		
Sex (male vs female)	0.91 [0.62–1.33]		
Well or very well informed (vs neutral, poor, or very poor)	5.27 [3.03–9.32]*		
Worried about becoming depressed (worried, very worried, or	0.37 [0.23–0.58]*		
extremely worried vs little or not at all worried)			
Worried about getting sick with COVID-19 (own health) (worried, very	0.96 [0.65–1.42]		
worried, or extremely worried vs little or not at all worried)			
Outcome: Overall difficulty with quarantine measures			
Age (per 10 years, reference age 18)	0.84 [0.75–0.95]*		
Sex (male vs female)	0.50 [0.34–0.72]*		
Working (possible or partially possible vs not possible)	1.00 [0.67–1.47]		
Household contact (vs index case not in the same household)	1.39 [0.95–2.04]		
Children in household (yes vs no)	1.49 [0.96–2.31]		
Outcome: More time to relax			
Age (per 10 years, reference age 18)	0.98 [0.87–1.10]		
Sex (male vs female)	0.92 [0.63–1.34]		
Working (possible or partially possible vs not possible)	0.66 [0.44–0.98]*		
Taking care of index case (yes vs no or not applicable)	1.08 [0.69–1.69]		
Children in household (yes vs no)	0.59 [0.37–0.92]*		
Outcome: Change in depression score (during quarantine minus before)			
Age (per 10 years, reference age 18)	1.02 [0.99–1.06]		
Sex (male vs female)	4.68 [1.67–13.12]*		
Quarantine duration (per additional day)	1.01 [0.81–1.27]		
Living alone (vs living with others)	0.91 [0.20–4.23]		
Children in household (yes vs no)	2.64 [0.75–9.31]		
Working (possible or partially possible vs not possible)	1.19 [0.38–3.64]		
Baseline depression score (per additional point)	1.23 [1.10–1.36]*		

437

439 Figure Legends

- 440 **Figure 1**: Participant flow through recruitment. Recruitment was performed in two stages. Potential
- 441 participants had to complete an online form to be contacted by the study team or give oral consent
- to have the online form completed for them.
- 443 Figure 2: Timing of conversions during quarantine and duration of quarantine
- A) Cumulative probability of conversion from close contact to case with positive test (i.e., secondary
- 445 attack rate), from the day of last contact with the index case. Most conversions occur within the first
- 446 5 days since last contact. Conversion on day 0 means that the sample for the PCR test was taken on
- the day of last contact. The person was reported as a close contact and most likely received the
- 448 positive test result one day later.
- B) Duration of quarantine for household contacts and non-household contacts. This includes
- 450 potential self-quarantine before the mandated quarantine and extensions due to other household
- 451 members in quarantine converting to cases.
- 452 Figure 3: Participants' responses regarding motivation for quarantine measures, reasons for
- 453 motivations, worries, and positive experiences during quarantine.
- 454 **Figure 4**: Change in proportion of persons with symptoms of depression, anxiety, or stress from
- 455 before to during mandated quarantine. This analysis is restricted to participants who completed both
- 456 questionnaires.

457 Figures



- Figure 1: Participant flow through recruitment. Recruitment was performed in two stages. Potential
 participants had to complete an online form to be contacted by the study team or give oral consent
- to have the online form completed for them.



Figure 2: Timing of conversions during quarantine and duration of quarantine

A) Cumulative probability of conversion from close contact to case with positive test (i.e., secondary
attack rate), from the day of last contact with the index case. Most conversions occur within the first
5 days since last contact. Conversion on day 0 means that the sample for the PCR test was taken on
the day of last contact. The person was reported as a close contact and most likely received the
positive test result one day later.



- 470 Figure 3: Participants' responses regarding motivation for quarantine measures, reasons for
- 471 motivations, worries, and positive experiences during quarantine.



472

473 **Figure 4**: Change in proportion of persons with symptoms of depression, anxiety, or stress from

- 474 before to during mandated quarantine. This analysis is restricted to participants who completed both
- 475 questionnaires.









Severity of symptoms — Any severity - - Mild - - Moderate - (Extremely) severe