

1 **Screening of Thalassemia Carriers, Hemoglobin Variants, and** 2 **Comparison of Hematological Parameters between Children of** 3 **Bangladesh and Forcibly Displaced Myanmar Nationals**

4 Rumana Mahtarin^{1,3}, Kasrina Azad¹, Mohabbat Hossain^{1,9}, Mst. Sharmin Aktar Mukta¹,
5 Mohammad Tanbir Habib¹, Abu Bakar Siddik¹, Nishat Sultana¹, Zannat Kawser¹, Umme Kulsum¹,
6 Nusrat Sultana^{1,3,5}, Farjana Akther Noor⁶, Ahmad Zubair Mahdi⁷, Muhammad Asaduzzaman⁸,
7 Md. Ahashan Habib⁸, Emran Kabir Chowdhury³, Firdausi Qadri^{1,2}, Mst. Noorjahan Begum⁴,
8 A.H.M. Nurun Nabi^{3*}

9
10 ¹Institute for Developing Science and Health Initiatives (ideSHi), Dhaka-1206, Bangladesh

11 ² Mucosal Immunology and Vaccinology, Infectious Diseases Division, International Centre for
12 Diarrhoeal Disease Research, Bangladesh, Mohakhali, Dhaka, Bangladesh.

13 ³Department of Biochemistry and Molecular Biology, University of Dhaka, Dhaka 1000,
14 Bangladesh

15 ⁴Virology Laboratory, Infectious Diseases Division, International Centre for Diarrhoeal Disease
16 Research, Bangladesh, Mohakhali, Dhaka, Bangladesh.

17 ⁵Department of Virology, Dhaka Medical College, Dhaka 1000, Bangladesh

18 ⁶Department of Biochemistry and Molecular Biology, Tejgaon College, 16 Indira Road, Dhaka-
19 1215, Bangladesh

20 ⁷Department of Community Medicine & Public Health, Tairunnessa Memorial Medical College
21 & Hospital, Gazipur, Bangladesh

22 ⁸Upazila Health Complex, Matlab (North), Chandpur, Bangladesh

23 ⁹Department of Genetic Engineering and Biotechnology, University of Chittagong, Chittagong-
24 4331, Bangladesh

25
26 *Correspondence: nabi@du.ac.bd

41 **Abstract**

42 Thalassaemia is a hereditary blood disorder with different severity spectrums. This study aimed to
43 assess thalassaemia screening rates between children of Bangladesh and selected camps of Forcibly
44 Displaced Myanmar Nationals (FDMN) in Cox's Bazar in Bangladesh and compare the
45 hematological parameters among the screening groups. Complete blood count (CBC) analysis and
46 hemoglobin electrophoresis for each participant were performed by collecting venous blood.
47 Statistical analysis was employed for the comparison of parameters in blood. The thalassaemia
48 carrier and other hemoglobin variant rate in Bangladeshi children in selected areas have been found
49 to be 20.7% and in FDMN children, the rate is 8.2%. Hematological differences are visualized
50 among children of two nations. Intra-and interpopulation variances are highlighted in principal
51 component analysis where higher variance (94.87%) in Bangladeshi children than FDMN children
52 (80.68%). Receiver operating characteristics (ROC) and area under the curve (AUC) analyses
53 revealed, RBC (0.761, 0.902, and 0.791) and RDW-CV are better model (0.819, 0.925, and 0.858)
54 among the classifier of blood parameters. Pearson correlation shows distinguished covariation or
55 association among the parameters. The outcome of the study highlights the discrepancies in levels
56 of carriers in regions in Bangladesh and suggests further screening as well as population based
57 molecular research to ensure better treatment strategies.

58

59 **Keywords:** Thalassaemia Carrier Screening, Beta Thalassaemia Trait, Hemoglobin Variant,
60 Children

61

62

63

64

65

66

67

68

69

70

71

72

73

74

75

76

77

78

79

80

81

82 **1. Introduction**

83 Thalassaemia is an inherited hemolytic anemia which passes through autosomal recessive pattern
84 [1]. A fractional or complete absence in the synthesis of α -globin chains or β -globin chains due to
85 genetic defects in the *α -globin* gene cluster on chromosome 16 or the *β -globin* gene cluster on
86 chromosome 11 can lead to the synthesis of defective hemoglobin [2]. In β -thalassaemia, the main
87 consequence of unstable hemoglobin and iron overload is oxidative stress [4]. Iron triggers the
88 manufacture of reactive oxygen species, which in excess are toxic, contributing to oxidative
89 damage to red blood cells, leukocytes (recurrent infections), platelets (hypercoagulable state) [4],
90 cellular dysfunction, apoptosis, and necrosis in vital organs such as heart, liver and endocrine
91 system [1, 2]. People with β -thalassaemia major (β^0/β^0 , β^0/β^+ , and sometimes β^+/β^+) generally
92 get medical attention within the first two years of life and necessitate regular RBC transfusions to
93 survive [2]. This transfusion can cause iron overload [3]. Dysregulation of iron homeostasis
94 promotes iron deposition in the organs and influences disease severity, which is the main cause of
95 death in beta-thalassaemia patients [4, 5]. Each year worldwide more than 300,000 newborns are
96 detected with major hemoglobinopathies of whom 60,000 to 70,000 are β -thalassaemia major cases,
97 especially in the Mediterranean area, Middle East, Far East, and East Asia [3]. In Bangladesh, the
98 birth rate of 33/10,000 had been reported where 8.68% had HbE trait, 2.24% had β -thalassaemia
99 (β -thal) trait and nearly 2500 thalassaemia major cases are added every year in Bangladesh [6,7]
100 while in Myanmar, thalassaemia carrier rate was 4.3% [8]. Previously migration increased the
101 frequency of thalassaemia disease in lower-prevalence regions [10]. Over a million Rohingyas from
102 Myanmar were forcefully displaced to Bangladesh and other neighboring countries by January
103 2018. Over half of the displaced Rohingyas have sought shelter in Bangladesh and settled in
104 Rohingya camps in Cox's Bazar [11]. There are no published thalassaemia screening outcomes on
105 Forcefully Displaced Myanmar Nationals (FDMN) in Bangladesh. This study aimed to compare
106 the prevalence of thalassaemia carriers and hemoglobin variants to associate hematological
107 parameters among Bangladeshi children and Rohingya children who were forcibly displaced from
108 Myanmar in Bangladesh.

109 **2. Methods**

110 **2.1. Enrollment of Study Participants**

111 In screening campaigns, a total of 92 Bangladeshi children (49% male and 51% female), aged
112 between 1 and 15 years and 110 children of forcibly displaced Myanmar Nationals (58% male and
113 42% female), aged between 7 and 16 years were enrolled in the study from 2 May 2022 to 22
114 October 2022. The study (# PNR-22003) was ethically approved by Institutional Review Board of
115 Institute for Developing Science and Health Initiatives (ideSHi). Participants were provided
116 informed consent and written consent was taken from parents or legal guardians of each
117 participant.

118

119 **Sample size calculation:**

120 Required sample size has been calculated by the following formula [12, 13]

$$121 \quad n = \frac{(Z_{\alpha} + Z_{\beta})^2 p(1 - p)}{e^2}$$

122 Where,

123 n = required sample size,

124 $Z_{1-\alpha/2}$ = Standard normal variate for level of significance

125 Z_{β} = Standard normal variate for power

126 p = Expected prevalence,

127 e = Margin of error or acceptable difference

128 n_b = Sample size for Bangladeshi Population.

129 n_m = Sample size for Myanmar Population.

130 Considering, the level of significance = 5%, power of the test = 85%, expected prevalence rate for
131 Bangladeshi population $P_b = 10.2\%$ [14], margin of error $e = 10\%$ and including 15% attrition rate
132 the sample size is $n_b = 92$. Again, with the level of significance = 5%, power of the test = 96%,
133 expected prevalence rate for Myanmar population $P_m = 4.3\%$ [8], margin of error $e = 9\%$ and
134 including 15% attrition rate the sample size is $n_m = 110$.

135 **2.2. Analysis of Complete Blood Count of The Study Participants**

136 Three ml of venous blood was collected from each participant in EDTA containing vacutainer and
137 the collected blood samples were transported to the laboratory maintaining 4-8°C and preserved at

138 4°C until analysis. The complete blood count (CBC) was performed for each sample by using XS-
139 800i Hematology Analyzer (Sysmex, Japan) following the manufacturers' instruction. Major
140 hematological parameters like hemoglobin (HGB), hematocrit (HCT), RBC count and Red Cell
141 Indices including mean corpuscular volume (MCV), mean corpuscular hemoglobin (MCH), mean
142 corpuscular hemoglobin concentration (MCHC) and Red cell distribution width (RDW) were
143 considered for analysis.

144 **2.3. Hemoglobin Electrophoresis**

145 Hemoglobin electrophoresis was performed on automated capillary electrophoresis system (Sebia,
146 France) using Capillary Hemoglobin (E) kit to measure HbA, HbA2, HbF, and other abnormal Hb
147 variants following manufacturer's instructions.

148 **2.4 Statistical Analysis**

149 Statistical analysis was performed using SPSS and GraphPad Prism software (version 9.0) and R
150 programming. $P < 0.05$ was considered statistically significant for all the tests. One-way ANOVA
151 was performed for the analysis of hematological differences, and the distribution pattern was
152 visualized. Principle component analysis was performed for understanding intra and inter
153 population variation. Covariation of association among the major hematological parameters was
154 visualized by Pearson correlation. Receiver operating characteristics (ROC) and area under the
155 curve (AUC) analyses were also conducted for diagnostic assessment.

156

157

158 **3. Results**

159 **3.1. Thalassemia Carrier and Hemoglobin Variant Screening** 160 **Outcomes**

161 In the study, the screening showed that the frequency of thalassemia traits and other hemoglobin
162 variants in Bangladeshi children was 20.7%, where beta thalassemia trait (BTT) and HbE trait
163 (HbET) were 5.43% and 13.04%, respectively. The highest carrier frequency was found among
164 the children in Dhaka and Rangpur (only HbE, 23.5%) divisions, while it is comparatively lower
165 in Chattogram (17.65%, BTT-11.76%, HbET-5.89%), and Barishal (17.65%, BTT-11.76%,
166 HbET- 5.89%). The lowest carrier frequency was observed among the children in Sylhet (only

167 HbET-5.8%), Mymensingh (only HbET-5.8%) and Khulna (only BTT-5.8%) divisions. The low
168 HbA2 had been detected in Rajshahi (1.1%).

169 On the other hand, thalassemia carrier and other hemoglobin variant frequency was 8.2% among
170 the children from the FDMNs, where BTT and HbE were 2.73% and 5.45%, respectively. In
171 addition, HbD variant was also found in the Bangladeshi children (1.09%) and FDMN children
172 (0.9%). The distribution of different groups in screening was displayed in Fig 1.

173
174 **Fig 1.** The alluvial plot represents distribution of screening outcomes in BD and FDMN children. The color codes
175 indicate different groups: Beta Thalassemia Trait (purple), HbD Trait (blue), HbE Trait (lemon), Low Hb A2 (orange).
176

177 **3.2. Analysis of Hemoglobin Variants**

178 Hemoglobin electrophoresis among the healthy participants of Bangladeshi children showed the
179 levels of HbA and HbA2 were $97.14 \pm 0.31\%$ and $2.80 \pm 0.25\%$, respectively. In beta thalassemia
180 trait (BTT), the levels of HbA, HbA2 and HbF were $93.78 \pm 1.5\%$, $5.40 \pm 0.36\%$, and $2.05 \pm 2.33\%$
181 respectively. In HbE trait the levels of HbA, HbA2 and HbE were $70.80 \pm 1.4\%$, $3.72 \pm 0.48\%$,
182 and $25.39 \pm 1.08\%$, respectively. In addition, HbA (54%), HbA2 (3.7%), and HbD (42.3%) were
183 found in HbD trait. In low HbA2 participant, HbA was 98.1% and HbA2 was 1.9%, and suspected
184 to have alpha thalassemia trait (HGB 8.7 g/dL, RBC $4.68 \times 10^{12}/L$, MCV 55.8 fL, MCH 18.6
185 pg) according to Mentzer index (MCV/RBC <13).

186 On the other hand, among the normal or non-carrier children of FDMN, the levels of HbA and
187 HbA2 were $97.3 \pm 0.29\%$ and $2.67 \pm 0.21\%$ respectively. In beta thalassemia trait (BTT), the level
188 of HbA was $93.6 \pm 0.47\%$, HbA2 was $5.3 \pm 0.47\%$ and HbF was $1.17 \pm 0.5\%$. In HbE trait, the
189 level of HbA was $72.9 \pm 3.12\%$, HbA2 was $3.8 \pm 0.35\%$, HbE was $23.0 \pm 3.4\%$, and HbF was
190 1.1%. In addition, a child of FDMN with HbD trait showed the level of HbA, HbA2, and HbD at
191 56.4%, 2.9%, and 41%, respectively. Alpha thalassemia trait was not found in FDMN children
192 from study population.

193

194 **3.3. Principal Component Analysis**

195 In this study, the principal component analysis was considered as feature extractor to compare the
196 participants of Bangladesh (BD) and Forcibly Displaced Children of Myanmar Nationals. The

197 principal components, PC1 and PC2 accounted for most of the variation in the participants based
198 on gender and age distribution. In Fig 2A among Bangladeshi children, PC1 and PC2 showed
199 63.0% and 31.7% variance, respectively (in total 94.87%). On the contrary, the FDMN children
200 showed the variance of PC1 and PC2 at 50% and 30.7%, respectively, while the total variance was
201 80.68% (Fig 2B).

202 **Fig 2.** Intra-and inter-population variances are highlighted in Principal component analysis. (A) PC scores in BD
203 children's groups and (B) PC scores in FDMN children's groups. The color codes represent different groups in BD
204 and FDMN children, Normal (blue), Beta Thalassemia Trait (BTT) (red), HbE (green), HbD (purple), Low HbA2
205 (orange).

206 **3.4. Analysis of Hematological Differences between Bangladeshi and** 207 **Forcibly Displaced Children of Myanmar Nationals**

208 The column bar graphs exhibited a diverse distribution of major hematological parameters among
209 the prevalent groups in children of the two countries.

210
211 **Fig 3.** Distribution pattern of major hematological parameters.

212 Normal, Beta Thalassemia Trait (BTT), and Hemoglobin E Trait (HbET) groups in both (A) BD and (B) FDMN
213 children.

214
215 According to the one-way ANOVA test, RBC, MCV, MCH, and RDW-CV were substantially
216 different among the normal, BTT, and HbET groups in Bangladeshi children (Fig 3A), with a p-
217 value of 0.05. Among the BTT children of Bangladesh, RBC and RDW-CV were found
218 significantly higher, whereas MCV and MCH were significantly lower compared to the normal
219 value (Table 1). In the FDMN children, all the hematological parameters except MCHC showed
220 significant results among the normal, BTT and HbET groups (Fig 3B) with a p-value of <0.05.
221 RBC and RDW-CV were significantly higher, while HGB, HCT, MCV, and MCH were
222 significantly lower compared to the normal level in BTT in the FDMN children (Table 2).

223 However, the hematological parameters of both Bangladeshi children and FDMN children showed
224 significant results for RBC, HGB, MCV, MCH, HCT, and RDW-CV among the normal, BTT and
225 HbET groups. RBC and RDW-CV were significantly higher, while HGB, HCT, and MCH were
226 significantly lower in BTT of both populations (Table 3).

227 **Table 1. Comparative analysis of major hematological parameters of Bangladeshi children**
228 **in major groups**

229

Hematological parameters	Normal	BTT	HbET	P value
RBC (X10 ¹² /L)	4.34 ± 0.47	5.83 ± 1.14	4.76 ± 0.50	0.000
HGB (g/dL)	11.53 ± 1.25	10.87 ± 1.68	10.79 ± 1.21	0.107
HCT (%)	31.59 ± 3.27	30.50 ± 4.55	29.56 ± 2.59	0.096
MCV (fL)	72.97 ± 4.98	52.73 ± 3.54	62.61 ± 6.06	0.000
MCH (pg)	26.62 ± 1.89	18.77 ± 1.17	22.68 ± 2.46	0.000
MCHC (g/dL)	36.49 ± 1.01	35.60 ± 0.26	36.19 ± 1.21	0.232
RDW-CV (%)	13.52 ± 1.01	17.73 ± 1.94	15.19 ± 2.24	0.000

230 *Here, RBC means Red Blood Corpuscle, HGB means Hemoglobin, HCT means Hematocrit, MCV means Mean*
 231 *Corpuscular Volume, MCH means Mean Corpuscular Hemoglobin, MCHC means Mean Corpuscular*
 232 *Hemoglobin Concentration, and RDW-CV means Red Cell Distribution Width - Coefficient of Variation*
 233

234 **Table 2. Comparative analysis of major hematological parameters of Forcibly Displaced**
 235 **Children of Myanmar Nationals (FDMN) children in major groups**

Hematological parameters	Normal	BTT	HbET	P value
RBC (X10 ¹² /L)	4.46 ± 0.33	5.45 ± 0.17	4.89 ± 0.26	0.000
HGB (g/dL)	11.41 ± 0.90	9.43 ± 0.65	10.90 ± 0.67	0.001
MCV (fL)	73.92 ± 4.03	51.03 ± 0.81	63.76 ± 3.85	0.000
HCT (%)	32.92 ± 2.39	27.83 ± 1.03	31.12 ± 1.70	0.001
MCH (pg)	25.62 ± 1.71	17.30 ± 0.66	22.34 ± 1.45	0.000
MCHC (g/dL)	34.64 ± 0.99	33.90 ± 1.15	35.02 ± 0.66	0.299
RDW-CV (%)	13.40 ± 0.80	19.50 ± 0.95	14.32 ± 0.30	0.000

236 *Here, RBC means Red Blood Corpuscle, HGB means Hemoglobin, HCT means Hematocrit, MCV means Mean*
 237 *Corpuscular Volume, MCH means Mean Corpuscular Hemoglobin, MCHC means Mean Corpuscular*
 238 *Hemoglobin Concentration, and RDW-CV means Red Cell Distribution Width - Coefficient of Variation*
 239

240 **Table 3. Comparative analysis of major hematological parameters of both Bangladeshi and**
 241 **Forcibly Displaced Children of Myanmar Nationals (FDMN) children in major groups**

Hematological parameters	Normal	BTT	HbET	P value
RBC (X10 ¹² /L)	4.41 ± 0.40	5.62 ± 0.64	4.71 ± 0.37	0.000
HGB (g/dL)	11.46 ± 1.06	10.09 ± 1.18	10.93 ± 1.09	0.000
MCV (fL)	72.98 ± 5.36	58.51 ± 9.04	66.58 ± 5.44	0.000
HCT (%)	32.36 ± 2.90	29.80 ± 3.04	30.79 ± 2.85	0.013
MCH (pg)	26.04 ± 1.85	17.96 ± 1.01	23.16 ± 1.47	0.000
MCHC (g/dL)	35.42 ± 1.35	34.96 ± 1.10	35.91 ± 1.26	0.206
RDW-CV (%)	13.45 ± 0.89	18.03 ± 1.80	14.81 ± 2.01	0.000

242 *Here, RBC means Red Blood Corpuscle, HGB means Hemoglobin, HCT means Hematocrit, MCV means Mean*
 243 *Corpuscular Volume, MCH means Mean Corpuscular Hemoglobin, MCHC means Mean Corpuscular*
 244 *Hemoglobin Concentration, and RDW-CV means Red Cell Distribution Width - Coefficient of Variation*
 245

246 **3.5. Correlation among Hematological Parameters in Different** 247 **Groups of Children in Bangladesh and Forcibly Displaced Myanmar** 248 **Nationals (FDMN)**

249 Pearson correlation showed covariation or association among the hematological parameters.
250 Diverse correlation pattern in CBC parameters was persistent in both BD and FDMN children.
251 Among Bangladeshi children, normal or non-carriers and HbET showed quite similar type of
252 strong positive correlation pattern among RBC, HGB, HCT, MCV and MCH, while RDW-CV
253 exhibited negative correlation with the other hematological parameters. BTT showed strong
254 positive relationship among RBC, HGB, HCT, and RDW-CV, MCV, and MCH. In FDMN,
255 variations were more prominent among the different groups. BTT exhibited strong positive
256 correlation among most of the hematological parameters except HCT and RDW-CV (Fig 4).
257

258 **Fig 4.** Correlation pattern in major hematological parameters in both BD and FDMN children in major groups.
259 The groups in (A) BD children and (B) FDMN children. BTT indicates Beta Thalassemia Trait and HbET indicates
260 Hemoglobin E Trait. The major hematological parameters are numbered as 1 to 7 and the correlation is measured as -
261 1.0 to 1.0. Color code indicates correlation pattern. Red color indicates negative correlation (-1.0 to 0) where -1.0
262 indicates least correlation and blue color indicates positive correlation (0 to 1.0) where 1.0 indicates highest correlation.
263

264 **3.6. ROC Curve Analysis**

265 ROC curve analysis was performed for the assessment of diagnostic value of thalassemia carrier
266 and hemoglobin variant screening. ROC curves were plotted for children of Bangladesh and
267 FDMN children (Fig 5A, Fig 5B) and for both nations (Fig 5C). AUC was measured to visualize
268 the classifier of blood parameter model. RBC (0.761, 0.902, and 0.791) and RDW-CV (0.819,
269 0.925, and 0.858) covered high AUC and good classifier for carrier screening. High AUC indicated
270 more accuracy for diagnostic performance which might be supportive to hemoglobin
271 electrophoresis test for proper diagnosis of thalassemia and hemoglobinopathies.
272

273 **Fig 5.** ROC curve and area under curve display classifier of blood parameter model. (A) BD (B) FDMN (C) both
274 BD and FDMN.
275

276
277
278

279 **4. Discussion**

280 This is the first study for assessment of thalassemia carrier and hemoglobin variant screening data
281 in children between Bangladesh and Forcibly Displaced Myanmar Nationals (FDMN). In the
282 study, the frequency of thalassemia carriers and hemoglobin variants in Bangladeshi children is
283 found 20.7%, which is double of the previous report (10.2%) [7, 14]. The highest carrier frequency
284 was found in Dhaka and Rangpur divisions while the lowest carrier frequency was observed in
285 Sylhet, Mymensingh, and Khulna divisions. On the other hand, the carrier rate in FDMN children
286 was 8.2%, where beta thalassemia trait (BTT) rate is 2.73%, and HbE trait (HbET) rate is 5.45%.
287 Combining the rate for both study groups, the cumulative carrier rate is around 14% which is also
288 demonstrate the increasing trend compared to previous record [7, 14]. Principle component
289 analysis highlights the marked variations in Bangladesh and Forcibly Displaced Myanmar
290 Nationals for their genetic ancestry conformation. Higher variance has been exposed in
291 Bangladeshi children (94.9%) than in FDMN children (80.7%) for intra-and inter-population
292 variances of different groups. Divergent distribution pattern of major hematological parameters
293 was visualized for the children of both Bangladesh and FDMN. The variations were higher in BTT
294 group of both populations while wider distribution in Bangladesh and narrower distribution
295 patterns are visible in FDMN children. ROC curve evaluates the diagnostic performance of carrier
296 and hemoglobin variant screening by demonstrating the sensitivity and specificity of the test. AUC
297 under ROC curve was measured to visualize the classifier of blood parameter model. RBC (0.761,
298 0.902, and 0.791) and RDW-CV (0.819, 0.925, and 0.858) display high AUC which indicates
299 better classifier for the screening. Comparing all the parameters, RBC and RDW-CV are
300 supportive to hemoglobin electrophoresis to evaluate the screening outcomes. There was
301 association of RDW-CV and thalassemia disease severity in previous study [9]. Correlation
302 matrixes represent covariation or association among the parameters. Distinguished correlation
303 was revealed in both Bangladesh and FDMN children. BTT groups of both Bangladesh and FDMN
304 demonstrate the higher number of strong positive correlation among parameters although the
305 parameters differ in their association. Mostly RDW-CV demonstrates quite negative association
306 with other hematological parameters in the FDMN children. All the analyses demonstrate the
307 similarities and discrepancies in the major hematological parameters of the groups from
308 Bangladesh and Forcibly Displaced Myanmar Nationals. The screening in children has been

309 conducted to determine the carrier and hemoglobin variant rates of existing populations with two
310 different genetic background in Bangladesh. However, a drawback is that we could screen only a
311 limited number of children since this was conducted within a thalassemia awareness program.
312 More extensive thalassemia carrier screening and awareness programs, prenatal diagnosis, and
313 compulsory carrier screening before marriage should be implemented to prevent the spread of
314 thalassemia disease and other hemoglobinopathies. This comparative study will be helpful for
315 track the rates of thalassemia carrier and hemoglobin variant in children to prevent the disease
316 burden. As well as this population-based study will expedite molecular analysis for the
317 improvement of thalassemia treatment strategies by initiating personalized medicine. Finally, such
318 studies and programs will help in creating awareness of thalassemia, its prevention and in policy
319 changes in the country.

320 **5. Conclusion**

321 The frequency of carrier and hemoglobin variant is much higher in Bangladeshi children than the
322 displaced population from Myanmar (FDMN) into Bangladesh. This comparative study highlights
323 the discrepancies in thalassemia carrier and hemoglobin variant status on children which will be
324 helpful to understand the burden of thalassemia and hemoglobinopathies. Increasing the number
325 of screening and being aware of devastating health consequences are vital for prevention of
326 disease.

327

328 **Acknowledgement**

329 The authors are grateful to the First Security Islami Bank, Bangladesh for their contribution and
330 support to conduct the thalassemia screening program. We are thankful to the study participants,
331 the clinicians and staff for their assistance in specimen collection, and the thalassemia screening
332 team members of ideSHi.

333

334 **References**

- 335 [1] Fibach, E. and E.A. Rachmilewitz, *Pathophysiology and treatment of patients with beta-*
336 *thalassemia - an update*. F1000Res, 2017. **6**: p. 2156.
337 [2] Taher, A.T. and A.N. Saliba, *Iron overload in thalassemia: different organs at different*
338 *rates*. Hematology Am Soc Hematol Educ Program, 2017. **2017**(1): p. 265-271.

- 339 [3] Taher, A., et al., in *Guidelines for the Management of Non Transfusion Dependent*
340 *Thalassaemia (NTDT)*, D. Weatherall, Editor. 2013, Thalassaemia International Federation
341 © 2013 Thalassaemia International Federation.: Nicosia (Cyprus).
- 342 [4] Sposi, N., *Oxidative Stress and Iron Overload in β -Thalassaemia: An Overview*. 2019.
- 343 [5] Leecharoenkiat, K., et al., *Iron dysregulation in beta-thalassaemia*. Asian Pac J Trop Med,
344 2016. **9**(11): p. 1035-1043.
- 345 [6] Hossain, M.S., et al., *Thalassemys in South Asia: clinical lessons learnt from Bangladesh*.
346 Orphanet Journal of Rare Diseases, 2017. **12**(1): p. 93.
- 347 [7] Noor, F.A., et al., *Nationwide carrier detection and molecular characterization of β -*
348 *thalassaemia and hemoglobin E variants in Bangladeshi population*. Orphanet J Rare Dis,
349 2020. **15**(1): p. 15.
- 350 [8] Win, S., *Epidemiology and current status of case management of thalassaemia in Myanmar*
351 *- Country Report*. Burma medical journal, 2016. **58**: p. 43-51.
- 352 [9] Islam, M.T., et al., Association of diverse population of red blood cells with different
353 disease manifestations in patients with beta-thalassaemia. Meta Gene, 2021. **27**: p. 100846.
- 354 [10] Kattamis, A., et al., *Changing patterns in the epidemiology of β -thalassaemia*. Eur J
355 Haematol, 2020. **105**(6): p. 692-703.
- 356 [11] Rashid, R., et al., *A descriptive study of Forcefully Displaced Myanmar Nationals (FDMN)*
357 *presenting for care at public health sector hospitals in Bangladesh*. Glob Health Action,
358 2021. **14**(1): p. 1968124.
- 359 [12] Pourhoseingholi, M.A., M. Vahedi, and M. Rahimzadeh, *Sample size calculation in*
360 *medical studies*. Gastroenterol Hepatol Bed Bench, 2013. **6**(1): p. 14-7.
- 361 [13] Charan, J. and T. Biswas, *How to calculate sample size for different study designs in*
362 *medical research?* Indian J Psychol Med, 2013. **35**(2): p. 121-6.
- 363 [14] Khan, W.A., et al. *Prevalence of Beta thalassaemia trait and Hb E trait in Bangladeshi*
364 *school children and health burden of thalassaemia in our population*. 2005.
- 365
- 366

medRxiv preprint doi: <https://doi.org/10.1101/2023.09.19.23295805>; this version posted September 21, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/).




Fig 1. The alluvial plot represents distribution of screening outcomes in BD and FDMN children. The color codes indicate different groups: Beta Thalassemia Trait (purple), HbD Trait (blue), HbE Trait (lemon), Low Hb A2 (orange).




Fig 2. Intra- and inter-population variances are highlighted in Principal component analysis. (A) PC scores in BD children's groups and (B) PC scores in FDMN children's groups. The color codes represent different groups in BD and FDMN children, Normal (blue), Beta Thalassemia Trait (BTT) (red), HbE (green), HbD (purple), Low HbA2 (orange).

medRxiv preprint doi: <https://doi.org/10.1101/2023.09.19.23295805>; this version posted September 21, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/).




Fig 3. Distribution pattern of major hematological parameters.

Normal, Beta Thalassemia Trait (BTT), and Hemoglobin E Trait (HbET) groups in both (A) BD and (B) FDMN children.





Fig 4. Correlation pattern in major hematological parameters in both BD and FDMN children in major groups.

The groups in (A) BD children and (B) FDMN children. BTT indicates Beta Thalassemia Trait and HbET indicates Hemoglobin E Trait. The major hematological parameters are numbered as 1 to 7 and the correlation is measured as -1.0 to 1.0. Color code indicates correlation pattern. Red color indicates negative correlation (-1.0 to 0) where -1.0 indicates least correlation and blue color indicates positive correlation (0 to 1.0) where 1.0 indicates highest correlation.




medRxiv preprint doi: <https://doi.org/10.1101/2023.09.19.23295805>; this version posted September 21, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Area Under the Curve

Test Result Variable(s)	Area	95% Confidence Interval	
		Lower Bound	Upper Bound
RBC (X10 ¹² /L)	0.761	0.801	1.000
HGB (g/dL)	0.325	0.054	0.386
HCT (%)	0.297	0.000	0.037
MCV (fL)	0.053	0.039	0.319
MCH (pg)	0.060	0.000	0.094
MCHC (g/dL)	0.418	0.294	0.718
RDW-CV (%)	0.819	0.866	0.983

B




medRxiv preprint doi: <https://doi.org/10.1101/2023.09.10.23295805>; this version posted September 21, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Area Under the Curve

Test Result Variable(s)	Area	95% Confidence Interval	
		Lower Bound	Upper Bound
RBC (X10 ¹² /L)	0.902	0.801	1.000
HGB (g/dL)	0.220	0.054	0.386
HCT (%)	0.016	0.000	0.037
MCV (fL)	0.179	0.039	0.319
MCH (pg)	0.046	0.000	0.094
MCHC (g/dL)	0.506	0.294	0.718
RDW-CV (%)	0.925	0.866	0.983

C



medRxiv preprint doi: <https://doi.org/10.1101/2023.09.19.23295805>; this version posted September 21, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

Area Under the Curve
It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Test Result Variable(s)	Area	95% Confidence Interval	
		Lower Bound	Upper Bound
RBC (X10 ¹² /L)	0.791	0.682	0.899
HGB (g/dL)	0.299	0.178	0.420
HCT (%)	0.222	0.141	0.302
MCV (fL)	0.443	0.334	0.552
MCH (pg)	0.072	0.032	0.113
MCHC (g/dL)	0.561	0.444	0.677
RDW-CV (%)	0.858	0.788	0.929

Fig 5. ROC curve and area under curve display classifier of blood parameter model. (A) BD (B) FDMN (C) both BD and FDMN.