1 Title

2 A Natural History of NAA15-related Neurodevelopmental Disorder Through Adolescence

- 3 Rikhil Makwana^{1*}, Carolina Christ^{1*}, Elaine Marchi¹, Randie Harpell¹, Gholson J. Lyon^{1,2,3**}
- ⁴ ¹Department of Human Genetics, New York State Institute for Basic Research in Developmental
- 5 Disabilities, Staten Island, New York, United States of America
- ²George A. Jervis Clinic, New York State Institute for Basic Research in Developmental
- 7 Disabilities, Staten Island, New York, United States of America
- ³Biology PhD Program, The Graduate Center, The City University of New York, New York,
- 9 United States of America
- 10

11 Abstract

NAA15 is a member of the NatA N-terminal acetyltransferase complex which acts by 12 interacting with the NAA10 enzymatic sub-unit. Individuals with variants in the NAA15 coding 13 region develop NAA15-related neurodevelopmental syndrome which presents with a wide array 14 of manifestations that affect the heart, brain, musculoskeletal system, and behavioral and 15 16 cognitive development. We tracked a cohort of 26 participants (8 females and 18 males) over time, each with a pathogenic NAA15 variant, and administered the Vineland-3 assessment to 17 them to assess their adaptive functioning. We found that the cohort performed significantly 18 19 worse compared to the normalized Vineland values. On average, females performed better than males across all domains. They performed significantly better on the Motor Domain and Fine 20 Motor Sub-Domain portions of the assessment. Over time, females showed a decrease in 21 22 adaptive functioning with the decline being especially strongly correlated at the Coping sub-23 domain, Domestic sub-domain, and Fine motor sub-domains. It is difficult to determine the

strength of these correlations due to limited power. Males (after excluding one outlier) showed a
moderate positive correlation between age and ABC standard score. Ultimately, additional
longitudinal data should be collected to determine the validity of the between sex-differences and
to better understand the change in adaptive behavioral outcomes of individuals with NAA15neurodevelopmental disorder as they age.

29 Introduction

N-terminal acetylation is a prevalent protein modification that involves the addition of an 30 31 acetyl group to the alpha-amino group of the N-terminal amino acid of a protein (Liszczak et al., 2013). This enzymatic process is catalyzed by a group of N-terminal acetyltransferases (NATs) 32 and plays a pivotal role in various cellular processes, including protein stability, protein-protein 33 interactions, and cellular localization (Aksnes et al., 2015; Bienvenut et al., 2012; Dikiy & 34 Eliezer, 2014). There are seven known NAT types, with NatA being the most prominent, believed 35 36 to be responsible for modifying 40-50% of the human proteome(Deng et al., 2019; Feng & Ma, 2016; Van Damme, 2021). The NatA complex is composed of the catalytic sub-unit NAA10, and 37 the auxiliary sub-units NAA15 and HYPK (Deng et al., 2019; Dörfel et al., 2017; Feng et al., 38 39 2016; Knorr et al., 2019; Weidenhausen et al., 2021).

Dysregulation of NAA10 and NAA15 expression or activity has been associated with the pathogenesis of various diseases, including cancer (Kuhns et al., 2018; Le et al., 2023; Zhang et al., 2020; Zhu et al., 2024), neurodevelopmental disorders (Cheng et al., 2018; Lyon et al., 2023; Makwana et al., 2024; Sidhu et al., 2017; Wu & Lyon, 2018), and cardiovascular diseases (Belbachir et al., 2023; Huth et al., 2023; Ritter et al., 2021; Støve et al., 2018; Ward et al., 2021). More notably, mutations in the X-linked gene NAA10 lead to Ogden Syndrome, which has numerous effects on individual's development and overall health. NAA15-related

47	neurodevelopmental syndrome has overlapping symptoms with Ogden Syndrome, including
48	intellectual disability, atypical or delayed development, heart anomalies, and more features
49	previously described (Cheng et al., 2020; Dörfel & Lyon, 2015; Lyon et al., 2023; McTiernan et
50	al., 2018; Myklebust et al., 2015; Patel et al., 2024; Pesz et al., 2018; Stessman et al., 2017;
51	Støve et al., 2018; Umlai et al., 2021; Wong et al., 2019; Zhao et al., 2018). Additionally,
52	NAA15 variants are associated with musculoskeletal and neuromuscular abnormalities that can
53	appear in childhood or may have a delayed appearance until adulthood (Danti et al., 2024;
54	Monestier et al., 2019; Pesz et al., 2018; Tian et al., 2022; Yubero et al., 2022).
55	The aim of the current study is to further describe adaptive functioning in individuals
56	with NAA15-related neurodevelopmental syndrome by analyzing prospective data and
57	understand how this disorder progresses over time.
58	Methods
58 59	Methods <u>Participants</u>
58 59 60	Methods Participants All families that participated in the investigation signed Institutional Review Board
58 59 60 61	Methods Participants All families that participated in the investigation signed Institutional Review Board (IRB)-approved consents and HIPAA forms. The scope of the project was explained in lay
58 59 60 61 62	Methods Participants All families that participated in the investigation signed Institutional Review Board (IRB)-approved consents and HIPAA forms. The scope of the project was explained in lay language. The participant population included children with NAA15 pathogenic variants that the
58 59 60 61 62 63	Methods Participants All families that participated in the investigation signed Institutional Review Board (IRB)-approved consents and HIPAA forms. The scope of the project was explained in lay language. The participant population included children with NAA15 pathogenic variants that the principal investigator has previously worked with for other research projects. Participants were
58 59 60 61 62 63 64	Methods Participants All families that participated in the investigation signed Institutional Review Board (IRB)-approved consents and HIPAA forms. The scope of the project was explained in lay language. The participant population included children with NAA15 pathogenic variants that the principal investigator has previously worked with for other research projects. Participants were not compensated for their time. Overall, there were 26 participants who were administered the
58 59 60 61 62 63 64 65	Methods Participants All families that participated in the investigation signed Institutional Review Board (IRB)-approved consents and HIPAA forms. The scope of the project was explained in lay language. The participant population included children with NAA15 pathogenic variants that the principal investigator has previously worked with for other research projects. Participants were not compensated for their time. Overall, there were 26 participants who were administered the assessment.
58 59 60 61 62 63 64 65 66	Methods Participants All families that participated in the investigation signed Institutional Review Board (IRB)-approved consents and HIPAA forms. The scope of the project was explained in lay language. The participant population included children with NAA15 pathogenic variants that the principal investigator has previously worked with for other research projects. Participants were not compensated for their time. Overall, there were 26 participants who were administered the assessment. Cognitive Assessment

68 Communication Daily Living Skills, and Socialization. These domains are further split into sub-

domains that are used to assess an individual's competency in tasks of personal and social 69 sufficiency (Perry et al., 2009). The scores for each core domain are norm referenced and 70 71 summed to generate an Adaptive Behavior composite (ABC) score that provides a holistic 72 picture of an individual's adaptive behavior across domains. The Vineland 3 tools were 73 administered by three trained assessors at various timepoints in the participants' lives ranging 74 from 1-23 years old. Individual assessor scores were cross validated with one another to ensure equivalency in how assessments were administered and scored. Participant caregivers were those 75 interviewed due to their proximity and knowledge of the patients. Participants were given access 76 77 to the Vineland assessment scores after they had been scored.

78 <u>Analysis</u>

The Principal Investigator and associated staff collected at least one Vineland 3 score for 79 each participant over multiple years. Some participants were able to take the assessment more 80 81 than once providing a longitudinal outlook. Natural history analysis and visualization was then performed using the GraphPad Prism software. All pathogenic variants were analyzed together 82 and graphed by comparing their ABC standard score against the age at which the assessment was 83 84 administered. Further analysis was performed by filtering graphs by sex. All pathogenic variants, male participants, and female participants were also graphed according to their Communication 85 (com), Daily Living Skills (dls), Social (soc), and Motor (mot) domain standard scores. The 86 87 domain scores were further broken down by sub-domain standard score to allow for more granular identification of participant strength and weakness over time. The Communication sub-88 domain scores were Receptive (rec), Expressive (exp), and Written (wrn). The Daily Living 89 90 Skills sub-domains analyzed were Personal (per), Domestic (dom), and Community (cmm). The 91 Socialization sub-domains of interest were Interpersonal Relationships (ipr), Play and Leisure

(pla), and Coping Skills (cop). The Motor sub-domains of Interest were Fine Motor (fmo) and
Gross Motor (gmo). Scores were only collected and included for this motor domain if patients
were under the age of nine to stay within the validated age range of the exam (Perry et al., 2009).
Lastly, the Internalizing (int) and Externalizing (ext) sub-domain components of the Maladaptive
Trait domain were also included in analysis, however the Critical Items component of the
domain were not.

The scores were compared in Microsoft Excel with two-tailed equal variance t-tests to 98 compare the adaptive behavior outcomes between males and females to determine if there were 99 sex differences. The p-value was set at .05 for all calculations. Correlation coefficients were also 100 101 calculated for each of the graphs made using Excel. A second round of calculations for the t-test and the correlation coefficients in the male cohort was performed to exclude NAA15-011 due to 102 being three standard deviations above the mean in age compared to the other participants and 103 having performed two standard deviations below the mean in Vineland ABC, Motor, Social, and 104 105 Daily Living Skills standard scores. Data presented in tables includes calculations performed including participant NAA15-011 unless otherwise specified. 106

107 **Results**

There were 26 participants who were administered the Vineland and included in this study. A breakdown of the different pathogenic variants in each sex can be seen in **Table 1**. Males and females each had unique pathogenic variants. Twenty-four participants developed their variant *de novo* and 2 participants have variants of unknown origin (due to the parents not being tested). Of the 26 participants, 8 are female and 18 are male with the age range being from ~0 to ~25 years old (mean = 8.8 years, standard deviation (SD) = 4.6).

114 Compared to the general population, individuals with NAA15 variants scored below 115 average on the Vineland 3 Assessment (where average is mean=100, sd=15). This was true for 116 ABC standard scores as well as the main domain scores. The average ABC score among NAA15 117 variants was 66.7 (SD = 21.1). Females tended to outperform males across adaptive behavior 118 domains. However, except for the Motor domain score these differences were not significant. A 119 summary of the Vineland 3 scores can be seen in **Table 2**.

120 Graphical representations of the ABC standard scores over time can be seen in Figure 1. Before stratifying by sex, there appears to be an overall downward trend in score with age. 121 Figure 2 shows the ABC standard scores separated by male and female. Separation by sex shows 122 123 a moderate linear increase in adaptive function in males over time, when excluding NAA15-011, 124 whereas females appear to show a moderately strong linear decay. There is also a strong linear 125 decay in the female Daily Living Skills domain over times whereas the males showed a moderate positive linear growth when excluding NAA15-011. Table 3 showcases the remaining calculated 126 127 correlation coefficient values for the Vineland ABC and Domain standard scores for the females, males, and males excluding NAA15-011. 128

The Communication, Daily Living Skills, Social, and Motor standard score graphs can be seen in **Figure 3** and follow similar trends for females and males, when excluding NAA15-011, as the ABC scores over time. It should be noted that excluding NAA15-011 did not significantly change the t-test statistics of the standard scores when comparing the males and the females. The magnitude at which the females outperformed the males decreased. However, except for the Motor domain score, their differences were still not significant.

Dividing the Motor domain into its Fine Motor and Gross Motor sub-domains elucidated
the difference between sexes. Males, with and without the inclusion of NAA15-011, on average

scored 7.0 (SD = 2.5) which is significantly less than the females who had an average score of 137 14.0 (SD = 3.6). On average for Gross Motor subdomain scores, Males (mean = 8.5, SD = 3.5) 138 139 also performed worse than their female counterparts (mean = 10.8, SD = 1.8). This difference was not significant. There were no other significant differences between how the sexes 140 performed in the various sub-domains. A summary of the breakdown of each domain into its sub-141 142 domain components can be seen in Table 4. Each domain sub-domain score can be seen in Figure 4. There are strong negative linear 143 associations between the Coping, Fine Motor, and Domestic sub-domains over time in the 144 NAA15 females. There are also moderately strong negative linear associations in the Receptive 145

146 and Personal sub domains. The correlation coefficients in the males were not as strong as those

147 in the females. However, when excluding NAA15-011, the Personal sub-domain had a moderate

148 positive linear relationship over time. Calculated correlation coefficients for each sub-domain

149 can be seen in **Table 5.**

150 Discussion

Individuals with NAA15-related neurodevelopmental syndrome performed significantly 151 152 worse than the mean on the Vineland 3. This difference was expected due to the expression of the 153 NatA complex being ubiquitous across tissue types and playing a prominent role in development (Cheng et al., 2020; Liszczak et al., 2013; Lyon et al., 2023). When comparing performance by 154 155 sex, the females on average performed better than the males. Except for two participants with an 156 unknown inheritance pattern, the NAA15 pathogenic variants in the study cohort occurred denovo. Without evidence that the disease is maternally inherited, the difference in sex is 157 unexpected. A possible explanation could be that the NAA10 catalytic subunit of the NatA 158 complex is present on the X-chromosome (Bader et al., 2020; Lyon et al., 2023; Rope et al., 159

2011; Wu & Lyon, 2018). An increase in NAA10 dosage due to partial inactivation or a different 160 X-chromosome escape mechanism could partially offset the loss of active site conformational 161 162 change of NAA10 by NAA15 (Carrel & Willard, 2005; Cotton et al., 2013; Schultz et al., 2015; Tukiainen et al., 2017). The mechanism of NAA15-related neurodevelopmental syndrome 163 appearing to be a milder form of NAA10-related neurodevelopmental syndrome (Ogden 164 165 Syndrome) also provides some support for this hypothesis as the NAA10 catalytic unit has function outside of the NatA complex but is unable to acetylate NatA specific sites without 166 NAA15 (Cheng et al., 2020; Lyon et al., 2023; Makwana et al., 2024; Patel et al., 2024; Wu & 167 Lyon, 2018). However, the differences seen between sexes could also be due to small sample 168 sizes. Also, because of a lack of overlap between pathogenic variants in males and females, it 169 was not possible to do a direct comparison in function while controlling for sex. Further genetic 170 171 analysis and cognitive testing needs to be performed to determine if the differences between sexes are significant. 172

173 While females performed on average better than males across all adaptive domains, they performed significantly better on the motor domain and fine motor sub-domain portions of the 174 175 Vineland. Pathogenic variants in NAA15 have been associated with infant onset dystonia in a 176 few individuals (Cheng et al., 2018; Danti et al., 2024; Lyon et al., 2023; Yubero et al., 2022) which could be one reason why the cohort performed worse on motor tasks compared to the 177 norm population. However, there did not appear to be a difference between the sexes in the 178 referenced literature in how musculoskeletal manifestations presented. Furthermore, there is a 179 180 case where a NAA15 variant led to the adult-onset presentation of parkinsonism in a man presenting with other manifestations of NAA15-related neurodevelopmental syndrome such as 181 182 speech delay and intellectual disability (Straka et al., 2022). Continued genetic testing of

individuals with adult-onset dystonia and longitudinal monitoring of the current cohort for motor
disorders will be necessary to help determine if the difference in motor function between male
and females is due to chance, sex differences, or pathogenic variant type.

In aggregate, it is difficult to determine if there is an overall increase or decrease in 186 function over time in the participants. However, when comparing performance by sex, there is a 187 188 strong negative correlation between adaptive behavior in females and age. The decline with age has the strongest correlation between the coping, domestic, and fine motor sub-domains. With a 189 decline in fine motor capacity, it would stand to reason that an individual's ability to take care of 190 oneself (domestic) would also decline. However, both the domestic and fine motor sub-domains 191 have age restrictions leading to unequal administration of those portions of the exam creating an 192 193 even smaller population size to compare from (Farmer et al., 2020; Perry et al., 2009, p. 3). Furthermore, this decline in function contradicts the mild recovery in gross motor, personal-194 social, and language development function previously reported in the literature (Cheng et al., 195 196 2018, 2020; Tian et al., 2022). The decline in the coping sub-domain in the females could also be partially explained by the decrease in motor skills as there is evidence that similar motor declines 197 198 present in those with Autism Spectrum Disorder (ASD) have led to decreased coping and 199 socialization skills overall (Peyre et al., 2024; Raditha et al., 2023). The overall decline could also be partially explained by the upper half of the age range being from 11-17, a time period that 200 201 is traditionally tumultuous for adolescents due to various physical and hormonal changes, and is associated with increased stress for those with ASD (Cridland et al., 2014; Steward et al., 2018; 202 203 Tager-Flusberg & Kasari, 2013). Ultimately, follow up studies need to be performed on this 204 cohort to determine if they are continuing to show decline over time, if there is a bottoming out 205 effect, or if they begin to improve after navigating adolescence.

In contrast to the female decline over time, males when analyzed without NAA15-011 are 206 correlated to have a moderate increase in adaptive function with age. However, there are not 207 208 strong correlations present in any subsequent sub-domain analysis performed. Additionally, when including NAA15-011, there were no or small correlations between Vineland score and age. 209 Patient NAA15-011 was removed from the analysis due to their having performed much 210 211 more poorly than the rest of their cohort. They had a de novo p.Asn864Ser pathogenic variant 212 which is located in part of the region (residues 500-866 (Arnesen et al., 2010)) that interacts with the HYPK chaperone like protein which is important for NatA mediated acetylation and 213 Huntingtin protein aggregation (Arnesen et al., 2010; Ghosh & Ranjan, 2022; Miklánková et al., 214 215 2022; Wever et al., 2017). In addition to dysfunctional N-terminal acetylation being associated 216 with neurodevelopmental delay, abnormal Huntingtin protein aggregation also can lead to cognitive decline (Alberti & Hyman, 2021; Liu et al., 2023; Soltani Khaboushan et al., 2023). 217 The stark difference in function between NAA15-011 and the rest of the male cohort could be 218 219 explained by a substantial decrease in HYPK interaction. However, additional molecular testing would need to be conducted to be certain due to others in the cohort having mutations in this 220 221 cohort without having similar decreases in function. If abnormally aggregated Huntingtin protein 222 is associated with the steeper decline in function in patient NAA15-011, then future potential therapies that are effective in decreasing Huntingtin aggregation could possible rescue partial 223 224 function in these individuals (Im et al., 2015; Jain & Roy, 2023; Posey et al., 2018; Roth et al., 2023). The increase in age could also be the reason for the sharp decline in function as NAA15-225 226 011 is 10 years older than the second oldest male. However, additional longitudinal data needs to 227 be collected before definitive conclusions can be drawn.

228	The overall decrease in female adaptive behavior scores and the overall significantly
229	decreased male adaptive scores suggest that there is an urgent need for additional study of and
230	the development of interventions for NAA15-related neurodevelopmental disorder. Future
231	research should aim to complete the clinical timeline as best as possible. The present cohort was
232	only able to include individuals up to the age of \sim 20-24 years due to the rarity of the disease.
233	Without insight into the later stages of the disorder, it will be more difficult for clinicians to help
234	provide expectations of what to expect at different life stages to caregivers.
235	NAA15-related neurodevelopmental disorder is thought to be caused by
236	haploinsufficiency (Cheng et al., 2018; Tian et al., 2022) of which there are several therapies in
237	development that show promise in cell lines and animal models that have been used in similar
238	disorders such as Rett Syndrome and Angelman Syndrome to help restore protein function and
239	dosage to adequate levels (Albadri et al., 2017; Burbano et al., 2022; Hill & Meisler, 2021;
240	Milazzo et al., 2021; Palmieri et al., 2023; Protic et al., 2019). If interventions are implemented
241	early enough, it may be possible to restore protein function and improve adaptive behavior skills.
242	Conclusion
243	NAA15-related neurodevelopmental disorder is closely related to Ogden Syndrome
244	(NAA10-related neurodevelopmental disorder) due to both the NAA15 and NAA10 proteins
245	being part of the NatA complex. NAA15 plays a supporting role in complex with the NAA10
246	enzymatic subunit, with dysfunction leading to clinical manifestations in the heart,
247	musculoskeletal system, and brain. The pathogenic variants in NAA15-related
248	neurodevelopmental disorder are heterozygous, so there is still one functioning copy of NAA15,
249	meaning that the NatA complex can still form, but likely with haploinsufficiency. As such,
250	NAA15 pathogenic variants usually present less severely than those persons with pathogenic

251	NAA10 variants. Repeated Vineland-3 administration to a cohort of 26 patients with unique de
252	novo NAA15 pathogenic variants has revealed the natural history of the disease, showcasing an
253	overall decrease in adaptive functioning compared to the norm from infancy till adolescence.
254	Females tended to perform on average better than males. However, over time, females tended to
255	show a decline in function whereas males showed an increase. Additional natural history studies
256	need to be performed to understand disease outcomes outside of adolescence, difference between
257	sexes, and to identify outliers who perform significantly better or worse than their peers so
258	genetic or environmental factors influencing disease can be identified.
259	
260	Ethical Approval
261	Both oral and written patient consent were obtained for research and publication, with
262	approval of protocol #7659 for the Jervis Clinic by the New York State Psychiatric Institute -
263	Columbia University Department of Psychiatry Institutional Review Board.
264	
265	Funding
266	This work is supported by New York State Office for People with Developmental
267	Disabilities (OPWDD) and NIH NIGMS R35-GM-133408.
268	
269	Competing Interests
270	The authors declare that they have no competing interests or personal relationships that
271	could have influenced the work reported in this paper.
272	
273	

274 **References**

- 275 Aksnes, H., Hole, K., & Arnesen, T. (2015). Molecular, cellular, and physiological significance of N-
- terminal acetylation. *International Review of Cell and Molecular Biology*, *316*, 267–305.
- 277 https://doi.org/10.1016/bs.ircmb.2015.01.001
- 278 Albadri, S., Del Bene, F., & Revenu, C. (2017). Genome editing using CRISPR/Cas9-based knock-in
- approaches in zebrafish. *Methods (San Diego, Calif.)*, 121–122, 77–85.
- 280 https://doi.org/10.1016/j.ymeth.2017.03.005
- Alberti, S., & Hyman, A. A. (2021). Biomolecular condensates at the nexus of cellular stress, protein
- aggregation disease and ageing. *Nature Reviews. Molecular Cell Biology*, 22(3), 196–213.
- 283 https://doi.org/10.1038/s41580-020-00326-6
- 284 Arnesen, T., Starheim, K. K., Van Damme, P., Evjenth, R., Dinh, H., Betts, M. J., Ryningen, A.,
- 285 Vandekerckhove, J., Gevaert, K., & Anderson, D. (2010). The Chaperone-Like Protein HYPK
- 286 Acts Together with NatA in Cotranslational N-Terminal Acetylation and Prevention of Huntingtin
- Aggregation. *Molecular and Cellular Biology*, *30*(8), 1898–1909.
- 288 https://doi.org/10.1128/MCB.01199-09
- 289 Bader, I., McTiernan, N., Darbakk, C., Boltshauser, E., Ree, R., Ebner, S., Mayr, J. A., & Arnesen, T.
- 290 (2020). Severe syndromic ID and skewed X-inactivation in a girl with NAA10 dysfunction and a
- 291 novel heterozygous de novo NAA10 p.(His16Pro) variant—A case report. BMC Medical
- 292 *Genetics*, 21(1), 153. https://doi.org/10.1186/s12881-020-01091-1
- 293 Belbachir, N., Wu, Y., Shen, M., Zhang, S. L., Zhang, J. Z., Liu, C., Knollmann, B. C., Lyon, G. J., Ma,
- 294 N., & Wu, J. C. (2023). Studying Long QT Syndrome Caused by NAA10 Genetic Variants Using
- 295 Patient-Derived Induced Pluripotent Stem Cells. *Circulation*, 148(20), 1598–1601.
- https://doi.org/10.1161/CIRCULATIONAHA.122.061864
- Bienvenut, W. V., Sumpton, D., Martinez, A., Lilla, S., Espagne, C., Meinnel, T., & Giglione, C. (2012).
- 298 Comparative large scale characterization of plant versus mammal proteins reveals similar and

- 299 idiosyncratic N-α-acetylation features. *Molecular & Cellular Proteomics: MCP*, 11(6),
- 300 M111.015131. https://doi.org/10.1074/mcp.M111.015131
- 301 Burbano, L. E., Li, M., Jancovski, N., Jafar-Nejad, P., Richards, K., Sedo, A., Soriano, A., Rollo, B., Jia,
- 302 L., Gazina, E. V., Piltz, S., Adikusuma, F., Thomas, P. Q., Kopsidas, H., Rigo, F., Reid, C. A.,
- 303 Maljevic, S., & Petrou, S. (2022). Antisense oligonucleotide therapy for KCNT1 encephalopathy.
- 304 *JCI Insight*, 7(23), e146090. https://doi.org/10.1172/jci.insight.146090
- Carrel, L., & Willard, H. F. (2005). X-inactivation profile reveals extensive variability in X-linked gene
 expression in females. *Nature*, 434(7031), 400–404. https://doi.org/10.1038/nature03479
- 307 Cheng, H., Dharmadhikari, A. V., Varland, S., Ma, N., Domingo, D., Kleyner, R., Rope, A. F., Yoon, M.,
- 308 Stray-Pedersen, A., Posey, J. E., Crews, S. R., Eldomery, M. K., Akdemir, Z. C., Lewis, A. M.,
- 309 Sutton, V. R., Rosenfeld, J. A., Conboy, E., Agre, K., Xia, F., ... Lyon, G. J. (2018). Truncating
- 310 Variants in NAA15 Are Associated with Variable Levels of Intellectual Disability, Autism
- 311 Spectrum Disorder, and Congenital Anomalies. *American Journal of Human Genetics*, 102(5),
- 312 985–994. https://doi.org/10.1016/j.ajhg.2018.03.004
- 313 Cheng, H., Gottlieb, L., Marchi, E., Kleyner, R., Bhardwaj, P., Rope, A. F., Rosenheck, S., Moutton, S.,
- 314 Philippe, C., Eyaid, W., Alkuraya, F. S., Toribio, J., Mena, R., Prada, C. E., Stessman, H., Bernier,
- 315 R., Wermuth, M., Kauffmann, B., Blaumeiser, B., ... Lyon, G. J. (2020). Phenotypic and
- 316 biochemical analysis of an international cohort of individuals with variants in NAA10 and
- 317 NAA15. Human Molecular Genetics, 29(5), 877–878. https://doi.org/10.1093/hmg/ddz173
- 318 Cotton, A. M., Ge, B., Light, N., Adoue, V., Pastinen, T., & Brown, C. J. (2013). Analysis of expressed
- 319 SNPs identifies variable extents of expression from the human inactive X chromosome. *Genome* 320 *Biology*, *14*(11), R122. https://doi.org/10.1186/gb-2013-14-11-r122
- 321 Cridland, E. K., Jones, S. C., Caputi, P., & Magee, C. A. (2014). Being a girl in a boys' world:
- 322 Investigating the experiences of girls with autism spectrum disorders during adolescence. *Journal*
- 323 of Autism and Developmental Disorders, 44(6), 1261–1274. https://doi.org/10.1007/s10803-013-
- 324 1985-6

- 325 Danti, F. R., Sarmiento, I. J. K., Moloney, P. B., Colangelo, I., Graziola, F., Garavaglia, B., Zorzi, G.,
- 326 Mencacci, N. E., & Lubbe, S. J. (2024). Childhood-Onset Lower Limb Focal Dystonia Due to a
- NAA15 Variant: A Case Report. *Movement Disorders: Official Journal of the Movement Disorder Societv.* https://doi.org/10.1002/mds.29732
- 329 Deng, S., Magin, R. S., Wei, X., Pan, B., Petersson, E. J., & Marmorstein, R. (2019). Structure and
- 330 Mechanism of Acetylation by the N-Terminal Dual Enzyme NatA/Naa50 Complex. *Structure* 331 (*London, England: 1993*), 27(7), 1057-1070.e4. https://doi.org/10.1016/j.str.2019.04.014
- 332 Dikiy, I., & Eliezer, D. (2014). N-terminal Acetylation Stabilizes N-terminal Helicity in Lipid- and
- 333 Micelle-bound α-Synuclein and Increases Its Affinity for Physiological Membranes. *Journal of* 334 *Biological Chemistry*, 289(6), 3652–3665. https://doi.org/10.1074/jbc.M113.512459
- 335 Dörfel, M. J., Fang, H., Crain, J., Klingener, M., Weiser, J., & Lyon, G. J. (2017). Proteomic and genomic
- characterization of a yeast model for Ogden syndrome. *Yeast (Chichester, England)*, *34*(1), 19–37.
 https://doi.org/10.1002/yea.3211
- Dörfel, M. J., & Lyon, G. J. (2015). The biological functions of Naa10—From amino-terminal acetylation
 to human disease. *Gene*, 567(2), 103–131. https://doi.org/10.1016/j.gene.2015.04.085
- 340 Farmer, C., Adedipe, D., Bal, V. H., Chlebowski, C., & Thurm, A. (2020). Concordance of the Vineland
- Adaptive Behavior Scales, second and third editions. *Journal of Intellectual Disability Research: JIDR*, 64(1), 18–26. https://doi.org/10.1111/jir.12691
- Feng, J., Li, R., Yu, J., Ma, S., Wu, C., Li, Y., Cao, Y., & Ma, L. (2016). Protein N-terminal acetylation is
 required for embryogenesis in Arabidopsis. *Journal of Experimental Botany*, 67(15), 4779–4789.
 https://doi.org/10.1093/jxb/erw257
- Feng, J., & Ma, L. (2016). NatA is required for suspensor development in Arabidopsis. *Plant Signaling & Behavior*, *11*(10), e1231293. https://doi.org/10.1080/15592324.2016.1231293
- 348 Ghosh, D. K., & Ranjan, A. (2022). HYPK coordinates degradation of polyneddylated proteins by
- 349 autophagy. *Autophagy*, 18(8), 1763–1784. https://doi.org/10.1080/15548627.2021.1997053

350	Hill, S. F., & Meisler, M. H. (2021). Antisense Oligonucleotide Therapy for Neurodevelopmental
351	Disorders. Developmental Neuroscience, 43(3-4), 247-252. https://doi.org/10.1159/000517686
352	Huth, E. A., Zhao, X., Owen, N., Luna, P. N., Vogel, I., Dorf, I. L. H., Joss, S., Clayton-Smith, J., Parker,
353	M. J., Louw, J. J., Gewillig, M., Breckpot, J., Kraus, A., Sasaki, E., Kini, U., Burgess, T., Tan, T.
354	Y., Armstrong, R., Neas, K., Scott, D. A. (2023). Clinical exome sequencing efficacy and
355	phenotypic expansions involving anomalous pulmonary venous return. European Journal of
356	Human Genetics: EJHG, 31(12), 1430-1439. https://doi.org/10.1038/s41431-023-01451-4
357	Im, W., Ban, JJ., Chung, JY., Lee, ST., Chu, K., & Kim, M. (2015). Multidrug resistance protein 1
358	reduces the aggregation of mutant huntingtin in neuronal cells derived from the Huntington's
359	disease R6/2 model. Scientific Reports, 5(1), 16887. https://doi.org/10.1038/srep16887
360	Jain, S., & Roy, I. (2023). Aptamer Reduces Aggregation of Mutant Huntingtin and Rescues Proteostasis
361	Network in Non-Neuronal and Neuronal Cells. ACS Chemical Neuroscience, 14(12), 2385–2395.
362	https://doi.org/10.1021/acschemneuro.3c00226
363	Knorr, A. G., Schmidt, C., Tesina, P., Berninghausen, O., Becker, T., Beatrix, B., & Beckmann, R. (2019).
364	Ribosome-NatA architecture reveals that rRNA expansion segments coordinate N-terminal
365	acetylation. Nature Structural & Molecular Biology, 26(1), 35–39.
366	https://doi.org/10.1038/s41594-018-0165-y
367	Kuhns, K. J., Zhang, G., Wang, Z., & Liu, W. (2018). ARD1/NAA10 acetylation in prostate cancer.
368	Experimental & Molecular Medicine, 50(7), 1-8. https://doi.org/10.1038/s12276-018-0107-0
369	Le, MK., Vuong, H. G., Nguyen, T. T. T., & Kondo, T. (2023). NAA10 overexpression dictates distinct
370	epigenetic, genetic, and clinicopathological characteristics in adult gliomas. Journal of
371	Neuropathology and Experimental Neurology, 82(7), 650–658.
372	https://doi.org/10.1093/jnen/nlad037
373	Liszczak, G., Goldberg, J. M., Foyn, H., Petersson, E. J., Arnesen, T., & Marmorstein, R. (2013).
374	Molecular basis for N-terminal acetylation by the heterodimeric NatA complex. Nature Structural
375	& Molecular Biology, 20(9), 1098–1105. https://doi.org/10.1038/nsmb.2636

- Liu, L., Tong, H., Sun, Y., Chen, X., Yang, T., Zhou, G., Li, X.-J., & Li, S. (2023). Huntingtin Interacting
- 377 Proteins and Pathological Implications. *International Journal of Molecular Sciences*, 24(17),
- 378 13060. https://doi.org/10.3390/ijms241713060
- 379 Lyon, G. J., Vedaie, M., Beisheim, T., Park, A., Marchi, E., Gottlieb, L., Hsieh, T.-C., Klinkhammer, H.,
- 380 Sandomirsky, K., Cheng, H., Starr, L. J., Preddy, I., Tseng, M., Li, Q., Hu, Y., Wang, K.,
- 381 Carvalho, A., Martinez, F., Caro-Llopis, A., ... Herr-Israel, E. (2023). Expanding the phenotypic
- 382 spectrum of NAA10-related neurodevelopmental syndrome and NAA15-related
- neurodevelopmental syndrome. *European Journal of Human Genetics: EJHG*, *31*(7), 824–833.
- 384 https://doi.org/10.1038/s41431-023-01368-y
- 385 Makwana, R., Christ, C., Marchi, E., Harpell, R., & Lyon, G. J. (2024). Longitudinal Adaptive Behavioral
- 386 Outcomes in Ogden Syndrome by Seizure Status and Therapeutic Intervention. *medRxiv: The*
- 387 *Preprint Server for Health Sciences*, 2024.02.23.24303144.
- 388 https://doi.org/10.1101/2024.02.23.24303144
- 389 McTiernan, N., Støve, S. I., Aukrust, I., Mårli, M. T., Myklebust, L. M., Houge, G., & Arnesen, T. (2018).
- 390 NAA10 dysfunction with normal NatA-complex activity in a girl with non-syndromic ID and a de
- 391 novo NAA10 p.(V111G) variant—A case report. *BMC Medical Genetics*, 19(1), 47.
- 392 https://doi.org/10.1186/s12881-018-0559-z
- 393 Miklánková, P., Linster, E., Boyer, J.-B., Weidenhausen, J., Mueller, J., Armbruster, L., Lapouge, K., De
- 394 La Torre, C., Bienvenut, W., Sticht, C., Mann, M., Meinnel, T., Sinning, I., Giglione, C., Hell, R.,
- 395 & Wirtz, M. (2022). HYPK promotes the activity of the Nα-acetyltransferase A complex to
- determine proteostasis of nonAc-X2/N-degron-containing proteins. *Science Advances*, 8(24),
- 397 eabn6153. https://doi.org/10.1126/sciadv.abn6153
- 398 Milazzo, C., Mientjes, E. J., Wallaard, I., Rasmussen, S. V., Erichsen, K. D., Kakunuri, T., van der Sman,
- A. S. E., Kremer, T., Miller, M. T., Hoener, M. C., & Elgersma, Y. (2021). Antisense
- 400 oligonucleotide treatment rescues UBE3A expression and multiple phenotypes of an Angelman
- 401 syndrome mouse model. JCI Insight, 6(15), e145991. https://doi.org/10.1172/jci.insight.145991

- 402 Monestier, O., Landemaine, A., Bugeon, J., Rescan, P.-Y., & Gabillard, J.-C. (2019). Naa15 knockdown
- 403 enhances c2c12 myoblast fusion and induces defects in zebrafish myotome morphogenesis.
- 404 Comparative Biochemistry and Physiology. Part B, Biochemistry & Molecular Biology, 228, 61–
- 405 67. https://doi.org/10.1016/j.cbpb.2018.11.005
- 406 Myklebust, L. M., Van Damme, P., Støve, S. I., Dörfel, M. J., Abboud, A., Kalvik, T. V., Grauffel, C.,
- 407 Jonckheere, V., Wu, Y., Swensen, J., Kaasa, H., Liszczak, G., Marmorstein, R., Reuter, N., Lyon,
- 408 G. J., Gevaert, K., & Arnesen, T. (2015). Biochemical and cellular analysis of Ogden syndrome
- 409 reveals downstream Nt-acetylation defects. *Human Molecular Genetics*, 24(7), 1956–1976.
- 410 https://doi.org/10.1093/hmg/ddu611
- 411 Palmieri, M., Pozzer, D., & Landsberger, N. (2023). Advanced genetic therapies for the treatment of Rett
- 412 syndrome: State of the art and future perspectives. *Frontiers in Neuroscience*, 17, 1172805.
- 413 https://doi.org/10.3389/fnins.2023.1172805
- 414 Patel, R., Park, A. Y., Marchi, E., Gropman, A. L., Whitehead, M. T., & Lyon, G. J. (2024). Ophthalmic
- 415 Manifestations of NAA10-Related and NAA15-Related Neurodevelopmental Syndrome:
- 416 Analysis of Cortical Visual Impairment and Refractive Errors. *medRxiv: The Preprint Server for*
- 417 *Health Sciences*, 2024.02.01.24302161. https://doi.org/10.1101/2024.02.01.24302161
- 418 Perry, A., Flanagan, H. E., Dunn Geier, J., & Freeman, N. L. (2009). Brief report: The Vineland Adaptive
- 419 Behavior Scales in young children with autism spectrum disorders at different cognitive levels.
- 420 *Journal of Autism and Developmental Disorders*, *39*(7), 1066–1078.
- 421 https://doi.org/10.1007/s10803-009-0704-9
- 422 Pesz, K., Pienkowski, V. M., Pollak, A., Gasperowicz, P., Sykulski, M., Kosińska, J., Kiszko, M.,
- 423 Krzykwa, B., Bartnik-Głaska, M., Nowakowska, B., Rydzanicz, M., Sasiadek, M. M., & Płoski,
- 424 R. (2018). Phenotypic consequences of gene disruption by a balanced de novo translocation
- 425 involving SLC6A1 and NAA15. European Journal of Medical Genetics, 61(10), 596–601.
- 426 https://doi.org/10.1016/j.ejmg.2018.03.013

427	Peyre, H.,	Peries, M.,	Madieu, E	., David, A.,	Picot, MC.,	Pickles, A., &	Baghdadli, A. (2024).
	J / /	, , ,	,	, , , ,	, , ,	, ,		

- 428 Association of difficulties in motor skills with longitudinal changes in social skills in children
- 429 with autism spectrum disorder: Findings from the ELENA French Cohort. *European Child &*

430 *Adolescent Psychiatry*. https://doi.org/10.1007/s00787-023-02324-3

- 431 Posey, A. E., Ruff, K. M., Harmon, T. S., Crick, S. L., Li, A., Diamond, M. I., & Pappu, R. V. (2018).
- 432 Profilin reduces aggregation and phase separation of huntingtin N-terminal fragments by
- 433 preferentially binding to soluble monomers and oligomers. *Journal of Biological Chemistry*,
- 434 293(10), 3734–3746. https://doi.org/10.1074/jbc.RA117.000357
- 435 Protic, D., Salcedo-Arellano, M. J., Dy, J. B., Potter, L. A., & Hagerman, R. J. (2019). New Targeted
- 436 Treatments for Fragile X Syndrome. *Current Pediatric Reviews*, 15(4), 251–258.
- 437 https://doi.org/10.2174/1573396315666190625110748
- 438 Raditha, C., Handryastuti, S., Pusponegoro, H. D., & Mangunatmadja, I. (2023). Positive behavioral

439 effect of sensory integration intervention in young children with autism spectrum disorder.

440 *Pediatric Research*, 93(6), 1667–1671. https://doi.org/10.1038/s41390-022-02277-4

- 441 Ritter, A., Berger, J. H., Deardorff, M., Izumi, K., Lin, K. Y., Medne, L., & Ahrens-Nicklas, R. C. (2021).
- 442 Variants in NAA15 cause pediatric hypertrophic cardiomyopathy. *American Journal of Medical*443 *Genetics. Part A*, 185(1), 228–233. https://doi.org/10.1002/ajmg.a.61928
- 444 Rope, A. F., Wang, K., Evjenth, R., Xing, J., Johnston, J. J., Swensen, J. J., Johnson, W. E., Moore, B.,
- 445 Huff, C. D., Bird, L. M., Carey, J. C., Opitz, J. M., Stevens, C. A., Jiang, T., Schank, C., Fain, H.
- 446 D., Robison, R., Dalley, B., Chin, S., ... Lyon, G. J. (2011). Using VAAST to identify an X-linked
- disorder resulting in lethality in male infants due to N-terminal acetyltransferase deficiency.
- 448 American Journal of Human Genetics, 89(1), 28–43. https://doi.org/10.1016/j.ajhg.2011.05.017
- 449 Roth, J. R., Moraes, R. C. M. de, Xu, B. P., Crawley, S. R., Khan, M. A., & Melkani, G. C. (2023).
- 450 Rapamycin reduces neuronal mutant huntingtin aggregation and ameliorates locomotor
- 451 performance in Drosophila. *Frontiers in Aging Neuroscience*, 15.
- 452 https://doi.org/10.3389/fnagi.2023.1223911

- 453 Schultz, M. D., He, Y., Whitaker, J. W., Hariharan, M., Mukamel, E. A., Leung, D., Rajagopal, N., Nery,
- 454 J. R., Urich, M. A., Chen, H., Lin, S., Lin, Y., Jung, I., Schmitt, A. D., Selvaraj, S., Ren, B.,
- 455 Sejnowski, T. J., Wang, W., & Ecker, J. R. (2015). Human body epigenome maps reveal
- 456 noncanonical DNA methylation variation. *Nature*, *523*(7559), 212–216.
- 457 https://doi.org/10.1038/nature14465
- 458 Sidhu, M., Brady, L., Tarnopolsky, M., & Ronen, G. M. (2017). Clinical Manifestations Associated With
- 459 the N-Terminal-Acetyltransferase NAA10 Gene Mutation in a Girl: Ogden Syndrome. *Pediatric*
- 460 *Neurology*, *76*, 82–85. https://doi.org/10.1016/j.pediatrneurol.2017.07.010
- 461 Soltani Khaboushan, A., Moeinafshar, A., Ersi, M. H., Teixeira, A. L., Majidi Zolbin, M., & Kajbafzadeh,
- 462 A.-M. (2023). Circulating levels of inflammatory biomarkers in Huntington's disease: A
- 463 systematic review and meta-analysis. *Journal of Neuroimmunology*, 385, 578243.
- 464 https://doi.org/10.1016/j.jneuroim.2023.578243
- 465 Stessman, H. A. F., Xiong, B., Coe, B. P., Wang, T., Hoekzema, K., Fenckova, M., Kvarnung, M., Gerdts,
- 466 J., Trinh, S., Cosemans, N., Vives, L., Lin, J., Turner, T. N., Santen, G., Ruivenkamp, C., Kriek,
- 467 M., van Haeringen, A., Aten, E., Friend, K., ... Eichler, E. E. (2017). Targeted sequencing
- 468 identifies 91 neurodevelopmental-disorder risk genes with autism and developmental-disability
- 469 biases. *Nature Genetics*, 49(4), 515–526. https://doi.org/10.1038/ng.3792
- 470 Steward, R., Crane, L., Mairi Roy, E., Remington, A., & Pellicano, E. (2018). "Life is Much More
- 471 Difficult to Manage During Periods": Autistic Experiences of Menstruation. *Journal of Autism*472 *and Developmental Disorders*, 48(12), 4287–4292. https://doi.org/10.1007/s10803-018-3664-0
- 473 Støve, S. I., Blenski, M., Stray-Pedersen, A., Wierenga, K. J., Jhangiani, S. N., Akdemir, Z. C., Crawford,
- 474 D., McTiernan, N., Myklebust, L. M., Purcarin, G., McNall-Knapp, R., Wadley, A., Belmont, J.
- 475 W., Kim, J. J., Lupski, J. R., & Arnesen, T. (2018). A novel NAA10 variant with impaired
- 476 acetyltransferase activity causes developmental delay, intellectual disability, and hypertrophic
- 477 cardiomyopathy. *European Journal of Human Genetics: EJHG*, 26(9), 1294–1305.
- 478 https://doi.org/10.1038/s41431-018-0136-0

- 479 Straka, I., Švantnerová, J., Minár, M., Stanková, S., & Zech, M. (2022). Neurodevelopmental Gene-
- 480 Related Dystonia-Parkinsonism with Onset in Adults: A Case with NAA15 Variant. *Movement*
- 481 Disorders: Official Journal of the Movement Disorder Society, 37(9), 1955–1957.
- 482 https://doi.org/10.1002/mds.29125
- 483 Tager-Flusberg, H., & Kasari, C. (2013). Minimally Verbal School-Aged Children with Autism Spectrum
- 484 Disorder: The Neglected End of the Spectrum. *Autism Research : Official Journal of the*
- 485 *International Society for Autism Research*, 6(6), 10.1002/aur.1329.
- 486 https://doi.org/10.1002/aur.1329
- 487 Tian, Y., Xie, H., Yang, S., Shangguan, S., Wang, J., Jin, C., Zhang, Y., Cui, X., Lyu, Y., Chen, X., &
- 488 Wang, L. (2022). Possible Catch-Up Developmental Trajectories for Children with Mild
- 489 Developmental Delay Caused by NAA15 Pathogenic Variants. *Genes*, 13(3), 536.
- 490 https://doi.org/10.3390/genes13030536
- 491 Tukiainen, T., Villani, A.-C., Yen, A., Rivas, M. A., Marshall, J. L., Satija, R., Aguirre, M., Gauthier, L.,
- 492 Fleharty, M., Kirby, A., Cummings, B. B., Castel, S. E., Karczewski, K. J., Aguet, F., Byrnes, A.,
- 493 Lappalainen, T., Aviv Regev, Ardlie, K. G., Hacohen, N., & MacArthur, D. G. (2017). Landscape
- 494 of X chromosome inactivation across human tissues. *Nature*, *550*(7675), 244–248.
- 495 https://doi.org/10.1038/nature24265
- 496 Umlai, U.-K. I., Haris, B., Hussain, K., & Jithesh, P. V. (2021). Case Report: Phenotype-Gene Correlation
- 497 in a Case of Novel Tandem 4q Microduplication With Short Stature, Speech Delay and
- 498 Microcephaly. *Frontiers in Endocrinology*, *12*, 783235.
- 499 https://doi.org/10.3389/fendo.2021.783235
- 500 Van Damme, P. (2021). Charting the N-Terminal Acetylome: A Comprehensive Map of Human NatA
- 501 Substrates. *International Journal of Molecular Sciences*, *22*(19), 10692.
- 502 https://doi.org/10.3390/ijms221910692
- 503 Ward, T., Tai, W., Morton, S., Impens, F., Van Damme, P., Van Haver, D., Timmerman, E., Venturini, G.,
- 504 Zhang, K., Jang, M. Y., Willcox, J. A. L., Haghighi, A., Gelb, B. D., Chung, W. K., Goldmuntz,

505	E., Porter	, G. A., I	Lifton, R. P.,	Brueckner,	M., Yost,	H. J.,	Seidman.	J. G.	(2021)). Mechanisms
-----	------------	------------	----------------	------------	-----------	--------	----------	-------	--------	---------------

- 506 of Congenital Heart Disease Caused by NAA15 Haploinsufficiency. *Circulation Research*,
- 507 *128*(8), 1156–1169. https://doi.org/10.1161/CIRCRESAHA.120.316966
- 508 Weidenhausen, J., Kopp, J., Armbruster, L., Wirtz, M., Lapouge, K., & Sinning, I. (2021). Structural and
- 509 functional characterization of the N-terminal acetyltransferase Naa50. Structure (London,
- 510 *England: 1993*), 29(5), 413-425.e5. https://doi.org/10.1016/j.str.2020.12.004
- 511 Weyer, F. A., Gumiero, A., Lapouge, K., Bange, G., Kopp, J., & Sinning, I. (2017). Structural basis of
- 512 HypK regulating N-terminal acetylation by the NatA complex. *Nature Communications*, *8*,
- 513 15726. https://doi.org/10.1038/ncomms15726
- 514 Wong, W.-R., Brugman, K. I., Maher, S., Oh, J. Y., Howe, K., Kato, M., & Sternberg, P. W. (2019).
- 515 Autism-associated missense genetic variants impact locomotion and neurodevelopment in

516 Caenorhabditis elegans. *Human Molecular Genetics*, 28(13), 2271–2281.

- 517 https://doi.org/10.1093/hmg/ddz051
- 518 Wu, Y., & Lyon, G. J. (2018). NAA10-related syndrome. *Experimental & Molecular Medicine*, 50(7), 1–
 519 10. https://doi.org/10.1038/s12276-018-0098-x
- 520 Yubero, D., Martorell, L., Nunes, T., Lyon, G. J., & Ortigoza-Escobar, J. D. (2022). Neurodevelopmental
- Gene-Related Dystonia: A Pediatric Case with NAA15 Variant. *Movement Disorders: Official Journal of the Movement Disorder Society*, *37*(11), 2320–2321.
- 523 https://doi.org/10.1002/mds.29241
- 524 Zhang, Z.-Y., Zhang, J.-L., Zhao, L.-X., Yang, Y., Guo, R., Zhou, N., Liu, Y.-R., & Zheng, G.-P. (2020).
- 525 NAA10 promotes proliferation of renal cell carcinoma by upregulating UPK1B. *European*
- 526 *Review for Medical and Pharmacological Sciences*, *24*(22), 11553–11560.
- 527 https://doi.org/10.26355/eurrev 202011 23796
- 528 Zhao, J. J., Halvardson, J., Zander, C. S., Zaghlool, A., Georgii-Hemming, P., Månsson, E., Brandberg,
- 529 G., Sävmarker, H. E., Frykholm, C., Kuchinskaya, E., Thuresson, A.-C., & Feuk, L. (2018).
- 530 Exome sequencing reveals NAA15 and PUF60 as candidate genes associated with intellectual

- 531 disability. American Journal of Medical Genetics. Part B, Neuropsychiatric Genetics: The
- 532 *Official Publication of the International Society of Psychiatric Genetics*, 177(1), 10–20.

533 https://doi.org/10.1002/ajmg.b.32574

- 534 Zhu, R., Chen, M., Luo, Y., Cheng, H., Zhao, Z., & Zhang, M. (2024). The role of N-acetyltransferases in
- 535 cancers. *Gene*, 892, 147866. https://doi.org/10.1016/j.gene.2023.147866

537 Tables:

538 Table 1. NAA15 Pathogenic Variant Breakdown by Sex

Pathogenic Variant	Males	Females
c.0692-1G>T in intron 6	1	
c.139+2T>A: IVS2+2T>A in		1
intron 2		
c.2057-2A>G (splice site)		1
p.Ala585Thr	1	
p.Ala678Cysfs*3	1	
p.Arg027Gly	1	
p.Arg549X	1	
p.Arg782X		1
p.Asn113Ser		1
p.Asn362MetfsX36	1	
p.Asn623Ilefs*	1	
p.Asn864Ser	1	
p.Asp0335Glufs*7	1	
p.Cys484Arg		1
p.Gln0621*	1	
p.Glu173*	2	
p.Glu173NfsX54	1	
p.His080Argfs*17	2	
p.intronic		1
p.Lys0628Glu	1	
p.splice site variant	1	
p.Trp0083X		1
p.Tyr682*	1	
p.Y333*		1
Total: 24	18	8

540

541 Table 2. Vineland ABC and Domain Standard Scores by Sex

	Communication	Daily Living Skills	Social	Motor	ABC Standard
Total Average	61.2	68.9	70.9	69.1	66.7
SD	26.4	20.1	22.7	17.4	21.1
Female					
Average	65.3	72.3	75.3	85.2	71.5
SD	30.9	24.0	24.8	11.0	25.9
Male Average	59.4	67.5	68.9	61.1	64.5
SD	23.8	17.9	21.4	14.2	18.3
t test	0.53	0.51	0.43	0.003*	0.35

542 *Asterisk denotes significance (p<.05)

543

- 544
- 545

546

547 Table 3. Correlation Coefficients of Vineland Domain Standard Scores vs Age

		Daily Living			
	Communication	Skills	Social	Motor	ABC
Female Correlation					
Coefficient	-0.69	-0.83	-0.72	-0.61	-0.76
Male Correlation					
Coefficient	0.10	0.00	-0.09	0.46	-0.01
Male Correlation					
Coefficient					
excluding					
ID:NAA15-011	0.45	0.54	0.32	0.46	0.47

	Communi	cation Sub-D	omain	Daily Living Skills Sub-Domain			
	Receptive	Expressive	Written	Personal	Domestic	Community	
Total	7.97	8.03	8.63	8.28	10.6	8.66	
Average							
SD	4.86	5.48	4.78	5.16	3.46	3.35	
Female	9.58	8.42	7.50	9.50	10.8	7.90	
Average							
SD	5.19	6.33	5.20	5.39	4.47	3.30	
Male	7.26	7.85	9.08	7.74	10.5	8.96	
Average							
SD	4.53	5.05	4.52	4.96	2.95	3.32	
t-test	0.18	0.77	0.39	0.34	0.81	0.41	

549 Table 4. Vineland Sub-Domain Standard Scores by Domain and Sex

550

	Socialization Sub-Domain			Moto Don	r Sub- nain	Maladaptive Trait Sub- Domain		
	Interpersonal Relations	Play and Leisure	Coping	Gross Motor	Fine Motor	Internalizing	Externalizing	
Total	9.18	9.15	10.7	9.22	9.33	18.8	17.3	
Average								
SD	4.85	5.16	2.99	3.21	4.41	3.61	2.91	
Female	9.92	9.92	10.8	10.8	14.0	18.4	16.7	
Average								
SD	5.62	5.33	3.01	1.67	3.61	3.37	2.76	
Male	8.85	8.81	10.6	8.42	7.00	19.0	17.6	
Average								
SD	4.42	5.05	2.98	3.48	2.52	3.71	2.94	
t-test	0.54	0.55	0.84	0.15	0.003*	0.74	0.53	

551 *Asterisk denotes significance (p < .05) between Male and Female scores

552

554 Table 5. Correlation Coefficients of Vineland Sub-Domain Standard Scores vs Age

	Comn	nunication Domain	Sub-	Daily Living Skills Sub- Domain			
	Receptive	Expressive	Written	Personal	Domestic	Community	
Female Correlation Coefficient	-0.70	-0.54	-0.64	-0.70	-0.79	-0.57	
Male Correlation Coefficient	0.15	0.19	-0.22	0.24	-0.52	-0.17	
Male Correlation excluding							
NAA15-011	0.46	0.51	0.06	0.58	-0.12	0.28	

555

	Socialization Sub-Domain			Motor Sub-Domain	
	Interpersonal	Play and		Gross	Fine
	Relations	Leisure	Coping	Motor	Motor
Female Correlation Coefficient	-0.47	-0.69	-0.95	-0.16	-0.80
Male Correlation Coefficient	0.03	0.01	-0.27	0.18	-0.09
Male Correlation Excluding NAA15-011	0.38	0.31	0.09	0.18	-0.09

Maladaptive Trait Sub-Domain		
Internalizing	Externalizing	
0.66		0.62
0.42		0.21
0.42		-0.34
	Maladap Internalizing 0.66 0.42 0.42	Maladaptive Trait Sub-DomainInternalizingExternalizing0.660.420.420.42





Figure 2. ABC Standard Score vs Age at time of Vineland Administration by Sex. Each point on the graph represents a single participants ABC standard score at a given time point. Points connected by lines show change in single participant scores over time.

563



Figure 3. Vineland Domain Score vs Age at time of Vineland Assessment. Points in black represent Males. Points in Blue represent Females. Each point on the graph represents a single participants Vineland Domain standard score at a given time point. Points connected by lines show change in single participant scores over time.

565

566

567





Figure 4. Vineland Sub-Domain Score vs Age at time of Assessment. A) Communication Sub-Domain Standard Scores. B) Daily Living Skills Sub-Domain Standard Scores. C) Socialization Sub-Domain Standard Scores. D) Motor Sub-Domain Standard Scores E) Maladaptive Trait Standard Scores. Points in black represent Males. Points in Blue represent Females. Each point on the graph represents a single participants sub-domain standard score at a given time point. Points connected by lines show change in single participant scores over time.