

### Supplementary materials

**Table S1.** Engagement in avoidance and recommended behaviors due to the novel coronavirus outbreak

	Wuhan (N=510) N(%)	Shanghai (N=501) N(%)	P-value
Last week, I have ... because of the outbreak			
Avoided eating out	488 (95.7)	487 (97.2)	0.193
Avoided using public transport	487 (95.5)	467 (93.2)	0.117
Reduced going to public areas	479 (93.9)	479 (95.6)	0.229
Rescheduled travel plan	468 (91.8)	451 (90.0)	0.335
Increased surface cleaning	456 (89.4)	390 (77.8)	<0.001
Maintained better indoor ventilation	499 (97.8)	488 (97.4)	0.648

**Table S2.** Perceived severity compared with SARS by age groups

	Wuhan		P-value	Shanghai		P-value
	Aged 18-39 (N=250)	Aged 40 and above (N=260)		Aged 18-39 (N=251)	Aged 40 and above (N=250)	
	N (%)	N (%)		N (%)	N (%)	
Severity compared with SARS			0.136		<0.001	
Much lower	6 (2.4)	7 (2.7)		8 (3.2)	4 (1.6)	
Lower	25 (10.0)	41 (15.8)		52 (20.7)	23 (9.2)	
Even	75 (30.0)	56 (21.5)		81 (32.3)	55 (22.0)	
Higher	76 (30.4)	82 (31.5)		50 (19.9)	77 (30.8)	
Much higher	68 (27.2)	74 (28.5)		60 (23.9)	91 (36.4)	

**Table S3.** Perception variables associated with face mask wearing and hands washing, odds ratios (95% CI)


	Increased frequency in...		Longer handwashing duration	Purchased goggles
	Face mask wearing	Immediate handwashing		
GAD score (continuous)	1.0 (0.9 - 1.0)	1.0 (1.0 - 1.0)	1.0 (1.0 - 1.0)	1.0 (1.0 - 1.0)
Correct perceived efficacy	1.4 (0.5 - 4.0)	1.9** (1.2 - 3.2)	1.2 (0.8 - 1.7)	1.2 (0.8 - 1.8)
Perceived risk	0.9 (0.7 - 1.3)	1.0 (0.9 - 1.3)	1.2** (1.1 - 1.4)	1.1 (0.9 - 1.3)
Perceived severity	0.8 (0.6 - 1.0)	0.8 (0.7 - 1.0)	0.8* (0.7 - 1.0)	1.1 (0.9 - 1.3)
Relative transmissibility to SARS (in 2003)	1.2 (0.8 - 2.0)	1.3* (1.0 - 1.7)	1.3** (1.1 - 1.6)	1.0 (0.8 - 1.1)
Relative harm to body to SARS (in 2003)	1.1 (0.8 - 1.4)	0.8* (0.7 - 1.0)	1.0 (0.9 - 1.2)	1.2* (1.0 - 1.4)
Received and read information brochure	1.3 (0.5 - 3.4)	3.2** (1.4 - 7.1)	1.2 (0.7 - 1.9)	0.9 (0.5 - 1.5)
Sufficient information	1.2 (0.8 - 1.9)	1.1 (0.9 - 1.4)	1.2* (1.0 - 1.4)	0.9 (0.8 - 1.1)
Confused about information reliability	1.1 (0.8 - 1.4)	1.0 (0.8 - 1.2)	1.1 (0.9 - 1.2)	1.0 (0.9 - 1.2)
Self-confidence	1.0 (0.6 - 1.6)	1.2 (0.9 - 1.6)	1.0 (0.8 - 1.1)	1.0 (0.8 - 1.2)

Notes: Each column presents a separate multivariate logistic regression result. \* for significance at the 5% level; \*\* for significance at the 1% level.

**Table S4.** Perceived efficacy of various behaviors

	Wuhan N (%)	Shanghai N (%)	P-value
Washing hands more frequently			0.274
Effective	435 (85.5)	414 (82.6)	
Even	71 (13.9)	80 (16.0)	
Ineffective	3 (0.6)	7 (1.4)	
Wearing a face mask			0.314
Effective	471 (92.5)	450 (89.8)	
Even	35 (6.9)	47 (9.4)	
Ineffective	3 (0.6)	4 (0.8)	
Reducing going out			0.547
Effective	491 (96.5)	481 (96.0)	
Even	14 (2.8)	18 (3.6)	
Ineffective	4 (0.8)	2 (0.4)	
Washing mouth with salty water			<0.001
Effective	153 (30.1)	124 (24.8)	
Even	231 (45.4)	187 (37.3)	
Ineffective	125 (24.6)	190 (37.9)	
Having vitamin C or radix isatidis (a traditional Chinese medicine)			0.002
Effective	81 (15.9)	89 (17.8)	
Even	256 (50.3)	198 (39.5)	
Ineffective	172 (33.8)	214 (42.7)	

**Figure S1.** Participant recruitment flowcharts



**Figure S2.** Top reasons for wearing a face mask and not always wearing in the past week

