Supplementary material: Diffuse white matter abnormality in very preterm infants reflects reduced brain network efficiency

Julia E. Kline, PhD¹, Venkata Sita Priyanka Illapani, MS¹, Hailong Li, PHD¹, Lili He, PhD^{1,2}, Nehal A. Parikh, DO, MS^{*1,2,3}

¹Perinatal Institute, Cincinnati Children's Hospital Medical Center, Cincinnati, OH ²Department of Pediatrics, University of Cincinnati College of Medicine, Cincinnati, OH ³Center for Perinatal Research, The Research Institute at Nationwide Children's Hospital, Columbus, OH

*Corresponding author's contact information: Nehal A. Parikh, DO, MS Professor of Pediatrics Cincinnati Children's Hospital 3333 Burnet Ave, MLC 7009 Cincinnati, OH 45229 (513) 636-7584 (Business) (513) 803-0969 (Fax) Nehal.Parikh@cchmc.org



Supplementary Figure S1: Blue circles represent nodes in which brain injury score was negatively correlated with local efficiency (top row) and clustering coefficient (bottom row) at term equivalent age in the very preterm brain, after correcting for postmenstrual age at MRI, gestational age, sex, structural brain abnormality score, and birth hospital and applying false discovery rate correction. The same networks are shown from a sagittal (left) and an axial (right) orientation. There were no significant positive correlations.