

**Article type:** Research Letter

**Title:** Highlighting Educational Gaps in Hairstyling Practices amongst Dermatologists and Trainees

**Authors:** Erinolaoluwa F. Araoye, BS\*<sup>1</sup>, Taylor A. Jamerson, BA\*<sup>2</sup>; Lu Yin, BA<sup>3</sup>, Kristen Lo Sicco MD<sup>3</sup>; Crystal Aguh, MD<sup>4</sup>

<sup>1</sup>Johns Hopkins School of Medicine, Baltimore, Maryland

<sup>2</sup>University of Michigan Medical School, Ann Arbor, Michigan

<sup>3</sup>The Ronald O. Perelman Department of Dermatology, New York University Grossman School of Medicine, New York, New York

<sup>4</sup>Department of Dermatology, Johns Hopkins School of Medicine, Baltimore, Maryland

\*These authors contributed equally and are co-first authors.

**Corresponding author information:**

Crystal Aguh, MD

Johns Hopkins School of Medicine

Department of Dermatology

10710 Charter Drive

Shared Suite 420

Columbia, MD 21044

Email: [cagi1@jhmi.edu](mailto:cagi1@jhmi.edu)

ORCID ID: 0000-0002-3590-9682

**Funding sources:** None

**Declarations of interests:** None

**COI:** The authors have no conflicts of interests.

**Manuscript word count:** 600

**Table/Figures:** 2

**Presentations:** None

**Keywords:** Hair practices, hair styling, ethnic hair, dermatology training, medical education

## **Abstract:**

**Background** Hairstyling practices are associated with the development and/or exacerbation of various forms of alopecia. Exposure to various hairstyling practices ranges but is often insufficient in current dermatologic textbooks and training curricula. We therefore conducted a survey to establish dermatologists understanding of hairstyling practices, particularly those that have been implicated in alopecia.

**Methods:** A 34-item anonymous, electronic survey was distributed by email to 291 board-certified dermatologists and dermatology residents across the US between August 2020 and February 2021. Responses were rated on a 10-point scale to identify physician confidence in various styling practices

**Results:** Black providers were more confident in both the knowledge and counseling of all hair practices (chemical straightening, heat styling, braiding, weaving, and wigs) compared to non-Black providers ( $p < 0.001$ ), with the exception of counseling patients on hair dyes for which no significant difference was found ( $p = 0.337$ ). Female providers were only more likely to indicate confidence in knowledge regarding different heat styling methods and hair dyes, and counseling of heat styling methods compared to male providers (OR 15.72,  $p < 0.001$ ; OR 2.47,  $p = 0.022$ ; OR 3.78,  $p = 0.001$  respectively) across all hair practices surveyed. Overall, 63.8% of providers reported that the majority of their knowledge on hair practices was from personal experience as opposed to formal training.

**Limitations:** This survey is limited by its response rate and the inability to characterize non-responders due to anonymity.

**Conclusion:** Our study highlights educational gaps in dermatologic training on hair practices, especially those more common among Black patients. Interestingly, the majority of provider

knowledge came from personal experience rather than dermatologic training emphasizing the need for formalized curricula to enhance understanding among all dermatology providers.

## Introduction

Hair loss and scalp dermatoses are among the most common concerns of patients presenting to dermatology clinics, many developing or becoming exacerbated as a direct result of damaging hair practices.<sup>1</sup> To provide competent care, dermatologists must have a fundamental understanding of the common hairstyling practices across different ethnic and racial groups to inform adequate counseling and treatment recommendations.<sup>2</sup> Exposure to various hairstyling practices ranges but is often insufficient in current dermatologic textbooks and training curricula.<sup>2,3</sup> We therefore conducted a survey to establish dermatologists' understanding of hairstyling practices, particularly those that have been implicated in alopecia.<sup>4,5</sup>

## Methods

A 34-item anonymous, electronic survey was developed on Research Electronic Data Capture (REDCap) following approval from the Johns Hopkins Institutional Review Board. The survey was distributed by email to 291 board-certified dermatologists and dermatology residents across the US between August 2020 and February 2021. Responses were rated on a 10-point scale and dichotomized into two groups: not confident (confidence scale ratings 0-5) and confident (confidence scale ratings 6-10). Racial and gender differences in responses were compared using two-sided  $\chi^2$  with statistical significance determined at  $P < 0.05$ . A logistic regression was used to control for the effect of ethnicity and gender on comparisons.

## Results

Of the 291 dermatology providers surveyed, 149 responses were received (response rate, 51.2%). Demographics and characteristics of respondents are summarized in Table 1.

Black providers were more confident in both the knowledge and counseling of all hair practices (chemical straightening, heat styling, braiding, weaving, and wigs) compared to non-Black providers ( $p < 0.001$ ), with the exception of counseling patients on hair dyes for which no significant difference was found ( $p = 0.337$ ) (see Table 2). Black providers were also more comfortable providing counseling on hair practices to patients with a discordant hair type (81.8% vs 50.5%;  $p < 0.001$ ). Though there was no statistical difference in the proportion of Black and non-Black providers who consider treatment vehicles based on hair type, non-Black providers indicated less confidence in selecting the optimal agent (69.4% vs 97%;  $p = 0.001$ ).

Controlling for ethnicity, female providers were only more likely to indicate confidence in knowledge regarding different heat styling methods and hair dyes, and counseling of heat styling methods compared to male providers (OR 15.72,  $p < 0.001$ ; OR 2.47,  $p = 0.022$ ; OR 3.78,  $p = 0.001$  respectively) across all hair practices surveyed.

Overall, 63.8% of providers reported that the majority (>50%) of their knowledge on hair practices was from personal experience. Only 13.4% indicated that their knowledge came almost exclusively (>90%) from dermatologic training and 87.2% of providers indicated they would have benefited from formal training on hair practices.

## Discussion

Our study highlights educational gaps in dermatologic training on hair practices, especially those more common among Black patients.<sup>4</sup> This likely accounts for our finding that non-Black

providers are less confident discussing hairstyling practices with patients who have discordant hair types compared to Black providers. The largest knowledge gaps were identified in braiding practices and types of dreadlocks, hairstyles highly associated with higher prevalence of traction alopecia.<sup>4</sup> Though Black providers indicated more overall confidence in knowledge of hairstyling practices, the majority of provider knowledge came from personal experience rather than dermatologic training.

### *Limitations*

This survey is limited by its response rate and the inability to characterize non-responders due to anonymity.

### Conclusions

Hairstyling practices are associated with the development and/or exacerbation of various forms of alopecia. An understanding of these practices is critically important to competent dermatologic care. To promote standardized management and equitable care for all patients presenting with hair and scalp concerns, we strongly encourage integration of this information into standard dermatology training through curricular textbooks, resident and conference lectures, and when possible, patient-centered learning.

No. of respondents (%)
------------------------

References:

1. Davis SA, Narahari S, Feldman SR, et al. Top dermatologic conditions in patients of color: an analysis of nationally representative data. *Journal of Drugs in Dermatology : JDD*. 2012 Apr;11(4):466-473.
2. Aguh C, Okoye GA (eds): *Fundamentals of Ethnic Hair: The Dermatologist's Perspective*. 2017. Cham, Switzerland, Springer International Publishing, 2017
3. Nijhawan RI, Jacob SE, Woolery-Lloyd H. Skin of color education in dermatology residency programs: does residency training reflect the changing demographics of the United States? *J Am Acad Dermatol*. 2008 Oct;59(4):615-8. doi: 10.1016/j.jaad.2008.06.024.
4. Haskin A, Aguh C. All hairstyles are not created equal: What the dermatologist needs to know about black hairstyling practices and the risk of traction alopecia (TA). *J Am Acad Dermatol*. 2016 Sep;75(3):606-611. doi: 10.1016/j.jaad.2016.02.1162.
5. Gathers RC, Jankowski M, Eide M, Lim HW. Hair grooming practices and central centrifugal cicatricial alopecia. *J Am Acad Dermatol*. 2009 Apr;60(4):574-8. doi: 10.1016/j.jaad.2008.10.064.

Sex	
Male	44 (29.5)
Female	104 (69.8)
Prefer not to answer	1 (0.7)
Race/ethnicity	
Caucasian	70 (46.9)
Black	32 (21.5)
Asian (East, South, Pacific Islander)	25 (16.8)
Hispanic	12 (8.1)
Multi-racial/ethnic	5 (3.4)
Prefer not to answer	5 (3.4)
Level of training	
PGY1	4 (2.7)
PGY2	10 (6.7)
PGY3	9 (6.0)
PGY4	11 (7.4)
Board certified dermatologist	115 (77.2)
Region of dermatologic practice	
Northeast	72 (48.3)
Midwest	9 (6.0)
South	57 (38.3)
West	11 (7.4)
Geographic location of dermatologic practice	
Urban	114 (76.5)
Rural	1 (0.7)
Suburban	31 (20.8)
Unsure	3 (2.0)
Dermatology practice setting	
Private practice	18 (12.1)
Academic	116 (77.9)
Mixed	7 (4.7)
None of the above	8 (5.4)
Years in practice (average)	11.39

**Table 1.**  
**Demographics and characteristics of survey respondents**



**Table 2. Percentage of providers who indicated confidence in their knowledge and counseling of hairstyling practices, including ethnic and gender comparisons**

	% confident (all respondents)	Black providers vs non- Black providers*		Female providers vs male providers**	
		OR	P-value	OR	P-value
<b>Physician confidence in <i>knowledge</i> of hairstyling practices</b>					
Overall	60.4	8.95	0.001	1.91	0.101
Chemical straightening	36.9	7.09	<0.001	1.52	0.332
Heat styling methods	79.2	1	<0.001	15.72	<0.001
Types of hair dyes	48.3	1.48	0.337	2.47	0.022
Types of hair braiding	37.6	61.07	<0.001	0.86	0.759
Types of dreadlocks	26.2	42.90	<0.001	2.71	0.144
Weaving installation practices	36.9	24.63	<0.001	1.71	0.269
Wig types	28.2	11.85	<0.001	1.73	0.291
<b>Physician confidence in patient <i>counseling</i> on hairstyling practices</b>					
Overall	53.0	13.85	<0.001	1.34	0.479
Chemical straightening	36.2	8.74	<0.001	1.49	0.364
Heat styling methods	64.4	10.76	0.002	3.78	0.001

Types of hair dyes	41.6	2.64	0.019	1.59	0.241
Types of hair braiding	33.6	51.53	<0.001	0.75	0.568
Types of dreadlocks	26.2	54.65	<0.001	1.26	0.787
Weaving installation practices	33.6	18.66	<0.001	0.88	0.276
Wig types	26.8	14.31	<0.001	1.11	0.839

\* = Adjusted for sex

\*\* = Adjusted for ethnicity