

The magnitude of exercise-induced hypoalgesia in female can be improved and correlated with endogenous pain modulation following 4 weeks of treadmill running

Zi-Han Xu¹ (Corresponding author)

¹School of Sport Medicine and Rehabilitation, Beijing Sport University
Beijing, China
2290259311@qq.com

ORCID ID : 0000-0002-2090-3528

Nan An¹

¹School of Sport Medicine and Rehabilitation, Beijing Sport University
Beijing, China
616455768@qq.com

ORCID ID : 0000-0001-9197-4465

Shuang Xu²

²Division of Surgery and Interventional Science, University College London
London, UK
zemrehab@163.com

ORCID ID: 0009-0008-7642-6397

Ru-Yun Wang¹

¹School of Sport Medicine and Rehabilitation, Beijing Sport University
Beijing, China
2021011262@bsu.edu.cn

ORCID ID: 0009-0004-6264-0895

Xu, Z.H., An, N., and Xu, S. contributed equally to this manuscript.

Funding

This study is self-funded

Data availability statement

Data available on request from the authors

Disclosure statement

The authors declare no actual or potential conflicts of interest that could influence this study.

Abstract

Objective: We aimed to investigate changes in pain perception, acute exercise-induced hypoalgesia (EIH), and endogenous pain modulation responses following 4-week treadmill running exercises of different intensities in female.

Methods: Forty-two female participants included in this study performed exercise interventions 12 times within 4 weeks. All participants were randomly assigned to two experimental groups (TRL, low-intensity treadmill running and TRH, high-intensity treadmill running). All participants performed a single treadmill running session at different intensities based on their target heart rate (THR). A running assessment was administered to every participant one week before implementing the exercise interventions. The magnitudes of EIH, conditioned pain modulation (CPM), and temporal summation (TS) responses following long-term treadmill running were assessed. Pressure pain thresholds (PPT) or mechanical pain thresholds (MPT) were also determined following long-term treadmill running.

Results: Treadmill running only induced acute EIH responses, with all pre-running PPT and MPT remaining unaltered. However, EIH responses were significantly positively and negatively correlated with running sessions in the TRL and TRH groups, respectively.

Conclusions: A 4-week low-intensity treadmill running improved acute EIH response by enhancing endogenous pain modulation in healthy females. Future studies should consider sex, behavior, and physiological factors to provide a comprehensive understanding of the changes in EIH following long-term exercises.

Keywords: exercise-induced hypoalgesia; endogenous pain modulation; treadmill running; training-induced hypoalgesia; moderate-intensity exercise; pain

Introduction

Acute reduction of pain perception in the body following a single bout of exercise, commonly called exercise-induced hypoalgesia (EIH), has been widely confirmed in healthy individuals and some patients with pain[1]. Usually, both the global aerobic[2] and local resistance exercise[3] with a certain intensity and duration can temporarily increase various pain thresholds (pressure pain thresholds, PPT or mechanical pain thresholds, MPT) and enhance emotional well-being[4]. However, the EIH response may be weakened (absence of hypoalgesia or hyperalgesia) in older adults [5] or patients with painful conditions[6] and contribute to the impairment of endogenous pain modulation [7] (pain sensitization or pain-related psychological syndrome) in these individuals.

Exercise has been recommended as a non-pharmacological intervention and overall health promotion for various patients with pain and older adults, and the attenuation of chronic pain syndrome and improvement of pain-related behavior following long-term exercise training, also known as training-induced hypoalgesia (TIH) have been reported in many studies [8-10]. However, there is no clear evidence proving that the magnitude of the analgesic effects following a single bout of exercise can be improved or restored by long-term training in healthy individuals or patients with pain. [11]

In both healthy individuals and patients with pain, the EIH magnitude is affected by conditioned pain modulation (CPM) [12, 13] or temporal summation (TS) [14], which refers to the function of endogenous pain modulation and usually changes in individuals with sensitization of pain perception. In healthy individuals, pain perception can be inhibited or facilitated by descending control of the midbrain [15] and cortex [16] when the thalamus [17] receives certain inputs from peripheral nociceptors such as C fibers, which might also be activated by exercise with sufficient loads, leading to EIH. Therefore, it is important to understand whether endogenous pain modulation can be improved through long-term exercise.

Additionally, the EIH magnitude can be modulated by the intensity of exercise [18]; high-intensity exercises often exacerbate pain in both human and rodent studies [19, 20], while moderate-intensity training increases the pain threshold in many conditions [21, 22]. Previous studies [23, 24] have shown that the relationship between EIH and exercise intensity is an inverted U-shaped curve in healthy individuals. Considering the possibility of pain exacerbation following high-intensity exercise [25], the long-term influence on pain perception and endogenous pain modulation may also differ between high- and moderate-intensity exercises.

Therefore, we aimed to compare the long-term effects of high- and moderate-intensity exercise on EIH and endogenous pain modulation in healthy individuals following 4-week high- and moderate-intensity exercises. We measured the effects of EIH in every exercise session and changes in CPM and TS responses before and after the 4-week training session. We hypothesized that (1) both high- and moderate-intensity exercises might elicit EIH responses (increase in PPT or MPT) in every exercise session, (2) the magnitude of EIH and CPM responses might gradually

improve following long-term moderate-intensity exercise with the attenuation of TS response, and (3) the magnitude of EIH might be correlated with the CPM and TS responses.

Methods

This study was approved by the Sports Science Experimental Ethics Committee of the Beijing Sport University (approval number: 2023023H) and was registered in the Chinese Clinical Trial Registry (registration number: ChiCTR2300074367).

Study design

Altogether, 42 healthy participants included in this study performed exercise interventions 12 times within 4 weeks. All participants provided written informed consent. Demographic data and baseline measurements (resting heart rate [HR_{rest}], PPT, MPT, and CPM responses) were collected. The maximum heart rate (HR_{max}) was estimated using the formula [26]: $HR_{max} = 202.5 - 0.53 * \text{age}$ and the reserved heart rate (HRR) was calculated as $HRR = HR_{max} - HR_{rest}$. Real-time HR was collected and recorded via the HR belt worn by the participants during running. The first running session was performed 48 h after baseline measurements to avoid potential long-lasting analgesic effects of the CPM test.

All participants were randomly assigned to two experimental groups (TRL, low-intensity treadmill running and TRH, high-intensity treadmill running). Randomized sequences were generated using Excel software.

The TRL and TRH groups performed low-intensity and high-intensity treadmill running with 55% and 70% HRR, respectively. Running speed was determined in accordance with the target heart rate (THR) during baseline measurements. All participants performed a single exercise session once a day, three times per week for 4 weeks (Figure 1).




Figure 1. Flowchart of the experiment

Participants

Based on previous studies [27], long-term aerobic exercise-induced effect size on PPT changes ranging from 0.20 to 0.38 was 0.20. Our study utilized G*Power software with an effect size of 0.38, an alpha level of 0.05, and a power of 0.80. Thus, a minimum sample size of 38 participants across the two groups was determined.

Forty-six healthy female students (aged 18–30 years) from Beijing Sports University were included in this study, of whom 42 were finally enrolled. The exclusion criteria were (1) had pain-related pathological or psychological syndrome within 3 months; (2) had injury history of lower extremities within 1 year; (3) had potential or confirmed heart disease, or recovered from a heart disease <1 year ago, (4) failed to maintain or tolerate the exercise intensity during the long-term treadmill running interventions; (5) showed serious exertion or fatigue in 24 h after any exercise session; (6) showed intolerable pain during the pain perception test; (7) currently menstruating; and (8) had exercises or training experiences previously.

Procedures

All participants performed 12 treadmill running sessions within 4 weeks at different intensities based on their THR. The participants wore an HR belt to monitor and record real-time HR during the test and running sessions. A running assessment was administered to every participant one week before implementing the exercise interventions. This assessment involved a progressive increase in speed until the THR was reached. Subsequently, the predetermined speed for each individual was established at the commencement of running. During the running session, the running speed was adjusted at any time according to the participant's heart rate changes.

Outcome measures

Outcome measures were assessed at multiple time points, where the PPT-arm, PPT-leg, and MPT were recorded 5 min before and 10 min after each running session. The CPM and TS responses were evaluated at baseline (48 h and 24 h before and after the first and last running sessions, respectively). All testing locations were marked with a sterile waterproof marker to ensure consistency in the repeated measures. The testing angle of the algometer was carefully adjusted perpendicular to the skin.

PPT

The PPT was evaluated using a quantitative sensory testing protocol [28] with a handheld pressure algometer (Baseline Dolorimeter, Fabrication Enterprises, USA) equipped with a 1 cm² metal probe. Pressure was applied at a rate of 0.5 kg/s over the right side of the two muscle groups: the extensor carpus radialis (PPT arm) and peroneus longus (PPT leg). The participants were instructed to indicate their perceived pain intensity using a visual analog scale (VAS) ranging from 0 to 100 cm. When the participants reported a pain intensity of 30 out of 100 cm (Pain30) during

pressure application, the pressure thresholds were recorded as PPT values.

MPT

The MPT was evaluated using a quantitative sensory testing protocol [29] with the handheld algometer equipped with a needle probe. Pressure was applied at 0.1 kg/s over the left side of the extensor carpus radialis. The participants were instructed to indicate their perceived pain intensity using the VAS ranging from 0 to 100 cm. When the participants reported a pain intensity of 30 out of 100 cm (Pain30) during pressure application, the pressure thresholds were recorded as MPT values.

CPM

The CPM response was measured using a quantitative sensory testing protocol [28], specifically, the cold pressor procedure. Pressure was applied as the test stimulation, and cold-water immersion served as the conditioned stimulation. Participants first received pressure stimulation at the ipsilateral extensor carpus radialis, and the PPT was recorded as a test stimulus when the pain intensity reached Pain30. Subsequently, participants were instructed to immerse the contralateral hand into cold water at 8°C for 1 min. The PPT at Pain30 was reassessed when the participants withdrew their hands from the immersion. The difference between the two PPTs was recorded in response to the CPM.

TS

The TS response was measured using a quantitative sensory testing protocol [29]. The needle probe of the algometer was applied to the left side of the extensor carpus radialis at an intensity of 1.25 times the participants' MPT. Subsequently, the mechanical stimulations were repeated 10 times at 0.5 Hz (1-second stimulus following a 1-second interval). Subsequently, the participants were instructed to report the pain perception of the first and last stimulations using the VAS score, and the differences in scores between these two mechanical stimulations were recorded as a response to the TS.

Statistical analysis

The normality of all data was assessed using the Shapiro–Wilk test. Differences in baseline data (height, weight, HR_{rest}, PPT, MPT, CPM, and TS) between both groups were analyzed using an independent t-test. The differences in the PPT and MPT values between pre- and post-running in each session were calculated as EIH responses, including EIH-A for changes in the PPT-arm, EIH-L for changes in the PPT-leg, and EIH-M for changes in the MPT.

A two-way repeated measures analysis of variance (ANOVA) was used to determine the differences between the two groups over time (running sessions), and independent t-tests (without post hoc comparisons) were applied to examine the EIH values of the PPT and MPT. An independent t-test was also applied for between-group comparison of the CPM and TS responses.

Linear regression analysis was used to investigate whether there was a linear correlation between the running sessions and EIH-A, EIH-L, and EIH-M values. The relationships between the CPM

and TS, EIH-A, EIH-L, and EIH-M values after the running intervention were analyzed using the Pearson correlation method. All statistical analyses were performed using SPSS Version 21.0, and a significance level of $p < 0.05$ was applied to all tests.

Results

Baseline characteristics

Four participants were excluded from this study because of myofascial pain syndrome that occurred one month before the experiments. Of the forty-two participants enrolled in this study, 20 in the TRL group completed 12 low-intensity running sessions, and 18 in the TRH group completed 12 high-intensity running sessions. Additionally, four participants withdrew from the study because of the onset of menstruation, failure to finish all running sessions, and loss to follow-up. No significant differences were observed in baseline characteristics between the groups ($p > 0.05$, Table 1).

Table 1. Baseline measurement (M±SD)¹

	Group TRL (n=20)	Group TRH (n=18)	P
PPT-arm (kg/cm ²)	2.19±0.27	2.13±0.30	0.532
PPT-leg (kg/cm ²)	3.83±0.42	3.98±0.48	0.324
MPT (kg/cm ²)	0.57±0.10	0.62±0.16	0.248
CPM (kg/cm ²)	0.64±0.30	0.58±0.23	0.545
TS score (cm)	31.99±6.40	28.42±6.10	0.087
Age (y)	21.30±2.77	21.22±1.92	0.921
Height (cm)	166.05±3.89	165.56±4.55	0.720
Weight (kg)	55.90±9.13	55.89±5.97	0.997

All data were presented as mean +/- standard deviation (M±SD)

¹: 1-way ANOVA, significant difference was set by $p \leq 0.05$.

Changes in EIH-A following running sessions

The two-way repeated-measures ANOVA revealed significant effects ($F=15.465$, $p < 0.001$) for the running sessions involving the EIH-A, indicating that the 4-week running intervention significantly increased global EIH responses in all participants. The interaction effect ($F=28.347$, $p < 0.001$) between running intensity and time for the EIH-A was also significant. However, independent t-tests showed that the EIH-A in the TRL group was significantly higher ($p < 0.001$) than that in the TRH group, which indicated that only the global EIH in the TRL group improved after the 4-week treadmill running. In addition, all pre-running PPT-arm values remained unaltered, indicating that long-term running may not change the baseline level of the PPT-arm (Figures 2 and 3).




Figure 2. Changes in PPT of arms following running sessions
 All data are presented as mean and standard deviation; PPT=pressure pain threshold;
 HRR=reserved heart rate




Figure 3. Changes in EIH-A following running sessions
 All data were presented as mean and standard deviation; PPT=pressure pain threshold;
 EIH=exercise-induced hypoalgesia; EIH-A=EIH value of PPT-arms; HRR=reserved heart rate
 *: PPT in TRL group significantly higher than TRH group

Changes in EIH-L following running sessions

Two-way repeated-measures ANOVA revealed significant main effects ($F=3.639$, $p<0.001$) for the running sessions involving the EIH-L, indicating that the 4-week running intervention significantly increased local EIH responses in all participants. The interaction effect ($F=34.274$, $p<0.001$) between running intensity and time for the EIH-L was also significant. However, independent t-tests showed that EIH-L in the TRL group was significantly higher ($p<0.001$) than that in the TRH group, indicating that only the local EIH in the TRL group improved after the 4-week treadmill running. All pre-running PPT-leg values remained unaltered, indicating that

long-term running may not change the baseline level of the PPT-leg (Figures 4 and 5).




Figure 4. Changes in PPT of legs following running sessions
All data were presented as mean and standard deviation; PPT=pressure pain threshold;
HRR=reserved heart rate




Figure 5. Changes in EIHL following running sessions
All data were presented as mean and standard deviation; PPT=pressure pain threshold;
EIH=exercise-induced hypoalgesia; EIHL= EIH value of PPT-legs; HRR=reserved heart rate
*: PPT in TRL group significantly higher than TRH group

Changes in EIHL following running sessions

The two-way ANOVA revealed significant main effects ($F=4.288$, $p=0.01$) for the running sessions involving the EIHL, which indicated that the 4-week running intervention significantly decreased EIHL responses in all participants. The interaction effect ($F=7.445$, $p<0.001$) between running intensity and time for the EIHL was also significant. Additionally, independent t-tests showed that the EIHL in the TRL group was significantly higher ($p<0.001$) than that in the TRH group in the first to third running sessions and significantly lower ($p<0.001$) than that in the TRH

group from the 9th to 12th sessions. This result indicated that the EIH-M in the TRH group gradually decreased following the 4-week treadmill running, whereas the EIH-M in the TRL group was unaltered during the intervention. Additionally, all pre-running MPT values remained unaltered, indicating that long-term running may not change the baseline level of the MPT (Figures 6 and 7).




Figure 6. Changes in MPT following running sessions

All data were presented as mean and standard deviation; MPT=mechanical pain threshold; HRR=reserved heart rate




Figure 7. Changes in EIH-M following running sessions

All data were presented as mean and standard deviation; MPT=mechanical pain threshold; EIH=exercise-induced hypoalgesia; EIH-M=EIH value of MPT; HRR=reserved heart rate

#: MPT in TRH group significantly higher than TRL group

*: MPT in TRL group significantly higher than TRH group

Relationship between running sessions and EIH magnitudes

Linear regression analysis showed a significant positive correlation between running sessions and EIH-A ($p<0.001$) and EIH-L ($p<0.001$) magnitudes in the 4-week low-intensity running intervention group. Figure 8 shows the regression formulae.

$$\text{EIH-A (low-intensity running)} = 0.564 + 0.084 * \text{running sessions}$$

$$\text{EIH-L (low-intensity running)} = 0.717 + 0.071 * \text{running sessions}$$




Figure 8. Linear regression analyses between running sessions and EIH magnitudes
EIH-A= EIH value of PPT-arms; EIH-L= EIH value of PPT-legs; EIH-M=EIH value of MPT;

Linear regression analysis also revealed a significant negative correlation between running sessions and the EIH-A ($p<0.001$), EIH-L ($p<0.001$), and EIH-M ($p<0.001$) magnitudes during the 4-week high-intensity running intervention. Figure 9 shows the regression formulae.

$$\text{EIH-A (high-intensity running)} = 0.515 - 0.013 * \text{running sessions}$$

$$\text{EIH-L (high-intensity running)} = 0.760 - 0.037 * \text{running sessions}$$

$$\text{EIH-M (high-intensity running)} = 0.217 - 0.015 * \text{running sessions}$$




Figure 9. Linear regression analyses between running sessions and EIH magnitudes
EIH-A= EIH value of PPT-arms; EIH-L= EIH value of PPT-legs; EIH-M=EIH value of MPT;

Changes in CPM and TS following running sessions

Independent t-tests revealed significant between-group differences in the CPM ($p < 0.001$) and TS responses ($p < 0.001$) after the 4-week treadmill running intervention. The CPM responses in the TRL group were significantly higher than those in the TRH group. In contrast, the TS scores of the TRL group were significantly lower than those of the TRH group, which showed no significant changes before and after the 4-week running intervention (Figure 10).




Figure 10. Changes in CPM after 4-week running

All data were presented as mean/standard deviation; CPM=conditioned pain modulation; HRR=reserved heart rate

*: CPM in TRL group significantly higher than TRH group




Figure 11. Changes in TS after 4-week running

All data were presented as mean/standard deviation; TS=temporal summation; HRR=reserved heart rate

#: TS in TRH group significantly higher than TRL group

Relationship between endogenous pain tests and EIH magnitudes

Pearson correlation analysis showed that there were significant positive relationships between CPM values and EIH-A ($r=0.830$, $p<0.001$), EIH-L ($r=0.866$, $p<0.001$), and EIH-M ($r=0.551$, $p<0.001$) magnitudes. There were also significant negative relationships between TS values and EIH-A ($r=-0.773$, $p<0.001$), EIH-L ($r=-0.714$, $p<0.001$), and EIH-M ($r=-0.419$, $p=0.008$) magnitudes (Figures 12 and 13).




Figure 12: Relationship between CPM and EIH magnitudes

CPM=conditioned pain modulation; EIH-A= EIH value of PPT-arms; EIH-L= EIH value of PPT-legs; EIH-M=EIH value of MPT.




Figure 13: Relationship between TS and EIH magnitudes

TS=temporal summation; EIH-A= EIH value of PPT-arms; EIH-L= EIH value of PPT-legs; EIH-M=EIH value of MPT.

Discussion

We aimed to investigate changes in pain perception, EIH, and endogenous pain modulation responses following 4-week treadmill running exercises of different intensities in females. Our results revealed the following: First, low- and high-intensity running may only induce short-term analgesic effects, including improvements in global and local PPT and MPT, within 24 h or less. Second, acute EIH responses following a running session varied according to the type of pain

perception and exercise intensity, where the EIH-A and EIH-L following low-intensity running were significantly increased along with exercise time, and the EIH-A and EIH-L following high-intensity running slightly decreased after the exercise intervention. Third, the EIH-M following high-intensity running significantly decreased with exercise time, and the EIH-M following low-intensity running remained unaltered. Finally, the function of endogenous pain inhibition was enhanced, and facilitation was decreased following the 4-week low-intensity running exercise, showing positive and negative correlations with EIH responses, respectively.

The baseline pain perception threshold is relatively constant in healthy individuals [30] and may only be affected by activated or impaired endogenous pain modulation rather than long-term exercise training. Recent studies have shown that a 24-week high-intensity interval training [31] and a 20-week resistant band exercise [32] have no significant effects on the PPT in healthy individuals. Tesarz et al. [33] investigated baseline pain perceptions in athletes and normally active individuals and observed that differences in pain thresholds between the groups were not significant.

However, as a response to endogenous pain modulation, EIH following exercise may change after a long-term exercise intervention. Song et al.[11] suggested that exercise training induces physiological changes leading to improved EIH. Ohlman et al.[5] observed a greater EIH response in individuals who performed moderate physical activity per week than in sedentary controls. Hansen et al.[34] also observed that the PPT and EIH in healthy individuals significantly increased after a 7-week military training. However, evidence from randomized trials with a pre-test-post-test design remains limited.

Exercise with a sufficient load can induce a short-term EIH response, whereas exercise with low or moderate intensity may elicit greater analgesic effects than high-intensity or exhaustive exercise [23, 24]. Running at low- or moderate-intensity may activate non-noxious C fibers [35] via repeated muscle contractions, induce descending inhibition, upregulate 5-HT receptors in the brainstem [36], and attenuate pressure or thermal pain perception. High-intensity exercise may trigger noxious [37] and non-noxious C fibers and potentially induce descending facilitation [38] with limited EIH responses. Additionally, the upregulation of cannabinoids and opioids expression following high-intensity exercise may decrease the perception of mechanical stimuli. [39]

Therefore, we hypothesized that long-term exercise may induce plastic changes in endogenous pain modulation. Low- or moderate-intensity exercise may enhance the central descending inhibition function and increase acute EIH responses with an increase in CPM responses. Lemley et al.[12] investigated EIH in healthy individuals and observed that those with greater CPM were more likely to experience greater EIH. Naugle et al.[40] also observed that healthy adults who self-reported increased total physical activity exhibited reduced TS and greater CPM.

In contrast, high-intensity exercise may induce the adaptation of endocannabinoids and opioid modulation, decreasing the EIH of the MPT with unaltered baseline pain perception. For example, athletes experiencing high-intensity training showed a partially decreased EIH response than did healthy controls. Siebers et al.[39] observed a downregulation of endocannabinoid levels

following long-term running training. However, differential changes in the PPT, MPT, and other pain perception measurements represent various pain modulation pathways.

Our study had several limitations. First, the indicators of the pain tests were limited. For instance, adding heat pain detection thresholds might provide a more complete description of the changes in pain perception. Second, the intervention period for running exercise was relatively short. Future studies should investigate the long-term (> 6 weeks) effects on pain perception and modulation following various types of exercises. Finally, all the participants in this study were female. Considering the potential sex differences in endogenous pain modulation function and exercise behaviors, future studies should consider sex, behavior, and physiological factors to provide a comprehensive understanding of the changes in EIH following long-term exercise.

Conclusion

In summary, a 4-week low-intensity treadmill running improved acute EIH responses by enhancing endogenous pain modulation in healthy females. CPM and TS may be correlated with EIH and changed after exercise training, indicating that treadmill running may induce TIH through functional changes in endogenous pain modulation. However, baseline pain thresholds may remain unaltered and may not be affected by long-term exercise interventions.

References

1. Rice, D., et al., *Exercise-Induced Hypoalgesia in Pain-Free and Chronic Pain Populations: State of the Art and Future Directions*. J Pain, 2019. **20**(11): p. 1249-1266.
2. Hviid, J.T., J.B. Thorlund, and H.B. Vaegter, *Walking increases pain tolerance in humans: an experimental cross-over study*. Scand J Pain, 2019. **19**(4): p. 813-822.
3. Vaegter, H.B., et al., *Exercise-Induced Hypoalgesia After Isometric Wall Squat Exercise: A Test-Retest Reliability Study*. Pain Med, 2019. **20**(1): p. 129-137.
4. Brellenthin, A.G., et al., *Psychosocial Influences on Exercise-Induced Hypoalgesia*. Pain Med, 2017. **18**(3): p. 538-550.
5. Ohlman, T., et al., *Physical Activity Levels Predict Exercise-induced Hypoalgesia in Older Adults*. Med Sci Sports Exerc, 2018. **50**(10): p. 2101-2109.
6. Vaegter, H.B., et al., *Impaired exercise-induced hypoalgesia in individuals reporting an increase in low back pain during acute exercise*. Eur J Pain, 2021.
7. Fingleton, C., K.M. Smart, and C.M. Doody, *Exercise-induced Hypoalgesia in People With Knee Osteoarthritis With Normal and Abnormal Conditioned Pain Modulation*. Clin J Pain, 2017. **33**(5): p. 395-404.
8. Wewege, M.A. and M.D. Jones, *Exercise-Induced Hypoalgesia in Healthy Individuals and People With Chronic Musculoskeletal Pain: A Systematic Review and Meta-Analysis*. J Pain, 2021. **22**(1): p. 21-31.
9. Belavy, D.L., et al., *Pain sensitivity is reduced by exercise training: Evidence from a systematic review and meta-analysis*. Neurosci Biobehav Rev, 2021. **120**: p. 100-108.
10. Tan, L., et al., *Does aerobic exercise effect pain sensitisation in individuals with*

- musculoskeletal pain? A systematic review. BMC Musculoskelet Disord*, 2022. **23**(1): p. 113.
11. Song, J.S., et al., *Can we improve exercise-induced hypoalgesia with exercise training? An overview and suggestions for future studies. Phys Ther Sport*, 2023. **63**: p. 67-72.
 12. Lemley, K.J., S.K. Hunter, and M.K. Bement, *Conditioned pain modulation predicts exercise-induced hypoalgesia in healthy adults. Med Sci Sports Exerc*, 2015. **47**(1): p. 176-84.
 13. Alsouhibani, A. and M. Hoeger Bement, *Impaired conditioned pain modulation was restored after a single exercise session in individuals with and without fibromyalgia. Pain Rep*, 2022. **7**(3): p. e996.
 14. Vaegter, H.B., G. Handberg, and T. Graven-Nielsen, *Isometric exercises reduce temporal summation of pressure pain in humans. Eur J Pain*, 2015. **19**(7): p. 973-83.
 15. Ossipov, M.H., *The perception and endogenous modulation of pain. Scientifica (Cairo)*, 2012. **2012**: p. 561761.
 16. Ong, W.Y., C.S. Stohler, and D.R. Herr, *Role of the Prefrontal Cortex in Pain Processing. Mol Neurobiol*, 2019. **56**(2): p. 1137-1166.
 17. You, H.J., J. Lei, and A. Pertovaara, *Thalamus: The 'promoter' of endogenous modulation of pain and potential therapeutic target in pathological pain. Neurosci Biobehav Rev*, 2022. **139**: p. 104745.
 18. Hoffman, M.D., et al., *Intensity and duration threshold for aerobic exercise-induced analgesia to pressure pain. Arch Phys Med Rehabil*, 2004. **85**(7): p. 1183-7.
 19. Damsgard, E., et al., *Activity-related pain in patients with chronic musculoskeletal disorders. Disabil Rehabil*, 2010. **32**(17): p. 1428-37.
 20. Yokoyama, T., et al., *Muscle fatigue increases the probability of developing hyperalgesia in mice. J Pain*, 2007. **8**(9): p. 692-9.
 21. Lazaridou, A., et al., *Impact of daily yoga-based exercise on pain, catastrophizing, and sleep amongst individuals with fibromyalgia. J Pain Res*, 2019. **12**: p. 2915-2923.
 22. Newcomb, L.W., et al., *Influence of preferred versus prescribed exercise on pain in fibromyalgia. Med Sci Sports Exerc*, 2011. **43**(6): p. 1106-13.
 23. Niwa, Y., et al., *Effects of Exercise-Induced Hypoalgesia at Different Aerobic Exercise Intensities in Healthy Young Adults. J Pain Res*, 2022. **15**: p. 3615-3624.
 24. Zi-Han, X., et al., *Modulation of pain perceptions following treadmill running with different intensities in females. Physiol Rep*, 2023. **11**(18): p. e15831.
 25. Sluka, K.A. and L.A. Rasmussen, *Fatiguing exercise enhances hyperalgesia to muscle inflammation. Pain*, 2010. **148**(2): p. 188-97.
 26. Lach, J., et al., *HR Max Prediction Based on Age, Body Composition, Fitness Level, Testing Modality and Sex in Physically Active Population. Front Physiol*, 2021. **12**: p. 695950.
 27. Kolle, T., W. Alt, and D. Wagner, *Effects of a 12-week home exercise therapy program on pain and neuromuscular activity in patients with patellofemoral pain syndrome. Arch Orthop Trauma Surg*, 2020. **140**(12): p. 1985-1992.
 28. Kovacevic, M., et al., *Test-retest reliability of pressure pain threshold and heat pain threshold as test stimuli for evaluation of conditioned pain modulation. Neurophysiol Clin*, 2021.
 29. Mailloux, C., et al., *Within-session test-retest reliability of pressure pain threshold and mechanical temporal summation in healthy subjects. PLoS One*, 2021. **16**(1): p. e0245278.
 30. SL, B. and K. I., *Cognitive mediators of pain perception and tolerance. J Pers Soc Psychol*, 1991. **61**(3): p. 504-510.

31. A, M., et al., *Body Image Relates to Exercise-Induced Antinociception and Mood Changes in Young Adults: A Randomized Longitudinal Exercise Intervention*. Int J Environ Res Public Health, 2020. **17**(18): p. 6801.
32. M, M., et al., *Self-administered physical exercise training as treatment of neck and shoulder pain among military helicopter pilots and crew: a randomized controlled trial*. . BMC Musculoskelet Disord, 2017. **18**(1): p. 147.
33. Tesarz, J., et al., *Pain perception in athletes compared to normally active controls: a systematic review with meta-analysis*. Pain, 2012. **153**(6): p. 1253-1262.
34. Hansen, S., et al., *Modulation of Exercise-Induced Hypoalgesia Following an Exercise Intervention in Healthy Subjects*. Pain Med, 2020. **21**(12): p. 3556-3566.
35. Adreani, C.M., J.M. Hill, and M.P. Kaufman, *Responses of group III and IV muscle afferents to dynamic exercise*. J Appl Physiol (1985), 1997. **82**(6): p. 1811-7.
36. Bobinski, F., et al., *Role of brainstem serotonin in analgesia produced by low-intensity exercise on neuropathic pain after sciatic nerve injury in mice*. Pain, 2015. **156**(12): p. 2595-2606.
37. Nagi, S.S. and D.A. Mahns, *C-tactile fibers contribute to cutaneous allodynia after eccentric exercise*. J Pain, 2013. **14**(5): p. 538-48.
38. Sluka, K.A., et al., *Exercise-induced pain requires NMDA receptor activation in the medullary raphe nuclei*. Med Sci Sports Exerc, 2012. **44**(3): p. 420-7.
39. Siebers, M., S.V. Biedermann, and J. Fuss, *Do Endocannabinoids Cause the Runner's High? Evidence and Open Questions*. Neuroscientist, 2022: p. 10738584211069981.
40. Naugle, K.M. and J.L. Riley, 3rd, *Self-reported physical activity predicts pain inhibitory and facilitatory function*. Med Sci Sports Exerc, 2014. **46**(3): p. 622-9.

Acknowledgments

We would like to thank Tian-Rui Wu, Zheng-Quan Shi, Meng-Fei Lei, Zhao-Xia Zhou, Hua-Lian Tang, and all the researchers who provided assistance and advice during our experiments. And we would also like to thank Editage (www.editage.cn) for English language editing.

Author Contribution

Zi-Han Xu, Nan An and Shuang Xu conceived and designed research, Zi-Han Xu, Nan An, and Ru-Yun Wang performed experiments, Zi-Han Xu and Shuang Xu analyzed data. Zi-Han Xu, Nan An, Shuang Xu and Ru-Yun Wang interpreted results of experiments. Zi-Han Xu prepared Tables and Figures and drafted manuscript. All authors edited and revised manuscript drafts and approved final manuscript.