

## **The Mental Health of Healthcare Staff during the COVID-19 Pandemic: It Depends on How Much They Work and Their Age**

**Xingzi Xu**

Doctoral Student, Tsinghua University, Beijing, China

**Stephen X. Zhang\***

Associate Professor, University of Adelaide, Adelaide, Australia

**Asghar Afshar Jahanshahi**

Associate Professor, Pontificia Universidad Católica del Perú, Peru

**Jizhen Li**

Professor, Tsinghua University, Beijing, China

**Afsaneh Bagheri**

Associate Professor, Faculty of Entrepreneurship, University of Tehran, Iran

**Khaled Nawaser**

Assistant Professor, Arvandan Non-profit Higher Education Institute, Iran

**\*Corresponding author:** Stephen X. Zhang, [stephen.x.zhang@gmail.com](mailto:stephen.x.zhang@gmail.com); Phone: +61 8831 39310; Address: 9-28 Nexus10 Tower, 10 Pulteney St, Adelaide SA 5000, Australia

## **The Mental Health of Healthcare Staff during the COVID-19 Pandemic: It Depends on How Much They Work and Their Age**

### **ABSTRACT**

#### **Background:**

Healthcare staff are the forefront of fight against COVID-19 and they are under enormous pressure due to the fast growth in the number and severity of infected cases. This creates their mental issues such as distress, depression and anxiety. Exploring healthcare staff's mental health during the pandemic contributes to improving their persistence in the growing challenges created by COVID-19 and enabling effective management of their mental health.

#### **Methods:**

An online survey of 280 healthcare staff in all the 31 provinces of Iran was conducted during April 5–20, 2020. The survey assessed staff's distress, depression and anxiety in the COVID-19 pandemic.

#### **Results:**

Nearly a third of healthcare staff suffered from distress, depression and anxiety. Females and more educated healthcare staff were more likely to experience distress. Compared to personnel who did not have COVID-19, those who were unsure whether they had COVID-19 were more likely to experience distress and depression. The number of COVID-19 cases among the staff's colleagues or friends positively predicted their anxiety. Compared to radio technologists, doctors were less likely to experience distress and anxiety. Technicians and obstetrics experienced less anxiety. Analysis the interaction between weekly working days and age of the staff indicated the chance of experiencing distress and depression varied greatly by working days among younger but not older healthcare staff.

#### **Conclusion:**

Exploring the potential predictors of mental health issues assists healthcare organizations to develop approaches to prevent and decrease the issues among staff

and improve their task performance in sever crises such as COVID-19, so they can stay motivated in the COVID-19 pandemic. It is useful to identify more working characteristic predictors for healthcare staff at different ages.

**Funding:**

This work was supported by Tsinghua University-INDITEX Sustainable Development Fund (No. TISD201904).

**Keywords:** Healthcare staff; Mental health; Working days; Age; COVID-19; Pandemic; Iran

## 1. Introduction

Healthcare staff have been under enormous physical and mental pressures since the beginning of the COVID-19 pandemic. In the pandemic, healthcare staff face tremendous workloads, long and irregular working hours, and shortages of personal protective equipment (PPE) [1]. They also experience emotional and moral dilemmas from being isolated from their families during the pandemic [2, 3]. All of these factors may eventually lead to mental health issues [4, 5]. For instance, one nurse committed suicide because of the overload stress in Italy [6]. “Staff are being exposed to high levels of personal risk, long hours in difficult environments clad in PPE, and also the possibility of something known as moral injury, which is the distressing awareness you may feel when you know you can't meet all the needs of the people you are trying to care for” [7]. While such issues are acute in many countries [8, 9], this paper aims to specify some unique predictors of healthcare staff's mental health disorders in Iran as the second country that experienced COVID-19 and the virus spread in all of the country very fast.

In predicting mental health issues created and/or developed by COVID-19, this current research focused on socio-demographic [10, 11] and clinical predictors of healthcare staffs [12]. This survey is one of the very first attempts that investigated working characteristic predictors of healthcare staff's mental health during the COVID-19 pandemic. Furthermore, this study is the first, to the best of our knowledge, to examine comprehensive predictors of healthcare staff's mental health by measuring socio-demographic, clinical and working variables.

While previous studies have suggested the risk factors as the predictors of mental health among healthcare staff [13, 14, 15], our study particularly investigated working characteristics [16] such as job function, working institute and the number of working days per week to predict distress, depression and anxiety of healthcare staff in Iran. We also tested how the interaction between weekly working days and age of the staff predicts these mental health issues. These unique predictors provide evidence to help psychological service to identify healthcare staff who may need support.

## **2. Methods**

Our study surveyed healthcare staff in Iran during April 5–20, 2020 when Iran was experiencing a high outbreak of COVID-19. On April 5, the date the survey started, a total of 58,226 cumulative COVID-19 cases were diagnosed in Iran and 2,483 were diagnosed as new cases. Of the diagnosed cases, 3,603 total deaths reported in a day, 43 of whom were healthcare staff [17]. On April 20, the date the survey ended, a total of 83,505 cumulative COVID-19 cases were reported in Iran with 1,294 daily new cases and 5,209 total deaths. This caused extreme strain in medical resources and great psychological pressures on Iranian healthcare staff.

### **2.1 Study Design**

This study followed the American Association for Public Opinion Research (AAPOR) reporting guideline. Written informed approval was provided by all survey participants prior to their enrolment. In order to access more participants under the constraints created by the COVID-19 pandemic, we shared the survey links on the social media channels (Instagram, Whatsapp and Telegram) which are very popular

among Iranian and asked the responding healthcare staff to share the survey with their colleagues. All of the participants completed the survey voluntarily and anonymously, and they were free to leave the survey at any time. The final sample comprised 280 healthcare staff across all the 31 provinces of Iran.

## **2.2 Measures**

### *2.2.1 Predictor variables*

Our predictor variables included socio-demographic variables, clinical variables and working characteristic variables. Socio-demographic characteristics were self-reported by all participants including *age* (that ranged from 20–30, 30–40, 40–50, to 50–60 years old), *gender* (male or female), *education level* (categorical variable ranging from 1 = under diploma, 2 = diploma with 12 years of education, 3 = student or graduate with 2 academic years, 4=student or graduate from university, 5 = student or graduate with a master degree, 6 = student or graduate with a doctoral degree).

Clinical variables included: *chronic disease*, *infected situation* and *number of colleagues or friends diagnosed with COVID-19*. Drawing on previous research finding that people who have comorbidity may experience more mental health disorder than others in the COVID-19 pandemic [18], we asked the participants in this survey whether they had chronic disease using a categorical variable (1=yes, 2=not sure, 3=no). The participants were also asked whether they had been infected with COVID-19 within the past two weeks (1=yes, 2=not sure, 3=no). Lastly, the participants reported the number of their colleagues or friends diagnosed with COVID-19 within the past two weeks.

Working characteristic variables consisted of three items. Because medical protective equipment are different by the job function [19], we asked the healthcare staffs to indicate their *job function* (a categorical variable ranging from 1 to 9, where 1= a doctor, 2= a nurse, 3= a technician, 4= a radio technologist, 5= a medical student and an intern, 6= a healthcare administrator, 7= a supporting staff such as facility or cleaning staff, 8= a volunteer and 9= a obstetric staff). We also asked the participants their *working institute* (private, public, or government sector) and how many days a week they worked as a proxy for *working days* (1 day to 5 days) because the variables were suggested to affect healthcare workers' mental health.

### 2.2.2 Outcome variables

The three outcome variables measured in this study were distress, depression, and anxiety. The Kessler Psychological Distress Scale (K6) with Cronbach's alpha of 0.79 [20] was used to assess the participants' distress during the COVID-19. The 4-item Patient Health Questionnaire (PHQ-4, with a score of 0–12) was also employed to measure the severity of depression and anxiety among the participants. PHQ-4 is an ultra-brief self-report questionnaire consisting of a 2-item depression scale (PHQ-2) and a 2-item anxiety scale (GAD-2). The cutoff scores for detecting symptoms of distress, depression and anxiety were 13, 3 and 3 respectively. Participants who scored higher than the cutoff threshold were characterized as having the symptoms of the diseases. We translated the measures from English into Farsi, the official language of Iran.

## 2.3 Statistical Analysis

Our empirical analysis was performed using Stata 16.1 software. The statistical significance level was assessed by  $p < .05$ , 2-tailed. Multivariable logistic regression was used to determine potential risk threats of distress, depression and anxiety. In the statistical regression results, we presented descriptive analysis with odds ratios (ORs) and 95% confidence intervals (CIs) to show the potential association between risk predictors and outcome variables. We also performed a margin analysis to further investigate the interaction between the participants' working days and age that predict their distress, depression and anxiety.

### **3. Results**

#### **3.1 Descriptive statistics**

Table 1 displays the descriptive characteristics of the 280 healthcare staff participated in this survey. Of the staff, 21.4% (60) had distress disorder, 30.0% (84) had depression disorder, and 32.9% (92) had anxiety disorder. The majority of the participants (60.0%, 168) were female. Most of the participants (70.5%, 196) were under 40 years old, and 79.4% (239) of the participants had a university degree or 2-year diploma. Most of the participants (80.3%, 226) did not have chronic diseases, 2.5% (7) reported they were diagnosed as COVID-19 positive, while 27.1% (76) reported that they were unsure if they were infected by the virus. Over half of the participants (53.0%, 161) reported that their colleagues or friends were infected by COVID-19 with a mean of 3.86 (min: 0; max: 150). Because the number of colleagues or friends infected by the virus is a count, we transformed it by taking its log.

Regarding the participants' job functions, 15.0% (42) were doctors, 20.4% (57) were nurses, 9.0% (25) were technicians, 22.1% (62) were radio technologists, 3.6% (10) were medical students and interns, 13.2% (37) were healthcare administrators or interns, 3.6% (10) were supporting staff (i.e., facility or cleaning staff), 2.8% (8) were volunteers and 10.3% (29) were obstetrics staff. Most the healthcare staff were from a public healthcare organization (74.7%, 223) and worked at least three days a week (72.2%, 202).

### **3.2 Logistic regression results**

Table 1 presents the multivariate logistic regression results. Firstly, our analysis revealed that females were 2.23 times more likely to have distress disorder (OR=2.23; 95% CI: 1.02 to 4.86;  $p=0.043$ ) than their male counterparts. Highly educated healthcare staff experienced a higher level distress disorder (OR=1.63; 95% CI: 1.05 to 2.44;  $p=0.017$ ) than less-educated staff. Compared to healthcare staff who were not infected by COVID-19, those who were unsure if they were infected by the virus had a higher level of distress (OR=3.23; 95% CI: 1.59 to 6.60;  $p=0.001$ ) and depression disorder (OR=2.42; 95% CI: 1.33 to 4.37;  $p=0.004$ ). The number of COVID-19 positive cases among colleagues or friends also positively predicted anxiety disorder (OR=2.11; 95% CI: 1.11 to 4.01;  $p=0.023$ ). Compared to radio technologists – the biggest job function category in this study, medical doctors were less likely to experience distress (OR=0.09; 95% CI: 0.02 to 0.53;  $p=0.007$ ) and anxiety disorder (OR=0.12; 95% CI: 0.03 to 0.48;  $p=0.003$ ) while technicians (OR=0.27; 95% CI: 0.08 to 0.89;  $p=0.031$ ) and obstetrics staff (OR=0.21, 95% CI: 0.06 to 0.81;  $p=0.023$ ) were

less likely to experience anxiety disorder. Our analysis also showed that existing chronic health issues and the type of healthcare organization (private, public or government sector) did not significantly predict distress, depression and anxiety. Significant results are presented in figure 1.

Secondly, the results demonstrated that the interaction between the participants' working days and age significantly predict their distress (OR=1.03; 95% CI: 1.00 to 1.06;  $p=0.047$ ) and depression (OR=1.03, 95% CI: 1.00 to 1.06;  $p=0.026$ ). Figure 1 shows the participants' working days per week and age predict their distress and depression. By further conducting a margin analysis, we found that 41% of the healthcare staff who were 20 years old and worked only one day per week had the highest likelihood of distress (95% CI: 23% to 65%;  $p=0.000$ ) and 61% of them experienced depression (95% CI: 40% to 82%;  $p=0.000$ ). Furthermore, the healthcare staff who were 20 years old but worked five days a week had 19% (95% CI: 19% to 36%;  $p=0.023$ ) chance of having distress disorder and 24% (95% CI: 7% to 40%;  $p=0.004$ ) chance of depression disorder. The chance of having distress and depression varied significantly by the number of working days among younger participants but not their older counterparts. Margin analysis results are presented in figure 2.

#### **4. Discussion**

The findings of our study revealed that nearly a third of all of the healthcare staff in Iran reported symptoms of distress, depression and anxiety during fighting with COVID-19 pandemic. Additionally, gender and education qualification were significant predictors of the healthcare staffs' distress. The participants who were

unsure if they have been infected by COVID-19 reported a more risk of distress and depression compared to those who were sure that they have not been infected by the virus. In addition, our survey suggested that the number of COVID-19 cases among colleagues or friends positively predicted anxiety disorder of the participants. Compared to radio technologists, doctors had less possibility of distress and anxiety while technicians and obstetrics staff had less possibility of anxiety. Our study demonstrated that the interaction between the number of working days a week and age predicts the respondents' distress and depression, with the chance of having distress and depression varying greatly by the number of working days among younger but not older healthcare staff.

In contrast to previous research, our study failed to find a significant effect of some demographic variables on the mental health of the healthcare staff. While, age, gender and education qualification predicted mental health among healthcare staff in China [21, 22, 23], in Italy [24], in Brazil [25] and in other countries [26] during the COVID-19 pandemic, we could not find a significant association between the variables. In addition, previous studies indicated having a chronic disease was shown to be a risk factor that significantly affects mental health of staff [27, 28], but we did not find a significant relationship between the variables. Our study also revealed that doctors were less likely to suffer from distress and anxiety than radio technologists, because radio technologists were more likely exposed to the risk of COVID-19 infection due to their frequent and close engagement in the process of diagnosing patients. These findings showed that the possible predictors of the mental health of

healthcare staff during a pandemic from socio-demographic variables, clinical variables and working characteristic variables vary across countries [29, 30, 31]. These predictors of the mental health of healthcare staff mentioned previously vary for different mental health symptoms in one pandemic.

Moreover, our study showed that the participants' age significantly predicted distress and depression differently when we focused on their working days per week. Our findings suggest that hospitals and psychological service organizations should pay more attention to their younger healthcare staff. Previous studies have shown that younger adults experienced greater mood swings during the SARS outbreak due to their inadequate experience of facing a changing environment [32]. During the COVID-19 pandemic, younger healthcare staff exhibited the same condition when they worked fewer days a week. Hence, hospitals in Iran may consider training more working days a week for younger healthcare staff.

## **5. Limitations**

Our study has several limitations that open new agendas for future studies. First, we focused on healthcare staff in Iran who were very busy during the COVID-19 pandemic. Therefore, we examined limited variables related to mental health of healthcare staff using K6 and PHQ-4 to measure distress, depression and anxiety. Second, our sample size is also relatively small. Future studies should examine the factors that determine mental health of healthcare staff using a larger sample size. Finally, we attempted to include samples from different provinces of Iran. However, our sample is limited to a small proportion of healthcare staff in each province of Iran.

Therefore, future studies should examine if geographic location or the actual density of COVID-19 cases in each province affect the healthcare staff's experience of mental diseases. This survey is also cross-sectional. Therefore, future longitudinal studies to examine the factors that affect healthcare staff's mental health highly contribute to our knowledge of healthcare mental health during pandemic crises.

## **6. Conclusion**

Identifying potential risk predictors for healthcare staff assists psychological service institutions better alleviate the mental health issues of staff during the COVID-19 pandemic. Importantly, this study demonstrated that it is imperative to pay more attention to female and highly educated healthcare staff, and some clinical predictors should also be recognized. The interaction of healthcare staff's number of working days a week and age play an important role in predicting their distress and depression disorders. Therefore, it is imperative to further investigate healthcare staff's potential working characteristics during the COVID-19 pandemic and help the staff maintain their professional and personal strength in the prolonged COVID-19 pandemic.

## References

- [1] Woolley, K., Smith, R., & Arumugam, S. (2020). Personal Protective Equipment (PPE) Guidelines, Adaptations and Lessons During the COVID-19 Pandemic. *Ethics, Medicine and Public Health*, 100546. <https://doi.org/10.1016/j.jemep.2020.100546>
- [2] Lee, S. (2020). Coronavirus Anxiety Scale: A brief mental health screener for COVID-19 related anxiety. *Death Studies*, 44, 1–9. <https://doi.org/10.1080/07481187.2020.1748481>
- [3] Roy, D., Tripathy, S., Kar, S. K., Sharma, N., Verma, S. K., & Kaushal, V. (2020). Study of knowledge, attitude, anxiety & perceived mental healthcare need in Indian population during COVID-19 pandemic. *Asian Journal of Psychiatry*, 51, 102083. <https://doi.org/10.1016/j.ajp.2020.102083>
- [4] Lai, J., Ma, S., Wang, Y., Cai, Z., Hu, J., Wei, N., Wu, J., Du, H., Chen, T., Li, R., Tan, H., Kang, L., Yao, L., Huang, M., Wang, H., Wang, G., Liu, Z., & Hu, S. (2020). Factors Associated With Mental Health Outcomes Among Health Care Workers Exposed to Coronavirus Disease 2019. *JAMA Network Open*, 3, e203976. <https://doi.org/10.1001/jamanetworkopen.2020.3976>
- [5] Afshar Jahanshahi, A., Dinani, M. M., Madavani, A. N., Li, J., & Zhang, S. X. (2020). The distress of Iranian adults during the Covid-19 pandemic – More distressed than the Chinese and with different predictors. *Brain, Behavior, and Immunity*, 87, 124–125. <https://doi.org/10.1016/j.bbi.2020.04.081>
- [6] Ruiz, M. (2020, March 26). *Italian nurse who fell ill commits suicide over fears of spreading coronavirus, nursing group says*. <https://www.foxnews.com/world/italian-nurse-suicide-fears-spreading-coronavirus>
- [7] Jane Kirby. Covid-19 mental health hotline set up for NHS staff. (n.d.). *MSN*. Retrieved July 10, 2020, from <https://www.msn.com/en-gb/news/uknews/covid-19-mental-health-hotline-set-up-for-nhs-staff/ar-BB12i92V>
- [8] Dai, H., Zhang, S. X., Looi, K. H., Su, R., & Li, J. (2020). Perception of Health Conditions and Test Availability as Predictors of Adults' Mental Health during the COVID-19 Pandemic: A Survey Study of Adults in Malaysia. *International Journal of Environmental Research and Public Health*, 17(15), 5498. <https://doi.org/10.3390/ijerph17155498>
- [9] Zhang, S. X., Sun, S., Jahanshahi, A. A., Alvarez-Risco, A., Ibarra, V. G., Li, J., & Patty-Tito, R. M. (2020). Developing and testing a measure of COVID-19

- organizational support of healthcare workers – results from Peru, Ecuador, and Bolivia. *Psychiatry Research*, 291, 113174. <https://doi.org/10.1016/j.psychres.2020.113174>
- [10] Byrne, L. (2020). A role for lived experience mental health leadership in the age of Covid-19. *Journal of Mental Health*, 29, 1–4. <https://doi.org/10.1080/09638237.2020.1766002>
- [11] Henry, B. M., & Lippi, G. (2020). Chronic kidney disease is associated with severe coronavirus disease 2019 (COVID-19) infection. *International Urology and Nephrology*, 52(6), 1193–1194. <https://doi.org/10.1007/s11255-020-02451-9>
- [12] Banerjee, D. (2020). ‘Age and ageism in COVID-19’: Elderly mental health-care vulnerabilities and needs. *Asian Journal of Psychiatry*, 51, 102154. <https://doi.org/10.1016/j.ajp.2020.102154>
- [13] Chen, J., Zhang, S. X., Wang, Y., Jahanshahi, A. A., Dinani, M. M., Madavani, A. N., & Nawaser, K. (2020). The curvilinear relationship between the age of adults and their mental health in Iran after its peak of COVID-19 cases. *MedRxiv*, 2020.06.11.20128132. <https://doi.org/10.1101/2020.06.11.20128132>
- [14] Chen, X., Zhang, S., Afshar Jahanshahi, A., Alvarez-Risco, A., Dai, H., Li, J., & Ibarra, V. (2020). *Belief in Conspiracy Theory about COVID-19 Predicts Mental Health and Well-being—A Study of Healthcare Staff in Ecuador*. <https://doi.org/10.1101/2020.05.26.20113258>
- [15] Zhang, S. X., Graf-Vlachy, L., Su, R., Li, J., & Looi, K. H. (2020). Who do not wash their hands during the Covid-19 pandemic? Social media use as a potential predictor. *MedRxiv*, 2020.06.01.20119230. <https://doi.org/10.1101/2020.06.01.20119230>
- [16] Zhang, S. X., Wang, Y., Rauch, A., & Wei, F. (2020). Unprecedented disruption of lives and work: Health, distress and life satisfaction of working adults in China one month into the COVID-19 outbreak. *Psychiatry Research*, 288, 112958. <https://doi.org/10.1016/j.psychres.2020.112958>
- [17] Gharebaghi, R., & Fatemeh, H. (2020). COVID-19 and Iran: Swimming with hands tied! *Swiss Medical Weekly: Official Journal of the Swiss Society of Infectious Diseases, the Swiss Society of Internal Medicine, the Swiss Society of Pneumology*, 150. <https://doi.org/10.4414/smw.2020.20242>
- [18] Fawad, M., Rawashdeh, F., Parmar, P., & Ratnayake, R. (2020). Simple ideas to mitigate the impacts of the COVID-19 epidemic on refugees with chronic diseases. *Conflict and Health*, 14. <https://doi.org/10.1186/s13031-020-00277-x>

- [19] Rimmer, A. (2020, March 13). Covid-19: GPs call for same personal protective equipment as hospital doctors. *BMJ*. <https://www.bmj.com/content/368/bmj.m1055>
- [20] Kessler, R. C., Andrews, G., Colpe, L. J., Hiripi, E., Mroczek, D. K., Normand, S.-L. T., Walters, E. E., & Zaslavsky, A. M. (2002). Short screening scales to monitor population prevalences and trends in non-specific psychological distress. *Psychological Medicine*, 32(6), 959–976. <https://doi.org/10.1017/S0033291702006074>
- [21] Li, H., Wang, S., Zhong, F., Bao, W., Li, Y., Liu, L., Wang, H., & He, Y. (2020). Age-Dependent Risks of Incidence and Mortality of COVID-19 in Hubei Province and Other Parts of China. *Frontiers in Medicine*, 7, 190. <https://doi.org/10.3389/fmed.2020.00190>
- [22] Xiao, X., Zhu, X., Fu, S., Hu, Y., Li, X., & Xiao, J. (2020). Psychological impact of healthcare workers in China during COVID-19 pneumonia epidemic: A multi-center cross-sectional survey investigation. *Journal of Affective Disorders*. <https://doi.org/10.1016/j.jad.2020.05.081>
- [23] Wang, Y., Zhu, L.-Y., Ma, Y.-F., Bo, H.-X., Deng, H.-B., Cao, J., Wang, Y., Wang, X.-J., Xu, Y., Lu, Q.-D., Wang, H., & Wu, X.-J. (2020). Association of insomnia disorder with sociodemographic factors and poor mental health in COVID-19 inpatients in China. *Sleep Medicine*. <https://doi.org/10.1016/j.sleep.2020.06.011>
- [24] Casagrande, M., Favieri, F., Tambelli, R., & Forte, G. (2020). The enemy who sealed the world: Effects quarantine due to the COVID-19 on sleep quality, anxiety, and psychological distress in the Italian population. *Sleep Medicine*. <https://doi.org/10.1016/j.sleep.2020.05.011>
- [25] Zhang, S. X., Wang, Y., Jahanshahi, A. A., Li, J., & Schmitt, V. G. H. (2020). Mental distress of adults in Brazil during the COVID-19 crisis. *MedRxiv*, 2020.04.18.20070896. <https://doi.org/10.1101/2020.04.18.20070896>
- [26] Zhang, S. X., Sun, S., Jahanshahi, A. A., Alvarez-Risco, A., Ibarra, V. G., Li, J., & Patty-Tito, R. M. (2020). Developing and testing a measure of COVID-19 organizational support of healthcare workers – results from Peru, Ecuador, and Bolivia. *Psychiatry Research*, 291, 113174. <https://doi.org/10.1016/j.psychres.2020.113174>
- [27] Sodhi, R., & Manju, D. (2013). Spiritually and mental health among normal and chronic disease group. *International Journal of Research Studies in Psychology*, 2. <https://doi.org/10.5861/ijrsp.2012.106>

- [28] Hüfner, K., Fuchs, D., Blauth, M., & Sperner-Unterweger, B. (2019). How acute and chronic physical disease may influence mental health—An Analysis of neurotransmitter precursor amino acid levels. *Psychoneuroendocrinology*, *106*, 95–101. <https://doi.org/10.1016/j.psyneuen.2019.03.028>
- [29] Zhang, S. X., Liu, J., Afshar Jahanshahi, A., Nawaser, K., Yousefi, A., Li, J., & Sun, S. (2020a). At the height of the storm: Healthcare staff's health conditions and job satisfaction and their associated predictors during the epidemic peak of COVID-19. *Brain, Behavior, and Immunity*, *87*, 144–146. <https://doi.org/10.1016/j.bbi.2020.05.010>
- [30] Zhang, S. X., Sun, S., Jahanshahi, A. A., Wang, Y., Madavani, A. N., & Dinani, M. M. (2020). Beyond predicting the number of infections: Predicting who is likely to be COVID negative or positive. *MedRxiv*, 2020.04.30.20086348. <https://doi.org/10.1101/2020.04.30.20086348>
- [31] Tang, P. M., Zhang, S. X., Li, C. H., & Wei, F. (2020). Geographical identification of the vulnerable groups during COVID-19 crisis: The typhoon eye effect and its boundary conditions. *MedRxiv*, 2020.04.28.20083667. <https://doi.org/10.1101/2020.04.28.20083667>
- [32] Yeung, D., & Fung, H. (2007). Age differences in coping and emotional responses toward SARS: A longitudinal study of Hong Kong Chinese. *Aging & Mental Health*, *11*, 579–587. <https://doi.org/10.1080/13607860601086355>

Table 1 Descriptive findings and predictors of healthcare staff's mental health by logistic regression (N=280)

Variables	Description	Distress		Depression		Anxiety	
		OR(95%CI)	p-value	OR (95%CI)	p-value	OR (95%CI)	p-value
<b>SOCIO-DEMOGRAPHIC VARIABLES</b>							
<b>Age</b>							
<i>20–30 years old</i>	101 (36.3%)	0.87(0.80 to 0.96)	0.004	0.87(0.79 to 0.96)	0.004	0.91(0.83 to 1.00)	0.052
<i>30–40 years old</i>	95 (34.2%)						
<i>40–50 years old</i>	63 (22.7%)						
<i>50–60 years old</i>	19 (6.8%)						
<b>Gender</b>							
<i>Male</i>	112 (40.0%)	Reference group		Reference group		Reference group	
<i>Female</i>	168 (60.0%)	2.23 (1.02 to 4.86)	0.043	1.41 (0.75 to 2.66)	0.283	1.70 (0.91 to 3.17)	0.095
<b>Education qualification</b>							
<i>Under diploma</i>	4 (1.4%)	1.63 (1.05 to 2.44)	0.017	1.14 (0.83 to 1.56)	0.433	1.40 (0.99 to 1.97)	0.055
<i>Diploma (with 12 years of education)</i>	19 (6.8%)						
<i>Student or graduate with 2 academic years</i>	34 (12.1%)						
<i>Student or graduate from university</i>	135 (48.2%)						
<i>Student or graduated with a master degree</i>	40 (14.3%)						
<i>Student or graduated with a doctoral degree</i>	48 (17.2%)						
<b>CLINICAL VARIABLES</b>							
<b>Chronic disease</b>							
<i>No</i>	226 (80.7%)	Reference group		Reference group		Reference group	
<i>Unsure</i>	36 (12.9%)	0.48 (0.15 to 1.56)	0.224	0.98 (0.39 to 2.44)	0.958	0.64 (0.26 to 1.61)	0.346
<i>Yes</i>	18 (6.4%)	0.83 (0.21 to 3.30)	0.791	0.67 (0.22 to 2.04)	0.486	1.20 (0.39 to 3.66)	0.750
<b>COVID-19 infected status</b>							

<i>No</i>	197 (79.4%)	Reference group		Reference group		Reference group	
<i>Unsure</i>	76 (27.1%)	3.23 (1.59 to 6.60)	0.001	2.42 (1.33 to 4.37)	0.004	1.72 (0.93 to 3.18)	0.084
<i>Yes</i>	7 (2.5%)	6.19 (0.82 to 46.64)	0.077	1.36 (0.18 to 9.98)	0.765	1.83 (0.28 to 11.96)	0.527
<b>Number of COVID-19 cases among colleagues or friends</b>	Mean=3.86	0.98 (0.40 to 2.37)	0.930	1.26 (0.62 to 2.58)	0.523	2.11 ( 1.11 to 4.01)	0.023
<b>WORKING CHARACTERISTIC VARIABLES</b>							
<b>Job function</b>							
<i>Radio technologist</i>	62 (22.1%)	Reference group		Reference group		Reference group	
<i>Doctor</i>	42 (15.0%)	0.09 (0.02 to 0.53)	0.007	0.83 (0.24 to 2.90)	0.776	0.12 (0.03 to 0.48)	0.003
<i>Nurse</i>	57 (20.4%)	0.58 (0.23 to 1.49)	0.260	0.79 (0.35 to 1.79)	0.572	0.61 (0.27 to 1.38)	0.236
<i>Technician</i>	25 (9.0%)	0.31 (0.09 to 1.11)	0.072	0.55 (0.16 to 1.85)	0.334	0.27 (0.08 to 0.89)	0.031
<i>Obstetrics</i>	29 (10.3%)	0.27 (0.06 to 1.21)	0.087	0.49 (0.14 to 1.74)	0.273	0.21 (0.06 to 0.81)	0.023
<i>Healthcare administrator</i>	47 (15.8%)	0.62 ( 0.23 to 1.68)	0.343	0.68 (0.27 to 1.67)	0.404	0.72 (0.31 to 1.71)	0.465
<i>Supporting staff</i>	10 (3.6%)	0.44 (0.02 to 7.96)	0.575	0.58 (0.06 to 5.34)	0.634	0.20 (0.02 to 1.86)	0.157
<i>Volunteer</i>	8 (2.8%)	0.32(0.03 to 3.46)	0.345	0.24 (0.02 to 2.84)	0.258	0.54 (0.08 to 3.52)	0.522
<b>Public or private institution</b>							
<i>Private</i>	71 (25.4%)	Reference group		Reference group		Reference group	
<i>Public</i>	209 (74.6%)	1.79 (0.82 to 3.89)	0.143	0.86 (0.43 to 1.74)	0.674	0.94 (0.48 to 1.87)	0.870
<b>Work days a week</b>							
<i>1 day</i>	41 (14.6%)	0.38 (0.13 to 1.10)	0.075	0.37 (0.15 to 0.91)	0.030	0.61 (0.25 to 1.51)	0.283
<i>2 days</i>	37 (13.2%)						
<i>3 days</i>	81 (28.9%)						
<i>4 days</i>	61 (21.8%)						
<i>5 days</i>	60 (21.5%)						
<b>INTERACTION</b>							
<i>Working days * Age</i>	280 (100%)	1.03 (1.00 to 1.06)	0.047	1.03 (1.01 to 1.06)	0.018	1.01 (0.99 to 1.04)	0.251




Figure 1 Predicted value and 95% confidence intervals (CIs) of distress, depression and anxiety by healthcare staff's socio-demographic variables, clinical variables and working characteristic variables




Figure 2 Predicted value and 95% confidence intervals (CIs) of distress, depression and anxiety by healthcare staff's number of work days a week and age