

1 **FULL TITLE:** Spatial epidemiology of Japanese encephalitis virus and other infections of the central
2 nervous system in Lao PDR (2003 – 2011): a retrospective analysis

3
4 **SHORT TITLE:** Spatial epidemiology of central nervous system infection in Lao PDR

5
6 Sayaphet Rattanavong¹, Audrey Dubot-Pérés^{1,2,3}, Mayfong Mayxay^{1,2,4}, Manivanh Vongsouvath¹, Sue J
7 Lee^{2,5}, Julien Cappelle^{6,7,8,9}, Paul N. Newton^{1,2,5}, Daniel M. Parker^{10,11}

8
9
10
11 Sayaphet Rattanavong: Sayaphet@tropmedres.ac

12 Audrey Dubot-Pérés: Audrey@tropmedres.ac

13 Mayfong Mayxay: Mayfong@tropmedres.ac

14 Manivanh Vongsouvath: Manivanh@tropmedres.ac

15 Sue J Lee: Sue@tropmedres.ac

16 Julien Cappelle: Julien.cappelle@cirad.fr

17 Paul N. Newton: Paul.Newton@tropmedres.ac

18 *Daniel M. Parker: Dparker1@uci.edu

19
20 *corresponding author

21
22 1. Lao-Oxford-Mahosot Hospital-Wellcome Trust Research Unit (LOMWRU), Microbiology Laboratory,
23 Mahosot Hospital, Vientiane, Lao PDR

24 2. Centre for Tropical Medicine and Global Health, Nuffield Department of Clinical Medicine, University
25 of Oxford, Churchill Hospital, Oxford, U.K.

26 3. Unité des Virus Émergents (UVE: Aix-Marseille Univ – IRD 190 – Inserm 1207 – IHU Méditerranée
27 Infection), Marseille, France

28 4. Institute of Research and Education Development, University of Health Sciences, Vientiane, Lao PDR

29 5. Mahidol-Oxford Tropical Medicine Research Unit, Faculty of Tropical Medicine, Mahidol University,
30 Thailand

31 6. Epidemiology and Public Health Unit, Institut Pasteur du Cambodge, Phnom Penh, Cambodia.

32 7. CIRAD, UMR ASTRE, F-34398, Montpellier, France

33 8. UMR ASTRE, CIRAD, INRA, Montpellier University, F-34398, Montpellier, France.

34 9. UMR EpiA, INRA, VetAgro Sup, F-69280, Marcy l’Etoile, France.

35 10. Department of Population Health and Disease Prevention, University of California, Irvine, U.S.A.

36 11. Department of Epidemiology, School of Medicine, University of California, Irvine, U.S.A.

37

38

39

40

41

42

43

44

45

46

47

48 **ABSTRACT**

49 **Background.** Central nervous system (CNS) infections are important contributors to morbidity and
50 mortality and the causative agents for ~50% patients are never identified. The causative agents of some
51 CNS infections have distinct spatial and temporal patterns.

52 **Methodology/Principal Findings.** Here we present the results of a spatial epidemiological and
53 ecological analysis of CNS infections in Lao PDR (2003 – 2011). The data came from hospitalizations for
54 suspected CNS infection at Mahosot Hospital in Vientiane. Out of 1,065 patients, 450 were assigned a
55 confirmed diagnosis. While many communities in Lao PDR are in rural and remote locations, most
56 patients in these data came from villages along major roads. Japanese encephalitis virus ((JEV); n=94)
57 and *Cryptococcus* spp. (n=70) were the most common infections. JEV infections peaked in the rainy
58 season and JEV patients came from villages with higher surface flooding during the same month as
59 admission. JEV infections were spatially dispersed throughout rural areas and were most common in
60 children. *Cryptococcus* spp. infections clustered near Vientiane (an urban area) and among adults.

61 **Conclusions/Significance.** The spatial and temporal patterns identified in this analysis are related to
62 complex environmental, social, and geographic factors. For example, JEV infected patients came from
63 locations with environmental conditions (surface water) that are suitable to support larger mosquito
64 vector populations. Most patients in these data came from villages that are near major roads; likely the
65 result of geographic and financial access to healthcare and also indicating that CNS diseases are
66 underestimated in the region (especially from more remote areas). As Lao PDR is undergoing major
67 developmental and environmental changes, the space-time distributions of the causative agents of CNS
68 infection will also likely change. There is a major need for increased diagnostic abilities; increased access
69 to healthcare, especially for rural populations; and for increased surveillance throughout the nation.

70

71

72 **AUTHOR SUMMARY**

73 Infections of the central nervous system (CNS) are important with regard to public health. However
74 many CNS infections are never diagnosed. In this analysis we investigated spatial and temporal patterns
75 in hospitalized patients with suspected CNS infections in Lao PDR. We found that patients were most
76 likely to come from villages located along major roads and highways. Patients from remote areas may
77 have more difficulty reaching healthcare facilities. The most commonly diagnosed infection in these
78 patients was Japanese encephalitis virus (JEV). Patients with this disease came from locations that were
79 optimal for the mosquito vectors that spread JEV, rural areas with surface water and during the rainy
80 season. Our analyses suggest that CNS infections should be a priority for public health workers in the
81 region. Diagnostic capabilities should be increased throughout the nation; surveillance efforts should be
82 broadened; and efforts should be increased toward providing easy access to healthcare for rural and
83 remote populations.

84

85

86

87

88

89

90

91

92

93

94

95

96 INTRODUCTION

97 Numerous illnesses go undiagnosed and the causative agents of many infections are never
98 identified. In regions where access to healthcare facilities is limited and where diagnostic capabilities are
99 few, a smaller proportion of diseases are objectively diagnosed. Diseases with mild symptoms may more
100 frequently go untreated, but in some areas even severe illnesses commonly go undiagnosed and
101 untreated. Infections of the central nervous system (CNS) can be particularly severe, affecting the brain
102 and/or spinal cord and/or the surrounding meninges, frequently resulting in death.

103 Pathogens that invade and infect the CNS include viruses, bacteria, fungi, parasites, and prions.
104 As with other infections, the causative agent(s) of many CNS infections are never determined
105 (frequently <50%) [1–3]. Symptoms of CNS infection can range widely, even for single causative agents,
106 leading to further difficulties with diagnosis. Diagnoses are frequently presumptive and non-specific (i.e.
107 meningitis is often presumed to be caused by bacteria whereas encephalitis is presumed to be caused
108 viruses [2,4–6]). Some causative agents are specific to regions (e.g. Japanese Encephalitis, Saint Louis
109 Encephalitis, Rift Valley Fever Viruses) and exhibit seasonal fluctuations (e.g. vector borne infections),
110 therefore geography and seasonality can facilitate presumptive diagnosis of CNS diseases [4,7].

111 In Southeast (SE) Asia, CNS infections are increasingly recognized as important contributors to
112 morbidity and mortality [1,8,9]. However, detailed medical and epidemiological data are frequently
113 lacking, especially for low income nations and from remote areas within middle-to-high income nations.
114 Known important viral CNS infections in SE Asia include Japanese encephalitis, dengue, and rabies
115 viruses [1]. Important bacterial CNS infections include *Streptococcus pneumoniae*, *Haemophilus*
116 *influenzae*, *S. suis*, *Mycobacterium tuberculosis* and *Neisseria meningitidis*. *Orientia tsutsugamushi*,
117 *Rickettsia typhi* and *Leptospira* spp. are increasingly recognized as important causes [9]. Detailed
118 analyses that confirm the cause of CNS related infections or assess their spatial and temporal
119 distribution in the region are rare [8–11,14].

120

121 We recently published the results of a study of the etiology and impact of CNS infections
122 diagnosed among 1,065 patients at Mahosot Hospital, Vientiane, Lao PDR [14]. The goal of this
123 secondary analysis was to investigate the spatial distribution(s) of CNS-related infections; to look for
124 differential spatial distributions for major causative agents; and to explore potential geographic,
125 demographic, and environmental correlates of these infections.

126 **DATA AND METHODS**

127 ***Data sources, processing and merging***

128 Data used in this research were compiled from four main sources (**Supporting Figure 1**). The
129 epidemiological data come from an 8-year research project on CNS infections in Lao PDR from all
130 patients who received diagnostic lumbar puncture (LP) at Mahosot Hospital in Vientiane, Lao PDR
131 between January 2003 and August 2011 and consenting to participate [14]. All patients were admitted
132 to the hospital because of suspected CNS infection and Mahosot Hospital is the only medical facility in
133 Lao PDR capable of performing diagnostic LP and cerebral spinal fluid (CSF) analysis. Tests for a large
134 panel of pathogens were performed at the Microbiology Laboratory following international standards
135 (details provided in **Supporting Materials I** and [14]). Demographic (age, gender, ethnicity) and
136 geographic (home village) characteristics of patients were recorded in the database.

137 The epidemiological data were used to create two separate datasets: One aggregated at the
138 village level (one row per location) and another was maintained at the individual level, with one row per
139 individual. The official Lao PDR censuses from 2005 and 2015 were used to geocode villages (based on
140 village name and administrative units listed in patient records) and to assign village population
141 estimates to each village (taking a mean population size between 2005 and 2015). Village location and
142 population sizes were then merged to both the individual- and village-level datasets.

143 A subset of villages within the geographic region of the home villages of included patients was
144 selected by overlaying a standard deviational ellipse with 3 standard deviations (calculations described
145 in **Supporting Materials II**) around the patient home villages and then selecting all villages within that
146 ellipse (**Supporting Figure 2**). These villages were then retained for village level comparisons between
147 villages populated, and not populated, with patients admitted with CNS disease needing an LP. This
148 subset of villages is hereafter referred to as the “study area”.

149 Major road network data was taken from OpenStreetMaps (<http://www.openstreetmap.la>),
150 selecting “primary”, “secondary”, and all major connecting roads (downloaded in February 2017,
151 **Supporting Figure 3**). Primary and secondary roads are the two largest road classifications for the
152 nation. Primary roads link major towns and cities and secondary roads link mid-sized towns. Primary and
153 secondary link roads are ramps or slip roads that connect other roads to primary or secondary roads.
154 Together, these types of roads are hereafter referred to as “major roads”. Smaller roads were not
155 included in this analysis as they are less likely to be accurately included in the OpenStreetMaps data. The
156 Euclidian distance was then calculated from all villages in the census to the nearest point along a major
157 road. These distances were merged to both the village- and the individual-level datasets.

158 Environmental predictor variables for vegetation and surface water were derived from
159 Moderate Resolution Imaging Spectroradiometer (MODIS) products (MOD13Q1/MYD13Q1 250 meter
160 AQUA/TERRA 16 day composites). Since many infectious diseases, especially vector borne diseases, are
161 strongly influenced by environmental factors we hypothesized that indicators of vegetation and surface
162 flooding would correlate with some specific diagnoses. Three environmental indices were downloaded
163 and considered in these analyses: a normalized flooding index (NFI) [15]; the normalized difference
164 vegetative index (NDVI); and the enhanced vegetation index (EVI). NFI is indicative of surface water,
165 NDVI is indicative of green surface vegetation, and EVI is an improved measure of green vegetation that
166 is intended to account for dense forest canopies and atmospheric conditions that can lead to error in

167 NDVI measurements. Data were downloaded for each of these environmental indices (EI) within each
168 16-day time period from February 2002 through December of 2011. The final analyses conducted in this
169 research retained only the EVI and NFI for environmental predictors (summary statistics for NDVI are
170 included) because NDVI and EVI were strongly correlated. The EIs are described in more detail in

171 **Supporting Materials III.**

172 The environmental raster data were then summarized and extracted based on varying buffer
173 sizes (2km, 5km, and 10km) for each village in the individual- and village-level datasets. Permanent
174 water bodies (such as the Mekong River and Nam Ngum Dam) were masked from the NFI calculations.
175 For the village-level datasets, mean values of each environmental variable was calculated for the study
176 period duration and used as an indicator of “average” vegetation or surface water characteristics of
177 each village. For the individual-level dataset the values were extracted based on the admission date,
178 using increasing durations of time prior to admission (within the same month, within the previous 2
179 months, within the previous year).

180 The final datasets include the village-level data, that is a subsample of 98% of all villages with
181 patients included in the study and the other census designated villages within the same region (the
182 study area), and the individual level dataset that includes all patients included in the study. Variables
183 used in this analysis and their descriptions are listed in **Table 1**.

184 ***Exploratory spatial data analysis***

185 Summary statistics (median; Q1:Q3; mean) were calculated for the distances between villages
186 and the nearest major road, and aggregated by whether or not the village was home to included
187 patients and by specific diagnoses. Summary statistics (mean and 95% confidence intervals) were also
188 calculated for all environmental variables, at each buffer size and temporal resolution, and for each of
189 the major diagnoses. Tukey’s post hoc range test was used to assess statistically significant differences in
190 environmental indices across diagnoses.

191 Standard distance deviations (SDDs) and standard deviation ellipses (SDEs) were calculated
192 (details in **Supporting Materials II**) and mapped to measure and visually analyze the central tendency
193 and spatial distributions for all patient home villages and by each of the major single (mono-infection)
194 diagnoses.

195 **Formal analyses**

196 Multivariable regressions were used to calculate model-adjusted odds ratios and confidence
197 intervals. The regressions at the village level focused on study patient villages and the home villages of
198 JEV diagnosed patients. A multivariable regression was also done at the individual level focusing on JEV
199 infected patients. Other diagnoses were not included in these analyses because of small numbers of
200 cases per village.

201 Logistic generalized additive models (GAMs) were used for variable selection and specification
202 (detailed in **Supporting Materials IV and in Table 1**) for the final models. The GAMs were used to
203 examine different specifications of the continuous environmental, geographic, and demographic
204 variables and for changes in model fit and strength of association across buffer sizes (i.e. 2km, 5km, or
205 10km buffers) and for different time durations for EI measurements (i.e. same month, 2 months prior,
206 12 months prior to hospital admission). The final model covariates were chosen based on a combination
207 of *a priori* hypotheses, model fit (using the Akaike information criterion), and strength of association
208 between the covariate and the model outcome variable.

209 The final model for the individual-level analysis was a generalized logistic mixed model with a
210 random effect for home village. The final model for the village-level analysis was a logistic regression.

211 **Software**

212 All maps were created using QGIS version 3.4.9. R cran version 3.5.2 was used for downloading,
213 processing, and wrangling MODIS data (using the “MODISStsp”; “raster”; “rgdal”; and “maptools”
214 packages) and for all regressions. The “mgcv” package was used for GAMs and the lme4 package was

215 used for the generalized mixed models. Euclidian distances between villages and major roads were
216 calculated using QGIS. ArcMap version 10.5.1 was used to calculate SDDs and SDEs.

217 ***Ethics approval***

218 Ethical clearance for the CNS study was granted by the Oxford University Tropical Ethics
219 Research Committee and by the Ethical Review Committee of the Faculty of Medical Sciences, National
220 University of Laos. Between 2003 and 2006, both Oxford Tropical Ethics Committee and the Ethical
221 Review Committee of the Faculty of Medical Sciences, National University of Laos, approved the use of
222 oral witnessed consent. Oral consent was confirmed by a signed witness statement. Verbal consent
223 (from 2003 – 2006) and written consent (from 2006 – 2011) were obtained from all recruited patients or
224 immediate relatives. All ages were included provided that they or their guardian gave witnesses oral
225 informed consent (2003 – 2006) and written informed consent since 2006.

226 **RESULTS**

227 ***Summary statistics***

228 A total of 1,065 patients were recruited with no LP contraindications and consented to have a
229 diagnostic LP; 450 (42%) were assigned a final laboratory diagnosis. The most common of these were
230 Japanese encephalitis virus (JEV) 94 individuals); followed by *Cryptococcus* spp. with 70 individuals (9
231 were *C. gattii*); scrub typhus (*Orientia tsutsugamushi*) 31; Dengue virus 27; *Leptospira* spp. 25; murine
232 typhus (*Rickettsia typhi*) 24; *Streptococcus pneumoniae* in 22 and 20 with *Mycobacterium tuberculosis*.
233 124 patients died prior to discharge (out of 893 with recorded discharge type recorded).

234 The majority (666, 63%) of patients were male, with the lowest sex bias in cryptococcal
235 infections (40/70, 57% male) and the highest among dengue infections (22/27, 82% male) (**Table 2**). Age
236 patterns were evident in JEV and cryptococcal infections, with median ages of 13 and 33 years,
237 respectively (**Table 2**). Patients were linked to 582 different villages (multiple patients could come from
238 the same village): 90 villages with JEV patients, 66 with cryptococcal patients, 31 with scrub typhus

239 patients, 27 with dengue patients, 24 with leptospiral patients, and 24 with murine typhus patients. The
240 majority (870, 82%) of patients came from within Vientiane Prefecture (678, 64%) or Vientiane Province
241 (192, 18%).

242 A total of 6,416 villages (of 10,520 recorded in 2005 [16]) were selected as the study area for
243 further village level analyses (**Table 3**). Villages that were home to study patients were 11 times (0.7km
244 versus 6.3km, from **Table 3**) closer to a major road when compared to other villages within the study
245 area. Scrub typhus and JEV infected patient homes were further from major roads than other patients,
246 but the difference was not statistically significant in univariate analyses.

247 The home villages of JEV patients were more broadly dispersed (**Figure 1B**) than for patients
248 with other etiologies (**Figure 1C**), evident from the circular, larger SDE and SDD. The distribution of these
249 JEV patient home villages was also relatively isotropic, with the SDE and SDD being nearly identical.
250 Conversely, patients with cryptococcal infections were clustered near Vientiane City and along the road
251 leading North/Northwest from the urban center (**Figure 1C**). Scrub typhus and murine typhus infections
252 were also both clustered around Vientiane City but showed perpendicular spatial distributions
253 (**Supporting Figures 4D and 4G**) a pattern previously described from IgG seropositivity data from
254 Vientiane City [17]).

255 ***Characteristics of patient home villages***

256 Mean NFI values for the 2km radius tended to be higher than for either the 5km or 10km radius
257 as surface flooding is heterogeneous and taking a mean across larger radii dilutes the measurement. The
258 opposite pattern was observed for both vegetation indices. The 2km radius for both mean NDVI and
259 mean EVI was usually smaller than at 5km or 10km radii (**Figures 2 and 3**).

260 Study patient villages had higher mean NFI values than non-study patient villages (**Figure 2A**)
261 (non-study patient villages are those in the same study area as study patient villages but were not home
262 to a study patient). The home villages of study patients diagnosed with *dengue virus* and cryptococcal

263 infections had high mean NFI values over the duration of the study period when compared to other
264 major diagnoses (**Figure 2A**).

265 Conversely, the home villages of study patients tended to have lower mean EVI values than non-
266 study patient villages (**Figure 2C**). Villages from which patients who were diagnosed with JEV were an
267 exception to this general pattern. JEV patient home villages had higher mean EVI when compared to
268 home villages of patients with dengue and cryptococcal infections (**Figure 2C**).

269 Home villages of patients with JEV diagnoses had lower mean NFI over the duration of the study
270 period, but had higher NFI than other major diagnoses when looking specifically at the month of
271 admission (especially when compared to cryptococcal infections and murine typhus (**Figure 3A**)). JEV
272 patient home villages also had higher EVI during the month of admission than most other mono-
273 infections, especially when compared to either cryptococcal infections or murine typhus (**Figure 3C**).
274 Scrub typhus infections had higher EVI than cryptococcal infections when the measurement was taken
275 at the 10km radius buffer (not detectable at smaller radii (**Figure 3C**)).

276 Home villages of patients with dengue infections had higher NFI than murine typhus or
277 *Cryptococcus* spp. patient home villages when considering the 2 months prior to admission (**Figure 3D**).
278 *Cryptococcus* spp. patient home villages had particularly low EVI in the two months leading up to
279 admission, especially when compared to JEV and *Leptospira* spp. patient home villages (**Figure 3F**).
280 At one year prior to admission both *Cryptococcus* spp. patient home and *dengue virus* patient home
281 villages had higher NFI than JEV patient home villages (**Figure 3G**).

282 ***Logistic regressions for geographic, environmental and demographic predictors of CNS infections***

283 In agreement with univariate analyses, villages from which study patients came tended to be
284 larger in population size (**Supporting Figure 4**), lower in elevation (**Supporting Figure 5**), and closer to a
285 major road when compared to other villages within the study area (**Table 4**). They also had higher mean
286 levels of surface flooding, with villages in the highest NFI quadrant having over two times the odds (AOR:

287 2.21; CI: 1.49 – 3.31) of being a home village for study patients when compared to neighboring villages
288 in the study area (**Table 4 and Figure 2**). Villages from which JEV patients originated had few defining
289 characteristics in the logistic regression, other than being larger in population size (AOR: 1.74; CI: 1.55 –
290 1.96) and at lower elevations (AOR: 0.69; CI: 0.46 – 0.97) than non-study patient villages (**Table 5**).

291 In the individual-level analysis (**Table 6**), age and season were the strongest predictors of JEV
292 infections among all patients. Patients who were admitted between July and September had over seven
293 times the odds (AOR: 7.40; CI: 1.45 – 37.67) of being diagnosed with JEV when compared to patients
294 who were admitted between January and March (**Table 6**). JEV was most common in children aged 5
295 through 14 (AOR: 2.74; CI: 1.31 – 5.69; ages 0 – 4 as the comparison group). NFI during the month of
296 admission (10km buffer used in the regression) was also a strong predictor of JEV infection. Individuals
297 who came from villages in the highest quadrant of NFI measurements had approximately 3 times the
298 odds being diagnosed with JEV (AOR: 3.06; CI: 1.04 – 8.96). While study patients came from villages with
299 lower mean elevations, patients who were diagnosed with JEV came from higher elevation villages in
300 comparison to the other patients (AOR: 1.36; CI: 1.11 – 1.66). EVI was a significant predictor in models
301 that did not include distance to road, village population, and elevation (**Table 6 M1 and M2**).

302 **DISCUSSION**

303 Patients from this study were recruited based on symptomology and a medical procedure that is
304 only available at a single location in the nation (diagnostic LP at Mahosot Hospital, Vientiane). The home
305 villages of all included study patients, regardless of diagnosis, were approximately centered on Vientiane
306 City and were closer to major roads than would be expected by chance alone. For many of the infections
307 studied in this analysis, this association is likely more related to geographic and financial access to
308 healthcare systems rather than exposure to environmental risk factors – especially for infections that
309 are more associated with rural areas (e.g. JEV). This finding also suggests that the results here may be a
310 vast underestimate of the true burden of CNS infections, with much of the Lao population not being in

311 near proximity to a major road (**Table 3**). The causative agents of CNS infections differ in biology,
312 ecology, and geography, and this is evident through the spatial distributions of the home villages of
313 patients. The geographic, environmental, and demographic patterns exhibited by patients needing a
314 diagnostic LP for suspected CNS infections, and for specific diagnoses, are the result of complex
315 overlapping factors.

316 A similar spatial pattern was described from an epidemiological analysis of CNS infections
317 among children admitted to Ho Chi Minh City hospitals in Vietnam – with most patients coming from
318 districts near the hospital [18]. While the majority of infections (55%) in the Vietnam study were
319 presumed to be bacterial in origin, in this study from Lao PDR bacterial infections were identified in only
320 38% (170 out of 450 patients with diagnoses).

321 JEV was the single largest identified cause of CNS infections in these data; it primarily affected
322 children (median 13 years of age, **Table 2**), occurred predominantly during the rainy season (likely
323 corresponding to peaks in mosquito vector populations), and in villages with recent high levels of
324 surface water [19]. JEV is a vaccine-preventable disease, but the vaccine has historically been expensive
325 and vaccine programs are frequently limited by access to remote communities. In 2013 the WHO
326 approved a less expensive vaccine (produced in China by the Chengdu Institute of Biological Products)
327 which has since been used in mass vaccination campaigns in Lao PDR in 2013 and 2015 [20]. The vaccine
328 is now routinely given to all children less than 9 months of age but coverage may be low in some areas.

329 The second largest contributor to CNS diseases were cryptococcal infections, which are
330 opportunistic fungal infections with high mortality [21]. Of the 70 patients with cryptococcal infections,
331 12 died prior to discharge and another 8 likely died at home after leaving the hospital. Cryptococcal
332 infections are generally acquired after inhalation of the yeast-like form of the fungus which has been
333 associated with several ecological habitats (*Cryptococcus gattii* has been associated with over 50 species
334 of trees; *Cryptococcus neoformans* has been associated with bird droppings but is also suspected to be

335 associated with plants [22,23]). This disease has a long incubation period [24,25] and while *C. gattii*
336 infections commonly occur among immunocompetent individuals, *C. neoformans* infections are
337 frequently associated with HIV infections [26,27]. In these data, 20 of the patients diagnosed with
338 cryptococcal infections also had HIV infections. While several studies have shown that cryptococcal
339 species exist in specific ecological habitats and have inferred environmental exposure, the long
340 incubation period and complex natural history likely obfuscate ecological correlations.

341 There are several limitations to this research. Diagnostic LPs are only conducted in Mahosot
342 Hospital in the national capital. Logistical and financial difficulties in accessing healthcare facilities, and
343 especially for etiological diagnostic capabilities, likely leads to severe under-reporting of meningitis,
344 encephalitis, or in the diagnosing the causative agent in these conditions when the patient does access
345 care. All of these factors ultimately lead to small case counts for numerous different causative agents.
346 The spatial patterns in points (villages) and ellipses exhibited in these data are likely influenced by the
347 shape of the nation and it is possible that the point patterns and ellipses would differ if we had data
348 from neighboring nations. Spatial and temporal patterns that differentiate different infections might be
349 more obvious if the surveillance system instead focused on any symptomatic infections (rather than only
350 suspected infections of the CNS). Some pathogens are neurotropic whereas others have tropism for
351 other organs, while being capable of occasionally infecting the CNS. This may partially explain the higher
352 case counts of JEV and why we were able to identify spatial, temporal, and environmental predictors for
353 this causative agent.

354 OpenStreetMaps data are volunteered data and may be prone to error. For this reason we
355 focused on major roads, whose routes have changed very little over the last decades. For the
356 regressions, the distances from all villages to the nearest major road was also rounded to the nearest
357 5km. Examination of satellite imagery in comparison with the major roads from OpenStreetMaps
358 suggests that where error does exist, it is on a scale of ± 100 meters, meaning that measurements of

359 distances, as used in this analysis, should not be strongly influenced. Some of these data now come from
360 over a decade ago. Surveillance systems of this type (based on relatively vague symptomology), with a
361 wide panel of possible contributing causative agents, and necessary intensive laboratory components
362 are extremely labor and time intensive.

363 Lao PDR is currently undergoing vast environmental, demographic, and economic changes. Road
364 networks are increasing in range and density and several areas (i.e. Vientiane, Savannakhet) are
365 undergoing expansive urbanization [28]. These environmental changes will most likely result in shifting
366 patterns of infectious diseases. As the region undergoes urbanization (including both a decrease in
367 urban landscape and movement of human populations to urban centers), pathogens that thrive in rural
368 areas (e.g. JEV) may undergo reduced transmission, especially if vaccine campaigns are more capable of
369 reaching rural populations. Conversely, infections that cluster in urban and peri-urban areas (such as
370 dengue and murine typhus) may increase in frequency.

371 Several environmental indices from remote sensing instruments have shown potential for
372 predicting disease risk, differentiating disease types, or for other surveillance efforts in SE Asia [29,30]
373 and globally [31–35]. This analysis, and others like it, illustrates the ability to differentiate some
374 infections (namely JEV when compared to other diagnoses) through the use of freely available data (i.e.
375 MODIS) and software (R and QGIS) and routinely collected healthcare data. Surveillance systems and
376 potentially diagnostic algorithms [36] in developing settings could benefit from inclusion of such
377 resources. A far-reaching surveillance system that is representative of the entire nation and includes
378 likely CNS infections would be beneficial in order to assess the true burden of CNS infections – many of
379 which would benefit from primary and secondary prevention through increased provision of vaccines,
380 vector control, and early diagnosis and treatment. Given the inherent difficulties in accurately
381 diagnosing and treating CNS infections, the predictors reported here and from other epidemiological
382 studies for major contributors to CNS diseases (i.e. age, seasonality, location, and environmental

383 characteristics) could be considered alongside clinical symptomology when presumptive diagnoses are
384 being made. However, it will be important to consider current and ongoing demographic,
385 environmental, and economic changes in Lao PDR.

386 Finally, increasing population access to vaccines, diagnosis, and treatment would have clear
387 benefits to overall population health. As with other parts of the developing world, a large fraction of the
388 Lao population must travel long distances in order to reach primary healthcare centers. In 2005 73% of
389 the Lao population was reported to live in rural areas, 21% without roads. By 2015 67% of the
390 population were reported to live in rural villages with 8% in villages without roads [37]. For many
391 communities, travel during the wet season remains difficult. Travel costs can also be prohibitive. Most of
392 the CNS infections in this analysis occurred or developed symptoms during the wet season. Public
393 health initiatives that help to decrease the distances between communities and the healthcare services
394 that they need are warranted.

395

396

397 **Acknowledgements**

398 We are very grateful to the patients and to Bounthaphany Bounxouei, past Director of Mahosot
399 Hospital; to Bounnack Saysanasongkham, Director of Department of Health Care, Ministry of Health; to
400 H.E. Bounkong Syhavong, Minister of Health, Lao PDR; to the staff of the Infectious Disease Center and
401 Microbiology Laboratory for their help and support.

402 **Funding**

403 This study was supported by the European Commission Innovate program (ComAcross project, grant no.
404 DCI-ASIE/2013/315-047). The work of LOMWRU, PN and MM are funded by the Wellcome Trust of Great
405 Britain, grant 106698/Z/14/Z. Travel to Lao PDR to support DMP for this research was obtained through
406 a University of California Council on Research, Computing, and Libraries (CORCL) Faculty Grant.

407

408 REFERENCES

- 409 1. Tarantola A, Goutard F, Newton P, de Lamballerie X, Lortholary O, Cappelle J, et al. Estimating the
410 Burden of Japanese Encephalitis Virus and Other Encephalitides in Countries of the Mekong
411 Region. *PLoS Neglected Tropical Diseases*. 2014;8: 4. doi:10.1371/journal.pntd.0002533
- 412 2. Glaser C a, Honarmand S, Anderson LJ, Schnurr DP, Forghani B, Cossen CK, et al. Beyond viruses:
413 clinical profiles and etiologies associated with encephalitis. *Clinical infectious diseases*: an official
414 publication of the Infectious Diseases Society of America. 2006;43: 1565–1577.
415 doi:10.1086/509330
- 416 3. Glaser CA, Gilliam S, Schnurr D, Forghani B, Honarmand S, Khetsuriani N, et al. In Search of
417 Encephalitis Etiologies: Diagnostic Challenges in the California Encephalitis Project, 1998–2000.
418 *Clinical Infectious Diseases*. 2003;36: 731–742. doi:10.1086/367841
- 419 4. Solomon T, Michael BD, Smith PE, Sanderson F, Davies NWS, Hart IJ, et al. Management of
420 suspected viral encephalitis in adults – Association of British Neurologists and British Infection
421 Association National Guidelines. *Journal of Infection*. 2012;64: 347–373.
422 doi:10.1016/j.jinf.2011.11.014
- 423 5. McGill F, Heyderman RS, Michael BD, Defres S, Beeching NJ, Borrow R, et al. The UK joint specialist
424 societies guideline on the diagnosis and management of acute meningitis and meningococcal
425 sepsis in immunocompetent adults. *Journal of Infection*. 2016;72: 405–438.
426 doi:10.1016/j.jinf.2016.01.007
- 427 6. Bharucha T, Vickers S, Ming D, Lee SJ, Dubot-Pérès A, de Lamballerie X, et al. Association between
428 reported aetiology of central nervous system infections and the speciality of study investigators—a
429 bias compartmental syndrome? *Trans R Soc Trop Med Hyg*. 2017;111: 579–583.
430 doi:10.1093/trstmh/try008
- 431 7. Kennedy PGE. Viral Encephalitis: Causes, Differential Diagnosis, and Management. *Journal of*
432 *Neurology, Neurosurgery & Psychiatry*. 2004;75: i10–i15. doi:10.1136/jnnp.2003.034280
- 433 8. Olsen SJ, Campbell AP, Supawat K, Liamsuwan S, Chotpitayasunondh T, Laptikulthum S, et al.
434 *Infectious Causes of Encephalitis and Meningoencephalitis*. 2015;21: 2003–2005.
- 435 9. Dittrich S, Rattanaovong S, Lee SJ, Panyanivong P, Craig SB, Tulsiani SM, et al. *Orientia*, *rickettsia*,
436 and *leptospira* pathogens as causes of CNS infections in Laos: A prospective study. *The Lancet*
437 *Global Health*. 2015;3: e104–e112. doi:10.1016/S2214-109X(14)70289-X
- 438 10. Mai NTH, Phu NH, Nhu LNT, Hong NTT, Hanh NHH, Nguyet LA, et al. Central Nervous System
439 Infection Diagnosis by Next-Generation Sequencing: A Glimpse Into the Future? *Open Forum Infect*
440 *Dis*. 2017;4. doi:10.1093/ofid/ofx046
- 441 11. Turner P, Suy K, Tan LV, Sar P, Miliya T, Hong NTT, et al. The aetiologies of central nervous system
442 infections in hospitalised Cambodian children. *BMC Infect Dis*. 2017;17. doi:10.1186/s12879-017-
443 2915-6

- 444 12. Ho NT, Hoang VMT, Le NNT, Nguyen DT, Tran A, Kaki D, et al. A spatial and temporal analysis of
445 paediatric central nervous system infections from 2005 to 2015 in Ho Chi Minh City, Vietnam.
446 *Epidemiology and Infection*. 2017;145: 3307–3317. doi:10.1017/S095026881700228X
- 447 13. Mawuntu AHP, Bernadus JBB, Dhenni R, Wiyatno A, Anggreani R, Feliana, et al. Detection of
448 central nervous system viral infections in adults in Manado, North Sulawesi, Indonesia. *PLOS ONE*.
449 2018;13: e0207440. doi:10.1371/journal.pone.0207440
- 450 14. Dubot-Pérès A, Mayxay M, Phetsouvanh R, Lee SJ, Rattnavong S, Vongsouvath M, et al.
451 Management of Central Nervous System Infections, Vientiane, Laos, 2003–2011 - Volume 25,
452 Number 5—May 2019 - *Emerging Infectious Diseases journal* - CDC. doi:10.3201/eid2505.180914
- 453 15. Boschetti M, Nutini F, Manfron G, Brivio PA, Nelson A. Comparative analysis of normalised
454 difference spectral indices derived from MODIS for detecting surface water in flooded rice
455 cropping systems. *PLoS ONE*. 2014;9. doi:10.1371/journal.pone.0088741
- 456 16. Lao Population and Housing Census 2015: Provisional Report [Internet]. Ministry of Planning and
457 Investment Lao Statistics Bureau; 2015 Dec p. 64. Available:
458 <https://lao.unfpa.org/sites/default/files/pub-pdf/Final%20report-editting-English1.pdf>
- 459 17. Vallé J, Thaojaikong T, Moore CE, Phetsouvanh R, Richards AL, Souris M, et al. Contrasting spatial
460 distribution and risk factors for past infection with scrub typhus and murine typhus in Vientiane
461 city, Lao PDR. *PLoS Neglected Tropical Diseases*. 2010;4: 1–10. doi:10.1371/journal.pntd.0000909
- 462 18. HO NT, HOANG VMT, LE NNT, NGUYEN DT, TRAN A, KAKI D, et al. A spatial and temporal analysis of
463 paediatric central nervous system infections from 2005 to 2015 in Ho Chi Minh City, Vietnam.
464 *Epidemiology and Infection*. 2017;145: 3307–3317. doi:10.1017/S095026881700228X
- 465 19. Campbell G, Hills S, Fischer M, Jacobson J, Hoke C, Hombach J, et al. Estimated global incidence of
466 Japanese encephalitis: *Bulletin of the World Health Organization*. 2011;89: 766–774.
467 doi:10.2471/BLT.10.085233
- 468 20. UNICEF. Japanese Encephalitis Vaccine Market and Supply Update. 2015;
- 469 21. Slavin MA, Chakrabarti A. Opportunistic fungal infections in the Asia-Pacific region. *Medical
470 Mycology*. 2012;50: 18–25. doi:10.3109/13693786.2011.602989
- 471 22. Springer DJ, Chaturvedi V. Projecting global occurrence of *Cryptococcus gattii*. *Emerging Infectious
472 Diseases*. 2010;16: 14–20. doi:10.3201/eid1601.090369
- 473 23. Ellis DH, Pfeiffer TJ. Ecology, life cycle, and infectious propagule of *Cryptococcus neoformans*. *The
474 Lancet*. 1990;336: 923–925. doi:10.1016/0140-6736(90)92283-N
- 475 24. Castrodale LJ, Gerlach RF, Preziosi DE, Frederickson P, Lockhart SR. Prolonged incubation period for
476 *cryptococcus gattii* infection in cat, Alaska, USA. *Emerging Infectious Diseases*. 2013;19: 1034–
477 1035. doi:10.3201/eid1906.130006

- 478 25. Georgi A, Schneemann M, Tintelnot K, Calligaris-Maibach RC, Meyer S, Weber R, et al.
479 *Cryptococcus gattii* meningoencephalitis in an immunocompetent person 13 months after
480 exposure. *Infection*. 2009;37: 370–373. doi:10.1007/s15010-008-8211-z
- 481 26. Wright P, Inverarity D. Human immunodeficiency virus (HIV) related cryptococcol meningitis in
482 rural Central Thailand-Treatment difficulties and prevention strategies. *SOUTHEAST ASIAN J TROP*
483 *MED PUBLIC HEALTH*. 2007;38: 4.
- 484 27. Inverarity D, Bradshaw Q, Wright P, Grant A. The spectrum of HIV-related disease in rural Central
485 Thailand. *SOUTHEAST ASIAN J TROP MED PUBLIC HEALTH*. 2002;33: 10.
- 486 28. Bank AD. Urban development in the Greater Mekong Subregion. Steinberg F, Hakim J, editors.
487 Mandaluyong City, Philippines: Asian Development Bank; 2016. doi:10.1111/j.1467-
488 9787.2010.00709.x
- 489 29. Ledien J, Sorn S, Hem S, Huy R, Buchy P, Tarantola A, et al. Assessing the performance of remotely-
490 sensed flooding indicators and their potential contribution to early warning for leptospirosis in
491 Cambodia. Schumann GJ-P, editor. *PLOS ONE*. 2017;12: e0181044.
492 doi:10.1371/journal.pone.0181044
- 493 30. Chadsuthi S, Chalvet-Monfray K, Wiratsudakul A, Suwancharoen D, Cappelle J. A remotely sensed
494 flooding indicator associated with cattle and buffalo leptospirosis cases in Thailand 2011–2013.
495 *BMC Infectious Diseases*. 2018;18. doi:10.1186/s12879-018-3537-3
- 496 31. Rogers DJ, Randolph SE, Snow RW, Hay SI. Satellite imagery in the study and forecast of malaria.
497 *Nature*. 2002;415: 710–715. doi:10.1038/415710a
- 498 32. Lambin EF, Tran A, Vanwambeke SO, Linard C, Soti V. Pathogenic landscapes: Interactions between
499 land, people, disease vectors, and their animal hosts. *International Journal of Health Geographics*.
500 2010;9: 54. doi:10.1186/1476-072X-9-54
- 501 33. Beck LR, Lobitz BM, Wood BL. Remote sensing and human health: new sensors and new
502 opportunities. *Emerg Infect Dis*. 2000;6: 217–227.
- 503 34. Ford TE, Colwell RR, Rose JB, Morse SS, Rogers DJ, Yates TL. Using Satellite Images of
504 Environmental Changes to Predict Infectious Disease Outbreaks. *Emerg Infect Dis*. 2009;15: 1341–
505 1346. doi:10.3201/eid1509.081334
- 506 35. Lleo MM, Lafaye M, Guell A. Application of space technologies to the surveillance and modelling of
507 waterborne diseases. *Curr Opin Biotechnol*. 2008;19: 307–312. doi:10.1016/j.copbio.2008.04.001
- 508 36. Pokharel S, White LJ, Aguas R, Celhay O, Pellé KG, Dittrich S. Algorithm in the diagnosis of febrile
509 illness using pathogen-specific rapid diagnostic tests. *Clin Infect Dis*. doi:10.1093/cid/ciz665
- 510 37. Results of the Population and Housing Census 2015 [Internet]. Lao Statistics Bureau; 2016 p. 282.
511 Available: https://lao.unfpa.org/sites/default/files/pub-pdf/PHC-ENG-FNAL-WEB_0.pdf
- 512

513 **Supporting information legends**

514

515 **Supporting information file 1: Detailed explanation of variables and analyses**

516

517

518

519

520

521

522

523

524

525

526

527

528

529

530

531

532

533

534

535

536

537

538 **Table 1:** List of variables, their spatial and temporal scales, and transformations

539

Variable	description	spatial scale	temporal scale	transformation
<i>Village population</i>	Population estimate of LP patient home village. This is calculated as a mean population value from the Lao PDR official census from years 2005 and 2015.			For multivariable regressions, this variable was centered on its mean and standardized by its standard deviation.
<i>Distance to major road</i>	This is the distance in meters from the LP patient home village and the nearest point on a major highway network, from the OpenStreetMaps map layer. This distance was transformed into kilometers and rounded to the nearest 5 kilometers in order to account for measurement error.	This distance was transformed into kilometers and rounded to the nearest 5 kilometers in order to account for measurement error.		For multivariable regressions, this variable was centered on its mean and standardized by its standard deviation.
<i>Village elevation</i>	This is the elevation of the LP patient home village, calculated from a digital elevation model. It is in meters above sea level.	at village point		For multivariable regressions, this variable was centered on its mean and standardized by its standard deviation.
<i>NFI</i>	This is the normalized flooding index, described in detail in the Supporting materials.	Mean values at 2km, 5km, and 10km buffers around each village.	For village level analysis: calculated as a mean value for the study duration. For individual level analysis: Calculated for the same month (same 16 day time period); the previous 2 months (mean of the previous 5 16 day intervals); and the previous year (mean of the previous 23 16 day intervals).	Aggregated into quartiles for multivariable regressions.
<i>NDVI</i>	This is the normalized difference vegetation index, detailed in the Supporting materials.	Mean values at 2km, 5km, and 10km buffers around each village.	For village level analysis: calculated as a mean value for the study duration. For individual level analysis: Calculated for the same month (same 16 day time period); the previous 2 months (mean of the previous 5 16 day intervals); and the previous year (mean of the previous 23 16 day intervals).	
<i>EVI</i>	This is the enhanced vegetation index, detailed in the Supporting materials.	Mean values at 2km, 5km, and 10km buffers around each village.	For village level analysis: calculated as a mean value for the study duration. For individual level analysis: Calculated for the same month (same 16 day time period); the previous 2 months (mean of the previous 5 16 day intervals); and the previous year (mean of the previous 23 16 day intervals).	Aggregated into quartiles for multivariable regressions.
<i>Gender</i>	Binary for male or female, self reported in hospital records			
<i>Age</i>	Self reported age in years.			Aggregated into age groups for multivariable
<i>Year</i>	The year of admission to the hospital			For multivariable regressions, this variable was centered on its mean and standardized by its standard deviation.
<i>Quarter</i>	The calendar quarter of admission (Jan - March; Apr - June; July - Sep; Oct - Dec)			

540

541

542

543

544

545

546

547 **Table 2:** Age and gender of study patients. (Q1 and Q3 indicate the first and third quartiles,
548 respectively).

549

	male/female	M/F ratio	median age in years (Q1 –Q3)	Total number
all patients	666/399	1.67	23 (8 - 38)	1065
JEV	55/39	1.41	13 (8 - 20)	94
<i>Cryptococcus</i> spp.	40/30	1.33	33 (27 - 41)	70
scrub typhus	22/9	2.44	16 (9 - 29)	31
<i>Dengue virus</i>	22/5	4.4	20 (9 - 30)	27
<i>Leptospirosis</i> spp.	17/8	2.13	25 (12 - 39)	25
murine typhus	17/7	2.43	32 (16 - 51)	24

550

551

552

553

554 **Table 3:** Distribution of distances (in km) to the nearest major road, by diagnosis type. Counts of villages
555 are from within 3 standard deviational ellipses (SDEs) of all LP villages (referred to as the “study area” in
556 text). In some cases, multiple patients came from the same village meaning that counts of villages will
557 be smaller than counts of total patients. (Q1 and Q3 indicate the first and third quartiles, respectively).

558

	n	median distance in km (Q1 - Q3)
All	6416	5.4 (0.5 - 15.2)
Villages without study patient	5847	6.3 (0.8 - 16.1)
Villages with study patient	569	0.7 (0.1 - 4.1)
JEV	88	0.6 (0.1 - 8.0)
<i>Cryptococcus</i> spp.	66	0.3 (0.1 - 1.4)
scrub typhus	31	0.6 (0.2 - 3.5)
<i>Dengue virus</i>	27	0.3 (0.1 - 1.1)
<i>Leptospirosis</i> spp.	22	0.4 (0.1 - 1.9)
murine typhus	24	0.3 (0.1 - 2.2)

559

560

561 **Table 4:** Logistic regression and model adjusted odds ratios (AOR) for village level analysis of LP villages

562

covariate	total	LP count	AOR (CI)
NDFI Q1	1604	68	reference group
NDFI Q2	1604	75	1.07 (0.73 - 1.56)
NDFI Q3	1604	130	1.48 (1.03 - 2.16)
NDFI Q4	1604	296	2.21 (1.49 - 3.31)
EVI Q1	1604	311	reference group
EVI Q2	1604	134	1.16 (0.86 - 1.57)
EVI Q3	1604	74	1.12 (0.76 - 1.66)
EVI Q4	1604	50	1.19 (0.74 - 1.91)
Village population			2.22 (2.03 - 2.42)
Elevation			0.52 (0.42 - 0.63)
Distance to major road			0.68 (0.57 - 0.80)

563

564 **Table 5:** Logistic regression and model adjusted odds ratios (AOR) for village level analysis of JEV villages

covariate	total	JEV count	AOR (CI)
NDFI Q1	1604	18	reference group
NDFI Q2	1604	15	0.83 (0.38 - 1.76)
NDFI Q3	1604	20	1.11 (0.53 - 2.35)
NDFI Q4	1604	35	1.26 (0.54 - 2.95)
EVI Q1	1604	35	reference group
EVI Q2	1604	23	1.81 (0.87 - 3.76)
EVI Q3	1604	17	1.85 (0.77 - 4.46)
EVI Q4	1604	13	1.76 (0.63 - 4.93)
Village population			1.74 (1.55 - 1.96)
Elevation			0.69 (0.46 - 0.97)
Distance to major road			0.88 (0.62 - 1.20)

565

566

567

568

569

570

571

572 **Table 6:** Mixed effects logistic regression and model adjusted odds ratios (AOR) for individual level
 573 analysis

			M1	M2	M3
covariate	total	JEV count	AOR (CI)	AOR (CI)	AOR (CI)
NDFI Q1	262	7	reference group	reference group	reference group
NDFI Q2	261	16	2.91 (1.06 - 7.97)	2.28 (0.86 - 6.01)	2.32 (0.87 - 6.19)
NDFI Q3	261	31	2.73 (1.02 - 7.28)	2.75 (1.07 - 7.07)	2.98 (1.13 - 7.85)
NDFI Q4	262	38	3.41 (1.17 - 9.94)	3.12 (1.09 - 8.90)	3.06 (1.04 - 8.96)
EVI Q1	262	8	reference group	reference group	reference group
EVI Q2	261	20	2.11 (0.84 - 5.28)	1.89 (0.76 - 4.74)	1.60 (0.63 - 4.04)
EVI Q3	261	24	2.04 (0.81 - 5.12)	1.65 (0.67 - 4.07)	1.33 (0.53 - 3.32)
EVI Q4	262	40	4.19 (1.62 - 10.87)	3.44 (1.35 - 8.73)	2.43 (0.91 - 6.46)
Jan - March	210	2	reference group	reference group	reference group
April - June	267	22	5.12 (1.10 - 23.89)	4.45 (0.95 - 20.76)	5.05 (1.07 - 23.73)
July - Sep	333	62	8.72 (1.75 - 43.46)	6.35 (1.26 - 31.89)	7.40 (1.45 - 37.67)
Oct - Dec	253	8	1.80 (0.35 - 9.25)	1.51 (0.29 - 7.81)	1.54 (0.29 - 8.17)
Year			1.31 (1.03 - 1.68)	1.34 (1.05 - 1.72)	1.27 (0.98 - 1.64)
0 through 4	208	13		reference group	reference group
5 through 14	150	37		3.00 (1.46 - 6.18)	2.74 (1.31 - 5.69)
15 through 24	192	28		1.74 (0.84 - 3.62)	1.37 (0.64 - 2.94)
25 through 34	186	10		0.66 (0.27 - 1.61)	0.61 (0.25 - 1.48)
35 through 44	133	4		0.26 (0.07 - 0.96)	0.26 (0.07 - 0.97)
45 plus	196	2		0.11 (0.02 - 0.52)	0.10 (0.02 - 0.47)
female	399	39		reference group	reference group
male	666	55		0.97 (0.60 - 1.58)	1.08 (0.66 - 1.79)
Village population					1.00 (0.77 - 1.29)
Elevation					1.36 (1.11 - 1.66)
Distance to major road					1.08 (0.88 - 1.32)

574
 575
 576
 577
 578
 579
 580
 581

582 **FIGURES**

583

584 **Figure 1:** Spatial distributions of the home villages of study patients, for A: all study patients, B: study
585 patients with JEV infections, and C: with cryptococcal infections, D: scrub typhus infections, E: with
586 *dengue virus* infections, F: with leptospiral infections, and G: with murine typhus infections. SDDs and
587 SDEs are weighted by case numbers, with some patients coming from the same village. Maps were
588 created using QGIS version 3.4.9. All layers were created by the authors of this manuscript.

589

590 **Figure 2:** Environmental indices for villages with study patient homes for the duration of the study
591 period (January 2003 through August 2011) for all study patient villages, non study patient villages in the
592 study area, and for major diagnoses (JEV = Japanese Encephalitis virus; Crypto = cryptococcal infection;
593 ST = scrub typhus; MT = murine typhus; dengue = *Dengue virus*; lepto = *Leptospira* spp. infection). Bar
594 values are mean values and the error bars are 95% confidence intervals, using the t-distribution. NFI
595 values here have a constant (0.25) added to them for visualization only.

596

597 **Figure 3:** Environmental indices for study patients by major diagnosis and at different times leading up
598 to the date of admission. JEV = Japanese Encephalitis virus; Crypto = cryptococcal infection; ST = scrub
599 typhus; MT = murine typhus; dengue = *Dengue virus*; lepto = *Leptospira* spp. infection. Bar values are
600 mean values and the error bars are 95% confidence intervals, using the t-distribution. NFI values here
601 have a constant (0.25) added to them for visualization only.

602

603