

## **A Systematic Review on Psychological and Biological Mediators Between Adversity and Psychosis: Potential Targets for Treatment**

Luis Alameda<sup>ab</sup>, Victoria Rodriguez<sup>a</sup>, Ewan Carr<sup>c</sup>, Monica Aas<sup>ad</sup>, Giulia Trotta<sup>a</sup>, Paolo Marino<sup>a</sup>, Natasha Vorontsova<sup>a</sup>, Andrés Herane-Vives<sup>ef</sup>, Edoardo Spinazzola<sup>a</sup>, Marta Di Forti<sup>ag</sup>, Craig Morgan<sup>h</sup>, Robin M Murray<sup>a</sup>

<sup>a</sup>Department of Psychosis Studies, Institute of Psychiatry, King's College London, London, UK.

<sup>b</sup>Service of General Psychiatry, Treatment and Early Intervention in Psychosis Program, Lausanne University Hospital (CHUV), CH-1008 Lausanne, Switzerland.

<sup>c</sup>Department of Biostatistics and Health Informatics, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London.

<sup>d</sup>Division of Mental Health and Addiction, Oslo University Hospital, Oslo, Norway.

<sup>e</sup> Centre for Affective Disorders, Department of Psychological Medicine, Institute of Psychiatry, Psychology & Neuroscience, King's College London

<sup>f</sup> Departamento de Clínicas, Facultad de Medicina, Universidad Católica del Norte, Coquimbo, Chile

<sup>g</sup>Social, Genetic and Developmental Psychiatry Centre, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK.

<sup>h</sup>Department of Health Services and Population Research, Institute of Psychiatry Psychology and Neuroscience, King's College London, London, UK

### **Abstract**

#### **Background:**

Various psychological and biological pathways have been proposed as mediators between childhood adverse events (CA) and psychosis. A systematic review of the evidence in this domain is needed.

#### **Aims:**

Systematically review the evidence on clinical and biological mediators between CA and psychosis across the psychosis spectrum.

#### **Methods**

This systematic review followed Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines. Articles published between 1979 and July 2019 were identified through a literature search in OVID (PsychINFO; Medline and Embase). The direction of the evidence by each analysis and each study results were presented by group of mediator categories found in the review; the percentage of total effect mediated was calculated.

## Results

47 studies were included, with a total of 79,668 general population (GP) subjects and 3,189 from clinical samples. The quality of studies was judged as “fair”. We found (i) solid evidence of mediation between CA and psychosis by negative cognitive schemas about the self, the world and others (NS); by dissociation and other PTSD symptoms; (ii) evidence of a partial mediation through an affective pathway (affective dysregulation, anxiety and depression) in GP; (iii) lack of studies exploring biological mediators.

## Conclusions

Experiences of adversity, along with relevant mediators such as PTSD and mood related symptoms and NS should be routinely assessed in patients with psychosis. Targeting such mediators through cognitive behavioural approaches using trauma-focused therapy and/or pharmacological means could be a useful addition to the traditional treatment of positive symptoms.

## Introduction

Evidence has accumulated over the past 15 years showing that exposure to childhood adversity (CA) – in the form of abuse, neglect and bullying – is associated with increased risk of psychosis(1) . Moreover, many studies have demonstrated that patients already suffering from psychosis who were exposed to CA experience higher severity of positive symptoms, particularly hallucinations and delusions, compared with their non-traumatised counterparts(2, 3). However, research to date is less clear on the mechanism involved. Therefore, a systematic review of the evidence about possible mediating mechanisms linking CA and psychosis is needed.

In terms of biology, substantial research on the underlying biological mechanism of the CA-psychosis association has been carried out in the last 15 years. This covers dysfunction in pathways such as the stress response system(4), dopaminergic neurotransmission(5), inflammation and redox dysregulation(6), and changes in stress related brain structures such as the amygdala or the hippocampus(7). For

example, it has been suggested that excessive exposure to stress might lead to an overactivation of the Hypothalamic-Pituitary-Adrenal (HPA) Axis. This could be toxic for hippocampal functioning (8) (9) which might in turn contribute to psychosis.

Furthermore, acute social stress also has been shown to increase striatal dopamine release in individuals at risk for psychosis, in patients with psychosis and in individuals with childhood trauma (5). Additionally, inflammatory dysfunctions as well as redox dysregulation conditions(6, 10) have been identified among individuals exposed to CA and among patients with psychosis.

Several psychological models have also been proposed to explain the relationship between CA and psychosis. One explanation relies on the so called “Affective Pathway to psychosis(11)”. This view postulates that CA may lead to psychosis through a pathway of heightened emotional distress, characterized by hypersensitivity to daily-life stressors, leading to anxiety and depression(11, 12). These symptoms, sometimes called “ancillary symptoms of psychosis” (12) might operate as mediators between adversity and positive symptoms of psychosis.

Secondly, another model proposes that severe forms of adversity might lead to cognitive bias, such as negative schema about the self and the world, which, in addition to a heightened tendency to attribute experiences to external causes, might give rise to paranoia, ideas of reference and misattributions of perceptual abnormalities(13, 14). A third explanatory model relies on Attachment Theory(15), which hypothesizes that adversity occurring within the context of caregiving relationships might disrupt the quality of attachment styles leading to insecure, fearful or anxious attachment. This might, in turn, play a role in psychosis itself, but also in processes contributing to treatment effectiveness, such as therapeutic alliance or engagement with services(16). Finally, another putative pathway linking

adversity with psychosis emphasises the mediating role of Post Traumatic Stress Disorder (PTSD) and related symptoms, such as dissociation, depersonalisation and intrusive memories, very frequently comorbid with psychotic phenomena(17). For example, It is suggested that these experienced could be interpreted as being externally generated, leading to hallucinatory experiences and hampering reality testing(13, 18).

Despite several narrative reviews covering potential biological pathways and psychological mediating pathways(7, 12, 19-23), to date only a single systematic review has been conducted(24). Williams et al.,(24) examined psychological mediators between CA and psychosis including papers up to September 2017, which consisted of a total of 37 studies. They found robust evidence for a mediation by cognitive bias, such as negative schemas about the self and the world; by symptoms of PTSD and by affective processes. However, this review is limited by the fact that (I) they did not analyze and discuss the direction of evidence of all analyses involving mediation within each study; (ii) no summary was given regarding the extent of mediation (e.g. partial or total – see method section for details); (iii) they did not extract the percentage of the total effect mediated; and (iv) their review did not include biological studies.

The present study overcomes those limitations and includes additional papers published up to July 2019, providing an additional period of 22 months of research in the field. Our aim is to systematically review the evidence on potential psychological and biological mediators between CA and psychosis (considered as presenting hallucinations, delusions or disorganisation) including, as an outcome: (i) psychosis onset; (ii) severity of positive symptoms in patients with a psychotic disorder (iii)

severity of attenuated psychotic symptoms in the general population. Results will be grouped based on existing proposed theoretical mechanisms and discussed in terms of potential treatment interventions.

## **Methods**

### ***Search strategy***

A systematic review of the literature was conducted following Preferred Reporting Items for Systematic Reviews and Meta-Analyses PRISMA(25) guidelines. The systematic review was registered in PROSPERO(26) in July 2018 (registration number: CRD42018100846). The main search was conducted on MEDLINE, EMBASE and PsycINFO, through Ovid provider in June 2018 and it was updated in July 2019. We searched Medical Subjects Headings (MeSH) and keywords related to: (1) childhood adversity, including exposure to sexual, physical, emotional abuse, physical and emotional neglect; bullying and early separation from parents; (2) mediation and (3) psychosis, using the Boolean operator 'AND' (full list of search terms provided in **Online Supplementary Material – Search Strategy**). In accord with others(1), the present analyses did not consider separation from parents, abandonment and parental loss.

Titles and abstracts of articles were screened independently by two reviewers (LA and VR) with 95% and 85% agreement (agreement was considered when equal quality assessment scores were) for articles on psychological and biological mediators, respectively. Discrepancies were resolved through discussion at a project group meeting.

### ***Inclusion and exclusion criteria***

Included studies were those that (1) examined psychological or biological mediators of the relationship between CA and (i) psychosis onset, (ii) severity of positive symptoms in patients with a psychotic disorder, or (iii) severity of attenuated psychotic symptoms in the general population; (2) employed a robust method for testing mediation in the analyses and fulfilling the Baron and Kenny criteria(27); (3) used diagnoses of psychosis according to Diagnostic and Statistical Manual of Mental Disorders (DSM-III, DSM-III-R, DSM-IV, DSM IV-TR(28)), Research Diagnostic Criteria or International Classification of Diseases, Ninth or Ten Revision (ICD-9), (ICD-10)(29); (4) assessed both mediators and outcomes using validated methods and scales; (5) included CA occurring before age 18 and involving exposure to sexual, physical and emotional abuse, physical and emotional neglect, bullying or equivalent experiences; (6) had been published as original research; and (7) were performed in humans.

Exclusion criteria were: (1) not being published in English language; (2) including more than 20% of participants aged 65 years or over, in accord with others(30); (3) being performed in homogeneous samples of specific populations such as pregnant women or samples from forensic settings; (4) including psychological mediators that cannot be directly targeted with current psychological or pharmacological interventions (e.g. being exposed to later life hassles, level of education or different aspects of social disadvantage).

### ***Quality Assessment and data extraction methodology***

Quality assessment was carried out using the Newcastle–Ottawa Scale(31) for cohort studies by two independent reviewers (LA and PM). Details on the instrument items and quality assessment procedures can be found in **Supplementary material**

**section.** The agreed quality grades of each study are presented in **Supplementary Material Table S1a, b and c.**

**Tables 1, 2, 3 and S2** show the direction of evidence for each mediator in clinical samples and in general population as follows: we enumerated the number of studies and analyses indicating (i) absence of mediation, (ii) partial mediation, or (iii) total mediation. These results were extracted from the text or tables of each paper. We included all mediational analyses (i.e. distinct pathways tested) from each paper, meaning the total number of analyses was greater than the number of papers.

Where the extent of mediation (absence, partial, total) could not be determined (i.e. authors suggested evidence of a mediating effect but did not quantify the percentage mediated) analyses were summarised as ‘suggested mediation’. In **Tables 1, 2, 3 and S2**, as well as in the text we provided the “percentage of analyses showing evidence of mediation”, across all categories of mediators. We considered that one analysis showed evidence of mediation when a significant partial, total or suggested mediation was found, and negative when null mediation was observed. This allowed us to provide the percentage of analyses supporting mediation per category. Where a mediating effect was reported but the “percentage mediated” was not given we derived an estimate by dividing the indirect effect by the total effect and multiplying it by 100(32). **Figure 2** represents the percentage of the total effect mediated by each analysis across the most meaningful categories described in the results section. The type of adversity and outcome are also shown to have a visual representation of the pathway. As an indicative measure, the median value of all the analyses per category is also highlighted in the figure and presented in the text. Further details on data extraction procedures can be found in **Supplementary materials** and in **Figure 2** footnote.

## Results

**Figure 1** summarizes how we selected the 47 studies included in our review from the 2310 studies found in the initial searches (2018 and 2019 combined).

Nineteen studies were conducted in clinical samples (four in First Episode of Psychosis (FEP) or early psychosis patients (33-36), three in Ultra High Risk (UHR) patients(37-39), 14 in non-FEP patients(40-51)); 26(52-77) in the general population. Our review included 79,668 subjects from the general population and 3,189 subjects from clinical studies, among which 551 were FEP or Early psychosis, 393 were UHR patients and 1,789 were non early psychosis patients. A total of 133 individuals with psychosis were included in biological papers. Participant ages ranged from 18.5 to 44.6 years old in clinical samples (30.4 in average) and between 9.8 to 51.7 in the general population (34.3 in average). In clinical samples, 35% of the participants were females, compared to 37% of the healthy volunteers.

This total number of analyses excludes two studies with extremely high numbers of analysis ((one used a network based approach exploring multiple connexions between adversity and symptoms(43), the other included up to 28(52)). These studies would have distorted the numerical summaries and therefore are described in the text only. Overall, 170 analyses were included in this review (ranging between 1 and 12 analysis per paper).

The quality check agreement rate between the two raters was 81.8%. Overall, quality was graded as 'fair' (between 4 and 7) for all studies except one, where quality was judged as 'good(58). Studies of biological mediators tended to be graded with lower scores failing to provide estimates of the indirect effects, relying on small samples, and/or cross-sectional data(78, 79). Overall, only 4/47 studies(39, 58, 63, 75) were



conducted with a prospective design. Eight(43, 52-55, 62, 65, 78) of the 47 studies reported a mediating effect but failed to provide estimates of the indirect and direct effects and thus were classified as 'suggested mediation', as described above.

**Table 1** summarises the direction of the evidence for each analysis by mediator category. A comprehensive table containing all extracted data including quality scores is available as online **Supplementary material (Table S1a, b, c)**. Given the high number of analyses and the heterogeneity found for the affective pathway, cognitive schema pathway and dissociation, three detailed tables were constructed for these groups (see **Table 2, Table 3** and **Table S2** in **Supplementary material** respectively) presenting the direction of the evidence for each subcategory.

The percentage of total effect mediated could be obtained for 103 out the 124 analyses showing evidence of mediation. **Figure 2** summarises the percentage of total effect mediated for 57 analyses included in 24 papers across the most relevant category of mediators found in our review (only categories being explored in at least three papers are shown in **Figure 2**, the percentage of total effect mediated of the remaining analyses are shown in **Tables S1a, b** and **c**).

The most relevant categories of mediators found in our review were: Dissociation (42 analyses, 12 papers), dysfunctional attachment (18 analyses, 5 papers), affective pathway to psychosis (32 analyses, 14 papers), loneliness (6 analyses, 5 papers), cognitive schemas (53 analyses, 20 papers), other PTSD symptoms (7 analyses, 4 papers), other psychological mediators (10 analyses, 6 papers), biological mediators (2 analyses, 2 papers).

As a whole, in terms of amount of mediation between CA and psychosis and the proportion of analyses showing evidence of mediation across all categories, we found (i) solid evidence of mediation between CA and psychosis by negative

cognitive schemas about the self, the world and others as well as solid evidence of mediation by PTSD symptoms, particularly dissociation. We found evidence of a partial mediation through an affective pathway (particularly for affective dysregulation, anxiety and depression) in the general population. Although only explored in a limited number of studies, the feeling of loneliness appeared to mediate the link between CA and psychosis. There was no evidence supporting mediating role of the remaining categories.

### ***Dissociation***

Despite only 43% of analyses showing evidence of mediation, 10/12 studies (36, 44, 45, 49, 53, 56, 60, 68, 70, 76, 77) were supportive of mediation, against 2/12 that were not (33, 80). The percentage of total effect mediated (which is the amount of the adversity-psychosis association that is explained by the mediator – see methods and **Figure 2** footnote for details) was consistently high across analyses within this category, with a median of around 50% of total effect explained. Moreover, as illustrated in **Figure 2**, most of the studies showing evidence of mediating effects used hallucinations as outcome and explored trauma as a composite score (44, 49, 53, 56, 60, 68, 76), while only 2 (36, 56) studies found positive effects with delusions as outcome (see also **Tables S1a** and **1b** for details). As shown in **Table S2**, due to high levels of heterogeneity for specific items within the dissociative category (depersonalisation, absorption, dissociative amnesia and defensive dissociation), it was not possible to draw conclusions about which dissociative symptom was more likely to mediate the adversity-psychosis association.

### ***Dysfunctional attachment***

Despite overall 43% of analyses showing evidence of mediation, 3/5(40, 71, 73) papers showed mixed findings, 1/5(48) showed no evidence of mediation and just 1/5(61) showed consistent evidence of mediation across all its analyses. As seen in **Figure 2**, the percentages of total effect mediated were the lowest among all categories, with a median value of around 12% of effect mediated. Studies examining different outcomes and types of traumas were not enough to allow conclusions about specific mediational pathways.

### ***Affective Pathway***

Overall, only one paper explored this pathway in clinical samples(39) showing no evidence of mediation. The remaining 13(43, 52, 57, 58, 60, 63-66, 70, 74-76) papers (including 29 analyses) were conducted in general population, with 66% of analyses supporting mediation (Ashford et al.,(52) and Isvoranu et al.,(43) not included in the analyses count as mentioned above). 11/13(43, 52, 57, 58, 60, 63-65, 70, 74, 75) of these general population studies showed evidence of mediation and 3 of them were prospective graded high in quality assessment(58, 63, 75) **Stable 1b**). This was the category with the highest number of analyses showing evidence of partial mediation (see **Table 1**).

As mentioned before, we divided the Affective Pathway into the following discrete subcategories: Anxiety, Depression, Affective dysregulation, Stress sensitivity and a residual composite measure (details on the direction of evidence can be found in **Table 2**). Briefly, in terms of proportion of analyses supporting mediation, results per subcategories showed that 67% of analyses for anxiety, 55.5% for depression, 62.5% for affective dysregulation and 100% for stress sensitivity showed evidence of mediation. As seen in **Figure 2**, the percentages of total effect mediated (the amount

of the adversity-psychosis association that is explained by the mediator) for anxiety and depression ranged to around 20% of total effect mediated; while affective dysregulation's percentages rose up to approximately 35%. As illustrated in **Figure 2**, the majority of studies examined a composite category of attenuated psychotic symptoms as outcome and tended to examine specific types of adversities such as abuse and bullying.

### ***Loneliness***

83% of analyses showed evidence of mediation. Only one study(46) was conducted in clinical samples and showed evidence for mediation. In general population, two studies showed evidence of mediation(54, 62) and another showed mixed findings(72). The low number of analyses did not allow us to draw consistent conclusions in terms of specific pathways between adversity-psychosis (see **Table S1a, S1b** for details).

### **Cognitive schemas**

53% of analyses overall showed evidence of mediation in this category. In terms of papers, 15/20 were supportive of mediation (33, 35, 37, 38, 42, 51-53, 55, 58, 60, 62, 67, 69, 74), 5/20 were not(34, 50, 57, 68, 76). Results were more likely to show evidence of mediation when conducted in the general population, compared to clinical samples (65% versus 38% of analyses were supportive of mediation respectively, see **Table 1**).

As displayed in **Table 3**, we divided this category into: Negative schemas about self, others and the world and External Locus of control (ELC). Briefly, the direction of the evidence for the former showed that 62.5% of analyses were supportive of

mediation); with a percentage of total effect mediated (the amount of the adversity-psychosis association that is explained by the mediator) of around 47% (median value across analyses in this category – see **Figure 2**). No studies in clinical samples explored the ELC and 2.3 studies in general population showed evidence of mediation(58, 60). Other biases such as maladaptive schemas(53, 55) or low self-esteem(58) showed interesting findings but were not explored specifically due to insufficient number of studies to draw conclusions. In terms of specific adversities and outcomes, studies were very heterogeneous thus not allowing us to draw conclusions about specific pathways.

### ***Other PTSD symptoms***

These included posttraumatic intrusions, avoidance & numbing or a general measure of posttraumatic symptoms. Overall, results showed evidence of mediation(35, 41, 42, 66). The percentage of total effect mediated (the amount of the adversity-psychosis association that is explained by the mediator) was also high, with a median value just below 50% (see **Stable 1a, 1b**)

### ***Other psychological mediators***

Other mediators including 4 studies containing 10 analyses that did not fit into the aforementioned categories were grouped in a residual group. Mediators such as Anomalous self-experiences(59) and Time perspective capabilities(47) showed promising results, while mania showed no mediation effects(39). MacCarthy-Jones et al.,(66) showed that while there was a positive mediating effect for compulsions, this was not the case for obsessions (see **Stable 1a, 1b**).

### ***Biological mediators***

Surprisingly, only 2 studies examining biological mechanism as potential mediators between CA and psychosis fulfilled the inclusion criteria for this systematic review (**Stable 1c**). One study found no evidence for a mediation of the inferior frontal gyrus activation between CA and the Positive and Negative Syndrome Scale (PANSS) positive scores(79); another showed that the grey matter density in Dorsolateral prefrontal Cortex (DLPFC) was mediating the link between emotional neglect and disorganized symptoms in patients with psychosis(81).

### **Discussion**

From this systematic review of 47 papers, we found consistent evidence of mediation between CA and psychosis particularly by negative cognitive schemas about the self, the world and others; and by dissociation and other PTSD symptoms. For general population samples there was good evidence for mediation through an affective pathway (affective dysregulation, anxiety and depression), but there were insufficient studies using clinical samples to allow conclusions to be drawn about its mediating role in patients.

Our systematic review found no evidence supporting the mediating role of dysfunctional attachment styles in the general population; there were too few studies addressing this mediator in patient samples to permit conclusions. Other mediators showing interesting findings included loneliness, stress sensitivity, or external locus of control, but generalisation of findings is limited due to the low number of studies, however they should be taken into account in future research. Contrary to our expectations, only two papers(78, 79) fulfilled our inclusion criteria examining biological mediators, showing an urgent need for more research in this field.

## **Evidence for mediational pathways between childhood adversity and psychosis**

### *Pathway 1: Dissociation and PTSD symptoms*

Mediation through this pathway was more common when hallucinations were used as an outcome. These two categories, intimately related, showed the highest percentages of total effect mediated (**Figure 2**).

The role of post-traumatic dissociation in psychosis was already the focus of research at the end of the eighteenth century, when Janet, among others, defined Hysterical Psychosis (HP) as characterised by its dissociative and stress related nature(82) . Despite these longstanding connections between dissociation and psychosis, current international classifications do not include dissociation among the diagnostic criteria for any form of psychosis. However, contemporary authors such as Ross(83) and Moskowitz(82) suggest the existence of a dissociative type of psychosis that could potentially be responsive to psychotherapy focused on trauma. Accordingly, our findings showing mediation by other PTSD symptoms in clinical samples(35, 41, 42), are consistent with reports suggesting that similar mechanisms could be involved in psychotic experiences and symptoms of PTSD. For instance, it has been suggested that some hallucinations represent a dissociated type of posttraumatic intrusion, which may not be recognized as such by people with psychosis(18, 82). A clinical implication is that if a person experiencing verbal auditory hallucinations recognized the identity or content of a heard voice as corresponding to the content of a previous traumatic event, the intrusion could be reappraised as a type of intrusive memory. It could then be understood in the same way as a as a re-experiencing symptom of PTSD (instead of an externally attributed

voice reflective of psychosis(13)), and could be addressed therapeutically using trauma-focused therapies, such as cognitive behavioural therapy (CBT). Our results therefore support the possibility of applying specific psychological interventions for patients with psychosis, dissociation and PTSD symptoms and with a history of clear-cut traumatic episode. In this regard, two recent systematic reviews exploring the safety and efficacy of trauma focused therapies in individuals with comorbid PTSD symptoms and psychosis have shown that trauma focused CBT is safe(84) and can reduce PTSD symptoms(85). Moreover, a meta-analysis of 12 studies also explored its impact on positive symptoms and found small improvements after treatment(86).

#### *Pathway 2: Cognitive schemas*

Cognitive schemas about the self, the world and others were the most consistent mediator between CA and psychosis in both clinical samples and general population. Moreover, this category contributed highly to the total effect between CA-psychosis, with median percentage mediated of around 46% across all analyses (**Figure 2**). This evidence supports previous cognitive models of the development of positive symptoms(13, 87): these models suggest that exposure to severe trauma might contribute to the development of cognitive bias such as negative schemas about the world, others and the self. Further exposure to stressors or to subtle perceptual abnormalities that are common in the general population(30), and even more common in genetically predisposed individuals, will lead to anomalous conscious self-experiences which will trigger the search for an explanation. The biased schemas, in combination with anomalous self-experience disrupt the appraisal process leading to a misinterpretation of reality, and subsequently delusional



ideas(13, 87, 88). Other factors are important in this process, such as the tendency to search for an explanation in an external source (otherwise called External locus of control(80)) which although not explored intensively in our review, has shown evidence for mediation, especially in the general population as detailed above(58, 60). It would be interesting for future research to explore if other cognitive biases such as jumping to conclusions and meta cognitive deficits can moderate or mediate the connexions between CA and psychosis in combination with biases such as negative schemas and External Locus of control.

### *Pathway 3: Affective pathway*

We found consistent evidence suggesting that anxiety, depression and affective dysregulation might partially mediate the link between CA and psychosis in general population. This was the category with the highest number of analyses showing partial mediation with median values of percentages of total effects mediated ranging from around 20 to 35 % (see **Figure 2**). This suggests that the affective component could be a mediational partner along with other mechanisms such as cognitive bias or PTSD related symptoms. For example, Fisher et al.,(58) report on a large prospective study, where 100% of the total effect was mediated only when anxiety and depression were added to the model in combination with cognitive bias and low self-esteem. Unfortunately, our interpretation is restricted to the general population (and thus to attenuated positive symptoms) due to the lack of studies performed in clinical settings, which limits the applicability to possible clinical interventions. Nevertheless, these results support previous claims that the association between social adversity and psychosis might be mediated by non-psychotic symptoms (otherwise the so called “ancillary symptoms of psychosis”(12) such as anxiety and

depression and affective dysregulation). Stress sensitivity related to daily life activities can lead to negative affect in vulnerable individuals previously exposed to CA, which in turn can affect mood and anxiety. These symptoms might be determinants of paranoid thinking, for example; anxiety might lead to anticipation of threat; and low mood might drive negatively biased interpretations of ongoing experience and impact self-esteem and negative schemas about the self, which in turn are precursors of psychotic symptoms, as illustrated in **Figure 3**.

#### *Pathway 4: Loneliness*

Although only explored in 4 studies(46, 54, 62, 72), we found good evidence suggesting that a feeling of loneliness might mediate the CA-psychosis relationship. We could hypothesize that social withdrawal and loneliness may increase an individual's sensitivity to potential stressors in daily life restrict access to balanced information from the environment, maintaining biased cognitive schemas. This could, in turn, predispose an individual to lower mood and anxiety, which would constitute a favourable ground for the emergence of psychotic symptoms. Thus, the feeling of loneliness and isolation must be taken very seriously as it might operate as a potential condition allowing other mediators to operate

#### *Pathway 5: Biological measures*

Despite the high emphasis put on the search for biomarkers during the last 20 years of psychiatric research, we could identify just two eligible papers exploring biological mechanisms using psychosis as an outcome(78, 79), which prevents us from drawing conclusions about specific pathways. This highlights the limited evidence in this field and need for future studies addressing different biomarkers comparing

between groups of individuals with or without trauma, and to conduct longitudinal designs allowing us to test the influence of biological processes on the onset and evolution of symptoms of psychosis.

### **Clinical implications**

Our systematic review provides evidence to support some potential treatments for traumatised individuals with psychosis, in addition to the treatment of positive symptoms, which often remain high in this vulnerable group(2, 3). **Figure 3** displays a model with the different potential treatment targets derived from the pathways mentioned above. Beyond the traditional treatment of psychotic symptoms (Target 1 in **Figure 3**) to which some traumatised patients do not respond, we propose targeting the relevant mediators found in our review, hypothesizing that an improvement in such targets would then have an indirectly beneficial effect on positive symptoms.

A common clinical picture corresponds to a situation where a traumatised patient with psychosis suffers from negative schemas about the self, others and the world and PTSD symptoms, including dissociation. If these are present (Target 2 A in **Figure 3**), a trauma focused therapy, such as trauma-focused cognitive behavioural therapy (CBT), could be appropriate, integrating treatment of negative schemas and traumatic intrusions, in addition to dissociative symptoms. Briefly, this could include grounding techniques, imaginal exposure, memory updating and cognitive restructuring, alongside more general CBT elements such as the use of behavioural experiments to gather new information about current safety(17). In some cases, dissociative symptoms may be very disruptive and prevent access to the other targets; in this case they can be prioritized and targeted using sensory grounding

techniques to help the individual reliably regain contact with present external stimuli(89, 90).

In addition to psychological interventions, some patients with highly distressing PTSD symptoms might respond to adjunctive pharmacological treatments such as Selective serotonin reuptake inhibitors (SSRIs) or other antidepressants blocking  $\alpha_1$  adrenoceptors commonly used in patients with PTSD(91). This integrated approach could be adapted depending on which mediators are present or predominant, as presented in **Figure 3**. For example, if there are no post-traumatic symptoms, the work on the schemas would be prioritised; similarly, the work on dissociative symptoms alone would be prioritised if these are predominant. Thus, clinicians should carefully address subthreshold anxiety and depressive symptoms in traumatised individuals with psychosis. These could be, in isolation or in combination with other features (such as cognitive bias and posttraumatic symptoms) further targets for treatment (Target 3 **Figure 3**) that could indirectly ameliorate positive symptoms as suggested in previous reviews(12). The use of antidepressants and CBT techniques such as behavioural activation and graded exposure(92), as well as relaxation techniques and mindfulness, could then be useful, targeting low mood and anxiety respectively(92).

Lastly, it is important to highlight the possible role of loneliness as another mediator that could be targeted for treatment. Promoting social inclusion and community membership through group interventions and vocational support could be beneficial, as well as improving existing interpersonal relationships using family therapy approaches. Also, as previously mentioned, loneliness might co-occur with other mediators.

In addition to pointing towards possible therapeutic interventions, our review indicates routine assessment of trauma history capitalises in patients suffering from psychosis. Furthermore, given the possible importance of mediators, it is also important to carry out a careful assessment of the cognitive schemas, PTSD, dissociative symptoms, anxiety, mood and feelings of loneliness. These mediators can then be targeted through tailored psychological and/or pharmacological means, depending on which of them is predominant in the clinical picture.

### **Strengths and limitations**

The findings of this review should be interpreted in the context of various strengths and limitations. A major strength is the large scale of this review including 47 studies, 79,668 subjects from the general population and 3,189 from clinical studies. This has allowed us to cover multiple mediator groups with a sufficient number of participants and show new pathways that did not appear in Williams et al., systematic review(24), such as the role of loneliness. Second, we have not limited our analysis to the description of studies but also examined the extent to which the mediation was total or partial, as well as presented the amount of mediation (percentage of total effect mediated) between pathways which provides a more accurate understanding of the mediational processes, for example suggesting the partial mediating role of the affective pathway. Moreover, we believe that our concrete clinical implications may be useful and could contribute to a better knowledge on trauma-informed care in services treating with individuals with psychosis.

Despite these positive aspects, some limitations must be mentioned: first, only three studies used a prospective design to estimate the indirect effects, while in the remaining studies it cannot be excluded that the mediator resulted as a consequence

of psychosis. Second, CA was measured retrospectively in all studies except two(58, 75) which were conducted in the general population. In patients this can constitute a risk of bias given the difficulties of patients to recall their experiences and to disclose these openly in assessment by research assistants, which is especially true in people with psychosis. Third, in 8 studies we recorded mediation as “suggested”, given that the significance of the direct effect after the introduction of the mediators was not reported, not allowing us to determine whether the mediation was total or partial. In the same way, the percentage of total effect could not be obtained nor calculated in 17% of the analyses that were supportive of mediation, thus our **Figure 2** is not totally representative of the total number of analyses included in this review. In addition, some mediators (i.e. attachment styles, anxiety and depression) were found to be highly explored in the general population but very little in clinical samples, which is an important limitation of current literature in the field and which limits the extent to which we can extrapolate our conclusions to clinical intervention. Lastly, contrary to what we had expected, studies examining mediation through biological variables were limited in number and tended to use less robust mediation analyses, which emphasise the need to further perform this type of method in biological settings.

## **Conclusions**

Our review suggests that cognitive schemas about the self and the world and post-traumatic symptoms (particularly dissociation) are key mechanistic variables linking childhood adversity with psychosis, other factors such as mood and feeling of loneliness seem to contribute partially to this link and interact with other mediators in contributing to the effect. These findings support the routine assessment of

experiences of childhood adversity in clinical settings, alongside with all the potential mediators. An integrated approach targeting mediators such as mood or PTSD symptoms with psychological and/or medical means could potentially be a beneficial add-on to the traditional treatment of positive symptoms.

## Funding

This work was supported by the Swiss National Science Foundation (Grant P2LAP3\_171804 to L.A.)

This paper represents independent research part funded by the National Institute for Health Research (NIHR) Biomedical Research Centre at South London and Maudsley NHS Foundation Trust and King's College London. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health and Social Care. South-Eastern Norway Health Authority (#2017060) and the NARSAD Young Investigator Award to Monica Aas (#22388).

## References

1. Varese F, Smeets F, Drukker M, Lieverse R, Lataster T, Viechtbauer W, et al. Childhood adversities increase the risk of psychosis: a meta-analysis of patient-control, prospective-and cross-sectional cohort studies. *Schizophrenia bulletin*. 2012; 38(4): 661-71.
2. Alameda L, Golay P, Baumann PS, Ferrari C, Do KQ, Conus P. Age at the time of exposure to trauma modulates the psychopathological profile in patients with early psychosis. *J Clin Psychiatry*. 2016; 77(5): e612-8.
3. Ajnakina O, Trotta A, Oakley-Hannibal E, Di Forti M, Stilo SA, Kolliakou A, et al. Impact of childhood adversities on specific symptom dimensions in first-episode psychosis. *Psychol Med*. 2016; 46(2): 317-26.
4. Ruby E, Polito S, McMahon K, Gorovitz M, Corcoran C, Malaspina D. Pathways Associating Childhood Trauma to the Neurobiology of Schizophrenia. *Frontiers in psychological and behavioral science*. 2014; 3(1): 1-17.
5. Howes OD, McCutcheon R, Owen MJ, Murray RM. The Role of Genes, Stress, and Dopamine in the Development of Schizophrenia. *Biological psychiatry*. 2017; 81(1): 9-20.
6. Steullet P, Cabungcal JH, Monin A, Dwir D, O'Donnell P, Cuenod M, et al. Redox dysregulation, neuroinflammation, and NMDA receptor hypofunction: A "central hub" in schizophrenia pathophysiology? *Schizophr Res*. 2016; 176(1): 41-51.

7. van Winkel R, van Nierop M, Myin-Germeys I, van Os J. Childhood trauma as a cause of psychosis: linking genes, psychology, and biology. *The Canadian Journal of Psychiatry*. 2013; 58(1): 44-51.
8. Aas M, Haukvik UK, Djurovic S, Tesli M, Athanasiu L, Bjella T, et al. Interplay between childhood trauma and BDNF val66met variants on blood BDNF mRNA levels and on hippocampus subfields volumes in schizophrenia spectrum and bipolar disorders. *Journal of psychiatric research*. 2014; 59: 14-21.
9. Teicher MH, Anderson CM, Polcari A. Childhood maltreatment is associated with reduced volume in the hippocampal subfields CA3, dentate gyrus, and subiculum. *Proceedings of the National Academy of Sciences of the United States of America*. 2012; 109(9): E563-72.
10. Alameda L, Fournier M, Khadimallah I, Griffa A, Cleusix M, Jenni R, et al. Redox dysregulation as a link between childhood trauma and psychopathological and neurocognitive profile in patients with early psychosis. *Proceedings of the National Academy of Sciences*. 2018; 115(49): 12495-500.
11. Myin-Germeys I, van Os J. Stress-reactivity in psychosis: evidence for an affective pathway to psychosis. *Clinical psychology review*. 2007; 27(4): 409-24.
12. Bebbington P. Unravelling psychosis: psychosocial epidemiology, mechanism, and meaning. *Shanghai Arch Psychiatry*. 2015; 27(2): 70-81.
13. Morrison AP, Frame L, Larkin W. Relationships between trauma and psychosis: A review and integration. *British Journal of Clinical Psychology*. 2003; 42(4): 331-53.
14. Garety PA, Bebbington P, Fowler D, Freeman D, Kuipers E. Implications for neurobiological research of cognitive models of psychosis: a theoretical paper. *Psychological medicine*. 2007; 37(10): 1377-91.
15. Gumley A, Taylor H, Schwannauer M, MacBeth A. A systematic review of attachment and psychosis: measurement, construct validity and outcomes. *Acta Psychiatrica Scandinavica*. 2014; 129(4): 257-74.
16. Read J, Gumley A. Can attachment theory help explain the relationship between childhood adversity and psychosis? *Attachment*. 2008; 2(1): 1-35.
17. Hardy A. Pathways from trauma to psychotic experiences: a theoretically informed model of posttraumatic stress in psychosis. *Frontiers in psychology*. 2017; 8: 697.
18. Allen JG, Coyne L, Console DA. Dissociative detachment relates to psychotic symptoms and personality decompensation. *Comprehensive psychiatry*. 1997; 38(6): 327-34.
19. Morgan C, Gayer Anderson C. Childhood adversities and psychosis: evidence, challenges, implications. *World Psychiatry*. 2016; 15(2): 93-102.
20. Read J, Fosse R, Moskowitz A, Perry B. The traumagenic neurodevelopmental model of psychosis revisited. *Neuropsychiatry*. 2014; 4(1): 65-79.
21. Bentall RP, de Sousa P, Varese F, Wickham S, Sitko K, Haarmans M, et al. From adversity to psychosis: pathways and mechanisms from specific adversities to specific symptoms. *Social psychiatry and psychiatric epidemiology*. 2014; 49(7): 1011-22.
22. Freeman D, Garety P. Advances in understanding and treating persecutory delusions: a review. *Social psychiatry and psychiatric epidemiology*. 2014; 49(8): 1179-89.
23. Misiak B, Krefft M, Bielawski T, Moustafa AA, Sasiadek MM, Frydecka D. Toward a unified theory of childhood trauma and psychosis: a comprehensive review of epidemiological, clinical, neuropsychological and biological findings. *Neuroscience & Biobehavioral Reviews*. 2017; 75: 393-406.
24. Williams J, Bucci S, Berry K, Varese F. Psychological mediators of the association between childhood adversities and psychosis: a systematic review. *Clinical psychology review*. 2018; 65: 175-96.



25. Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Annals of internal medicine*. 2009; 151(4): 264-9.
26. Moher D, Booth A, Stewart L. How to reduce unnecessary duplication: use PROSPERO. *BJOG: An International Journal of Obstetrics & Gynaecology*. 2014; 121(7): 784-6.
27. Hayes AF. Beyond Baron and Kenny: Statistical mediation analysis in the new millennium. *Communication monographs*. 2009; 76(4): 408-20.
28. Diagnostic APA. statistical manual of mental disorders. American Psychiatric Association. Washington, DC. 1994: 886.
29. Organization WH. The ICD-10 classification of mental and behavioural disorders: clinical descriptions and diagnostic guidelines. Geneva: World Health Organization, 1992.
30. Van Os J, Linscott RJ, Myin-Germeys I, Delespaul P, Krabbendam L. A systematic review and meta-analysis of the psychosis continuum: evidence for a psychosis proneness–persistence–impairment model of psychotic disorder. *Psychological medicine*. 2009; 39(2): 179-95.
31. Stang A. Critical evaluation of the Newcastle-Ottawa scale for the assessment of the quality of nonrandomized studies in meta-analyses. *European journal of epidemiology*. 2010; 25(9): 603-5.
32. Alwin DF, Hauser RM. The decomposition of effects in path analysis. *American sociological review*. 1975: 37-47.
33. Evans GJ, Reid G, Preston P, Palmier-Claus J, Sellwood W. Trauma and psychosis: The mediating role of self-concept clarity and dissociation. *Psychiatry Research*. 2015; 228(3): 626-32.
34. Morgan C, Reininghaus U, Fearon P, Hutchinson G, Morgan K, Dazzan P, et al. Modelling the interplay between childhood and adult adversity in pathways to psychosis: initial evidence from the AESOP study. *Psychological medicine*. 2014; 44(2): 407-19.
35. Peach N, Alvarez-Jimenez M, Cropper SJ, Sun P, Bendall S. Testing models of post-traumatic intrusions, trauma-related beliefs, hallucinations, and delusions in a first episode psychosis sample. *British Journal of Clinical Psychology*. 2019; 58(2): 154-72.
36. Sun P, Alvarez-Jimenez M, Simpson K, Lawrence K, Peach N, Bendall S. Does dissociation mediate the relationship between childhood trauma and hallucinations, delusions in first episode psychosis? *Comprehensive psychiatry*. 2018; 84: 68-74.
37. Appiah-Kusi E, Fisher H, Petros N, Wilson R, Mondelli V, Garety P, et al. Do cognitive schema mediate the association between childhood trauma and being at ultra-high risk for psychosis? *Journal of psychiatric research*. 2017; 88: 89-96.
38. McDonnell J, Stahl D, Day F, McGuire P, Valmaggia L. Interpersonal sensitivity in those at clinical high risk for psychosis mediates the association between childhood bullying victimisation and paranoid ideation: a virtual reality study. *Schizophrenia research*. 2018; 192: 89-95.
39. Thompson A, Marwaha S, Nelson B, Wood SJ, McGorry PD, Yung AR, et al. Do affective or dissociative symptoms mediate the association between childhood sexual trauma and transition to psychosis in an ultra-high risk cohort? *Psychiatry research*. 2016; 236: 182-5.
40. Chatziioannidis S, Andreou C, Agorastos A, Kaprinis S, Malliaris Y, Garyfallos G, et al. The role of attachment anxiety in the relationship between childhood trauma and schizophrenia-spectrum psychosis. *Psychiatry research*. 2019; 276: 223-31.
41. Choi JY, Choi YM, Kim B, Lee DW, Gim MS, Park SH. The effects of childhood abuse on self-reported psychotic symptoms in severe mental illness: Mediating effects of posttraumatic stress symptoms. *Psychiatry research*. 2015; 229(1-2): 389-93.


42. Hardy A, Emsley R, Freeman D, Bebbington P, Garety PA, Kuipers EE, et al. Psychological mechanisms mediating effects between trauma and psychotic symptoms: the role of affect regulation, intrusive trauma memory, beliefs, and depression. *Schizophrenia Bulletin*. 2016; 42(suppl\_1): S34-S43.
43. Isvoranu A-M, van Borkulo CD, Boyette L-L, Wigman JT, Vinkers CH, Borsboom D, et al. A network approach to psychosis: pathways between childhood trauma and psychotic symptoms. *Schizophrenia bulletin*. 2016; 43(1): 187-96.
44. Perona-Garcelán S, Carrascoso-López F, García-Montes JM, Ductor-Recuerda MJ, López Jiménez AM, Vallina-Fernández O, et al. Dissociative experiences as mediators between childhood trauma and auditory hallucinations. *Journal of Traumatic Stress*. 2012; 25(3): 323-9.
45. Schalinski I, Breinlinger S, Hirt V, Teicher MH, Odenwald M, Rockstroh B. Environmental adversities and psychotic symptoms: The impact of timing of trauma, abuse, and neglect. *Schizophrenia research*. 2017.
46. Steenkamp L, Weijers J, Gerrmann J, Eurelings-Bontekoe E, Selten J-P. The relationship between childhood abuse and severity of psychosis is mediated by loneliness: an experience sampling study. *Schizophrenia research*. 2019.
47. Styła R, Stolarski M, Szymanowska A. Linking childhood adversities with schizophrenia: A mediating role of the balanced time perspective. *Schizophrenia research*. 2019.
48. Van Dam D, Korver-Nieberg N, Velthorst E, Meijer C, de Haan L. Childhood maltreatment, adult attachment and psychotic symptomatology: a study in patients, siblings and controls. *Social psychiatry and psychiatric epidemiology*. 2014; 49(11): 1759-67.
49. Varese F, Barkus E, Bentall R. Dissociation mediates the relationship between childhood trauma and hallucination-proneness. *Psychological medicine*. 2012; 42(5): 1025-36.
50. Weijers J, Fonagy P, Eurelings-Bontekoe E, Termorshuizen F, Viechtbauer W, Selten J. Mentalizing impairment as a mediator between reported childhood abuse and outcome in nonaffective psychotic disorder. *Psychiatry research*. 2018; 259: 463-9.
51. Wickham S, Bentall R. Are specific early-life adversities associated with specific symptoms of psychosis?: A patient study considering just world beliefs as a mediator. *The Journal of nervous and mental disease*. 2016; 204(8): 606.
52. Ashford CD, Ashcroft K, Maguire N. Emotions, traits and negative beliefs as possible mediators in the relationship between childhood experiences of being bullied and paranoid thinking in a non-clinical sample. *Journal of Experimental Psychopathology*. 2012; 3(4): 624-38.
53. Bortolon C, Seillé J, Raffard S. Exploration of trauma, dissociation, maladaptive schemas and auditory hallucinations in a French sample. *Cognitive neuropsychiatry*. 2017; 22(6): 468-85.
54. Boyda D, McFeeters D. Childhood maltreatment and social functioning in adults with sub-clinical psychosis. *Psychiatry research*. 2015; 226(1): 376-82.
55. Boyda D, McFeeters D, Dhingra K, Rhoden L. Childhood maltreatment and psychotic experiences: Exploring the specificity of early maladaptive schemas. *Journal of clinical psychology*. 2018; 74(12): 2287-301.
56. Cole CL, Newman-Taylor K, Kennedy F. Dissociation mediates the relationship between childhood maltreatment and subclinical psychosis. *Journal of Trauma & Dissociation*. 2016; 17(5): 577-92.
57. Fisher HL, Appiah-Kusi E, Grant C. Anxiety and negative self-schemas mediate the association between childhood maltreatment and paranoia. *Psychiatry research*. 2012; 196(2-3): 323-4.

58. Fisher HL, Schreier A, Zammit S, Maughan B, Munafò MR, Lewis G, et al. Pathways between childhood victimization and psychosis-like symptoms in the ALSPAC birth cohort. *Schizophrenia bulletin*. 2012; 39(5): 1045-55.
59. Gawęda Ł, Göritz AS, Moritz S. Mediating role of aberrant salience and self-disturbances for the relationship between childhood trauma and psychotic-like experiences in the general population. *Schizophrenia research*. 2019; 206: 149-56.
60. Gibson LE, Reeves LE, Cooper S, Olinio TM, Ellman LM. Traumatic life event exposure and psychotic-like experiences: A multiple mediation model of cognitive-based mechanisms. *Schizophrenia research*. 2019; 205: 15-22.
61. Goodall K, Rush R, Grünwald L, Darling S, Tiliopoulos N. Attachment as a partial mediator of the relationship between emotional abuse and schizotypy. *Psychiatry research*. 2015; 230(2): 531-6.
62. Jaya ES, Ascone L, Lincoln TM. Social adversity and psychosis: the mediating role of cognitive vulnerability. *Schizophrenia bulletin*. 2016; 43(3): 557-65.
63. Lincoln TM, Marin N, Jaya ES. Childhood trauma and psychotic experiences in a general population sample: a prospective study on the mediating role of emotion regulation. *European Psychiatry*. 2017; 42: 111-9.
64. Marwaha S, Bebbington P. Mood as a mediator of the link between child sexual abuse and psychosis. *Social psychiatry and psychiatric epidemiology*. 2015; 50(4): 661-3.
65. Marwaha S, Broome MR, Bebbington PE, Kuipers E, Freeman D. Mood instability and psychosis: analyses of British national survey data. *Schizophrenia bulletin*. 2013; 40(2): 269-77.
66. McCarthy-Jones S. Post-traumatic symptomatology and compulsions as potential mediators of the relation between child sexual abuse and auditory verbal hallucinations. *Behavioural and cognitive psychotherapy*. 2018; 46(3): 318-31.
67. Murphy S, Murphy J, Shevlin M. Negative evaluations of self and others, and peer victimization as mediators of the relationship between childhood adversity and psychotic experiences in adolescence: the moderating role of loneliness. *British Journal of Clinical Psychology*. 2015; 54(3): 326-44.
68. Perona-Garcelán S, García-Montes JM, Rodríguez-Testal JF, López-Jiménez AM, Ruiz-Veguilla M, Ductor-Recuerda MJ, et al. Relationship between childhood trauma, mindfulness, and dissociation in subjects with and without hallucination proneness. *Journal of Trauma & Dissociation*. 2014; 15(1): 35-51.
69. Pinto-Gouveia J, Matos M, Castilho P, Xavier A. Differences between depression and paranoia: the role of emotional memories, shame and subordination. *Clinical psychology & psychotherapy*. 2014; 21(1): 49-61.
70. Rössler W, Ajdacic-Gross V, Rodgers S, Haker H, Müller M. Childhood trauma as a risk factor for the onset of subclinical psychotic experiences: Exploring the mediating effect of stress sensitivity in a cross-sectional epidemiological community study. *Schizophrenia research*. 2016; 172(1-3): 46-53.
71. Sheinbaum T, Kwapil TR, Barrantes-Vidal N. Fearful attachment mediates the association of childhood trauma with schizotypy and psychotic-like experiences. *Psychiatry research*. 2014; 220(1-2): 691-3.
72. Shevlin M, McElroy E, Murphy J. Loneliness mediates the relationship between childhood trauma and adult psychopathology: evidence from the adult psychiatric morbidity survey. *Social psychiatry and psychiatric epidemiology*. 2015; 50(4): 591-601.
73. Sitko K, Bentall RP, Shevlin M, Sellwood W. Associations between specific psychotic symptoms and specific childhood adversities are mediated by attachment styles: an analysis of the National Comorbidity Survey. *Psychiatry research*. 2014; 217(3): 202-9.


74. van Nierop M, Van Os J, Gunther N, Van Zelst C, De Graaf R, Ten Have M, et al. Does social defeat mediate the association between childhood trauma and psychosis? Evidence from the NEMESIS-2 S study. *Acta Psychiatrica Scandinavica*. 2014; 129(6): 467-76.
75. Wolke D, Lereya S, Fisher H, Lewis G, Zammit S. Bullying in elementary school and psychotic experiences at 18 years: a longitudinal, population-based cohort study. *Psychological medicine*. 2014; 44(10): 2199-211.
76. Yamasaki S, Ando S, Koike S, Usami S, Endo K, French P, et al. Dissociation mediates the relationship between peer victimization and hallucinatory experiences among early adolescents. *Schizophrenia Research: Cognition*. 2016; 4: 18-23.
77. Bortolon C, Raffard S. Dissociation Mediates the Relationship Between Childhood Trauma and Experiences of Seeing Visions in a French Sample. *The Journal of nervous and mental disease*. 2018; 206(11): 850-8.
78. Cancel A, Comte M, Truillet R, Boukezzi S, Rousseau P, Zendjidjian X, et al. Childhood neglect predicts disorganization in schizophrenia through grey matter decrease in dorsolateral prefrontal cortex. *Acta Psychiatrica Scandinavica*. 2015.
79. Quidé Y, O'Reilly N, Watkeys O, Carr V, Green M. Effects of childhood trauma on left inferior frontal gyrus function during response inhibition across psychotic disorders. *Psychological medicine*. 2018; 48(9): 1454-63.
80. Thompson A, Sullivan S, Lewis G, Zammit S, Heron J, Horwood J, et al. Association between locus of control in childhood and psychotic symptoms in early adolescence: results from a large birth cohort. *Cognitive neuropsychiatry*. 2011; 16(5): 385-402.
81. Cancel A, Comte M, Truillet R, Boukezzi S, Rousseau PF, Zendjidjian X, et al. Childhood neglect predicts disorganization in schizophrenia through grey matter decrease in dorsolateral prefrontal cortex. *Acta Psychiatrica Scandinavica*. 2015; 132(4): 244-56.
82. Moskowitz A, Schäfer I, Dorahy MJ. Psychosis, trauma and dissociation. *Emerging Perspectives on Severe Psychopathology*. 2008.
83. Ross CA. Dissociation and psychosis: The need for integration of theory and practice. 2006.
84. Sin J, Spain D. Psychological interventions for trauma in individuals who have psychosis: a systematic review and meta-analysis. *Psychosis*. 2017; 9(1): 67-81.
85. Swan S, Keen N, Reynolds N, Onwumere J. Psychological interventions for post-traumatic stress symptoms in psychosis: a systematic review of outcomes. *Frontiers in psychology*. 2017; 8: 341.
86. Brand RM, McEnery C, Rossell S, Bendall S, Thomas N. Do trauma-focussed psychological interventions have an effect on psychotic symptoms? A systematic review and meta-analysis. *Schizophrenia research*. 2018; 195: 13-22.
87. Garety PA, Kuipers E, Fowler D, Freeman D, Bebbington P. A cognitive model of the positive symptoms of psychosis. *Psychological medicine*. 2001; 31(2): 189-95.
88. Howes OD, Murray RM. Schizophrenia: an integrated sociodevelopmental-cognitive model. *The Lancet*. 2014; 383(9929): 1677-87.
89. Keen N, Hunter E, Peters E. Integrated trauma-focused cognitive-behavioural therapy for post-traumatic stress and psychotic symptoms: a case-series study using imaginal reprocessing strategies. *Frontiers in psychiatry*. 2017; 8: 92.
90. Steel C, Hardy A, Smith B, Wykes T, Rose S, Enright S, et al. Cognitive-behaviour therapy for post-traumatic stress in schizophrenia. A randomized controlled trial. *Psychological Medicine*. 2017; 47(1): 43-51.
91. Steckler T, Risbrough V. Pharmacological treatment of PTSD—established and new approaches. *Neuropharmacology*. 2012; 62(2): 617-27.

92. Waller H, Garety P, Jolley S, Fornells-Ambrojo M, Kuipers E, Onwumere J, et al. Low intensity cognitive behavioural therapy for psychosis: a pilot study. *Journal of behavior therapy and experimental psychiatry*. 2013; 44(1): 98-104.

**Figure 1. Flow chart**




**Figure 2**



We managed to obtain the percentage of total effect mediated for for 83.1% of the analyses that found a significant indirect effect (103 out of the 124 positive analyses, 44 of which were already calculated and 59 which were calculated as described in **Supplementary Material - Data extraction procedures** section). In this table are presented the percentage of the most meaningful categories of mediators found in our results. As presented in the right-hand side of the figure, the type of predictor is detailed by different symbols and the type of outcome is detailed by different colours.

Figure 3

medRxiv preprint doi: <https://doi.org/10.1101/2019.12.14.19014506>; this version posted December 18, 2019. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. All rights reserved. No reuse allowed without permission.



This figure displays the potential targets for treatment that could be addressed in individuals with psychosis who have been exposure to adversity. Beyond the traditional treatment of psychotic symptoms (**Target 1**) different clinical presentations with various targets for treatment can occur: **Target 2** represent targets from the posttraumatic spectrum such as Negative Schemas about the Self, Others and the World (NS) and PTSD related symptoms, including Dissociation. These can be combined in different forms as displayed in **Target 2A**, **2B** and **2C**. **Target 3** represents mood-related symptoms and **Target 4** represent the presence of feeling of loneliness. The treatment options could be adapted depending on which targets are present or predominant in the clinical picture. For **Target 2 A**: CBT-trauma focused therapy (for PTSD symptoms and NS) + sensory grounding techniques (for dissociation) + SSRI/ antagonists  $\alpha_1$  if required (for PTSD symptoms). The implementation of these interventions would be adapted according to the presence/absence of the aforementioned targets (**Target 2 B and C**). **Target 3**: CBT techniques such as behavioural activation and graded exposure, relaxation techniques and mindfulness + SSRI if needed for anxiety and depressive symptoms. **Target 4**: Promoting social inclusion and community membership through therapy/activity groups and vocational support. Improving existing interpersonal relationships using family therapy approaches. The thickness of the arrows shows the robustness of the evidence according to our results. The red arrows represent a hypothetical connexion not addressed in our review.



**Table 1.** Summary of evidence for mediators between adversity and psychosis

Category ( <i>N total studies</i> )	Clinical samples					General population					Total  % analyses supporting mediation (N total analyses)*
	Direction of evidence <i>Number of analyses (number of studies)</i>					Direction of evidence <i>Number of analyses (number of studies)</i>					
	Total mediation	Null mediation	Partial mediation	Suggested mediation	% analyses supporting mediation (N total analyses)*	Total mediation	Null mediation	Partial mediation	Suggested mediation	% analyses supporting mediation (N total analyses)*	
<b>Dissociation (12)</b>	4(3)	20(6)	2(1)	-	<b>23% (26)</b>	6(5)	5(1)	5(2)	1(1)	<b>70% (17)</b>	<b>42% (43)</b>
<b>Dysfunctional attachment (5)</b>	-	3(2)	1(1)	-	<b>25% (4)</b>	1(1)	7(2)	6(3)	-	<b>50% (14)</b>	<b>44% (18)</b>
<b>Affective pathway (14)<sup>ab</sup></b>	-	3(1)	-	-	<b>0% (3)</b>	10(5)	8(5)	8(6)	3(1)	<b>72% (29)</b>	<b>66% (32)</b>
<b>Loneliness (4)</b>	1(1)	-	-	-	<b>100% (1)</b>	2(1)	2(1)	1(1)	1(1)	<b>66% (6)</b>	<b>83% (6)</b>
<b>Cognitive schemas (20)<sup>b</sup></b>	5(4)	13(5)	3(2)	-	<b>38% (21)</b>	8(3)	12(6)	5(4)	7(4)	<b>62% (32)</b>	<b>53% (53)</b>
<b>Others PTSD symptoms (4)</b>	4(2)	1(1)	1(1)	-	<b>83% (6)</b>	-	-	1(1)	-	<b>100% (1)</b>	<b>86% (7)</b>
<b>Others<sup>c</sup> (6)</b>	1(1)	1(1)	-	-	<b>50% (2)</b>	4(1)	1(1)	3(2)	-	<b>87% (8)</b>	<b>80% (10)</b>

<sup>a</sup>Isvoranu et al., was only included in the articles count within but not in the analyses count. <sup>b</sup>Ashford et al., was only included in the articles count but not in the analyses count. <sup>c</sup>Others included anomalous self-experiences; mania; time-perspective capabilities and compulsions as mediators. \*We considered one analysis as “supportive of mediation” when a significant mediation (partial, total or “suggested”) was found, and negative when null mediation was found.

**Table 2.** Summary of evidence for mediators within the Affective Pathway category between adversity and psychosis

Category (N total studies)	Clinical samples					General population					Total  % analyses supporting mediation (N total analyses)*
	Direction of evidence					Direction of evidence					
	Number of analyses (number of studies)					Number of analyses (number of studies)					
	Total mediation	Null mediation	Partial mediation	Suggested mediation	% analyses supporting mediation (N total analyses)*	Total mediation	Null mediation	Partial mediation	Suggested mediation	% analyses supporting mediation (N total analyses)*	
<b>Affective pathway (14)<sup>ab</sup></b>	-	3(1)	-	-	<b>0% (3)</b>	10(5)	8(5)	8(6)	3(1)	<b>72% (29)</b>	<b>66% (32)</b>
<b>Anxiety (4)</b>	-	1(1)	-	-	<b>0% (1)</b>	3(2)	2(2)	1(1)	-	<b>67% (6)</b>	<b>67% (7)</b>
<b>Depression (6)</b>	-	1(1)	-	-	<b>0% (1)</b>	2(1)	4(3)	3(2)	-	<b>55% (9)</b>	<b>55.5% (10)</b>
<b>Emotional dysregulation (4)</b>	-	1(1)	-	-	<b>0% (1)</b>	-	2(2)	2(2)	3(1)	<b>71% (7)</b>	<b>62.5% (8)</b>
<b>Stress sensitivity/perception (2)</b>	-	-	-	-	-	4(2)	-	1(1)	-	<b>100% (5)</b>	<b>100% (5)</b>
<b>Overall (1)</b>	-	-	-	-	-	1	-	1	-	<b>100% (2)</b>	<b>100% (2)</b>

<sup>a</sup>Isvoranu et al., was only included in the articles count within but not in the analyses count. <sup>b</sup>Ashford et al., was only included in the articles count but not in the analyses count. \*We considered one analysis as “supportive of mediation” when a significant partial, total or suggested. We considered one analysis as “supportive of mediation” when a significant mediation (partial, total or “suggested”) was found, and negative when null mediation was found

**Table 3.** Summary of evidence for mediators within the Cognitive Schemas category between adversity and psychosis

Category (N total studies)	Clinical samples					General population					Total  % analyses supporting mediation (N total analyses)*
	Direction of evidence					Direction of evidence					
	Number of analyses (number of studies)					Number of analyses (number of studies)					
	Total mediation	Null mediation	Partial mediation	Suggested mediation	% analyses supporting mediation (N total analyses)*	Total mediation	Null mediation	Partial mediation	Suggested mediation	% analyses supporting mediation (N total analyses)*	
Cognitive schemas (21) <sup>b</sup>	5(4)	13(5)	3(2)	-	<b>38% (21)</b>	8(3)	12(7)	5(4)	7(4)	<b>62% (32)</b>	<b>53% (53)</b>
Negative schemas (8)	3(2)	2(1)	2(1)	-	<b>71% (7)</b>	3(2)	4(1)	1(1)	1(1)	<b>55% (9)</b>	<b>62.5% (16)</b>
ELC (3)	-	-	-	-	-	3(2)	1(1)	1(1)	-	<b>80% (5)</b>	<b>80% (5)</b>

<sup>b</sup>Ashford et al., was only included in the articles count but not in the analyses count. We considered one analysis as “supportive of mediation” when a significant mediation (partial, total or “suggested”) was found, and negative when null mediation was found