

1 **Doubling Time of the COVID-19 Epidemic by Chinese Province**

2 **Manuscript Number:** EID-20-0219.R3 (Accepted for publication) This paper will be available in the
3 journal Emerging Infectious Diseases:

4 Muniz-Rodriguez K, Chowell G, Cheung CH, Jia D, Lai PY, Lee Y, et al. Doubling Time of the COVID-
5 19 Epidemic by Chinese Province. Emerg Infect Dis. 2020 Aug. 26(8). In press.

6 **Running Title:** COVID-19 doubling time by Chinese province

7 **Keywords:** China; Coronavirus; COVID-19; Disease Transmission, Infectious; Epidemiology;
8 Respiratory Tract Infections

9 **Authors:** Kamalich Muniz-Rodriguez, MPH¹; Gerardo Chowell, PhD¹; Chi-Hin Cheung, MS; Dongyu
10 Jia, PhD; Po-Ying Lai, MS; Yiseul Lee, MPH, Manyun Liu, MS; Sylvia K. Ofori, MPH; Kimberlyn M.
11 Roosa, MPH; Lone Simonsen, PhD; Cecile Viboud, PhD; Isaac Chun-Hai Fung, PhD*

12 ¹ These authors contribute equally as co-first authors

13 Address for correspondence: Isaac Chun-Hai Fung, Department of Biostatistics, Epidemiology and
14 Environmental Health Sciences, Jiann-Ping Hsu College of Public Health, Georgia Southern University,
15 Statesboro, GA 30460, USA; email: cfung@georgiasouthern.edu

16 **Affiliations:**

17 Georgia Southern University, Statesboro, GA, USA (K. Muniz-Rodriguez, D. Jia, M. Liu, S. K. Ofori, I.
18 C.-H. Fung); Georgia State University, Atlanta, GA, USA (G. Chowell, Y. Lee, K. M. Roosa);
19 Independent researcher (C.-H. Cheung); Boston University, Boston, MA, USA (P.-Y. Lai); Roskilde
20 University, Roskilde, Denmark (L. Simonsen); The National Institutes of Health, Bethesda, MD, USA (G.
21 Chowell, C. Viboud)

22

23 **Abstract: 50 words**

24 COVID-19 epidemic doubling time by Chinese province was increasing from January 20 through
25 February 9, 2020. The harmonic mean of the arithmetic mean doubling time estimates ranged from 1.4
26 (Hunan, 95% CI, 1.2-2.0) to 3.1 (Xinjiang, 95% CI, 2.1-4.8), with an estimate of 2.5 days (95% CI, 2.4-
27 2.6) for Hubei.

28 **Main text – 791 words**

29 Our ability to estimate the basic reproduction number of emerging infectious diseases is often
30 hindered by the paucity of information about the epidemiological characteristics and transmission
31 mechanisms of new pathogens (1). Alternative metrics could synthesize real-time information about the
32 extent to which the epidemic is expanding over time. Such metrics would be particularly useful if they
33 rely on minimal and routinely collected data that capture the trajectory of an outbreak (2).

34 Epidemic doubling times characterize the sequence of intervals at which the cumulative incidence
35 doubles (3). If an epidemic is growing exponentially with a constant growth rate r , the doubling time
36 remains constant and equals to $(\ln 2)/r$. An increase in the doubling time indicates a slowdown in
37 transmission if the underlying reporting rate remains unchanged (Technical Appendix) (4).

38 Here we analyzed by province the number of times COVID-19 cumulative incidence doubled and
39 the evolution of the doubling times in mainland China (5), from January 20 (when nationwide reporting
40 began) through February 9, 2020. Province-level daily cumulative incidence data were retrieved from
41 provincial health commissions' websites. Two sensitivity analyses based on a longer and a shorter time
42 period respectively were conducted (Technical Appendix). Tibet was excluded from further analysis
43 because there had only been one case reported during the study period.

44 From January 20 through February 9, the harmonic mean of the arithmetic means of the doubling
45 times estimated from cumulative incidence ranged from 1.4 (95% CI, 1.2, 2.0) days (Hunan) to 3.1 (95%
46 CI, 2.1, 4.8) days (Xinjiang). In Hubei, it was estimated as 2.5 (95% CI, 2.4, 2.6) days. The cumulative
47 incidence doubled 6 times in Hubei. The harmonic mean of the arithmetic means of doubling times in all
48 of mainland China except Hubei was 1.8 (95% CI, 1.5, 2.3) days. Provinces with a harmonic mean of the
49 arithmetic means of doubling times $<2d$ included Fujian, Guangxi, Hebei, Heilongjiang, Henan, Hunan,
50 Jiangxi, Shandong, Sichuan, and Zhejiang (Figures 1 and S1).

51 As the epidemic progressed, it took longer for the cumulative incidence in mainland China
52 (except Hubei) to double itself, which indicated an overall sub-exponential growth pattern outside Hubei
53 (Figures S1, S2). In Hubei, the doubling time decreased and then increased. A gradual increase in the
54 doubling time coincided with the social distancing measures and intra-and-inter-provincial travel
55 restrictions imposed across China since the implementation of quarantine of Wuhan on January 23 (6).

56 Our estimates of doubling times are shorter than prior estimates of 7.4 days (95% CI, 4.2-14) (5),
57 6.4 days (95% CrI, 5.8-7.1) (7), and 7.1 days (95% CI, 3.0-20.5) (8) respectively. Li et al. covered cases
58 reported by January 22 (5). Wu et al. statistically inferred case counts in Wuhan by internationally
59 exported cases as of January 25 (7). Volz et al. identified a common viral ancestor on December 8, 2019
60 using Bayesian phylogenetic analysis and fitted an exponential growth model to provide the epidemic
61 growth rate (8). Our estimates are based on cumulative confirmed case count by reporting date by
62 province from January 20 through February 9.

63 Our study is subject to limitations, including underreporting of cases (9). One reason for
64 underreporting is underdiagnosis, due to lack of diagnostic tests, healthcare workers, and other resources.
65 Further, underreporting is likely heterogeneous across provinces. As long as reporting remains invariant
66 over time within the same province, the calculation of doubling times remains reliable; however, this is a
67 strong assumption. Growing awareness of the epidemic and increasing availability of diagnostic tests
68 might have strengthened reporting over time, which could have artificially shortened the doubling time.
69 Nevertheless, apart from Hubei and Guangdong (first case reported on January 19), nationwide reporting
70 only began on January 20, and at this point, Chinese authorities had openly acknowledged the magnitude
71 and severity of the epidemic. Due to a lack of detailed case data describing incidence trends for imported
72 and local cases, we focused our analysis on the overall trajectory of the epidemic without adjusting for the
73 role of imported cases on the local transmission dynamics. Indeed, it is likely that the proportion of
74 imported cases was significant for provinces that only reported a few cases; their short doubling times in
75 the study period could simply reflect rapid detection of imported cases. However, with the data until

76 February 9, only two provinces had a cumulative case count of <40 (Table S1). It would be interesting to
77 investigate the evolution of the doubling time after accounting for case importations if more detailed data
78 becomes available.

79 To conclude, we observed an increasing trend in the epidemic doubling time of COVID-19 by
80 Chinese province from January 20 through February 9, 2020. The harmonic mean of the arithmetic means
81 of doubling times of cumulative incidence in Hubei during the study period was estimated at 2.5 (95% CI,
82 2.4, 2.6) days.

83 **Acknowledgements**

84 GC acknowledges support from NSF grant 1414374 as part of the joint NSF-NIH-USDA Ecology and
85 Evolution of Infectious Diseases program. ICHF acknowledges salary support from the National Center
86 for Emerging and Zoonotic Infectious Diseases, Centers for Disease Control and Prevention
87 (19IPA1908208). This article is not part of ICHF's CDC-sponsored projects.

88 **Disclaimers**

89 This article does not represent the official positions of the Centers for Disease Control and Prevention, the
90 National Institutes of Health, or the United States Government.

91 **Author Bio**

92 Kamalich Muniz-Rodriguez, MPH, is a doctoral student at the Jiann-Ping Hsu College of Public Health,
93 Georgia Southern University. Her research interests include infectious disease epidemiology, digital
94 epidemiology and disaster epidemiology.

95 Gerardo Chowell, PhD, is Professor of Epidemiology and Biostatistics, and Chair of the Department of
96 Population Health Sciences at Georgia State University School of Public Health. As a mathematical
97 epidemiologist, Prof Chowell studies the transmission dynamics of emerging infectious diseases, such as
98 Ebola, MERS and SARS.

99 **References**

- 100 1. Anderson RM, May RM. Infectious diseases of humans. Oxford: Oxford University
101 Press; 1991.
- 102 2. Drake JM, Bakach I, Just MR, O'Regan SM, Gambhir M, Fung IC-H. Transmission
103 Models of Historical Ebola Outbreaks. Emerging Infectious Disease journal.
104 2015;21(8):1447.
- 105 3. Vynnycky E, White RG. An Introduction to Infectious Disease Modelling. Oxford:
106 Oxford University Press; 2010.
- 107 4. Muniz-Rodriguez K, Fung IC-H, Ferdosi SR, Ofori SK, Lee Y, Tariq A, et al.
108 Transmission potential of SARS-CoV-2 in Iran, 2020. Emerging Infectious Disease journal.
109 2020;26(8):DOI: <https://doi.org/10.3201/eid2608.200536>.
- 110 5. Li Q, Guan X, Wu P, Wang X, Zhou L, Tong Y, et al. Early Transmission Dynamics in
111 Wuhan, China, of Novel Coronavirus-Infected Pneumonia. N Engl J Med. 2020 Mar 26
112 382(13):1199-207.
- 113 6. Du Z, Wang L, Cauchemez S, Xu X, Wang X, Cowling BJ, et al. Risk for Transportation
114 of Coronavirus Disease from Wuhan to Other Cities in China. Emerging Infectious Disease
115 journal. 2020 May 17;26(5):1049-52.
- 116 7. Wu JT, Leung K, Leung GM. Nowcasting and forecasting the potential domestic and
117 international spread of the 2019-nCoV outbreak originating in Wuhan, China: a modelling
118 study. Lancet. 2020 Feb 29;395(10225):689-97.
- 119 8. Volz E, Baguelin M, Bhatia S, Boonyasiri A, Cori A, Cucunubá Z, et al. Report 5:
120 Phylogenetic analysis of SARS-CoV-2. 2020 [cited Apr 15, 2020]; Available from:
121 [https://www.imperial.ac.uk/media/imperial-college/medicine/sph/ide/gida-
122 fellowships/Imperial-College-COVID19-phylogenetics-15-02-2020.pdf](https://www.imperial.ac.uk/media/imperial-college/medicine/sph/ide/gida-fellowships/Imperial-College-COVID19-phylogenetics-15-02-2020.pdf)
- 123 9. Fang G, Li S, Liu Y, Xin N, Ma K. People beyond the statistics: Died of "normal
124 pneumonia"? [统计数字之外的人：他们死于“普通肺炎”？]. Caijing Zachi [财经杂志]
125 (Finance and Economics Magazine) 2020 [cited Feb 13, 2020]; Available from:
126 [https://web.archive.org/web/20200213190623/http://www.sohu.com/a/370032279_12
127 0094087](https://web.archive.org/web/20200213190623/http://www.sohu.com/a/370032279_120094087)

128

129


130

131 Address for correspondence: Isaac Chun-Hai Fung, Department of Biostatistics, Epidemiology and
132 Environmental Health Sciences, Jiann-Ping Hsu College of Public Health, Georgia Southern University,
133 Statesboro, GA 30460, USA; email: cfung@georgiasouthern.edu


134 **Figure legend**


135 **Figure 1.** The map of the harmonic mean of the arithmetic means of doubling time estimates (Panel A)
136 and the number of times the COVID-19 epidemic cumulative incidence has doubled (Panel B) by
137 province in mainland China, from January 20 through February 9, 2020.

138



Harmonic mean of the arithmetic means of COVID-19 epidemic doubling times





Doubling Time of the COVID-19 Epidemic by Chinese Province

Technical Appendix

Kamalich Muniz-Rodriguez, MPH¹; Gerardo Chowell, PhD¹; Chi-Hin Cheung, MS; Dongyu Jia, PhD; Po-Ying Lai, MS; Yiseul Lee, MPH, Manyun Liu, MS; Sylvia K. Ofori, MPH; Kimberlyn M. Roosa, MPH; Lone Simonsen, PhD; Cecile Viboud, PhD; Isaac Chun-Hai Fung, PhD

¹ These authors contribute equally as co-first authors

Author affiliations: Georgia Southern University (K. Muniz-Rodriguez, D. Jia, M. Liu, S. K. Ofori, I. C.-H. Fung); Georgia State University (G. Chowell, Y. Lee, K. M. Roosa); Independent researcher (C.-H. Cheung); Boston University (P.-Y. Lai), Roskilde University (L. Simonsen), The National Institutes of Health (G. Chowell, C. Viboud)

Please address correspondence to Isaac Chun-Hai Fung, PhD, Department of Biostatistics, Epidemiology and Environmental Health Sciences, Jiann-Ping Hsu College of Public Health, Georgia Southern University, Statesboro, GA 30460, USA. +1 912 478 5079. Email: cfung@georgiasouthern.edu

Technical appendix

Additional information on our motivation, scope and methods

Motivation. R_0 is a widely used indicator of transmission potential in a totally susceptible population and is driven by the average contact rate and the mean infectious period of the disease (1). Yet, it only characterizes transmission potential at the onset of the epidemic and varies geographically for a given infectious disease according to local healthcare provision, outbreak response, as well as socioeconomic and cultural factors. Furthermore, estimating R_0 requires information about the natural history of the infectious disease. Thus, our ability to estimate reproduction numbers for novel infectious diseases is hindered by the paucity of information about their epidemiological characteristics and transmission mechanisms. More informative metrics could synthesize real-time information about the extent to which the epidemic is expanding over time. Such metrics would be particularly useful if they rely on minimal data on the outbreak's trajectory (2).

Scope and definition. Our analysis is restricted to mainland China in this paper. A 'province' herein encompasses three different types of political sub-divisions of mainland China, namely, a province, a centrally (literally, 'directly') administered municipality (Beijing, Chongqing, Shanghai, and Tianjin) and an 'ethnic minority' autonomous region (Guangxi, Inner Mongolia, Ningxia, Tibet, and Xinjiang). Our analysis does not include the Hong Kong Special Administrative Region and the Macau Special Administrative Region, which are under the effective rule of the People's Republic of China through the so-called 'One Country, Two Systems' political arrangements. Likewise, our analysis does not include Taiwan, which is *de facto* governed by a different government (the Republic of China).

Data sources. Daily cumulative incidence data were retrieved from provincial health commissions' websites (Table S8). Data were double-checked against the cumulative national total published by the National Health Commission (3), data compiled by the Centre for Health Protection, Hong Kong, when

available (4) and by John Hopkins University Center for Systems Science and Engineering (5). Whenever discrepancies arose, provincial government sources were deemed authoritative.

Doubling time calculation and its relationship with growth rate of an epidemic

As the epidemic grows, the times at which cumulative incidence doubles are given by t_{d_i} such that

$2C(t_{d_i}) = C(t_{d_{i+1}})$ where $t_{d_0} = 0$, $C(t_{d_0}) = C_0$, and $i = 0, 1, 2, 3, \dots, n_d$ where n_d is the total number of

times cumulative incidence doubles. The actual sequence of “doubling times” are defined as follows:

$$d_j = \Delta t_{d_j} = t_{d_j} - t_{d_{j-1}} \text{ where } j = 1, 2, 3, \dots, n_d.$$

To quantify parameter uncertainty, we used parametric bootstrapping with a Poisson error structure around the harmonic mean of doubling times d_j to obtain the 95% confidence interval. See references (6-8) for further details.

If we assume homogeneous mixing (equal probability of acquiring infection through contacts) and exponential growth, then, $C(t_2) = C(t_1)\exp(rt)$, and therefore, $\ln(C(t_2)/C(t_1)) = rt$. When $C(t_2)/C(t_1) = 2$ and thus t is the doubling time, i.e. $t = t_d$, $\ln 2 = rt_d$. Therefore, the doubling time, t_d , equals to $(\ln 2)/r$. See Vynnycky and White (9), panel 4.1, p.74 for further explanation.

Additional details on methods. Doubling time calculation was conducted using MATLAB R2019b (Mathworks, Natick, MA). Figures were created either using R version 3.6.2 (R Core Team) or MATLAB 2019b. Significance level in this manuscript was a priori decided to be $\alpha = 0.05$.

Additional information on our results and discussion

Cumulative incidence over time. From Figure S7 to Figure S10, we provided plots of cumulative incidence over time (left panel) and semi-log plots with \log_{10} -transformed cumulative incidence over time

(right panel) for a total of 8 provinces with a relatively high number of cases, namely, the epicenter Hubei, followed by (in alphabetical order) Fujian, Guangdong, Heilongjiang, Henan, Hubei, Hunan, Jiangxi and Shandong. If the epidemic is growing exponentially, the \log_{10} -transformed cumulative incidence over time will be a linear curve. If social distancing would have an impact, the slope of the semi-log plot would decrease, indicating a decreasing epidemic growth rate.

Harmonic mean of the harmonic mean. In this study, we also presented the harmonic mean of the harmonic means of the estimates of the epidemic doubling times. The harmonic means of the epidemic doubling times are shorter than their arithmetic means. From January 20 through February 9, the harmonic mean of the harmonic means of the doubling times estimated ranged from 0.5 (95% CI, 0.2, 1.3) days for Guangxi, to 2.3 (95% CI, 2.3, 2.4) days for Hubei. The harmonic mean of the harmonic means of doubling times in mainland China except Hubei were 1.2 (95% CI, 1.0, 1.4) days.

Further discussion. The slowing-down of the epidemic as represented in increasing epidemic doubling times in our study is also consistent with a study by Benjamin F. Maier and Dirk Brockmann, “Effective containment explains sub-exponential growth in confirmed cases of recent COVID-19 outbreak in Mainland China” (pre-print available at arXiv. 2020:2002.07572). They also identified sub-exponential growth of the outbreak across provinces, as mass quarantine and restriction of travels across mainland China began since January 23, 2020.

Sensitivity analysis #1. We performed a sensitivity analysis by expanding our data analysis to the data since December 31, 2019, when Hubei first reported a cluster of pneumonia cases with unexplained etiology that turned out to be COVID-19. The only difference between the sensitivity analysis and the main analysis is the inclusion of Hubei and Guangdong data from December 31, 2019, through January 19, 2020, because nationwide reporting started on January 20, 2020. The only differences in results were found for Hubei and Guangdong. For Hubei, the harmonic mean of the arithmetic mean of the doubling

times was 4.06 (95% CI, 3.85-4.33); the harmonic mean of the harmonic means of the doubling times for Hubei was 2.28 (95% CI, 2.08-2.56); and the cumulative incidence in Hubei doubled nine times from December 31, 2019, through February 9, 2020 (Table S5, Figures S3, S4, S12, S13, S14). The first doubling time of Hubei (Figure S3) was high, reflecting that real-time data was unavailable before mid-January. It was only by January 17, 2020, onwards when data reporting become increasingly transparent and timely.

Sensitivity analysis #2. We also performed a sensitivity analysis by restricting our data analysis to the data from January 23, 2020 through February 9, 2020, to allow for the time that all the other provinces to ramp up their testing. January 23 was also the day when the Chinese authorities to put the city of Wuhan on ‘lockdown’ and major inter-provincial travel restrictions were put in place. When we changed the start date of our study period from January 20 (main analysis) to January 23, 2020 (sensitivity analysis #2), the epidemic doubling time of the aggregate cumulative incidence of mainland China (except Hubei) increased from 1.79 (95% CI, 1.52, 2.25) to 2.90 (95% CI, 2.62, 3.24) (harmonic mean of the arithmetic means), and from 1.18 (95% CI, 0.96, 1.42) to 1.98 (95% CI, 1.82, 2.17) (harmonic mean of the harmonic means) (Table S7, Figure S5, S6). Apart from the epidemic doubling time of the aggregate cumulative incidence of mainland China (except Hubei), we did not observe significant differences by province between results in the main analysis and sensitivity analysis #2. Therefore, our results should be robust for the purpose of this study.

Table S1. Confirmed cases of COVID-19 (December 31, 2019 to January 19, 2020) by province in mainland China extracted from official government sources used for the sensitivity analysis.

| Locations ¹ | Dec | January | | | | | | | | | | | | | | | | | | | |
|--|-----|---------|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|-----|-----|----|
| | 31 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | |
| Mainland China (Excluding Hubei) (Sum of provincial reports) | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | 1 |
| Mainland China (Including Hubei) (Sum of provincial reports) | 27 | NR | NR | 44 | NR | 59 | NR | NR | NR | NR | 41 | 41 | 41 | 41 | 41 | 41 | 45 | 62 | 121 | 199 | |
| Mainland China (Including Hubei) (Sum by NCH) ² | NA | NA | NA | NA | NA | NA | NA | NA | NA | NA | NA | NA | NA | NA | NA | NA | NA | NA | NA | NA | NA |
| Hubei | 27 | NR | NR | 44 | NR | 59 | NR | NR | NR | NR | 41 | 41 | 41 | 41 | 41 | 41 | 45 | 62 | 121 | 198 | |
| Guangdong | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | 1 |

Note. NA, not applicable. NR, not reported. ¹Observations were collected directly from government official sites from each province in mainland China. If a press release included data reported at midnight and early morning, they were considered to belong to the previous day the data was reported. ²Official national tally of cumulative case count of confirmed cases was first published by the National Health Commission of China on January 21, 2020 for January 20, 2020 (3).

Table S2. Confirmed cases of COVID-19 (January 20 to 31, 2020) by province in mainland China extracted from official government sources used for the main analysis and sensitivity analysis.

| Locations ¹ | January | | | | | | | | | | | |
|--|---------|-----|-----|-----|------|------|------|------|------|------|------|-------|
| | 20 | 21 | 22 | 23 | 24 | 25 | 26 | 27 | 28 | 29 | 30 | 31 |
| Mainland China (Excluding Hubei) (Sum of provincial reports) | 26 | 71 | 145 | 291 | 585 | 923 | 1321 | 1802 | 2386 | 3126 | 3885 | 4637 |
| Mainland China (Including Hubei) (Sum of provincial reports) | 296 | 446 | 589 | 840 | 1314 | 1975 | 2744 | 4516 | 5940 | 7712 | 9691 | 11790 |
| Mainland China (Including Hubei) (Sum by NCH) ² | 291 | 440 | 571 | 830 | 1287 | 1975 | 2744 | 4515 | 5974 | 7711 | 9692 | 11791 |
| Hubei | 270 | 375 | 444 | 549 | 729 | 1052 | 1423 | 2714 | 3554 | 4586 | 5806 | 7153 |
| Anhui | 0 | 1 | 9 | 15 | 39 | 60 | 70 | 106 | 152 | 200 | 237 | 297 |
| Beijing | 5 | 10 | 14 | 26 | 36 | 49 | 68 | 80 | 91 | 111 | 132 | 156 |
| Chongqing | 0 | 5 | 9 | 27 | 57 | 75 | 110 | 132 | 147 | 165 | 206 | 238 |
| Fujian | 0 | 0 | 1 | 5 | 10 | 18 | 35 | 59 | 82 | 101 | 120 | 144 |
| Gansu | 0 | 0 | 0 | 2 | 4 | 7 | 14 | 19 | 24 | 26 | 29 | 35 |
| Guangdong | 14 | 26 | 32 | 53 | 78 | 98 | 146 | 188 | 241 | 311 | 393 | 520 |
| Guangxi | 0 | 0 | 2 | 13 | 23 | 33 | 46 | 51 | 58 | 78 | 87 | 100 |
| Guizhou | 0 | 0 | 0 | 3 | 5 | 5 | 7 | 9 | 9 | 12 | 15 | 29 |
| Hainan | 0 | 0 | 4 | 8 | 11 | 20 | 27 | 33 | 40 | 46 | 49 | 57 |
| Hebei | 0 | 0 | 1 | 2 | 8 | 13 | 18 | 33 | 48 | 65 | 82 | 96 |
| Heilongjiang | 0 | 0 | 1 | 4 | 9 | 15 | 21 | 30 | 37 | 43 | 59 | 80 |
| Henan | 0 | 1 | 5 | 9 | 32 | 83 | 128 | 168 | 206 | 278 | 352 | 422 |
| Hunan | 0 | 1 | 4 | 9 | 43 | 69 | 100 | 143 | 221 | 277 | 332 | 389 |
| Inner Mongolia | 0 | 0 | 0 | 1 | 2 | 7 | 11 | 13 | 16 | 18 | 20 | 23 |
| Jiangsu | 0 | 0 | 1 | 9 | 18 | 31 | 47 | 70 | 99 | 129 | 168 | 202 |
| Jiangxi | 0 | 2 | 3 | 7 | 18 | 36 | 48 | 72 | 109 | 162 | 240 | 286 |
| Jilin | 0 | 0 | 1 | 3 | 4 | 4 | 6 | 8 | 9 | 14 | 14 | 17 |
| Liaoning | 0 | 0 | 2 | 4 | 12 | 19 | 22 | 30 | 36 | 41 | 45 | 60 |
| Ningxia | 0 | 0 | 1 | 2 | 3 | 4 | 7 | 11 | 12 | 17 | 21 | 26 |
| Qinghai | 0 | 0 | 0 | 0 | 0 | 1 | 4 | 6 | 6 | 6 | 8 | 9 |
| Shaanxi | 0 | 0 | 0 | 0 | 7 | 15 | 22 | 46 | 56 | 63 | 87 | 101 |
| Shandong | 0 | 1 | 1 | 1 | 21 | 39 | 63 | 87 | 87 | 145 | 178 | 202 |
| Shanghai | 2 | 9 | 16 | 20 | 33 | 40 | 53 | 66 | 80 | 101 | 128 | 153 |
| Shanxi | 0 | 0 | 1 | 1 | 6 | 9 | 13 | 20 | 27 | 35 | 39 | 47 |
| Sichuan | 0 | 2 | 5 | 15 | 28 | 44 | 69 | 90 | 108 | 142 | 177 | 207 |
| Tianjin | 0 | 2 | 4 | 5 | 8 | 10 | 14 | 23 | 25 | 27 | 32 | 32 |
| Tibet | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 1 |
| Xinjiang | 0 | 0 | 0 | 2 | 3 | 4 | 5 | 10 | 13 | 14 | 17 | 18 |
| Yunnan | 0 | 1 | 1 | 2 | 5 | 11 | 19 | 26 | 51 | 70 | 80 | 91 |
| Zhejiang | 5 | 10 | 27 | 43 | 62 | 104 | 128 | 173 | 296 | 428 | 537 | 599 |

Note. ¹Observations were collected directly from government official sites from each province in mainland China. If a press release included data reported at midnight and early morning, they were considered to belong to the previous day the data was reported. ²Data was collected of press releases of the National Health Commission of China (3).

Table S3. Confirmed cases of COVID-19 (February 1 to 9, 2020) by province in mainland China extracted from official government sources used for the main analysis and sensitivity analysis.

| Locations ¹ | February | | | | | | | | |
|--|----------|-------|-------|-------|-------|-------|-------|-------|-------|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 |
| Mainland China (Excluding Hubei) (Sum of provincial reports) | 5396 | 6031 | 6910 | 7646 | 8352 | 9049 | 9614 | 10098 | 10507 |
| Mainland China (Including Hubei) (Sum of provincial reports) | 14381 | 17208 | 20432 | 24324 | 28017 | 31161 | 34567 | 37198 | 40138 |
| Mainland China (Including Hubei) (Sum by NCH) ² | 14380 | 17205 | 20438 | 24324 | 28018 | 31161 | 34546 | 37198 | 40171 |
| Hubei | 9074 | 11177 | 13522 | 16678 | 19665 | 22112 | 24953 | 27100 | 29631 |
| Anhui | 340 | 408 | 480 | 530 | 591 | 665 | 733 | 779 | 830 |
| Beijing | 183 | 212 | 228 | 253 | 274 | 297 | 315 | 326 | 337 |
| Chongqing | 262 | 300 | 337 | 366 | 389 | 411 | 426 | 446 | 468 |
| Fujian | 159 | 179 | 194 | 205 | 215 | 224 | 239 | 250 | 261 |
| Gansu | 40 | 51 | 55 | 57 | 62 | 67 | 71 | 79 | 83 |
| Guangdong | 604 | 683 | 797 | 870 | 944 | 1018 | 1075 | 1120 | 1131 |
| Guangxi | 111 | 127 | 139 | 150 | 168 | 172 | 183 | 195 | 210 |
| Guizhou | 38 | 46 | 56 | 64 | 69 | 77 | 89 | 96 | 99 |
| Hainan | 63 | 70 | 79 | 89 | 100 | 111 | 122 | 128 | 136 |
| Hebei | 104 | 113 | 126 | 135 | 157 | 171 | 195 | 206 | 218 |
| Heilongjiang | 95 | 118 | 155 | 190 | 227 | 277 | 295 | 307 | 331 |
| Henan | 493 | 566 | 675 | 764 | 851 | 914 | 981 | 1033 | 1073 |
| Hunan | 463 | 521 | 593 | 661 | 711 | 772 | 803 | 838 | 879 |
| Inner Mongolia | 27 | 34 | 35 | 42 | 46 | 50 | 52 | 54 | 58 |
| Jiangsu | 236 | 271 | 308 | 341 | 373 | 408 | 439 | 468 | 492 |
| Jiangxi | 333 | 391 | 476 | 548 | 600 | 661 | 698 | 740 | 771 |
| Jilin | 23 | 31 | 42 | 54 | 59 | 65 | 69 | 78 | 80 |
| Liaoning | 64 | 73 | 74 | 81 | 89 | 94 | 99 | 105 | 108 |
| Ningxia | 28 | 31 | 34 | 34 | 40 | 43 | 45 | 45 | 49 |
| Qinghai | 9 | 13 | 15 | 17 | 18 | 18 | 18 | 18 | 18 |
| Shaanxi | 116 | 128 | 142 | 165 | 173 | 184 | 195 | 208 | 213 |
| Shandong | 225 | 246 | 270 | 298 | 343 | 379 | 407 | 435 | 466 |
| Shanghai | 177 | 193 | 208 | 233 | 254 | 269 | 281 | 292 | 295 |
| Shanxi | 56 | 66 | 74 | 81 | 90 | 96 | 104 | 115 | 119 |
| Sichuan | 231 | 254 | 282 | 301 | 321 | 344 | 363 | 386 | 405 |
| Tianjin | 45 | 48 | 60 | 67 | 69 | 81 | 88 | 90 | 94 |
| Tibet | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| Xinjiang | 21 | 24 | 29 | 32 | 36 | 39 | 42 | 45 | 49 |
| Yunnan | 99 | 109 | 117 | 122 | 128 | 135 | 138 | 140 | 141 |
| Zhejiang | 661 | 724 | 829 | 895 | 954 | 1006 | 1048 | 1075 | 1092 |

Note. ¹Observations were collected directly from government official sites from each province in mainland China. If a press release included data reported at midnight and early morning, they were considered to belong to the previous day the data was reported. ²Data was collected of press releases of the National Health Commission of China (3).

Table S4. Main analysis: Doubling times of COVID-19 cumulative incidence and their harmonic mean of the arithmetic means of the doubling times and harmonic mean of the harmonic means of the doubling times (95% Confidence interval) by province in mainland China from January 20 through February 9, 2020.

| | | Mainland China (Except Hubei) | Hubei | Anhui | Beijing | Chongqing | Fujian | Gansu | Guangdong | Guangxi | Guizhou | Hainan |
|---------------------------------------|----|--|---------------------|---------------------|---------------------|-----------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|
| Harmonic mean of arithmetic means | | 1.79 (1.52-2.25) | 2.54 (2.44-2.64) | 2.56 (2.16-3.11) | 2.49 (1.89-3.38) | 2.22 (1.53-3.22) | 1.71 (1.15-2.52) | 2.56 (2.00-3.78) | 2.47 (1.97-3.20) | 1.92 (1.45-3.09) | 2.71 (1.90-3.90) | 2.91 (1.91-3.89) |
| Harmonic mean of harmonic means | | 1.18 (0.96-1.42) | 2.34 (2.27-2.41) | 1.72 (1.13-2.67) | 1.48 (0.63-2.70) | 1.23 (0.67-1.96) | 0.82 (0.46-1.41) | 1.36 (0.76-2.86) | 2.01 (1.53-2.54) | 0.48 (0.22-1.34) | 1.88 (0.81-3.28) | 1.52 (0.65-2.99) |
| Times doubled | 1 | 0.59 | 2.91 | 2.12 | 1.00 | 2.05 | 0.25 | 1 | 1.33 | 0.18 | 2.5 | 1.00 |
| | 2 | 0.86 | 2.16 | 0.75 | 1.5 | 0.56 | 0.5 | 1.14 | 1.79 | 0.36 | 3.5 | 1.55 |
| | 3 | 0.98 | 1.5 | 2.18 | 1.8 | 0.82 | 0.85 | 1.26 | 2.17 | 0.76 | 1.64 | 2.28 |
| | 4 | 1 | 2.17 | 1.77 | 2.7 | 1.71 | 1.15 | 4.1 | 2.38 | 1.6 | 2.56 | 5.31 |
| | 5 | 1.3 | 3.03 | 4.03 | 4.14 | 3.58 | 1.07 | 5.9 | 2.76 | 3.4 | 5.8 | 6.86 |
| | 6 | 1.98 | 3.43 | 4.9 | 7.31 | 4.82 | 1.39 | | 4.92 | 4.78 | | |
| | 7 | 2.55 | | | | | 3.12 | | | | | |
| | 8 | 4.53 | | | | | 9.21 | | | | | |
| | | Hebei | Heilongjiang | Henan | Hunan | Inner Mongolia | Jiangsu | Jiangxi | Jilin | Liaoning | Ningxia | Qinghai |
| Harmonic mean of arithmetic means | | 1.88 (1.57-2.72) | 1.93 (1.76-2.36) | 1.81 (1.35-2.05) | 1.42 (1.24-2.04) | 2.37 (1.80-3.67) | 2.43 (1.77-3.26) | 1.68 (1.45-2.33) | 2.64 (2.13-3.50) | 2.10 (1.45-3.30) | 2.54 (1.76-4.33) | 2.50 (1.50-5.00) |
| Harmonic mean doubling time | | 1.04 (0.67-1.93) | 1.08 (0.62-1.98) | 0.81 (0.56-1.12) | 0.71 (0.47-1.13) | 1.17 (0.67-2.67) | 1.93 (1.35-2.63) | 1.13 (0.71-1.71) | 1.48 (0.66-3.03) | 1.05 (0.54-1.94) | 1.59 (0.73-3.07) | 1.00 (0.38-3.87) |
| Times doubled | 1 | 1 | 0.33 | 0.25 | 0.33 | 1 | 2 | 1.25 | 0.5 | 1 | 1 | 0.33 |
| | 2 | 0.33 | 0.67 | 0.5 | 0.67 | 0.4 | 1.31 | 0.84 | 2.5 | 0.5 | 2 | 0.67 |
| | 3 | 0.67 | 0.8 | 1 | 0.8 | 0.85 | 1.75 | 0.72 | 2 | 1.07 | 1.25 | 4 |
| | 4 | 1.6 | 1.36 | 0.55 | 0.4 | 2.75 | 2.32 | 0.96 | 3.66 | 2.76 | 2.55 | 4.5 |
| | 5 | 1.33 | 2.12 | 0.7 | 0.47 | 4.71 | 4.07 | 1.89 | 2.43 | 4.67 | 4.53 | |
| | 6 | 2.01 | 2.95 | 0.62 | 1.13 | | | 1.69 | 3.74 | | | |
| | 7 | 5.28 | 3.04 | 1.38 | 1.85 | | | 1.99 | | | | |
| | 8 | | 3.31 | 2.69 | 1.97 | | | 4.16 | | | | |
| | 9 | | | 3.57 | 4.22 | | | | | | | |
| | 10 | | | 6.56 | | | | | | | | |
| | | Shaanxi | Shandong | Shanghai | Shanxi | Sichuan | Tianjin | Tibet | Xinjiang | Yunnan | Zhejiang | |
| Harmonic mean of the arithmetic means | | 2.82 (2.12-9.97) | 1.68 (1.42-2.39) | 2.19 (1.88-2.68) | 2.31 (1.67-3.25) | 1.83 (1.39-2.70) | 2.78 (2.07-4.06) | Not applied | 3.05 (2.06-4.75) | 2.05 (1.34-2.72) | 1.91 (1.60-2.51) | |
| Harmonic mean doubling time | | 2.04 (1.28-3.01) | 0.48 (0.28-1.15) | 0.77 (0.34-1.73) | 1.22 (0.68-2.51) | 0.96 (0.51-1.75) | 1.69 (0.80-3.55) | Not applied | 1.91 (0.83-4.46) | 1.25 (0.89-1.81) | 1.20 (0.74-1.70) | |
| Times doubled | 1 | 1.33 | 2.05 | 0.28 | 1.2 | 0.66 | 1 | | 2 | 2 | 1 | |
| | 2 | 2.24 | 0.1 | 0.57 | 0.4 | 0.64 | 2 | | 1.6 | 0.66 | 0.58 | |
| | 3 | 3.76 | 0.19 | 1.15 | 1.06 | 0.77 | 2.22 | | 3.06 | 0.84 | 1.23 | |
| | 4 | | 0.41 | 1.92 | 1.76 | 1.18 | 4.78 | | 5.34 | 1.12 | 1.61 | |
| | 5 | | 0.86 | 2.92 | 2.2 | 1.55 | 3.57 | | | 1.61 | 2.29 | |
| | 6 | | 1.43 | 3.16 | 4.18 | 2.78 | | | | 1.45 | 1.47 | |
| | 7 | | 2.66 | 6.13 | | 4.49 | | | | 7.32 | 3.48 | |
| | 8 | | 4.71 | | | | | | | | | |

Table S5. Sensitivity analysis #1 (1 of 2 tables): Doubling times of COVID-19 cumulative incidence and their harmonic mean of the arithmetic means of the doubling times and harmonic mean of the harmonic means of the doubling times (95% Confidence interval) by province in mainland China from December 31, 2019 through February 9, 2020: Mainland China (Except Hubei), Hubei, and from Anhui to Qinghai.

| | | Mainland China (Except Hubei) | Hubei | Anhui | Beijing | Chongqing | Fujian | Gansu | Guangdong | Guangxi | Guizhou | Hainan |
|-----------------------------------|----|----------------------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|
| Harmonic mean of arithmetic means | | 1.34 (1.28-1.52) | 4.06 (3.85-4.33) | 2.57 (2.12-3.00) | 2.51 (1.99-3.26) | 2.22 (1.60-3.23) | 1.82 (1.18-2.55) | 2.55 (1.83-3.79) | 1.88 (1.74-2.19) | 1.93 (1.47-2.96) | 2.78 (2.00-3.97) | 2.92 (1.97-4.25) |
| Harmonic mean of harmonic means | | 0.29 (0.15-0.59) | 2.28 (2.08-2.56) | 1.76 (1.21-2.40) | 1.60 (0.93-2.70) | 1.23 (0.74-1.88) | 0.83 (0.47-1.42) | 1.33 (0.70-2.62) | 0.44 (0.25-1.13) | 0.49 (0.22-1.29) | 1.98 (1.09-3.53) | 1.55 (0.60-3.29) |
| Times doubled | 1 | 0.04 | 17.33 | 2.12 | 1.00 | 2.05 | 0.25 | 1.00 | 0.07 | 0.18 | 2.50 | 1.00 |
| | 2 | 0.08 | 1.22 | 0.75 | 1.5 | 0.56 | 0.5 | 1.14 | 0.16 | 0.36 | 3.5 | 1.55 |
| | 3 | 0.15 | 2 | 2.18 | 1.8 | 0.82 | 0.85 | 1.26 | 0.3 | 0.76 | 1.64 | 2.28 |
| | 4 | 0.33 | 3.04 | 1.77 | 2.7 | 1.71 | 1.15 | 4.1 | 0.63 | 1.6 | 2.56 | 5.31 |
| | 5 | 0.53 | 2.11 | 4.03 | 4.14 | 3.58 | 1.07 | 5.9 | 1.84 | 3.4 | 5.8 | 6.86 |
| | 6 | 0.73 | 1.23 | 4.9 | 7.31 | 4.82 | 1.39 | | 1.44 | 4.78 | | |
| | 7 | 0.92 | 2.61 | | | | | 3.12 | 2.18 | | | |
| | 8 | 0.98 | 3.13 | | | | | 9.21 | 2.59 | | | |
| | 9 | 1.01 | 3.87 | | | | | | 2.72 | | | |
| | 10 | 1.48 | | | | | | | 6.17 | | | |
| | 11 | 2.17 | | | | | | | | | | |
| | 12 | 2.88 | | | | | | | | | | |
| | 13 | 5.55 | | | | | | | | | | |
| | | Hebei | Heilongjiang | Henan | Hunan | Inner Mongolia | Jiangsu | Jiangxi | Jilin | Liaoning | Ningxia | Qinghai |
| Harmonic mean of arithmetic means | | 1.89 (1.55-2.74) | 1.96 (1.76-2.26) | 1.80 (1.31-2.10) | 1.41 (1.26-1.99) | 2.37 (1.82-3.57) | 2.45 (1.75-3.31) | 1.72 (1.44-2.36) | 2.67 (2.13-3.50) | 2.16 (1.49-3.53) | 2.58 (1.72-4.43) | 2.64 (1.79-5.00) |
| Harmonic mean of harmonic means | | 1.07 (0.66-1.90) | 1.12 (0.66-1.97) | 0.77 (0.48-1.14) | 0.73 (0.48-1.15) | 1.15 (0.65-2.71) | 1.92 (1.31-2.68) | 1.17 (0.81-1.74) | 1.60 (0.70-3.11) | 1.06 (0.50-2.45) | 1.67 (0.87-3.65) | 0.96 (0.39-3.69) |
| Times doubled | 1 | 1.00 | 0.33 | 0.25 | 0.33 | 1.00 | 2.00 | 1.25 | 0.50 | 1.00 | 1.00 | 0.33 |
| | 2 | 0.33 | 0.67 | 0.5 | 0.67 | 0.4 | 1.31 | 0.84 | 2.5 | 0.5 | 2 | 0.67 |
| | 3 | 0.67 | 0.8 | 1 | 0.8 | 0.85 | 1.75 | 0.72 | 2 | 1.07 | 1.25 | 4 |
| | 4 | 1.6 | 1.36 | 0.55 | 0.4 | 2.75 | 2.32 | 0.96 | 3.66 | 2.76 | 2.55 | 4.5 |
| | 5 | 1.33 | 2.12 | 0.7 | 0.47 | 4.71 | 4.07 | 1.89 | 2.43 | 4.67 | 4.53 | |
| | 6 | 2.01 | 2.95 | 0.62 | 1.13 | | | 1.69 | 3.74 | | | |
| | 7 | 5.28 | 3.04 | 1.38 | 1.85 | | | 1.99 | | | | |
| | 8 | | 3.31 | 2.69 | 1.97 | | | 4.16 | | | | |
| | 9 | | | 3.57 | 4.22 | | | | | | | |
| | 10 | | | 6.56 | | | | | | | | |

Table S6. Sensitivity analysis #1 (2 of 2 tables): Doubling times of COVID-19 cumulative incidence and their harmonic mean of the arithmetic means of the doubling times and harmonic mean of the harmonic means of the doubling times (95% Confidence interval) by province in mainland China from December 31, 2019 through February 9, 2020: from Shaanxi to Zhejiang.

| | | Shaanxi | Shandong | Shanghai | Shanxi | Sichuan | Tianjin | Tibet | Xinjiang | Yunnan | Zhejiang |
|---------------|-----------------------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|--------------|---------------------|---------------------|---------------------|
| | Harmonic mean of arithmetic means | 2.77 (2.06-3.93) | 1.68 (1.41-2.36) | 2.21 (1.91-2.78) | 2.12 (1.67-3.00) | 1.79 (1.40-2.65) | 2.75 (2.10-3.89) | Not applied | 3.09 (2.12-4.89) | 2.10 (1.42-2.78) | 1.90 (1.59-2.55) |
| | Harmonic mean of harmonic means | 2.03 (1.27-2.93) | 0.48 (0.30-1.11) | 0.82 (0.40-1.83) | 1.26 (0.68-2.60) | 0.96 (0.62-1.73) | 1.67 (0.78-3.38) | Not applied | 1.98 (0.80-4.69) | 1.28 (0.80-1.93) | 1.23 (0.77-1.72) |
| Times doubled | 1 | 1.33 | 2.05 | 0.28 | 1.20 | 0.66 | 1.00 | | 2.00 | 2 | 1.00 |
| | 2 | 2.24 | 0.1 | 0.57 | 0.4 | 0.64 | 2 | | 1.6 | 0.66 | 0.58 |
| | 3 | 3.76 | 0.19 | 1.15 | 1.06 | 0.77 | 2.22 | | 3.06 | 0.84 | 1.23 |
| | 4 | | 0.41 | 1.92 | 1.76 | 1.18 | 4.78 | | 5.34 | 1.12 | 1.61 |
| | 5 | | 0.86 | 2.92 | 2.2 | 1.55 | 3.57 | | | 1.61 | 2.29 |
| | 6 | | 1.43 | 3.16 | 4.18 | 2.78 | | | | 1.45 | 1.47 |
| | 7 | | 2.66 | 6.13 | | 4.49 | | | | 7.32 | 3.48 |
| | 8 | | 4.71 | | | | | | | | |

Table S7. Sensitivity analysis #2: Doubling times of COVID-19 cumulative incidence and their harmonic mean of the arithmetic means of the doubling times and harmonic mean of the harmonic means of the doubling times (95% Confidence interval) by province in mainland China from January 23, 2020 through February 9, 2020.

| | | Mainland China (Except Hubei) | Hubei | Anhui | Beijing | Chongqing | Fujian | Gansu | Guangdong | Guangxi | Guizhou | Hainan |
|-----------------------------------|---|----------------------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|
| Harmonic mean of arithmetic means | | 2.9 (2.62-3.24) | 2.46 (2.37-2.55) | 2.54 (2.12-2.99) | 3.46 (2.77-4.57) | 3.11 (2.38-4.17) | 2.03 (1.29-3.10) | 2.54 (1.80-3.89) | 2.91 (2.40-3.61) | 3.26 (2.37-4.22) | 2.67 (1.85-3.92) | 3.43 (2.57-4.62) |
| Harmonic mean of harmonic means | | 1.98 (1.82-2.17) | 2.25 (2.18-2.33) | 1.47 (0.90-2.29) | 3.03 (2.23-3.99) | 1.87 (1.28-2.80) | 1.26 (0.69-2.01) | 1.27 (0.66-2.84) | 2.65 (2.14-3.10) | 2.23 (1.33-3.29) | 1.73 (0.67-3.40) | 2.31 (1.40-3.71) |
| Times doubled | 1 | 1.01 | 2.12 | 0.62 | 2.15 | 0.90 | 1.00 | 1.00 | 2.16 | 1.30 | 2.50 | 1.55 |
| | 2 | 1.59 | 1.47 | 1.38 | 3.50 | 2.04 | 1.11 | 1.14 | 2.29 | 2.84 | 3.50 | 2.28 |
| | 3 | 2.30 | 2.22 | 2.30 | 4.21 | 4.37 | 1.09 | 1.26 | 2.79 | 4.22 | 1.64 | 5.31 |
| | 4 | 3.21 | 3.03 | 3.74 | | 7.99 | 1.71 | 4.10 | 4.45 | 8.50 | 2.56 | 6.86 |
| | 5 | 6.41 | 3.45 | 3.96 | | | 4.13 | 5.90 | | | 5.80 | |
| | | Hebei | Heilongjiang | Henan | Hunan | Inner Mongolia | Jiangsu | Jiangxi | Jilin | Liaoning | Ningxia | Qinghai |
| Harmonic mean of arithmetic means | | 1.91 (1.42-2.83) | 2.21 (1.74-2.81) | 1.87 (1.50-2.40) | 1.89 (1.48-2.77) | 2.39 (1.84-4.00) | 2.31 (1.80-3.10) | 1.89 (1.44-2.52) | 3.01 (2.14-4.06) | 2.44 (1.49-4.00) | 2.68 (1.70-4.50) | 3.21 (2.25-5.67) |
| Harmonic mean of harmonic means | | 0.81 (0.39-1.78) | 1.47 (0.77-2.51) | 0.85 (0.48-1.37) | 0.75 (0.41-1.27) | 1.16 (0.66-3.02) | 1.60 (1.06-2.36) | 1.18 (0.62-1.86) | 2.44 (1.29-3.73) | 0.99 (0.37-2.46) | 1.88 (0.94-3.96) | 1.8 (0.80-5.08) |
| Times doubled | 1 | 0.33 | 0.80 | 0.39 | 0.26 | 1.00 | 1.00 | 0.63 | 3.00 | 0.50 | 2.00 | 2.33 |
| | 2 | 0.67 | 1.36 | 0.68 | 0.53 | 0.40 | 1.31 | 0.92 | 2.59 | 1.07 | 1.25 | 0.67 |
| | 3 | 1.60 | 2.12 | 0.71 | 1.30 | 0.85 | 1.75 | 1.78 | 3.53 | 2.76 | 2.55 | 4.00 |
| | 4 | 1.33 | 2.95 | 1.62 | 1.92 | 2.75 | 2.32 | 1.72 | 2.38 | 4.67 | 4.53 | 4.50 |
| | 5 | 2.01 | 3.04 | 2.73 | 2.19 | 4.71 | 4.07 | 1.74 | | | | |
| | 6 | 5.28 | 3.31 | 3.96 | 4.56 | | | 3.88 | | | | |
| | | Shaanxi | Shandong | Shanghai | Shanxi | Sichuan | Tianjin | Tibet | Xinjiang | Yunnan | Zhejiang | |
| Harmonic mean of arithmetic means | | 3.44 (2.76-4.40) | 1.43 (1.18-2.14) | 3.08 (2.52-4.07) | 1.93 (1.50-3.09) | 2.61 (1.89-3.60) | 3.17 (2.16-4.59) | Not applied | 3.05 (2.10-4.67) | 1.82 (1.20-2.87) | 2.37 (1.87-3.14) | |
| Harmonic mean of harmonic means | | 2.84 (1.82-4.05) | 0.24 (0.14-0.60) | 2.61 (1.72-3.59) | 0.71 (0.32-1.88) | 1.82 (1.28-2.57) | 2.12 (0.79-4.34) | Not applied | 1.86 (0.83-4.40) | 1.03 (0.56-1.77) | 1.98 (1.73-2.41) | |
| Times doubled | 1 | 3.33 | 0.05 | 2.00 | 0.20 | 1.12 | 2.00 | | 2.00 | 0.66 | 1.57 | |
| | 2 | 2.24 | 0.10 | 3.00 | 0.40 | 1.51 | 1.66 | | 1.60 | 0.84 | 2.4 | |
| | 3 | 3.76 | 0.20 | 3.29 | 1.06 | 2.72 | 4.95 | | 3.06 | 1.12 | 1.39 | |
| | 4 | | 0.40 | | 1.76 | 4.04 | 5.30 | | 5.34 | 1.61 | 4.06 | |
| | 5 | | 0.86 | | 2.20 | | | | | 1.45 | | |
| | 6 | | 1.43 | | 4.18 | | | | | 7.32 | | |
| | 7 | | 2.66 | | | | | | | | | |
| | 8 | | 4.71 | | | | | | | | | |

Table S8. Websites of national and provincial health commissions in mainland China

| Health commission | URL | Notes |
|---|---|---|
| National Health Commission of the People's Republic of China | http://www.nhc.gov.cn | |
| Provincial health commissions | | |
| Anhui | http://wjw.ah.gov.cn/ | |
| Beijing | http://wjw.beijing.gov.cn/ | |
| Chongqing | http://wsjkw.cq.gov.cn/ | |
| Fujian | http://fjwsjk.fjsen.com/ | |
| Gansu | http://wsjk.gansu.gov.cn/ | |
| Guangdong | http://wsjkw.gd.gov.cn/ | |
| Guangxi | http://wsjkw.gxzf.gov.cn/ | |
| Guizhou | http://www.gzhfpc.gov.cn/ | |
| Hainan | http://wst.hainan.gov.cn/ | |
| Hebei | http://www.hebwtst.gov.cn/ | Our team members found it often inaccessible from Statesboro, GA, USA. |
| Heilongjiang | http://wsjkw.hl.gov.cn/ | |
| Henan | http://www.hnwsjsw.gov.cn/ | |
| Hubei | http://wjw.hubei.gov.cn/ | |
| Hunan | http://wjw.hunan.gov.cn/ | |
| Inner Mongolia | http://wjw.nmg.gov.cn/ | |
| Jiangsu | http://wjw.jiangsu.gov.cn/ | |
| Jiangxi | http://hc.jiangxi.gov.cn/ | |
| Jilin | http://www.jl.gov.cn/ | |
| Liaoning | http://www.shenyang.gov.cn/ | |
| Ningxia | http://wsjkw.nx.gov.cn/index.htm | |
| Qinghai | https://wsjkw.qinghai.gov.cn/ | |
| Shaanxi | http://sxwjw.shaanxi.gov.cn/ | |
| Shandong | http://wsjkw.shandong.gov.cn | Our team members found it persistently inaccessible from Statesboro, GA, USA. |
| Shanghai | http://www.shanghai.gov.cn/ | |
| Shanxi | http://wjw.shanxi.gov.cn/ | |
| Sichuan | http://wsjkw.sc.gov.cn/ | |
| Tianjin | http://www.tj.gov.cn/ | |
| Tibet | http://wjw.xizang.gov.cn/ | |
| Xinjiang | http://www.xjhfp.gov.cn/ | |
| Yunnan | http://ynswsjkw.yn.gov.cn/ | |
| Zhejiang | https://www.zjwjw.gov.cn | |

Notes: If our team was unable to directly retrieve the press release from a provincial health commissions, we will use mainland Chinese media reports that directly reported on the provincial health commissions' announcements. Please note that mainland Chinese media are controlled by the Chinese Communist Party and they could not deviate from the government's announcements.

Figure S1. Main analysis: The harmonic mean of the arithmetic means of COVID-19 epidemic doubling times (red circle) with 95% confidence interval (red bar) of the doubling times (days), and their values (black diamond) by the number of times the reported cumulative incidence doubles by province within mainland China from January 20, 2020 through February 9, 2020. Each panel represents a province except the panel representing “Mainland China (except Hubei)” that is the aggregate of all other provinces in mainland China, except Hubei. Doubling time for Tibet is not available, because there had only been 1 confirmed case in Tibet as of February 9, 2020.

Figure S2. Main analysis: The harmonic mean of the harmonic means of COVID-19 epidemic doubling times (red circle) with 95% confidence interval (red bar) of the doubling times (days), and their values (black diamond) by the number of times the reported cumulative incidence doubles by province within mainland China from January 20, 2020 through February 9, 2020. Each panel represents a province except the panel representing “Mainland China (except Hubei)” that is the aggregate of all other provinces in mainland China, except Hubei. Doubling time for Tibet is not available, because there had only been 1 confirmed case in Tibet as of February 9, 2020.

Figure S3. Sensitivity analysis #1: The harmonic mean of the arithmetic means of COVID-19 doubling times (red circle) with 95% confidence interval (red bar) of the doubling times (days), and their values (black diamond) by the number of times the reported cumulative incidence doubles by province within mainland China from December 31, 2019 through February 9, 2020. Each panel represents a province except the panel representing “Mainland China (except Hubei)” that is the aggregate of all other provinces in mainland China, except Hubei. Doubling time for Tibet is not available, because there had only been 1 confirmed case in Tibet as of February 9, 2020.

Figure S4. Sensitivity analysis #1: The harmonic mean of the harmonic means of COVID-19 doubling times (red circle) with 95% confidence interval (red bar) of the doubling times (days), and their values (black diamond) by the number of times the reported cumulative incidence doubles by province within mainland China from December 31, 2019 through February 9, 2020. Each panel represents a province except the panel representing “Mainland China (except Hubei)” that is the aggregate of all other provinces in mainland China, except Hubei. Doubling time for Tibet is not available, because there had only been 1 confirmed case in Tibet as of February 9, 2020.

Figure S5. Sensitivity analysis #2: The harmonic mean of the arithmetic means of COVID-19 doubling times (red circle) with 95% confidence interval (red bar) of the doubling times (days), and their values (black diamond) by the number of times the reported cumulative incidence doubles by province within mainland China from January 23, 2020 through February 9, 2020. Each panel represents a province except the panel representing “Mainland China (except Hubei)” that is the aggregate of all other provinces in mainland China, except Hubei. Doubling time for Tibet is not available, because there had only been 1 confirmed case in Tibet as of February 9, 2020.

Figure S6. Sensitivity analysis #2: The harmonic mean of the harmonic means of COVID-19 doubling times (red circle) with 95% confidence interval (red bar) of the doubling times (days), and their values (black diamond) by the number of times the reported cumulative incidence doubles by province within mainland China from January 23, 2020 through February 9, 2020. Each panel represents a province except the panel representing “Mainland China (except Hubei)” that is the aggregate of all other provinces in mainland China, except Hubei. Doubling time for Tibet is not available, because there had only been 1 confirmed case in Tibet as of February 9, 2020.

Figure S7. Cumulative incidence and \log_{10} cumulative incidence over time (date) for Hubei (upper panel) and Fujian (lower panel).

Figure S8. Cumulative incidence and \log_{10} cumulative incidence over time (date) for Guangdong (upper panel) and Heilongjiang (lower panel).

Figure S9. Cumulative incidence and \log_{10} cumulative incidence over time (date) for Henan (upper panel) and Hunan (lower panel).

Figure S10. Cumulative incidence and \log_{10} cumulative incidence over time (date) for Jiangxi (upper panel) and Shandong (lower panel).

Figure S11. Main analysis: The map of the harmonic mean of the harmonic means of COVID-19 by province in mainland China, from January 20, 2020 through February 9, 2020.

Figure S12. Sensitivity analysis #1: The map of the harmonic mean of the arithmetic means of COVID-19 by province in mainland China, from December 31, 2019 through February 9, 2020.

Figure S13. Sensitivity Analysis #1: The map of the harmonic mean of the harmonic means of COVID-19 by province in mainland China, from December 31, 2019 through February 9, 2020.

Figure S14. Sensitivity analysis #1: The map of the number of times the COVID-19 outbreak has doubled by province in mainland China, from December 31, 2019 through February 9, 2020.

Authors' contributions

Project management: Dr. Gerardo Chowell, Dr. Isaac Chun-Hai Fung and Ms. Kamalich Muniz-Rodriguez

Manuscript writing: Dr. Isaac Chun-Hai Fung and Dr. Gerardo Chowell

Manuscript editing and data interpretation: Ms. Kamalich Muniz-Rodriguez, Dr. Gerardo Chowell, Dr. Isaac Chun-Hai Fung, Dr. Lone Simonsen, and Dr. Cecile Viboud

MATLAB code and methods of doubling time estimation: Dr. Gerardo Chowell

Doubling time calculation using MATLAB and presentation of results: Ms. Kamalich Muniz-Rodriguez, Dr. Gerardo Chowell and Dr. Isaac Chun-Hai Fung

Statistical analysis in R: Dr. Isaac Chun-Hai Fung, Ms. Kamalich Muniz-Rodriguez

Data management and quality check of epidemic data entry: Ms. Kamalich Muniz-Rodriguez, Dr. Isaac Chun-Hai Fung

Curation of epidemic data for countries and territories outside mainland China (including Hong Kong, Macao and Taiwan): Ms. Kamalich Muniz-Rodriguez and Ms. Sylvia K. Ofori

Curation of epidemic data for provinces in mainland China: Ms. Manyun Liu (from the early reports, up to Jan 24, 2020 data), Ms. Po-Ying Lai (since Jan 25, 2020 data to today), Mr. Chi-Hin Cheung (since Jan 27, 2020 data to today), and Ms. Kamalich Muniz-Rodriguez and Dr. Isaac Chun-Hai Fung (whenever there is a back-log).

Retrieval of epidemic data from official websites (downloading and archiving of China's national and provincial authorities' press releases): Ms. Manyun Liu and Dr. Dongyu Jia (at the very beginning of our project)

Retrieval of statistical data from the official website of National Bureau of Statistics of the People's Republic of China: Mr. Chi-Hin Cheung

Retrieval of publicly available statistical data from various sources: Ms. Yiseul Lee, Dr. Isaac Chun-Hai Fung


Table S8: Ms. Manyun Liu, Dr. Isaac Chun-Hai Fung


Map creation: Ms. Kimberlyn M. Roosa


Assistance provided to Dr. Fung and Ms. Muniz-Rodriguez: Ms. Sylvia K. Ofori


References


1. Anderson RM, May RM. Infectious diseases of humans. Oxford: Oxford University Press; 1991.
2. Drake JM, Bakach I, Just MR, O'Regan SM, Gambhir M, Fung IC-H. Transmission Models of Historical Ebola Outbreaks. *Emerging Infectious Disease journal*. 2015 Aug;21(8):1447-50.
3. National Health Commission of the People's Republic of China. 2020 [cited Feb 2, 2020]; Available from: <http://www.nhc.gov.cn/>
4. Centre for Health Protection, Department of Health, The Government for the Hong Kong Special Administrative Region. 2020 [cited Feb 2, 2020]; Available from: <https://www.chp.gov.hk/en/index.html>
5. John Hopkins University Center for Systems Science and Engineering. 2019 Novel Coronavirus COVID-19 (2019-nCoV) Data Repository by Johns Hopkins CSSE. 2020 [cited Feb 13, 2020]; Available from: <https://github.com/CSSEGISandData/COVID-19>
6. Banks HT, Hu S, Thompson WC. Modeling and inverse problems in the presence of uncertainty: CRC Press; 2014.
7. Chowell G, Ammon CE, Hengartner NW, Hyman JM. Transmission dynamics of the great influenza pandemic of 1918 in Geneva, Switzerland: Assessing the effects of hypothetical interventions. *J Theor Biol*. 2006 Jul 21;241(2):193-204.
8. Chowell G, Shim E, Brauer F, Diaz-Duenas P, Hyman JM, Castillo-Chavez C. Modelling the transmission dynamics of acute haemorrhagic conjunctivitis: application to the 2003 outbreak in Mexico. *Stat Med*. 2006 Jun 15;25(11):1840-57.
9. Vynnycky E, White RG. An Introduction to Infectious Disease Modelling. Oxford: Oxford University Press; 2010.

















Hubei Cumulative Incidence




Hubei Semi-log Cumulative Incidence




Fujian Cumulative Incidence




Fujian Semi-log Cumulative Incidence




Guangdong Cumulative Incidence




Guangdong Semi-log Cumulative Incidence




Heilongjiang Cumulative Incidence




Heilongjiang Semi-log Cumulative Incidence




Henan Cumulative Incidence




Henan Semi-log Cumulative Incidence




Hunan Cumulative Incidence




Hunan Semi-log Cumulative Incidence




Jiangxi Cumulative Incidence




Jiangxi Semi-log Cumulative Incidence





Shandong Cumulative Incidence




Shandong Semi-log Cumulative Incidence









Harmonic mean of the harmonic means of COVID-19 epidemic doubling times

- 0.4 - 0.69
- 0.7 - 0.99
- 1.0 - 1.19
- 1.2 - 1.49
- 1.5 - 1.89
- 1.9 - 2.29

