1	Estimation of Rift Valley fever virus spillover to humans
2	during the Mayotte 2018-2019 epidemic
3	
4	Raphaëlle Métras ^{a,b,c} PhD, W John Edmunds ^d PhD, Chouanibou Youssouffi ^e BSc, Laure
5	Dommergues ^f DVM, Guillaume Fournié ^g PhD, Anton Camacho ^{d,h} PhD, Sebastian Funk ^d PhD,
6	Eric Cardinale ^{a,i} PhD, Gilles Le Godais ^j MSc, Soihibou Combo ^j BSc, Laurent Filleul ^k PhD,
7	Hassani Youssouf ^{k*} PhD, Marion Subiros ^{k*} MSc.
8	*these authors share last authorship
9	
10	^a CIRAD, UMR ASTRE, Campus International de Baillarguet, 34398 Montpellier, France
11	^b ASTRE, Univ Montpellier (I-MUSE), CIRAD, INRA, 34398 Montpellier, France
12	^c Inserm, Sorbonne Université, Institut Pierre Louis d'Épidémiologie et de Santé Publique (iPLesp), 27
13	rue de Chaligny, 75012 Paris, France.
14	^d Centre for the Mathematical Modelling of Infectious Diseases, Department of Infectious Disease
15	Epidemiology, London School of Hygiene & Tropical Medicine, Keppel Street WC1E 7HT, London,
16	United Kingdom
17	°GSD Mayotte-Coopérative Agricole des Eleveurs Mahorais, 97670 Coconi, Mayotte, France
18	^f La Coopération Agricole, 43 rue Sedaine, F-75538 Paris, France
19	^g Veterinary Epidemiology, Economics and Public Health group, Department of Pathobiology and
20	Population Sciences, The Royal Veterinary College, Hatfield, United Kingdom
21	^h Epicentre, 14-34 avenue Jean Jaurès, 75019 Paris, France
22	ⁱ CIRAD, UMR ASTRE, F-97490 Sainte Clotilde, La Réunion, France
23	^j Direction de l'Alimentation, de l'Agriculture et de la Forêt de Mayotte, Mamoudzou, France
24	^k Santé Publique France, Mamoudzou, Mayotte, France

25 Abstract

Background. Rift Valley fever (RVF) is an emerging, zoonotic, arboviral haemorrhagic fever 26 threatening livestock and humans mainly in Africa. In the absence of a human vaccine, 27 estimating the transmission potential of RVF virus from livestock to humans is key to 28 assessing the impact of livestock disease control measures on preventing human disease. 29 Methods. We combined a unique RVF dataset, with livestock and human surveillance data of 30 the 2018-2019 RVF epidemic in Mayotte, and used them in a mathematical model. Using 31 Bayesian inference, we quantified the transmission amongst livestock and spillover to 32 humans, and we assessed the impact of livestock vaccination on reducing human disease risk. 33 Findings. Vaccination scenarios indicate that early livestock vaccination (December 2018) 34

would have reduced the human epidemic size by a third, whilst vaccinating one month later required using 50 % more vaccine doses for a similar impact. In addition, the likelihood of virus re-emergence in the next rainy season (2019-2020) was estimated very low, with 55.8 % (90 Credible Interval [27.1-59.5]) of the livestock population being immune in August 2019.

Interpretation. Human and livestock health surveillance, early detection, and timely vaccination in livestock are crucial to reducing disease risk in humans. We present the first study quantifying RVF virus spillover using livestock and human data, and use this quantitative information to inform on the impact of potential control programmes. This demonstrates the value of a One Health approach to surveillance and control of this emerging infectious disease. Funding. ARS Océan Indien, EAFRD, RITA Mayotte, VEEPED, Wellcome.

45 Introduction

Controlling zoonotic and vector-borne infections is complex, and requires getting an accurate 46 understanding of pathogen transmission within animal populations, and pathogen spillover to 47 humans, whilst accounting for environmental factors impacting on vectors.^{1,2} Rift Valley fever 48 (RVF) is an emerging zoonotic arbovirosis and an haemorrhagic fever that is threatening 49 animal and human health, mainly in Africa.³ Livestock (cattle, sheep and goats) are RVF virus 50 51 amplifying hosts, aquiring infection through the bites of infectious mosquitoes (mainly Aedes spp. and *Culex* spp.).⁴ Humans aquire infection by direct contact with infectious animal tissues 52 (upon abortions or animal slaughter), although vector transmission may also play a role.^{4,5} 53 Since 2015, RVF has been listed as a priority emerging disease by the WHO R&D Blueprint⁶ 54 and is today of global health concern,⁷ particularly as it has significantly expanded its 55 geographical range over recent decades.⁵ Current disease control options for reducing disease 56 risk in humans heavily rely on controlling virus transmission in animal populations. The 57 impact of disease control measures in livestock on reducing RVF risk in humans has not yet 58 been assessed, and doing so requires estimating key transmission parameters between 59 livestock and from livestock to humans; using animal and human epidemiological data. 60

Mayotte, an island located in the South Western Indian Ocean region, reported a RVF 61 epidemic in 2007-2008.8 In a previous paper, we used longitudinal livestock seroprevalence 62 data to model RVF virus emergence in the livestock population, and we estimated that the 63 likelihood of re-emergence was very low in a closed ecosystem (i.e. without the import of 64 infectious animals). However, a few imported infectious animals would be sufficient to 65 trigger another large epidemic, especially if neighbouring countries were affected and levels 66 of herd immunity had declined due to livestock population turnover.⁹ About ten years later, in 67 2018, RVF outbreaks were reported in several East African countries (e.g. Kenya, South 68 Sudan, Uganda, Rwanda).^{10,11} In Mayotte, between November 2018 and August 2019, a total 69

of 143 human cases (RVF virus RT-PCR confirmed) were reported (figure 1). The Veterinary Services of Mayotte, the regional health authorities (Agence de Santé Océan Indien) and the French Public Health Agency (Santé Publique France) did further epidemiological investigations to track the level of virus transmission in the animal population and study the possible route of human exposure to RVF virus. These investigations produced a uniquely well documented RVF epidemic, with both human and livestock datasets, including incidence and prevalence data.

We present these data and use them to extend the mathematical model previously developed to study RVF virus transmission in the livestock population in Mayotte,⁹ by explicitly modelling the human population. We fit this new mathematical model to both livestock and human epidemiological datasets from the 2018-2019 epidemic, allowing for the first time (i) to estimate the probability of RVF virus transmission to humans (virus spillover), (ii) to estimate the likelihood of another epidemic, and (iii) to assess the impact of different vaccination strategies in livestock on the disease risk reduction in humans.

84

85 Methods

86 Data: the course of the epidemic in both humans and livestock

87 The first human case, defined as a patient showing dengue-like symptoms and testing positive to RVF virus RT-PCR¹² was reported at the end of November 2018. The number of reported 88 cases rose quickly after the start of the rainy season (figure 1). The epidemic peaked around 89 week 7 of 2019 (February 11-17, 2019), with 18 confirmed cases. Between November 2018 90 and August 2019, a total of 143 confirmed RVF human cases were reported. From these 143 91 human cases, 127 were investigated and 16 were lost to follow-up. All investigated cases 92 93 were indigenous, most of them were male (male/female ratio=3), with a median age of 41 years-old [age range : 4-75] (13% were under 20, 80% were between 20 and 65 years-old, and 94

7% were over 65). About two-third of investigated cases (68%, n=86) reported having regular 95 direct contact with livestock, whilst 32% (n= 41) reported no previous contact with animals. 96 Symptoms were available for 98 cases, including fever, headache, arthralgia, asthenia, 97 myalgia, nausea and vomiting, and retro-orbital pain. In addition, between July 2018 and June 98 2019, a total of 1169 livestock sera, collected as part as the annual surveillance campaign, 99 were tested against RVF IgG (ID Screen RVF Competition ELISA, IDVet, Grabels, France). 100 RVF IgG antibodies can be detected in infected animals from about a week post-infection, 101 and up to several years.⁴ Date of birth was available for 493 of these sampled animals, and to 102 follow the emergence of the virus in the livestock population, we plotted quarterly age-103 stratified RVF IgG prevalence for the period July 2018 to June 2019 (figure 2A-D). In the 104 third quarter of 2018 (July - September 2018), that is before the report of the first human case, 105 the oldest animals were seropositive (figure 2A) probably from the previous epidemic.⁹ The 106 107 IgG prevalence increased in all age groups during the first quarter (January-March 2019, figure 2C) and second quarter (April-June 2019, figure 2D) of 2019, coincident with the 108 109 reported cases in humans.

Moreover, sera from illegally imported livestock seized by the Veterinary Services between June and August 2018 tested positive to RVF IgM (indicative of recent infections) (table 1). Finally, ongoing phylogenetic analyses on human-derived samples suggest that the incriminated RVF virus lineage may belong to the Kenya-2 clade, and is closely related to the strains from recent outbreaks in Eastern Africa (Cardinale, pers comm).

115

116 Model assumptions

We used these human and livestock data to develop our mathematical model. The study period ranged from the first week of July 2018 (week 27, July 2nd-9th) until the first week of August 2019 (week 31, July 29th- August 4th 2019), capturing the period between the

120 assumed importing time-window of infectious animals and the last reported human case.
121 Based on the data from illegally imported livestock, we assumed that RVF virus was imported
122 in Mayotte in July 2018, at which time most indigenous livestock were susceptible.
123 Throughout the epidemic, we assumed humans were infected by direct contact with infectious
124 livestock.

125

126 Model structure

The SEIR mathematical model of emergence previously developed for RVF in the livestock 127 population of Mayotte⁹ was adapted to the current epidemiological context. A discrete-time 128 deterministic framework was used, with a daily time step. For full details on the model in 129 livestock, see Metras and colleagues.⁹ As a reminder, transmission of RVF virus amongst 130 livestock ($\beta_{livestock-livestock}(t)$), was assumed to be vector-borne and modelled as a function of 131 132 monthly NDVI (Normalized Difference Vegetation Index) values as a proxy for vector abundance. Here we used rainfall data¹³ instead of NDVI as the study period, the time step 133 134 and the human epidemic curve available for fitting had a smaller time resolution (appendix p2, equations S1a-S1c). To this livestock model, we added a module simulating RVF virus 135 spillover from livestock into the human population, that is, the transmission of RVF virus 136 from livestock to humans. In this model (noted Model 1), we assumed that humans acquired 137 infection by direct contact with livestock (figure S1). Humans were Susceptible (S_h) , became 138 infected by direct contact (E_{hc}) with infectious livestock; after which they were successively 139 infectious (I_{hc}) and immune (R_h) . The corresponding model equations, transmission parameter 140 $\beta_{livestock-human-c}$ and force of infection by direct contact with livestock $\lambda_{contact}(t)$ are presented in 141 the supplementary appendix (appendix p2, equations S2a-S2e). 142

143

145 Model parameterization and fitting

Model parameters related to the natural history of infection (in animals and humans), and to 146 the structure of livestock and human populations were fixed (appendix p4, table S1). Animals 147 from all age-groups were assumed susceptibles at time t_0 , except the age-groups 9 and 10, for 148 which the proportion of immune at t_0 were 5% and 20.5 %, respectively (as per the 2017-2018) 149 IgG seroprevalence campaign). The reporting fraction of human cases was set to $\rho=1.9\%$, as a 150 post-epidemic serological study in humans, conducted in 2011 in Mayotte, estimated that 151 3.5% (95 %CI [2.6 - 4.8]) of the human population was RVF IgG-positive.14 Assuming a 152 population size of 212,645 inhabitants in 2012,15 this corresponds on average to 7,442 persons 153 154 being seropositive. Assuming that the past epidemic size is similar to the present one, and based on 143 cases currently reported, gives a reporting fraction of 1.9% (95% CI [1.4 -155 2.5]). Finally, we seeded the model with 10 infectious animals at t_0 (t_0 =Monday 2nd July, 156 2018), representing the infectious imports. 157

Parameters related to rainfall-dependent and constant viral transmission (appendix p4, model 158 1, table S1) were estimated by fitting the model to data. Parameter estimation was done by 159 fitting simultaneously the (i) quarterly age-stratified simulated proportion of immune 160 livestock $(p_{a,q})$ to quarterly RVF IgG prevalence (appendix p5, equation S5a, and figure 2AD); 161 and (ii) the simulated weekly number of reported incident cases in humans (appendix p5, 162 equation S6c) to the weekly number of reported cases (figure 1, all cases). For fitting, we 163 sampled from the posterior distributions of three parameters, $\theta_1 = \{A, B, \beta_{livestock-human-c}\}$ 164 (appendix p6) using a Monte Carlo Markov Chain Metropolis-Hastings (MCMC-MH) 165 algorithm.¹⁶ For details on parameter estimation and model fitting, see appendix p5-6. 166

Finally, rainfall data were downloaded from Meteofrance website, as cumulated rainfall over
10-day periods.¹³ Daily rainfall, used as input data in the model, was calculated by dividing
these values by ten over each 10-day period.

170 Forecasting and vaccination scenarios

We did projections for six scenarios (table 2). For all scenarios, we simulated 2,500 stochastic 171 trajectories by sampling randomly from the posterior distributions of the estimated 172 parameters, and we seeded the model with 10 infectious (I_{liv}) livestock. Scenario 1 aimed at 173 estimating the likelihood of virus re-emergence, without disease control intervention, in the 174 following rainy season, that is 2019-2020, in a closed ecosystem, using the same rainfall 175 values as in the 2018-2019 rainy season. Scenarios 2-6 aimed at assessing the impact that 176 different livestock vaccination strategies could have had on the number of human cases in the 177 past epidemic (2018-2019). We assumed the use of a single-dose highly immunogenic 178 vaccine (90 % vaccine efficacy),¹⁷ and a 14-days lag between vaccination and build-up of 179 immunity. Figures of vaccination campaigns in Mayotte in 2017 (vaccination for other 180 livestock diseases conducted on the island), showed that about 3,000 vaccine doses are 181 182 routinely administered to livestock over a year by the local veterinarians (Data CoopADEM, not shown). Scenario 2 tested the impact of administrating all these 3,000 doses in one month, 183 184 in December 2018, immediately after the report of the first human case (joint animal-human alert date for response), corresponding to the current vaccinating capacity in Mayotte in an 185 emergency setting. Scenario 3 assumed that all 3,000 doses were administered in January 186 2019, one month following the report of first human case, allowing extra time for organising 187 the vaccination campaign. Scenario 4-6 assumed an extra-vaccine supply and an emergency 188 mass vaccination, allowing 6,000 doses to be administered in December 2018 (Scenario 4), 189 6,000 doses to be administered in January 2019 (Scenario 5), and 9,000 doses to be 190 191 administered in January 2019 (Scenario 6).

192 Sensitivity analysis

193 To test for the possibility of a viral transmission from livestock to humans by direct contact 194 and by vector transmission, and to assess the impact of rainfall on transmission, we also developed two further models and compared them (Model 2 and Model 3, table S2). Details
on models equations, parameters, fitting methods and model comparison are presented in the
appendix p3-7.

198

199 Ethics statement

The livestock data were collected under the under the Mayotte disease surveillance system 200 (Système d'Epidémiosurveillance Animale à Mayotte, SESAM) with the approval of the 201 Direction of Agriculture, Food and Forestry (DAAF) of Mayotte. For human data, according 202 to French law, only "research involving a human being" (research defined by article L. 1121-203 204 1 and article R. 1121-1 of the Code de la santé publique) are compelled to receive the approval of ethics committee. This study was based on anonymous data collected from health 205 professionals for public health purposes relating to the health surveillance mission entrusted 206 207 to Santé publique France by the French Law (article L. 1413-1 code de la santé publique). Therefore, the study did not meet the criteria for qualifying a study "research involving a 208 209 human being" and did not require the approval of an ethics committee. Furthermore, as the data were anonymous, it did not require an authorization of the French data protection 210 authority (Commission Nationale informatique et libertés). 211

212

213 Role of the funding sources

The funding sources have no role in study design; in the collection, analysis, and interpretation of data; in the writing of the report; and in the decision to submit the paper for publication.

218 **<u>Results</u>**

The median number of predicted human reported cases was 140 (90% Credible Interval CrI [9 219 - 265], table 2 and figure 3A) showing good agreement with the observed confirmed cases 220 (143 cases, figure 3A), and the livestock IgG prevalence data (figure 2AD). The epidemic 221 started in November 2018, shortly after the rainy season started. The epidemic peaked in 222 livestock during the 5th week of 2019 (week 31 of the model, January 28 - February 3, 2019) 223 and two weeks later in humans (week 33 of the model, February 11-17, 2019) (figure 3B and 224 225 table 2). The total median number of predicted human cases was 7,362 (90%CrI [458 -13,962]), corresponding to 3.0% (90%CrI [1.2 - 3.6]) of the total human population (table 2). 226 For livestock, the total number of animals affected at the end of the epidemic were estimated 227 to 18,801 (90%CrI [1,250 - 35,572]), corresponding to 55.8 % (90%CrI [27.1 - 59.5]) of the 228 total livestock population (figure 3C and table 2). 229

The timing-varying reproductive number $R_s(t)$ in the livestock population was a function of rainfall and varied along the study period (figure 3D). Its maximum value (median $R_s(t)=2.24$, 90%CrI [1.77 – 3.10]) was reached on the first week of 2019 (December 31, 2018 - January 6th, 2019, 87.4 mm of cumulated rain on that week), that is four and six weeks prior to the peaks in livestock and humans, respectively. Finally, the constant transmission rate from livestock to humans was estimated at 2.2 per 10 million inhabitants per day (95%CrI [1.8 – 2.7]) (appendix p8, table S3).

The probabilistic forecasts showed that RVF re-emergence in the next rainy season (2019-2020) was very unlikely (figure 2A). Figure 4AD presents the results of the impact of the different livestock vaccination strategies on the number of human reported incident cases (figure 4AB) and livestock incident cases (figure 4CD). The implementation of massive vaccination campaign immediately after the first human case (table 2, Scenario 4, i.e. 6,000 doses in December 2018) allowed reducing the number of human cases reported by a third

(median= 94 cases), while waiting one more month would have required 50 % more vaccine 243 doses for a similar impact (figure 4AD and table 2, Scenario 6, i.e. 9,000 doses in January 244 2019, n=93 cases). The two alternative models tested as sensitivity analysis (Model 2 and 3) 245 did not fit the data as well as Model 1 (appendix p7, figure S2AB, and table S2). For Model 2, 246 the spillover to humans via direct contact was slightly lower than in Model 1, estimated at 1.3247 per 10 million inhabitants per day (95%CrI $[2 \cdot 1 - 1 \cdot 9]$) (appendix p8, table S3). In this model, 248 the maximum $R_s(t)$ values in livestock was $R_s(t) = 2.48$ (90%CrI [1.96 - 3.46]), and the 249 intensity of the vector transmission from livestock to humans represented 0.6 % (95%CrI 250 [0.12 - 2.1]) of the between livestock vector transmission (scaling factor X, appendix p8, 251 table S3). 252

253

254 **Discussion**

This paper presents a dataset of a large RVF epidemic, combining prospectively collected 255 livestock and human data, which represents a reference case study for RVF. Testing of human 256 257 cases of dengue-like syndrome with RVF RT-PCR has been systematically performed in Mayotte for the last ten years (since 2008), giving confidence that RVF had been absent from 258 the Island for a decade, and that the shape of the epidemic curve accurately reflected its actual 259 260 timing. This was cross-validated by the observed changes in the seroprevalence in livestock, exhibiting a clear pattern of viral emergence. Finally, the routine IgM testing of illegally 261 imported livestock provided an estimate of the likely time-window of virus importation, 262 which also coincided with the timing of RVF outbreaks on the East African mainland.¹¹ 263

We used these data to fit a mathematical model to quantify transmission amongst livestock and spillover to the human population, to estimate the likelihood of another emergence in the following rainy season (2019-2020), and to assess the impact of livestock vaccination on preventing disease risk in humans. At the end of the modelled epidemic wave, in August 2019, the proportion of immune livestock was estimated at $55 \cdot 8 \%$ (90%CrI [27·1 - 59·5]), and the likelihood of RVF re-emergence in Mayotte in the 2019-2020 rainy season was very low. We also demonstrated that reactive vaccination (6,000 animals vaccinated in December 2018) would have reduced the number of human cases by a third, while 50 % more vaccine doses would have been necessary a month later for a similar impact. This highlights that animal vaccination can prevent human cases, as well as cases in livestock, but that early detection and rapid vaccination are critical to RVF control at the early stage of the epidemic.

Rainfall is a known driver for RVF virus transmission,¹⁸ and was used as a proxy for vector 275 abundance. Temperature above 26°C may also drive RVF virus transmission.¹⁹⁻²¹ The 276 temperatures of Mayotte varying annually between 25°C and 35°C,⁹ we therefore assumed 277 that temperature would not be a driver for transmission in this setting. In areas with cooler 278 climates, temperature should be explicitly modelled.²² The highest estimated $R_s(t)$ value was 279 2.24 (90%CrI [1.77 – 3.10]), in line with previous estimations of R_0 .²³⁻²⁵ Finally, previous 280 models parameterised the transmission parameter from livestock to humans as a fixed 281 parameter at 1.7 per 10 thousand persons per day.^{24,26,27} For the first time, our model estimated 282 that value, which was lower, at 2.2 per 10 million persons per day (95%CrI [1.8 - 2.7]). This 283 can be used as benchmark for future modelling work. 284

RVF human cases without previous contact with animals or animal products have been 285 reported in other settings, although this fraction had not been documented during an epidemic. 286 ^{28,29} Here, the epidemiological investigations documented the fraction of cases that had no 287 contact with animals, and were therefore probably infected via mosquito bite. Regarding 288 livestock data, sera were tested as part of the national surveillance campaign, and some 289 samples were taken in areas reporting human cases. This may have led to an over-estimate the 290 proportion of IgG positive animals. However, most animal sampling was conducted from 291 January 2019 onwards, when RVF virus had already spread across the whole island. 292

A limitation of the model related to the parameterization of the reporting rate in humans, used 293 as an input value. For this, we used past data,¹⁴ assuming the 2007-2008 wave was of similar 294 size to the 2018-2019 epidemic. Whilst there is no human data to support this assumption, our 295 previous work⁹ estimated an overall post-epidemic livestock prevalence at about 50 %, which 296 is in line with our current estimates. Further data collection estimating human post-epidemic 297 prevalence would allow an accurate estimation of this reporting rate. In addition, the limited 298 total number of human reported cases led to wide confidence intervals for the forecasts. 299 Collecting more epidemic data at a smaller spatio-temporal resolution would have also 300 allowed spatial stratification, and testing for finer vaccination protocols. Finally, as in our 301 previous model, we assumed homogeneous mixing. However, Mayotte is a small island 302 (374km²), the ecosystem shows limited spatial variations, livestock production systems are 303 extensive with animals raised outdoor, and therefore potentially similarly exposed to RVF 304 mosquito vectors.9 305

As long as no vaccination in humans is available, disease surveillance in animals, contingency planning, and the timely implementation of livestock vaccination, are key for reducing human disease risk. During this epidemic, livestock were not vaccinated due to a lack of vaccine supply, and limited human resources, and the vaccination scenarios tested would require emergency funds and vaccine doses.

311

This is a uniquely detailed investigation into an outbreak of an emerging arbovirus, combining animal and human data, with a mathematical model for RVF. This work represents a collaboration between public health agency, animal health surveillance network, farmers' association, and researchers, from the beginning of the epidemic, and conducted in real-time, as the epidemic unfolded. Delays in getting serological data were inherent to climatic conditions (storms) and field work constraints in remote areas. Nevertheless, we addressed in

318 practice the challenges of a One Health approach,³⁰ and this work demonstrates its value to 319 surveillance and control of zoonotic emerging infectious diseases.

320

321 Acknowledgements

The authors wish to thank the Agence de Santé océan Indien that has participated in collecting 322 human cases data, the laboratory of Centre Hospitalier de Mayotte which has performed the 323 virological analyses on human samples, the animal SESAM surveillance system, the 324 CoopADEM (Coopérative agricole des éleveurs mahorais), the Cirad-CYROI, the Veterinary 325 Services, and the LVAD (Laboratoire Vétérinaire d'Analyses Départemental de Mayotte) for 326 the data collection and the serological analyses on livestock samples. Finally, we thank 327 Harold Noël (MD) from Santé publique France for facilitating human data access in the early 328 stage of the epidemic. 329

330

331 **Declaration of interests**

332 The authors declare no conflict of interest.

333

334 Funding sources

RVF RT-PCR were conducted as part as the surveillance system on dengue-like syndrome 335 since 2008, funded by Agence de Santé océan Indien. The animal sampling and analyses were 336 funded by EAFRD (European Agricultural Fund for Rural Development) and RITA (Réseau 337 d'Innovation et de Transfert Agricole) Mayotte. WJE and AC were funded by the Department 338 of Health and Social Care using UK Aid funding managed by the NIHR (VEEPED: PR-OD-339 1017-20007). The views expressed in this publication are those of the authors and not 340 necessarily those of the Department of Health and Social Care. SF was funded by a Wellcome 341 Senior Research Fellowship (210758/Z/18/Z). 342

343 **Contributions**

The corresponding author (RM) had access to all the data in the study and had final reponsibility for the decision to submit for publication. RM, WJE, LD, GF, YH, MS conceptualized and designed the study. CY, LD, SC, YH, MS collated the data and did data management. RM, GF performed the analyses. RM, WJE, CY, LD, GF, AC, SF, GLG, CS, EC, LF, YH, MS interpreted and discussed the data and results. RM was reponsible for drafting the manuscript. All authors reviewed and approved the final manuscript.

350

351 **<u>References</u>**

Woolhouse MEJ, Dye C. Preface. *Philos Trans Roy Soc London Ser B* 2001; **356**: 981–
 982.

354

Jones BA, Grace D, Kock R, et al. Zoonosis emergence linked to agricultural
intensification and environmental change. *Proc Natl Acad Sci U S A* 2013;110:8399-404.

3. Clark MHA, Warimwe GM, Di Nardo A, Lyons NA, Gubbins S. Systematic literature 358 review of Rift Valley fever virus seroprevalence in livestock, wildlife and humans in Africa 359 from 1968 to 2016. PLoS Negl Trop Dis 2018;**12**:e0006627. 360 doi:10.1371/journal.pntd.0006627 361

362

4. Bird BH, Ksiazek TG, Nichol ST and MacLachlan NJ. Rift Valley fever virus. *J Am Vet Med Assoc* 2009;234:883–893.

365

5. Nanyingi MO, Munyua P, Kiama SG, et al. A systematic review of Rift Valley Fever
epidemiology 1931–2014. *Infect Ecol Epidemiol* 2015; 5: 10.3402/iee.v5.28024

369 6. World Health Organization. 2018. Epidemic and pandemic-prone diseases, List of
370 Blueprint priority diseases. Available at <u>http://www.emro.who.int/fr/pandemic-epidemic-</u>
371 <u>diseases/news/list-of-blueprint-priority-diseases.html</u> Accessed on Nov 26, 2019.

372

- 7. Hatchett R and Lurie N. Outbreak responses as an essential component of vaccine
 development. *Lancet Infect Dis* 2019;19:e399-e403. doi:10.1016/S1473-3099(19)30305-6
- 8. Métras R, Cavalerie L, Dommergues L, et al. The Epidemiology of Rift Valley Fever in
- 377 Mayotte: Insights and Perspectives from 11 Years of Data. PLoS Negl Trop Dis 2016;10:
- 378 e0004783. https://doi.org/10.1371/journal.pntd.0004783
- 379
- 9. Métras R, Fournié G, Dommergues L, et al. Drivers for Rift Valley fever emergence in
 Mayotte: A Bayesian modelling approach. *PLoS Negl Trop Dis* 2017;11:e0005767.
 <u>https://doi.org/10.1371/journal.pntd.0005767</u>
- 383
- 10. ProMED. Rift Valley fever Kenya (02): (Wajir). Published Date: 2018-06-09. Archive
 Number: 20180609.5847216. Accessed on Dec 5th 2019.
- 386
- 11. Food and Agriculture Organization of the United Nations. EMPRES-i. Global Animal
 Disease Information System. Available at : <u>http://empres-i.fao.org/eipws3g/</u>. Accessed on
 Dec 11th, 2019.

391	12. Bird BH, Bawiec DA, Ksiazek TG, Shoemaker TR, Nichol ST. Highly sensitive and
392	broadly reactive quantitative reverse transcription-PCR assay for high-throughput detection of
393	Rift Valley fever virus. J Clin Microbiol 2007;45:3506–3513. doi:10.1128/JCM.00936-07
394	
395	13. Meteofrance. 2019. Donnees decadaires agrometeorologiques. Available at :
396	https://donneespubliques.meteofrance.fr/?fond=produit&id_produit=113&id_rubrique=37

397 Accessed on Sept 23, 2019.

398

14. Lernout T, Cardinale E, Jego M, et al. Rift Valley Fever in Humans and Animals in
Mayotte, an Endemic Situation? *PLoS ONE* 2013; 8: e74192.
<u>https://doi.org/10.1371/journal.pone.0074192</u>

402

- 403 15. Institut National de la statistique et des etudes economiques (Insee). Habitants a Mayotte.
- 2017 Available at : <u>https://www.insee.fr/fr/statistiques/3286558#documentation</u>, Accessed on
 Oct 5, 2019.

406

407 16. Camacho A and Funk S. fitR: Tool box for fitting dynamic infectious disease models to408 time series. R package version 0.1.

409

410 17. Dungu B, Lubisi BA, Ikegami T. Rift Valley fever vaccines : current and future needs.
411 *Curr Opin Virol* 2018;**29**:8-15. doi: 10.1016/j.coviro.2018.02.001

- 413 18. Sang R, Lutomiah J, Said M, et al. Effects of Irrigation and Rainfall on the Population
- 414 Dynamics of Rift Valley Fever and Other Arbovirus Mosquito Vectors in the Epidemic-Prone
- 415 Tana River County, Kenya. J Med Entomol 2017;54:460–470. doi:10.1093/jme/tjw206

416

417 19. Turell M J, Rossi CA and Bailey CL. Effect of extrinsic incubation temperature on the
418 ability of Aedes taeniorhynchus and Culex pipiens to transmit Rift Valley fever virus. *Am J*419 *Trop Med Hyg* 1985;**3**4:1211–1218.

420

20. Brubaker JF and Turell MJ. Effect of environmental temperature on the susceptibility of
Culex pipiens (Diptera: Culicidae) to Rift Valley fever virus. *J Med Entomol* 1998;**35**:918–
921.

424

425 21. Lo Iacono G, Cunningham AA, Bett B, Grace D, Redding DW, Wood JLN.
426 Environmental limits of Rift Valley fever revealed using ecoepidemiological mechanistic
427 models. *Proc Natl Acad Sci U S A* 2018;115:E7448–E7456. doi:10.1073/pnas.1803264115
428

429 22. Esser HJ, Mögling R, Cleton NB, et al. Risk factors associated with sustained circulation
430 of six zoonotic arboviruses: a systematic review for selection of surveillance sites in non431 endemic areas. *Parasit Vectors* 2019;**12**:265. doi:10.1186/s13071-019-3515-7

432

433 23. Barker CM, Niu T, Reisen WK, Hartley DM. Data-Driven Modeling to Assess
434 Receptivity for Rift Valley Fever Virus. *PLoS Negl Trop Dis* 2013; 7: e2515.
435 https://doi.org/10.1371/journal.pntd.0002515

436

437 24. Xue L, Scott MH, Cohnstaedt LW and Scoglio C. A network-based meta-population
438 approach to model Rift Valley fever epidemics. *J Theor Biol* 2012;**306**:129–144.

439 doi: 10.1016/j.jtbi.2012.04.029

441	25. Danzetta ML, Bruno R, Sauro F, Savini F and Calistri P. Rift Valley fever transmission
442	dynamics described by compartmental models. Prev Vet Med 2010;134:197-210.
443	https://doi.org/10.1016/j.prevetmed.2016.09.007
444	
445	26. Mpeshe SC, Haario H and Tchuenche JM. A Mathematical Model of Rift Valley Fever
446	with Human Host. Acta Biotheor 2011; 59: 231. https://doi.org/10.1007/s10441-011-9132-2
447	
448	27. Lugoye J, Wairimu J, Alphonce CB, Ronoh M. Modeling Rift Valley fever with treatment
449	and trapping control strategies. Appl Math 2016;7:556-568
450	
451	28. Archer BN, Thomas J, Weyer J, et al. Epidemiologic Investigations into Outbreaks of Rift
452	Valley Fever in Humans, South Africa, 2008-2011. Emerg Infect Dis 2013;19:1918-1925.
453	doi:10.3201/eid1912.121527
454	
455	29. Shoemaker TR, Nyakarahuka L, Balinandi S, et al. First Laboratory-Confirmed Outbreak
456	of Human and Animal Rift Valley Fever Virus in Uganda in 48 Years. Am J Trop Med Hyg
457	2019; 100 :659–671. doi:10.4269/ajtmh.18-0732
458	
459	30. Scoones I, Jones K, Lo Iacono G, Redding DW, Wilkinson A, Wood JLN. Integrative
460	modelling for One Health: pattern, process and participation. Philos Trans R Soc Lond B Biol
461	Sci 2017;372:20160164. doi:10.1098/rstb.2016.0164
462	
463	
464	
465	



Figure 1. Weekly number of reported human cases and average daily rainfall pattern (mm). Between November 2018 and July 2019, 143 cases were reported. A total of 86 cases reported a direct exposure to livestock or their fluids (red), 41 did not report any direct exposure to livestock (green), and 16 were lost to follow-up (grey).





539

Figure 3AD. Model results. (A) Weekly reported incident human cases. Observed data (red dots), predicted median (red solid line), interquartile range (dashed lines) and 90 % CrI (red envelope). (B) Predicted (reported and unreported) number of infectious livestock (blue) and humans (red). (C) Median (solid lines) and 90%CrI envelopes of the predicted proportion of Susceptible (green) and Immune (black) livestock over the course of the epidemic. (D) Value of Rs(t) over the course of the epidemic, and 90%CrI. In (A), (B) and (C), the vertical blue and red vertical lines correspond to the predicted epidemic peaks in livestock and humans, respectively. The vertical black line corresponds to the end of August 2019 (end of the fitting period).

- 547
- 548
- 549
- 550
- 551
- 552
- 553



Figure 4AD. Comparison of the vaccination strategies versus no intervention. (A) Median weekly number of incident human cases and (B) Predicted number of human reported cases. (C) Median weekly number of incident infected livestock and (D) Predicted number of infected livestock. (A) and (C) present the scenario with no intervention (red solid line, Scenario 1), the vaccinations in December 2008 (the black solid line represents the 3,000 doses, and the dashed black line the 6,000 doses), and in January 2019 (the blue solid line represents the 3,000 doses, the dashed blue line the 6,000 doses ; and the dotted blue line the 9,000 doses scenario).

576	Table 1. Number of	illegally imported liv	estock seized by the	Veterinary Se	ervices and	positive to RVF
577	IgM, between May ar	nd October 2018 (Sou	rce : Mayotte Veteri	nary Services)		

	No. of IgM positive	No. animals seized	Proportion of IgM positive [95 % CI]
May 2018	0	8	0 % [0-40]
June 2018	10	31	32% [17-51]
August 2018	2	18	11% [2-36]
September 2018	0	1	0 % [0-90]
October 2018	0	5	0 % [0-54]

579 Table 2. Predicted median and 90%CrI (Credibility Interval) of the epidemic size, post-epidemic prevalences, and timing of the epidemic peaks under the different scenarios tested (Model 1)

Scenario number	Scenario 1	Scenario 2 (2.1)	Scenario 3 (2.2)	Scenario 4 (2.4)	Scenario 5 (2.3)	Scenario 6 (2.5)
Assumptions	No intervention	3,000 doses, Dec 2018	3,000 doses, Jan 2019	6,000 doses, Dec 2018	6,000 doses, Jan 2019	9,000 doses, Jan 2019
Epidemic size						
Total human cases	7,362 [458-13,962]	6,088[542-11,564]	6,410 [0-12,074]	4,931 [28-9,240]	5,552[317-10,611]	4,895 [135-9,122]
Human reported cases	140 [9-265]	116 [10-220]	122 [0-230]	94 [1-176]	105 [6-202]	93 [3-173]
Total livestock cases	18,801 [1,250-35,572]	15,650[1519-29,387]	16,459 [10 - 30,890]	12,586 [167-23,276]	14,182 [871 -26,859]	12,540 [389-23,045]
Post-epidemic						
Human prevalence	3.0 [1.2-3.6]	$2 \cdot 1 \ [1 \cdot 0 - 3 \cdot 0]$	2.7 [0.0-3.2]	$2 \cdot 0 [0 \cdot 1 - 2 \cdot 6]$	2.3 [0.5 -2.9]	2.0 [0.2 - 2.7]
Livestock prevalence	55.8 [27.1-59.5]	46.6 [21.6 - 51.1]	$48 \cdot 7[1 \cdot 0 - 53 \cdot 9]$	37.7 [1.7 - 43.3]	42.1 [10.8-49.1]	36.9 [3.9 - 45.2]
Timing epidemic peaks						
Peak in humans	Week 33	Week 33	Week 33	Week 32	Week 32	Week 32
Peak in livestock	Week 31	Week 31	Week 31	Week 31	Week 31	Week 31