

# Brief Report: Genomic epidemiology of a densely sampled COVID19 outbreak in China

Erik M Volz<sup>1\*</sup>, Han Fu<sup>1</sup>, Haowei Wang<sup>1</sup>, Xiaoyue Xi<sup>2</sup>, Wei Chen<sup>3</sup>, Dehui Liu<sup>3</sup>,  
Yingying Chen<sup>3</sup>, Mengmeng Tian<sup>3</sup>, Wei Tan<sup>4</sup>, Junjie Zai<sup>5</sup>, Wanying Sun<sup>6</sup>, Jiandong  
Li<sup>6</sup>, Junhua Li<sup>6</sup>, Xingguang Li<sup>7†\*</sup>, Qing Nie<sup>3†\*</sup>

**\*For correspondence:**

[nieqing0454@163.com](mailto:nieqing0454@163.com) (QN);  
[xingguanglee@hotmail.com](mailto:xingguanglee@hotmail.com) (XL)

†These authors contributed equally  
to this work

**Present address:** <sup>†</sup>Department of  
Microbiology, Weifang Center for  
Disease Control and Prevention,  
Weifang 261061, China. Tel:  
+86-0536-8098503; <sup>‡</sup>Hubei  
Engineering Research Center of  
Viral Vector, Wuhan University of  
Bioengineering, Wuhan, 430415,  
China. Tel: +86-027-89648139

<sup>1</sup>Department of Infectious Disease Epidemiology and MRC Centre for Global Infectious  
Disease Analysis, Imperial College London, Norfolk Place, W2 1PG, United Kingdom;  
<sup>2</sup>Department of Mathematics, Imperial College London, London SW7 2AZ, United  
Kingdom; <sup>3</sup>Department of Microbiology, Weifang Center for Disease Control and  
Prevention, Weifang 261061, China.; <sup>4</sup>Department of Respiratory Medicine, Weifang  
People's Hospital, Weifang 261061, China.; <sup>5</sup>Immunology Innovation Team, School of  
Medicine, Ningbo University, Ningbo 315211, China.; <sup>6</sup>Shenzhen Key Laboratory of  
Unknown Pathogen Identification, BGI-Shenzhen, Shenzhen 518083, China.; <sup>7</sup>Hubei  
Engineering Research Center of Viral Vector, Wuhan University of Bioengineering, Wuhan,  
430415, China.

**Abstract** Analysis of genetic sequence data from the pandemic SARS Coronavirus 2 can provide  
insights into epidemic origins, worldwide dispersal, and epidemiological history. With few  
exceptions, genomic epidemiological analysis has focused on geographically distributed data sets  
with few isolates in any given location. Here we report an analysis of 20 whole SARS-CoV 2 genomes  
from a single relatively small and geographically constrained outbreak in Weifang, People's Republic  
of China. Using Bayesian model-based phylodynamic methods, we estimate the reproduction  
number for the outbreak to be 1.99(95% CI:1.48-3.14). We further estimate the number of  
infections through time and compare these estimates to confirmed diagnoses by the Weifang  
Centers for Disease Control. We find that these estimates are consistent with reported cases and  
there is unlikely to be a large undiagnosed burden of infection over the period we studied.

## Introduction

We report a genomic epidemiological analysis of one of the first geographically concentrated  
community transmission samples of SARS-CoV 2 genetic sequences collected outside of the initial

32 outbreak in Wuhan, China. These data comprise 20 whole genome sequences from confirmed  
33 COVID19 infections in Weifang, Shandong Province, People's Republic of China. The data were  
34 collected over the course of several weeks up to February 10, 2020 and overlap with a period  
35 of intensifying public health and social distancing measures. Phylodynamic analysis allows us to  
36 evaluate epidemiological trends after seeding events which took place in mid to late January, 2020.

37 The objective of our analysis is to evaluate epidemiological trends based on national surveillance  
38 and response efforts by Weifang Centers for Disease Control (CDC). This analysis provides an  
39 estimate of the initial rate of spread and reproduction number in Weifang City. In contrast to the  
40 early spread of COVID19 in Hubei Province of China, most community transmissions within Weifang  
41 took place after public health interventions and social distancing measures were put in place. We  
42 therefore hypothesize that genetic data should reflect a lower growth rate and reproduction number  
43 than was observed in Wuhan. A secondary aim is to estimate the total numbers infected and to  
44 evaluate the possibility that there is a large unmeasured burden of infection due to imperfect case  
45 ascertainment and a large proportion of infections with mild or asymptomatic illness.

46 To analyze the Weifang sequences, we have adapted model-based phylodynamic methods  
47 which were previously used to estimate growth rates and reproduction numbers using sequence  
48 data from Wuhan and exported international cases(Volz *et al.*, 2020). This analysis has several  
49 constraints and requirements:

50

51 *Importation of lineages from Wuhan.* The outbreak in Weifang was seeded by multiple lineages  
52 imported at various times from the rest of China. We use a phylodynamic model that accounts for  
53 location of sampling. Migration is modeled as a bi-directional process with rates proportional to  
54 epidemic size in Weifang. The larger international reservoir of COVID19 cases serves as a source of  
55 new infections and is assumed to be growing exponentially over this period of time.

56 *Nonlinear epidemiological dynamics in Weifang.* The maximum number of daily confirmed COVID19  
57 cases occurred on February 5, but it is unknown when the maximum prevalence of infection oc-  
58 curred. We use a susceptible-exposed-infectious-recovered (SEIR) model(Keeling and Rohani, 2011)  
59 for epidemic dynamics in Weifang. The model accounts for a realistic distribution of generation  
60 times and can potentially capture a nonlinear decrease in cases following epidemic peak.

61 *Variance in transmission rates(Lloyd-Smith *et al.*, 2005).*To estimate total numbers infected, the  
62 phylodynamic model must account for epidemiological variables which are known to significantly  
63 influence genetic diversity. Foremost among these is the variance in offspring distribution (number  
64 of transmissions per primary case). We draw on previous evidence based on the previous SARS  
65 epidemic which indicates that the offspring distribution is highly over-dispersed. High variance of  
66 transmission rates will reduce genetic diversity of a sample and failure to account for this factor  
67 will lead to highly biased estimates of epidemic size(Li *et al.*, 2017). Recent analyses of sequence  
68 data drawn primarily from Wuhan has found that high over-dispersion was required for estimated  
69 cases to be consistent with the epidemiological record(Volz *et al.*, 2020). Models assuming low  
70 variance in transmission rates between people would generate estimates of cases that are lower  
71 than the known number of confirmed cases. Separately, Grantz *et al.*(Grantz *et al.*, ????) have

72 found that high over-dispersion is required to reconcile estimated reproduction numbers with  
73 the observed frequency of international outbreaks. In this study, we elaborate the SEIR model to  
74 include a compartment( $J$ ) with higher transmission rates. The variance of the implied offspring  
75 distribution is calibrated to give similar overdispersion from the SARS epidemic.

## 76 Results

77 Despite an initial rapid increase in confirmed cases in Weifang in late January and early February, the  
78 number of confirmed cases by Weifang CDC show that outbreak peaked quite early and maximum  
79 number of cases occurred on February 5. Phylodynamic analysis supports the interpretation that  
80 control efforts reduced epidemic growth rates and contributed to eventual control. **Figure 1A**  
81 shows the estimated time scaled phylogeny (maximum clade credibility) including 20 lineages  
82 sampled from distinct patients in Weifang and 33 genomes sampled from Wuhan and internationally.  
83 **Figure 1B** illustrates the phylodynamic model which was co-estimated with the phylogeny which  
84 provides estimates of epidemiological parameters summarized in **Table 1**.

85 The estimated number of infections is shown Figure 1C. The time series of confirmed cases  
86 should lag the estimated number of infected because of delays from infection to appearance of  
87 symptoms and delays from symptoms to diagnosis. We also expect that an unknown proportion of  
88 infections will be missed by the surveillance system due very mild, subclinical, or asymptomatic  
89 infection. Our estimates do not support the hypothesis that there was a very large hidden burden  
90 of infection in Weifang over the period that the sequence data were sampled. Indeed, our central  
91 estimate for the number infected on 10th February is 53, which is only slightly in excess of the 44  
92 cumulative confirmed cases at the end of February.


**Table 1.** Summary of primary epidemiological and evolutionary parameters, including Bayesian prior distributions and estimated posteriors. Posterior uncertainty is summarized using a 95% highest posterior density interval.

Parameter	Prior	Posterior mean	95% HPD
Initial infected	Exponential(mean=1)	2.3	0.18-6.9
Initial susceptible	Log-normal(mean log=6, sd log=1)	787	102-3235
Migration rate <sup>1</sup>	Exponential(mean=10)	1.67	0.96-2.0
Reproduction number	Log-normal(mean log=1.03, sd log =0.5)	1.99	1.48-3.14
Molecular clock rate <sup>2</sup>	Uniform(0.0007,0.003)	0.0028	0.0024-0.003
Transition/transversion	Log-normal(mean log=1, sd log=1.25)	7.0	4.4-11.1
Gamma shape	Exponential(mean=1)	0.05	0.003-0.12

<sup>1</sup> Units: Migrations per lineage per year.

<sup>2</sup> Units: Substitutions per site per year.

93 While we do not have sufficient data to detect a large decrease in epidemic growth rates as the  
94 epidemic progressed, we do find that the growth rates are somewhat lower than estimated in other  
95 settings and during the early epidemic in Wuhan. We estimate  $R_0 = 1.99$  (95% HPD:1.48-3.14) and  
96 the growth rate in cases was approximately 6% per day. The relatively low value of  $R_0$  corresponds  
97 to growth during a period when Weifang was implementing a variety of public health interventions



**Figure 1.** Phylodynamic estimates and epidemiological model. A. A time scaled phylogeny co-estimated with epidemiological parameters. The colour of tips corresponds to location sampling. Red tips were sampled from Weifang, China. The credible interval of TMRCA is shown as a blue bar for all nodes with more than 50% posterior support. B. A diagram representing the structure of the epidemiological SEIR model which was fitted in tandem with the time scaled phylogeny. Colours correspond to the state of individuals sampled and represented in the tree (A). Note that infected and infectious individuals may occupy a low transmission state (I) or a high transmission rate state (J) to account for high dispersion of the reproduction number. C. Cumulative estimated infections through time produced by fitting the SEIR model and the cumulative confirmed cases (points) reported by Weifan CDC. The shaded region shows the 95% HPD and the line shows the posterior median. D. A root to tip regression showing approximately linear increase in diversity with time of sampling.

**Figure 1-Figure supplement 1.** Maximum likelihood time tree.

**Figure 1-Figure supplement 2.** Tree posterior density plot.

**Figure 1-Figure supplement 3.** Tree posterior density plot.

98 and contact tracing to limit epidemic spread. These interventions included public health messaging,  
99 establishing phone hotlines, encouraging home isolation for recent visitors from Wuhan (January  
100 23-26), optimizing triage of suspected cases in hospitals (January 24), travel restrictions (January 26),  
101 extending school closures, and establishing 'fever clinics' for consultation and diagnosis (January  
102 27) (Mao, 2020).

103 As well as providing novel epidemiological estimates, our results point to the significance of  
104 realistic modeling for fidelity of phylogenetic inference. The use of a model-based structured coa-  
105 lescent prior had large influence over estimated molecular clock rates and inferred TMRCA. **Figure**  
106 **Supplement 1** shows that maximum likelihood inference of time-scaled phylogenies produces a  
107 distribution of TMRCA which are substantially different than the Bayesian model-based analysis.  
108 Choice of population genetic prior will have a large influence on phylogenetic inference based on  
109 sparse or poorly informative genetic sequence data. Among the 20 Weifang sequences included in  
110 this analysis, there is mean pairwise difference of only three single nucleotide polymorphisms and  
111 only approximately twice as much diversity observed among the remainder of the sequences we  
112 studied. There is correspondingly low confidence in tree topology (**Figure Supplement 2**), and only  
113 three clades had greater than 50% posterior support including one clade which had a monophyletic  
114 composition of 13 Weifang lineages. The earliest Weifang sequence was sampled on January 25  
115 from a patient who showed first symptoms on January 16. These dates cover a similar range as the  
116 posterior TMRCA of all Weifang sequences (**Figure Supplement 3**).

## 117 Discussion

118 Our analysis of 20 SARS-CoV 2 genomes from Weifang, China has confirmed independent observa-  
119 tions regarding the rate of spread and burden of infection in the city. Surveillance of COVID19 is  
120 rendered difficult by high proportions of illness with mild severity and an unknown proportion of  
121 asymptomatic infection (Guan et al., 2020). The extent of under-reporting and case ascertainment  
122 rates has been widely debated. Analysis of genetic sequence data provides an alternative source of  
123 information about epidemic size which can be more robust to imperfect case ascertainment. We do  
124 not find evidence for a large hidden burden of infection within Weifang. The relatively low estimate  
125 of  $R_0$  is consistent with a slower rate of spread outside of Wuhan and effective control strategies  
126 implemented in late January.

127 While the value of pathogen genomic analyses is widely recognized for estimating dates of  
128 emergence (Verity Hill, 2020; Gire et al., 2014) and identifying animal reservoirs (Zhou et al., 2020;  
129 Dudas et al., 2018), analysis of pathogen sequences also has potential to inform epidemic surveil-  
130 lance and intervention efforts. With few exceptions (Stadler, 2020; Bedford, 2020), this potential  
131 is currently not being realized for the international response to COVID19. It is worth noting that  
132 the analysis described in this report was accomplished in approximately 48 hours and drew on  
133 previously developed models and packages for BEAST2 (Bouckaert et al., 2019; Volz and Siveroni,  
134 2018). It is therefore feasible for phylodynamic analysis to provide a rapid supplement to epidemio-  
135 logical surveillance, however this requires rapid sequencing and timely sharing of data as well as  
136 randomized concentrated sampling of the epidemic within localities such as individual cities.

## 137 Methods and Materials

### 138 Epidemiological investigation, sampling, and genetic sequencing.

139 As of 10 February 2020, 136  
140 suspected cases, and 214 close contacts were diagnosed by Weifang Center for Disease Control and  
141 Prevention. 28 cases were detected positive with SARS-CoV-2. Viral RNA was extracted using Maxwell  
142 16 Viral Total Nucleic Acid Purification Kit (Promega AS1150) by magnetic bead method and RNeasy  
143 Mini Kit (QIAGEN 74104) by column method. RT-qPCR was carried out using 2019 novel coronavirus  
144 nucleic acid detection kit (BioGerm, Shanghai, China) to confirm the presence of SARS-CoV-2 viral  
145 RNA with cycle threshold (Ct) values range from 17 to 37, targeting the high conservative region  
146 (ORF1ab/N gene) in SARS-CoV-2 genome. Metagenomic sequencing: The concentration of RNA  
147 samples was measurement by Qubit RNA HS Assay Kit (Thermo Fisher Scientific, Waltham, MA, USA).  
148 DNase was used to remove host DNA. The remaining RNA was used to construct the single-stranded  
149 circular DNA library with MGIEasy RNA Library preparation reagent set (MGI, Shenzhen, China).  
150 Purified RNA was then fragmented. Using these short fragments as templates, random hexamers  
151 were used to synthesize the first-strand cDNA, followed by the second strand synthesis. Using  
152 the short double-strand DNA, a DNA library was constructed through end repair, adaptor ligation,  
153 and PCR amplification. PCR products were transformed into a single strand circular DNA library  
154 through DNA-denaturation and circularization. DNA nanoballs (DNBs) were generated with the  
155 single-stranded circular DNA library by rolling circle replication (RCR). The DNBs were loaded into  
156 the flow cell and pair-end 100bp sequencing on the DNBSEQ-T7 platform 8 (MGI, Shenzhen, China).  
157 20 genomes were assembled with length from 26,840 to 29,882 nucleotides. The median age of  
158 patients was 36 (range:6-75). Two of twenty patients suffered severe or critical illness. Weifang  
159 sequences were combined with a diverse selection of sequences from China outside of Weifang and  
160 other countries provided by GISAID *Elbe and Buckland-Merrett (2017)*. The new Weifang sequences  
are deposited in GISAID (XXXX-XXXX).

**Mathematical model.** The phylodynamic model is designed to account for nonlinear epidemic dynamics in Weifang, a realistic course of infection (incubation and infectious periods), migration of lineages in and out of Weifang, and variance in transmission rates which can influence epidemic size estimates. The model of epidemic dynamics within Weifang is based on a susceptible-exposed-infectious-recovered (SEIR) model. We elaborate the model with with an additional compartment  $J$  which has a higher transmission rate ( $\tau$ -fold higher) than the  $I$  compartment. Upon leaving the incubation period individuals progress to the  $J$  compartment with probability  $p_h$ , or otherwise to  $I$ . The model is implemented as a system of ordinary differential equations:

$$\dot{S}(t) = -\beta(\beta I(t) + \beta\tau J(t)) \frac{S(t)}{S(t) + I(t) + J(t) + R(t)} \quad (1)$$

$$\dot{E}(t) = \beta(\beta I(t) + \beta\tau J(t)) \frac{S(t)}{S(t) + I(t) + J(t) + R(t)} - \gamma_0 E(t) \quad (2)$$

$$\dot{I}(t) = \gamma_0(1 - p_h)E(t) - \gamma_1 I(t) \quad (3)$$

$$\dot{J}(t) = \gamma_0 p_h E(t) - \gamma_1 J(t) \quad (4)$$

$$\dot{R}(t) = \gamma_1(E(t) + J(t)) \quad (5)$$

We also model an exponentially growing reservoir  $Y(t)$  for imported lineages in to Weifang. The

equation governing this population is

$$\dot{Y}(t) = (\rho - \mu)Y(t). \quad (6)$$

161 Migration is modeled as a bidirectional process which only depends on the size of variables in  
162 the Weifang compartment and thus migration does not influence epidemic dynamics; it will only  
163 influence the inferred probability that a lineage resides within Weifang. For a compartment  $X$  (E, I,  
164 or J),  $\eta$  is the per lineage rate of migration out of Weifang and the total rate of migration in and out  
165 of Weifang is  $\eta X$ .

166 During phylodynamic model fitting  $\beta$  and  $\rho$  are estimated. Additionally, we estimate initial sizes  
167 of  $Y$ ,  $E$ , and  $S$ . Other parameters are fixed based on prior information. We fix  $1/\gamma_0 = 4.1$  days and  
168  $1/\gamma_1 = 3.8$  days. We set  $p_h = 0.20$  and  $\tau = 74$  which yields a dispersion of the reproduction number  
169 that matches a negative binomial distribution with  $k = 0.22$  if  $R_0 = 2$ , similar to values estimated for  
170 the 2003 SARS epidemic (*Lloyd-Smith et al., 2005*).

171 **Phylogenetic analysis.** We aligned the 20 Weifang sequences using MAFFT (*Katoh and Standley,*  
172 *2013*) with a previous alignment of 35 SARS-CoV 2 sequences from outside of Weifang (*Volz et al.,*  
173 *2020*). Maximum likelihood analysis was carried using IQTree (*Minh et al., 2019*) with a HKY+G4  
174 substitution model and a time-scaled tree was estimated using treedater 0.5.0 (*Volz and Frost, 2017*).  
175 Two outliers according to the molecular clock model were identified and removed using 'treedater'  
176 which was also used to compute the root to tip regression.

177 Bayesian phylogenetic analysis was carried out using BEAST 2.6.1 (*Bouckaert et al., 2019*) using a  
178 HKY+G4 substitution model and a strict molecular clock. The phylodynamic model was implemented  
179 using the PhyDyn package (*Volz and Siveroni, 2018*) using the QL likelihood approximation and the  
180 RK ODE solver. The model was fitted by running 8 MCMC chains in parallel and combining chains  
181 after removing 50% burn-in.

182 The *ggtree* package was used for all phylogeny visualizations (*Yu et al., 2017*).

183 Code to replicate this analysis and and BEAST XML files can be found at <https://github.com/emvolz/weifang-sarscov2>.  
184

## 185 Funding

186 This work was supported by Centre funding from the UK Medical Research Council under a con-  
187 cordat with the UK Department for International Development. NIHR. J-IDEA. This work was also  
188 supported by a grant from the Special Project for Prevention and Control of Pneumonia of New  
189 Coronavirus Infection in Weifang Science and Technology Development Plan in 2020 (2020YQFK015)  
190 to Associate Senior Technologist Qing Nie. Role of the Funders: All funders of the study had no role  
191 in study design, data analysis, data interpretation, or writing of the report.

## 192 Acknowledgements

193 We gratefully acknowledge China National GeneBank at Shenzhen, China for the sequencing strategy  
194 and capacity support. We also gratefully acknowledge the laboratories that have contributed  
195 publicly available genomes via GISAID: Shanghai Public Health Clinical Center & School of Public




196 Health, Fudan University, Shanghai, China, at the National Institute for Viral Disease Control and  
197 Prevention, China CDC, Beijing, China, at the Institute of Pathogen Biology, Chinese Academy of  
198 Medical Sciences & Peking Union Medical College, Beijing, China, at the Wuhan Institute of Virology,  
199 Chinese Academy of Sciences, Wuhan, China, at the Department of Microbiology, Zhejiang Provincial  
200 Center for Disease Control and Prevention, Hangzhou, China, at the Guangdong Provincial Center  
201 for Diseases Control and Prevention at the Department of Medical Sciences, at the Shenzhen Key  
202 Laboratory of Pathogen and Immunity, Shenzhen, China, at the Hangzhou Center for Disease and  
203 Control Microbiology Lab, Zhejiang, China, at the National Institute of Health, Nonthaburi, Thailand,  
204 at the National Institute of Infectious Diseases, Tokyo, Japan, at the Korea Centers for Disease  
205 Control & Prevention, Cheongju, Korea, at the National Public Health Laboratory, Singapore, at the  
206 US Centers for Disease Control and Prevention, Atlanta, USA, at the Institut Pasteur, Paris, France,  
207 at the Respiratory Virus Unit, Microbiology Services Colindale, Public Health England, and at the  
208 Department of Virology, University of Helsinki and Helsinki University Hospital, Helsinki, Finland,  
209 and at the University of Melbourne, Peter Doherty Institute for Infection and Immunity, Melbourne,  
210 Australia, at the Victorian Infectious Disease Reference Laboratory, Melbourne, Australia, at the  
211 Public Health Virology Laboratory, Brisbane, Australia and at the Institute of Clinical Pathology and  
212 Medical Research, University of Sydney, Westmead, Australia.

## 213 References


- 214 **Bedford T**, Cryptic transmission of novel coronavirus revealed by genomic epidemiology; 2020. Accessed:  
215 2020-3-8. <https://bedford.io/blog/ncov-cryptic-transmission/>.
- 216 **Bouckaert R**, Vaughan TG, Barido-Sottani J, Duchêne S, Fourment M, Gavryushkina A, Heled J, Jones G, Kühnert  
217 D, De Maio N, et al. BEAST 2.5: An advanced software platform for Bayesian evolutionary analysis. *PLoS*  
218 *computational biology*. 2019; 15(4):e1006650.
- 219 **Dudas G**, Carvalho LM, Rambaut A, Bedford T. MERS-CoV spillover at the camel-human interface. *Elife*. 2018  
220 Apr; 7.
- 221 **Elbe S**, Buckland-Merrett G. Data, disease and diplomacy: GISAID's innovative contribution to global health.  
222 *Global Challenges*. 2017; 1(1):33–46.
- 223 **Gire SK**, Goba A, Andersen KG, Sealfon RSG, Park DJ, Kanneh L, Jalloh S, Momoh M, Fullah M, Dudas G, Wohl S,  
224 Moses LM, Yozwiak NL, Winnicki S, Matranga CB, Malboeuf CM, Qu J, Gladden AD, Schaffner SF, Yang X, et al.  
225 Genomic surveillance elucidates Ebola virus origin and transmission during the 2014 outbreak. *Science*. 2014  
226 Sep; 345(6202):1369–1372.
- 227 **Grantz K**, Metcalf J, Lessler J, Dispersion vs. Control [Internet].[cited 2020 Feb 12];.
- 228 **Guan WJ**, Ni ZY, Hu Y, Liang WH, Ou CQ, He JX, Liu L, Shan H, Lei CL, Hui DSC, Du B, Li LJ, Zeng G, Yuen KY, Chen  
229 RC, Tang CL, Wang T, Chen PY, Xiang J, Li SY, et al. Clinical Characteristics of Coronavirus Disease 2019 in China.  
230 *N Engl J Med*. 2020 Feb; .
- 231 **Katoh K**, Standley DM. MAFFT multiple sequence alignment software version 7: improvements in performance  
232 and usability. *Molecular biology and evolution*. 2013; 30(4):772–780.
- 233 **Keeling MJ**, Rohani P. *Modeling Infectious Diseases in Humans and Animals*. Princeton University Press; 2011.
- 234 **Li LM**, Grassly NC, Fraser C. Quantifying Transmission Heterogeneity Using Both Pathogen Phylogenies and  
235 Incidence Time Series. *Mol Biol Evol*. 2017 Nov; 34(11):2982–2995.




- 236 **Lloyd-Smith JO**, Schreiber SJ, Kopp PE, Getz WM. Superspreading and the effect of individual variation on  
237 disease emergence. *Nature*. 2005 Nov; 438(7066):355–359.
- 238 **Mao H**. Weifang City announces fever clinics. Weifang News Network. 2020 Jan; .
- 239 **Minh BQ**, Schmidt H, Chernomor O, Schrempf D, Woodhams M, von Haeseler A, Lanfear R. IQ-TREE 2: New  
240 models and efficient methods for phylogenetic inference in the genomic era; 2019.
- 241 **Stadler T**, Phylodynamic Analyses based on 11 genomes from the Italian outbreak; 2020. Accessed: 2020-3-8.  
242 <http://virological.org/t/phylogenetic-analyses-based-on-11-genomes-from-the-italian-outbreak/426>.
- 243 **Verity Hill AR**, Phylodynamic analysis of SARS-CoV-2 | Update 2020-03-06; 2020. Accessed: 2020-3-8. <http://virological.org/t/phylogenetic-analysis-of-sars-cov-2-update-2020-03-06/420>.
- 244
- 245 **Volz EM**, Frost SDW. Scalable relaxed clock phylogenetic dating. *Virus Evol*. 2017 Jul; 3(2).
- 246 **Volz E**, Baguelin M, Bhatia S, Boonyasiri A, Cori A, Cucunubá Z, Cuomo-Dannenburg G, Donnelly CA, Dorigatti  
247 I, Fitzjohn R, Fu H, Gaythorpe K, Ghani A, Hamlet A, Hinsley W, Imai N, Laydon D, Nedjati-Gilani L, Gemma  
248 abd Okell, Riley S, van Elsland S, et al. Report 5: Phylogenetic analysis of SARS-CoV-2; 2020.
- 249 **Volz EM**, Siveroni I. Bayesian phylodynamic inference with complex models. *PLoS Comput Biol*. 2018 Nov;  
250 14(11):e1006546.
- 251 **Yu G**, Smith DK, Zhu H, Guan Y, Lam TTY. ggtree : an R package for visualization and annotation of phylogenetic  
252 trees with their covariates and other associated data. *Methods Ecol Evol*. 2017 Jan; 8(1):28–36.
- 253 **Zhou P**, Yang XL, Wang XG, Hu B, Zhang L, Zhang W, Si HR, Zhu Y, Li B, Huang CL, Chen HD, Chen J, Luo Y, Guo H,  
254 Jiang RD, Liu MQ, Chen Y, Shen XR, Wang X, Zheng XS, et al. A pneumonia outbreak associated with a new  
255 coronavirus of probable bat origin. *Nature*. 2020 Feb; .



**Figure 1-Figure supplement 1.** A time scaled phylogeny estimated using IQTree and treedater and using the same data as used for the Bayesian analysis.



**Figure 1-Figure supplement 2.** A tree density plot based on the posterior distribution of trees computed in BEAST2.



**Figure 1-Figure supplement 3.** The estimated posterior TMRCA among all Weifang lineages.