

A single holiday was the turning point of the COVID-19 policy of Israel

Klausner, Ziv PhD^{1 †}; Fattal, Eyal PhD^{1* †}; Hirsch, Eitan PhD²;
Shapira, Shmuel C., MD, MPH³

¹ Department of Applied Mathematics, Israel Institute for Biological Research, Ness-Ziona, Israel

² Environmental Sciences Division, Israel Institute for Biological Research, Ness-Ziona, Israel

³ Director general, Israel Institute for Biological Research, Ness-Ziona, Israel

* Correspondence to: Applied Math department, Israel Institute for Biological Research, P.O. Box 19, Ness-Ziona, 7410001, ISRAEL. Tel.: +972 8 9381794; fax: +972 8 9381432; e-mail: eyalf@iibr.gov.il

† These two authors contributed equally to the article

42 **Abstract**

43 **Background:**

44 The impact of COVID-19 has been profound, and the public health challenge
45 seem to be the most serious regarding respiratory viruses since the 1918 H1N1
46 influenza pandemic. In the absence of effective vaccine or biomedical treatment, the
47 basic rules of public health measures have not changed, namely public distancing.

48 **Methods:**

49 We analyzed epidemiological investigation reports during the first month of
50 the outbreak in Israel. In addition, we present a deterministic compartment model and
51 simulations of several scenarios emphasizing quarantine and isolation policies given
52 their efficiency.

53 **Results:**

54 We identify an abrupt change from controlled epidemic regime to an
55 exponential growth ($R_0 = 2.19$) in light of the actual policy-makers decisions and
56 public behavior in Israel. Our analysis show that before the abrupt change, the new
57 cases trend was due to returning citizens infected abroad. The abrupt change followed
58 a holiday in which social distancing was clearly inefficient and many public
59 gatherings were held. We further discuss three different modeled scenarios of
60 quarantine efficiency: high-, medium-, and low-efficiency.

61 **Conclusions:**

62 Israel early lessons show that there is no allowance to compromise with the
63 directive of social distancing. Even before the onset of the pandemic in Israel, fine-
64 tuned but determined early decisions were taken by policy makers to monitor flight
65 arrivals from Covid-19 affected regions and to limit public gatherings. Our analysis
66 show that one particular holiday has shifted the occurrence curve from controlled

67 regime to exponential growth. Therefore, even a short lapse in public responsiveness
68 can have a dramatic effect.

69

70

71

72 **1. Introduction**

73 Since its emergence the impact of COVID-19 has been profound, and the
74 public health challenge seem to be the most serious seen in a respiratory virus since
75 the 1918 H1N1 influenza pandemic (Soper 1919). In this study we present the results
76 of epidemiological data and modelling of one month since the onset of the outbreak in
77 Isreal, addressing public events occurring during this period and the sensitivity to a
78 number of public health measures focusing on social distancing (quarantine and
79 isolation).

80 The epidemiological data studied, consisting of 381 laboratory confirmed
81 COVID-19 cases, has been obtained from the epidemiological investigation reports
82 that were released by the Israeli Ministry of Health (Israeli Ministry of Health, 2020).
83 This allowed us to identify and separate the incoming from abroad (which policy
84 required to enter quarantine on arrival) to the local cases, thus analyzing the net
85 contribution of local infections.

86 In addition, we present an extended deterministic SEIR (Susceptible, Exposed,
87 Infectious, and Recovered) model to simulate disease outbreak scenarios. In
88 particular, the model includes quarantine of asymptomatic suspected population
89 (exposed) and isolation of symptomatic and infectious patients. The model takes into
90 account the efficiency of the quarantine and isolation measures. We discuss three
91 different quarantine efficiency scenarios: high-efficiency, medium-efficiency, and

92 low-efficiency. The resulting analysis from the epidemiological cases data are
93 discussed in light of public events and compared to model simulations. We analyze
94 and discuss an abrupt change from controlled epidemic regime to an exponential
95 growth regime in light of policy makers' decisions and public behavior.

96

97 **2. Methods**

98 The dynamics of spread of epidemics as well as the quarantine-isolation policy
99 of Israel was modeled using the SEQIJR model (e.g. Gumel, et al., 2004). This is a
100 deterministic compartmental model which allows the implicit inclusion of biological
101 epidemiological phases (including incubation period) as well as governmental
102 interventions such as quarantine and their actual efficiency of implementation. A
103 successful a posteriori implementation of this model to the transmission dynamics and
104 control of the SARS epidemics in Toronto, Hong Kong, Singapore and Beijing is
105 given in Gumel, et al., 2004. The model consists of a system of 7 dynamical equations
106 and 15 parameters. For details of the model and its parameters see the online
107 supplementary information.

108 The data analyzed in this study was obtained from the epidemiological
109 investigation reports that were released by the Israeli Ministry of Health (Israeli
110 Ministry of Health, 2020). From the total of 883 PCR laboratory confirmed COVID-
111 19 cases we analyzed the 384 cases that were investigated epidemiologically,
112 allowing us to separate the incoming from abroad (quarantined on arrival by policy)
113 to local cases. This allowed us to separate the imported cases (travelers arriving from
114 abroad) from the locally infected cases. The data spans over the first month of the
115 COVID-19 outbreak in Israel, beginning in February 21st 2020 and going until March
116 20th. We further note that during the examined period the number of PCR tests

117 performed rose daily, reaching 1869 at the end of the period. However, the proportion
118 of positive tests remained approximately the same, as 7.9% (Israeli Ministry of
119 Health, 2020).

120 The incidence curve was modelled as a fit to an exponential growth function
121 (de Silva, et al., 2009; Zhao, et al., 2020). Several serial interval distributions that
122 were estimated for COVID-19 were examined (Nishiura, et al., 2020; Ganyani, et al.,
123 2020; Zhao, et al., 2020).


124

125 **3. Results**

126 The dynamics of epidemic spread in Israel in terms of daily reported
127 confirmed cases is shown in fig. 1. There are two distinct regimes in the curve. The
128 first ended in March 12, 2020 in which the local to travel associated infection ratio
129 was around 1:3 (up to 1:2.5), i.e., for every three imported cases, there was one local
130 infection. The second period, began a day later. The first distinct sign for the change
131 was evident on March 14th where the ratio rose to 1:2, then continued to rise on March
132 17th to 1:1.5 and finally, after a month into the epidemic in Israel, on March 21st there
133 were more local infections in Israel than imported cases (the ratio was above 1:1).

134 March 12th is two days after the Jewish holiday of Purim, which this year was
135 celebrated on March 10th. Purim customs include wearing masks and costumes and
136 holding public celebrations, parades as well as religious gatherings. Because Purim is
137 also a school holiday, many celebration are held a day or two earlier.

138



139

140 FIG. 1. Accumulated reported confirmed COVID-19 cases in Israel (blue bars)

141 and the ratio of local infections to travel associated infections (orange line)

142

143 The effective reproduction number, R_0 , was estimated for the daily new cases

144 data of the first month of the COVID-19 epidemic. The lowest estimated R_0 was 2.08

145 (95%CI 1.93-2.6) for the Gamma distributed serial interval with mean 4.4 and SD 3

146 days (Zhao, et al., 2020). The highest estimate was 2.37 (95%CI 2.16-2.61) for the

147 Gamma distributed serial interval with mean 5.2 and SD 2.8 days (Ganyani, et al.,

148 2020). The mean R_0 overall the serial interval distributions examined, was 2.19.

149 The SEQJR model solutions are characterized by 3 locally stable equilibrium

150 points in parameters space. These refer to the following regimes or types of dynamics

151 of epidemic spread: controlled (decaying), flattened, uncontrolled (baseline SEIR


152 model; further details can be found in the supplementary material). In this study we

153 simulated three scenarios, the first corresponding to the first equilibrium and two

154 corresponding to the second:

- 155 1. A controlled high-efficiency quarantining (decaying green curve in
156 Fig. 2). This regime is characterized by an early entry of asymptomatic
157 suspected population to home quarantine. Moreover we assume the
158 infectiousness in home quarantine is one sixth compared to free
159 asymptomatic. The efficiency of isolation is 70%.
- 160 2. A flattened for medium-efficiency quarantining (purple curve in Fig.
161 2). This regime is characterized by a late entry of asymptomatic
162 suspected population to home quarantine. Moreover we assume the
163 infectiousness in home quarantine is one third compared to free
164 asymptomatic. The efficiency of isolation is 70%.
- 165 3. A flattened for low-efficiency quarantining (light-blue curve in Fig. 2).
166 This regime is characterized by a late entry of asymptomatic suspected
167 population to home quarantine. Moreover we assume the
168 infectiousness in home quarantine is similar to free asymptomatic. The
169 efficiency of isolation is 30%.

170 As discussed above, the dynamics of epidemic spread in Israel until March 8, 2020
171 corresponds to the controlled regime characterizes with R_0 of the order of half (green
172 curve in Fig. 2). On the other hand, after March 8, 2020 the regime corresponds to the
173 flattened regime with low-efficiency quarantining and $R_0 = 2.19$.



174

175

FIG. 2. Total number of symptomatic cases in Israel from SEQIJR model

176

simulations for 3 scenarios: green – high-efficiency quarantining, purple – medium-

177

efficiency quarantining, light blue – low-efficiency quarantining.

178

179 4. Discussion

180

Israel has begun early its mitigation policy against the COVID-19 epidemic.

181

From the beginning of February, it was decided: to close all border passages (via land,

182

sea and air) to people that are not Israeli citizens or resident that have visited China

183

recently; to stop all direct flights from Israel to China and to require Israeli citizens

184

returning from China or that were in touch with a confirmed COVID-19 patients to a

185

14 days home quarantine. A short while later, the quarantine requirement was

186

expanded to other Asian countries. The first COVID-19 patients in Israel were two

187

passengers that returned from the “Diamond Princess”, in 21 and 23 of February.

188 They entered directly to hospital isolation. A week later, a passenger from Italy was
189 diagnosed as a COVID-19 patient, and Italy was added to the list of countries that
190 require 14 days quarantine.

191 The quarantine-isolation policy succeeded in keeping the rate of daily new
192 cases small, up until March 9th. Then, 4 days later, a sudden change of regime has
193 occurred, which was manifested by the distinct change of the epidemic curve of Israel
194 towards exponential growth. March 15th marks the first time that the daily new cases
195 of locally infected were higher than the new travel associated cases.

196 The timing of this abrupt change is not of coincidence. Regarding the cases
197 arriving from abroad, a requirement for home quarantine affecting all travelers
198 arriving began on March 9th. Moreover, during the period between March 9th to 11th a
199 Jewish holiday, Purim, was celebrated. This holiday is characterized by big parades
200 organized by local municipalities, as well as religious gatherings and privately
201 organized parties. Although authorities cancelled the public parades, many privately
202 organized and religious crowding had occurred. Regrettably, these drove Israel from a
203 controlled, mitigated regime to an exponential growth, as described in the results
204 section. Therefore, despite its intense efforts, Israel's effective R_0 for the period
205 ending in March 20th stands on around 2.19, slightly smaller than the R_0 of 2.6-3.2
206 estimated for the republic of Korea and Italy, for the period ending in March 5th.
207 Such abrupt transition based on social behavior emphasizes the fragility of mitigation
208 policies.

209 We therefore emphasize the importance of early fine-tuned but intense
210 directives for social distancing and isolation measures. This study clearly
211 demonstrates the lesson learned from the Israeli policy, that even a short lapse in

212 public responsiveness can have a dramatic effect on public health during pandemic
213 outbreak.

214

215

216

217 **References**

- 218 1. Israel Ministry of Health. Information on confirmed patients and COVID-19
219 press releases. [https://govextra.gov.il/ministry-of-health/corona/corona-](https://govextra.gov.il/ministry-of-health/corona/corona-virus/spokesman-messages-corona/)
220 [virus/spokesman-messages-corona/](https://govextra.gov.il/ministry-of-health/corona/corona-virus/spokesman-messages-corona/). Accessed Mar 22, 2020.
- 221 2. De Silva UC, Warachit J, Waicharoen S, Chittaganpitch M. 2009. A
222 preliminary analysis of the epimemiology of Influenza A(H1N1)v virus
223 infection in Thailand from early outbreak data. *Eurosurveillance*: **14**(31).
- 224 3. Nishiura H, Linton NM, Akhmetzhanov AR. 2020. Serial interval of novel
225 coronavirus (COVID-19) infections. *International Journal of Infectious*
226 *Diseases* DOI: 10.1016/j.ijid.2020.02.060
- 227 4. Zhao S, Gao D, Zhuang Z, Chong MKC, Cai Y, Ran J, Cao P, Wang K, Lou
228 Y, Wang W, Yang L, He D. 2020. Estimating the serial interval of the novel
229 coronavirus disease (COVID-19): A statistical analysis using the public data in
230 Hong Kong from January 16 to February 15, 2020. preprint is under
231 consideration at *Infectious Diseases of Poverty*. DOI: 10.21203/rs.3.rs-
232 18805/v1
- 233 5. Ganyani T, Kremer C, Dongxuan C, Torneri A, Faes C, Wallinga J, Hens N.
234 Estimating the generation interval for COVID-19 based on symptom onset
235 data. medRxiv doi:10.1101/2020.03.05.20031815

- 236 6. Gumel AB, Ruan S, Day T, Watmough J, Brauer F, van den Driessche P,
237 Gabrielson D, Bowman C, Alexander ME, Ardal S, Wu J, Sahai BM.
238 Modelling strategies for controlling SARS outbreaks. *Proc Biol Sci.* 2004;
239 **271**(1554):2223-32. doi:10.1098/rspb.2004.2800
- 240 7. Soper GA. 1919. The lessons of the Pandemic. *Science* **49**(1274): 501-506.
241 DOI: 10.1126/science.49.1274.501
- 242 8. Zhuang Z, Zhao S, Lin Q, Cao P, Lou Y, Yang L, Yang S, He D, Xiao L.
243 2020. Preliminary estimating the reproduction number of the coronavirus
244 disease (COVID-19) outbreak in Republic of Korea and Italy by 5 March
245 2020. medRxiv preprint DOI: <https://doi.org/10.1101/2020.03.02.20030312>
246