

Reporting the life tracks of confirmed cases can effectively prevent and control the COVID-19 outbreak in China

Jie Zhang^{1†}, Tianjing Wang^{†1}, Jiaqi Wang^{1†}, Jingjing Chen¹, Hongwei Yan^{1*}, Lin Sun^{2*}

¹ Key Laboratory of Spine and Spinal Cord Injury Repair and Regeneration of Ministry of Education, Orthopaedic Department of Tongji Hospital, School of Medicine, Tongji University, Shanghai, China

² Shanghai Mental Health Center, Shanghai Jiao Tong University School of Medicine, Shanghai, China

[†] Contribute equally

*Correspondence to: Hongwei Yan, No. 389 Xincun Road, Shanghai, P. R. China. E-mail: zhenxin8855@126.com; Sun Lin, No. 600 South Wanping Road, Shanghai, China. E-mail: xiaosuan2004@126.com

Abstract

Since late December 2019, a novel coronavirus (COVID-19) has emerged in Wuhan city and rapidly spread throughout China. Fears were raised higher, so effective policies about the emergency prevention and control were concerned.

Till March 3, 91273 confirmed COVID-19 cases were included in. Using Mann-Whitney U test, the provinces that reported the life tracks of confirmed cases to public have lower increased in daily new confirmed cases. Compared with the paired province, Tianjin, Jilin, Gansu, Shanxi, Hainan and Guizhou have significant differences in the number of new confirmed cases and have lower mean rank ($P < 0.05$). Compared with the matched municipalities, Shanghai has not significant difference but has a lower mean rank ($P = 0.175$). Besides this, the success prevention and control work need other effective strategies, such as the appointment and removal of leaders in Hubei province, real time media coverage, isolating the suspected cases or even close contacts, encouraging people to avoid going outside, delaying the return of work, closing schools, wearing facial masks, and disinfecting the communities, et.al, have kept China's epidemic under control.

Strategies in China were adjusted in real time according to the epidemic which made a well prevention and control. They may be suggestions to other counties with an outbreak of COVID-19.

Keywords: COVID-19, epidemic, outbreak, SARS, coronavirus

Introduction

The current outbreak of 2019 novel coronavirus (COVID-19) was originated from Wuhan/Hubei province in central China. Just in merely a month, it fast spread across other provinces or cities(1, 2). After two months, other countries, such as Korean, Japan, Iran and Italy had severe situations. It has become a global health concern(3).

Evidence pointed out there was a strong person-to-person transmission in hospital and family settings(4-8). Despite of lockdown of the Wuhan city since 23 January 2020, the massive human movement during the period of Chinese traditional new year may have fueled the spread of the disease. As we know, general rules for the prevention and control of infectious diseases are to cut off the routes of transmission, to control the source of infection, and to protect vulnerable populations. To date, there is no effective antiviral treatment or vaccine specifically designed for this virus, which is the seventh member of enveloped RNA coronavirus(2). In this situation, some non-pharmaceutical interventions, closing the public gathering places, wearing facial masks and social distancing could slow the spread of the disease. However, just one intervention, may be ineffective for this disease. So, effective public health policies, issued or executed by government, are needed for the disease preventive and control. Based on the experience of SARS (Severe Acute Respiratory Syndrome) prevention and control in 2003(9-12), the Chinese government has carried out a rapid and effective prevention and control(13).

In this study, by collecting the published data from the Health Commission of 23 provinces, 5 autonomous regions, 4 municipalities directly under the Central Government and 2 special administrative regions, we compared the effect of epidemic prevent and control for each paired provinces or cities which were based on the population density of the administrative population and the administrative area.

Materials and Methods

Data collection

Public data on official websites of Health Commission of 23 provinces, 5 autonomous regions, 4 municipalities directly under the Central Government and 2 special administrative regions were retrieved manually. The published date, the number of confirmed COVID-19 cases and the detailed level of data description including symptoms, transmission routes, etc. were recorded. From January 21st to March 3rd, 80303 cases of China were included in. All information above was collected by two persons independently.

Population density

We calculated the population density of the administrative population and the administrative area. The population of administrative regions in various provinces and cities were based on the China Statistical Yearbook 2019 (<http://www.stats.gov.cn/tjsj/ndsj/2019/indexch.htm>), and the area of administrative regions in various provinces and cities were based on the China's second national land survey(14).

Statistical analysis

Time distribution of confirmed cases with the public health policies were compared between different areas in China by line charts. For testing hypothesis that whether the report on life tracks of confirmed cases to public is useful for the epidemic, the number of confirmed cases were compared by Mann-Whitney U test. A pair of provinces or cities were matched if they had similar population densities. A two-tailed $P < 0.05$ was considered as statistically significant. ALL statistical analyses were done using the SPSS software, version 20.0 (IBM Corporation, Armonk,

NY).

Results

Situation reports of COVID-19 in various provinces of China

Till March 3rd, there were 80303 confirmed cases, 2947 deaths and 47281 recurred cases in domestic (Table 1). The top five provinces had more than 1000 cases. Hubei province had 67214 confirmed cases. Guangdong, Henan, Zhejiang, Hunan, and Anhui had 1350, 1272, 1213 and 1018 confirmed cases respectively. The existing confirmed cases in Hubei, Guangdong, Shandong, Sichuan and Zhejiang were 28216, 259, 250, 146 and 130 respectively. Xizang and Qinghai had no existing confirmed cases.

Comparison of COVID-19 in Wuhan city, other cities in Hubei province and other provinces

Figure 1 described the trend of COVID-19 in Wuhan city, other cities in Hubei province and other provinces, and the main strategies in China. On December 1, first case was confirmed in Wuhan city. Since Jan 25th, the confirmed cases were increased in Wuhan city, other cities in Hubei province and other provinces, municipalities and special administrative regions. Wuhan city had the largest confirmed cases every day, then other cities in Hubei province and then other provinces and cities. On Feb. 12th, 15162 cases were confirmed nationwide. 15152 cases were diagnosed by clinical diagnosis criteria, including 1332 cases in Wuhan city. After Feb.20th, there were almost 400 confirmed cases in Wuhan city, no more than 100 cases in Hubei province (except Wuhan city), no more than 10 cases in other provinces.

Impacts of whether reporting the life tracks of confirmed cases in different provinces

There were different levels of situation reports in various provinces in domestic. In 34 units, seven provinces or municipalities directly under the central government had published the life tracks of confirmed cases on their official websites. Tianjin, Jilin, Gansu, Shanxi and Hainan reported the detail life tracks of confirmed since Jan.25th, Jan.22nd, Jan.23rd, Jan.22nd, and Jan 22nd, respectively. These five units had a milder outbreak, and up to March 3rd, the existing cases in the five units were no more than 15. Table 1 showed they ranked behind other provinces. Shanghai and Guizhou reported the detail life tracks of confirmed cases in the middle of the epidemic since Feb.9th and Feb.5th, respectively. Among the provinces that have never published their life tracks, other seven other provinces were matched to the seven provinces with the similar population density respectively. Then we got seven pairs of units, each pair had one unit that published the life tracks of confirmed cases and one unit did not publish such information. Through Mann-Whitney U test, it was found that the difference between the two units of six pairs was statistically significant (Table 2). Compared with the paired province, Tianjin, Jilin, Gansu, Shanxi, Hainan and Guizhou have significant differences in the number of new confirmed cases and have lower mean rank ($P < 0.05$). Compared with the matched municipalities, Shanghai has not significant difference but has a lower mean rank ($P = 0.175$).

Discussion

In this highly mobile world, the outbreak of COVID-19 is a challenging task to China. In particular, there is a strong person to person transmission(5) and the treatment and prevention options are limited. To control the novel pathogen could be a challenging task for China. Based on the prevention and control for MERS (Middle East Respiratory Syndrome)(15, 16) or SARS (Severe Acute Respiratory Syndrome)(9-12, 17-20), Chinese government had issued some effective policies or proposed some precautions in case detection and management, vulnerable population protection, and transmission routes cutting off. To date, these measures are very

effective to prevent and control the spread of COVID-19.

An outbreak requires attention in advance to clarity, collaboration, communication, coordination and capacity(21). During the outbreak of SARS, policies were issued by Chinese government. Such as media reporting in real time, built Xiaotangshan hospital for confirmed disease, and published the life tracks of confirmed cases, which may make people were more alert. The spread of SARS in 2003, also brought to light substantial weaknesses in the country's public-health system. After the SARS epidemic was brought under control, the government increased its commitment and leadership to tackle public-health problems and, among other efforts, increased public-health funding, revised laws that concerned the control of infectious diseases, implemented the world's largest internet-based disease reporting system, and started a program to rebuild local public-health facilities(22).

For the outbreak of epidemic in the whole country, the prevention and control system should be changed timely. Different levels of cases should be treated differently. Case detection and management, vulnerable population protection, and transmission routes cutting off are needed to decrease the rate of infection. The monitoring of the epidemic by Health Commission is necessary. Real time media are needed to attract the people's attention. In this study, we found the provinces that reported the life tracks for confirmed cases had mild epidemic and less confirmed cases (Table 2). Compared with the paired province, Tianjin, Jilin, Gansu, Shanxi, Hainan and Guizhou have significant differences in the number of new confirmed cases and have lower mean rank ($P<0.05$). Compared with the matched municipalities, Shanghai has not significant difference but has a lower mean rank ($P=0.175$). There may be two reasons for this result. Firstly, the population densities were the best matched, but there was a big gap between Beijing and Shanghai. Secondly, the reports on the life tracks for confirmed cases in Shanghai began later not at the beginning.

From our opinion, there were two advantages for the epidemic prevention and control for COVID-19. Firstly, in this period, most people stay at home. They always to read the daily media report, and talk about the number of confirmed cases in Wechat or other media social platforms. They can be alert if they find the confirmed case living near their house. Or they may avoid to go the places where the confirmed cases stayed by. So, for the public, reporting the life tracks of confirmed cases is an effective way to help them keep away from the 'infectious' things or the places. This may good alert for protect the public. Secondly, for tracking the detail information of life tracks would need more time and work for some of government administration departments. Whether the local governments are willing to put in the effort and time in it, to some extent, may showed how meticulous and important the local governments were in their work.

Although the report for the life tracks of confirmed cases had some controversial, it was proved that this kind of information was effective and feasible in our country. To date, except Wuhan city and other cities in Hubei province, we found the number of new COVID-19 cases increased slowly, even in some provinces, there were no new cases increased these days. Nine provinces had no more than 10 cases, and Xizang and Qinghai provinces had no confirmed case now. Even Wuhan city had a decreased number of confirmed cases every day. The information showed a controlled trend for this epidemic outbreak.

The epidemic was under control in China. Still, there were some limitations in our study. Firstly, the outbreak in China is not over yet. There may be another peak of infection rate with the resumption of work. Secondly, Wuhan city still had a severe situation now. Anyway, the number of daily confirmed cases were dropped sharply which showed the effective of epidemic prevention

and control.

Conclusion

COVID-19, with a strong person-to-person transmission made a pandemic in China and outbreak in Korean, Iran, Italy and Japan. Along with the decreased number of new confirmed cases and the increased number of recovered cases, there might be a turning corner of COVID-19 in China. This can be attributed to the success of a serial of emergence public measures and the confidence of the whole nation, which may be useful for the epidemic prevention and control in other countries. However, it remains a challenging task to improve the mechanism for major epidemic prevention and control and the national public health emergency management system.

Author contributions

JW and TW analyzed the data, JZ wrote the manuscript. JW, TW, JZ, LS, HY and JC collected the data, LS and JZ designed the research and monitored the quality of the analysis.

Conflict of interest

The authors declare that they have no conflict of interest.

References

1. Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet*. 2020;395(10223):497-506. doi:10.1016/S0140-6736(20)30183-5
2. Zhu N, Zhang D, Wang W, et al. A Novel Coronavirus from Patients with Pneumonia in China, 2019. *N Engl J Med*. 2020;382(8):727-33. doi:10.1056/NEJMoa2001017
3. Habibzadeh P, Stoneman EK. The Novel Coronavirus: A Bird's Eye View. *Int J Occup Environ Med*. 2020;11(2):65-71. doi:10.15171/ijoem.2020.1921
4. Chan JF, Yuan S, Kok KH, et al. A familial cluster of pneumonia associated with the 2019 novel coronavirus indicating person-to-person transmission: a study of a family cluster. *Lancet*. 2020;395(10223):514-23. doi:10.1016/S0140-6736(20)30154-9
5. Li Q, Guan X, Wu P, et al. Early Transmission Dynamics in Wuhan, China, of Novel Coronavirus-Infected Pneumonia. *N Engl J Med*. 2020. doi:10.1056/NEJMoa2001316
6. Phan LT, Nguyen TV, Luong QC, et al. Importation and Human-to-Human Transmission of a Novel Coronavirus in Vietnam. *N Engl J Med*. 2020. doi:10.1056/NEJMc2001272
7. Rothe C, Schunk M, Sothmann P, et al. Transmission of 2019-nCoV Infection from an Asymptomatic Contact in Germany. *N Engl J Med*. 2020. doi:10.1056/NEJMc2001468
8. Wu JT, Leung K, Leung GM. Nowcasting and forecasting the potential domestic and international spread of the 2019-nCoV outbreak originating in Wuhan, China: a modelling study. *Lancet*. 2020. doi:10.1016/S0140-6736(20)30260-9
9. From the Centers for Disease Control and Prevention. Updated interim surveillance case definition for severe acute respiratory syndrome (SARS)--United States, April 29, 2003. *JAMA*. 2003;289(20):2637-9. doi:10.1001/jama.289.20.2637-b
10. Chinese Medical A, China Preventive Medicine A, Chinese Association of Medical A. [Summary of the corss-strait seminar on prevention and control of SARS]. *Zhonghua Yi Xue Za Zhi*. 2003;83(9):708-12.
11. Liang TH. [The epidemiological findings and the experience in the prevention and control of SARS in the Hong Kong Special Administrative Region]. *Zhonghua Yi Xue Za Zhi*. 2003;83(11):902-5.
12. Wang LD. [Rethinking about the prevalence of SARS, recent work on SARS prevention and control]. *Zhonghua Yi Xue Za Zhi*. 2003;83(20):1753-6.
13. Wu Z, McGoogan JM. Characteristics of and Important Lessons From the Coronavirus Disease

- 2019 (COVID-19) Outbreak in China: Summary of a Report of 72314 Cases From the Chinese Center for Disease Control and Prevention. *JAMA*. 2020. doi:10.1001/jama.2020.2648
14. Land Resources and Utilization in China[M]. Beijing:Geological Publishing House. 2019:127.
 15. Al-Tawfiq JA, Memish ZA. Infection control measures for the prevention of MERS coronavirus transmission in healthcare settings. *Expert Rev Anti Infect Ther*. 2016;14(3):281-3. doi:10.1586/14787210.2016.1135053
 16. Madani TA, Althaqafi AO, Alraddadi BM. Infection prevention and control guidelines for patients with Middle East Respiratory Syndrome Coronavirus (MERS-CoV) infection. *Saudi Med J*. 2014;35(8):897-913.
 17. From the Centers for Disease Control and Prevention. Severe Acute Respiratory Syndrome (SARS) and coronavirus testing--United States, 2003. *JAMA*. 2003;289(17):2203-6. doi:10.1001/jama.289.17.2203
 18. Ammon A. From SARS to Ebola--10 years of disease prevention and control at ECDC. *Euro Surveill*. 2015;20(37). doi:10.2807/1560-7917.ES.2015.20.37.30016
 19. Vogel LN, Roberts A, Paddock CD, et al. Utility of the aged BALB/c mouse model to demonstrate prevention and control strategies for severe acute respiratory syndrome coronavirus (SARS-CoV). *Vaccine*. 2007;25(12):2173-9. doi:10.1016/j.vaccine.2006.11.055
 20. Yen MY, Lu YC, Huang PH, Chen CM, Chen YC, Lin YE. Quantitative evaluation of infection control models in the prevention of nosocomial transmission of SARS virus to healthcare workers: implication to nosocomial viral infection control for healthcare workers. *Scand J Infect Dis*. 2010;42(6-7):510-5. doi:10.3109/00365540903582400
 21. Naylor CD, Chantler C, Griffiths S. Learning from SARS in Hong Kong and Toronto. *JAMA*. 2004;291(20):2483-7. doi:10.1001/jama.291.20.2483
 22. Wang L, Liu J, Chin DP. Progress in tuberculosis control and the evolving public-health system in China. *Lancet*. 2007;369(9562):691-6. doi:10.1016/S0140-6736(07)60316-X

TABLES AND FIGURES

Table 1. Existing cases, confirmed cases, deaths and recurred cases of COVID-19 acute respiratory disease reported by provinces, regions and cities in China, Data as 03 March 2020.

Abbreviations: COVID-19: a novel coronavirus outbreak in Wuhan city in 2019

Table 2. Comparison between pair of regions that published the life tracks of confirmed cases or not based on Mann-Whitney U Test

Figure 1. The comparison of COVID-19 confirmed cases in Wuhan city, other cities in Hubei province and other provinces.


Abbreviations: COVID-19: a novel coronavirus outbreak in Wuhan city in 2019

Table 1. Existing cases, confirmed cases, deaths and recurred cases of COVID-19 acute respiratory disease reported by provinces, regions and cities in China, Data as 03 March 2020

Area	Existing cases	Confirmed cases	Deaths	Recurred cases
Hubei	28179	67217	2835	36203
Wuhan	24111	49426	2252	23063
Other cities	4068	17791	583	13140
Guangdong	242	1350	7	1101
Henan	19	1272	22	1231
Zhejiang	120	1213	1	1092
Hunan	113	1018	4	901
Anhui	56	990	6	928
Jiangxi	64	935	1	870
Shandong	241	758	6	511
Jiangsu	75	631	0	556
Chongqing	101	576	6	469
Sichuan	141	538	3	394
Heilongjiang	106	480	13	361
Beijing	118	414	8	288
Shanghai	41	338	3	294
Hebei	12	318	6	300
Fujian	36	296	1	259
Guangxi	48	252	2	202
Shaanxi	28	245	1	216
Yunnan	3	174	2	169
Hainan	12	168	5	151
Guizhou	30	146	2	114
Tianjin	15	136	3	118
Shanxi	13	133	0	120
Liaoning	20	125	1	104
Hong Kong SAR	62	100	2	36
Jilin	9	93	1	83
Gansu	4	91	2	85
Xinjiang	5	76	3	68
Inner Mongolia	15	75	1	59
Ningxia	5	74	0	69
Taipei and environs	29	42	1	12
Qinghai	0	18	0	18
Macao SAR	2	10	0	8
Xizang	0	1	0	1
Total	29964	80303	2948	47391

Table 2. Comparison between pair of regions that published the life tracks of confirmed cases or not based on Mann-Whitney U Test

Regions		Matched regions		<i>P</i>	Mean Rank	
Name	Population density (person/square kilometer)	Name	Population density (person/square kilometer)		Reported regions	Unreported regions
Tianjin	1309.05	Beijing	1312.93	0.002	33.23	49.77
Jilin	141.42	Sichuan	171.58	<0.001	29.20	53.80
Gansu	61.92	Heilongjiang	83.37	<0.001	30.96	52.04
Shanxi	237.27	Guangxi	207.32	0.032	35.94	47.06
Hainan	265.51	Jiangxi	278.43	0.001	33.26	49.74
Shanghai	2899.87	Beijing	1312.93	0.175	37.95	45.05
Guizhou	204.43	Guangxi	207.32	0.026	35.76	47.24



On December 1, 2019, the first case was confirmed in Wuhan.

① On January 23, 2020, Wuhan, Hubei Province was sealed at 10 am.

② Since January 24, 2020, medical teams from various provinces and cities across the country have been mobilized to carry out treatment in Hubei Province.

③ On January 25, 2020, the Party Central Committee set up a leading group for the response to Covid-19 to establish a centralized, unified and efficient leading chain of command and accurately resolve the frontline problems of the epidemic.

④ Since February 2, 2020, Wuhan, Hubei Province has focused on the centralized admission and quarantine of "four types of personnel", including unconditional centralized admission of diagnosed patients, centralized quarantine and treatment of suspected patients, and centralized quarantine and observation of febrile patients and close contacts.

⑤ On February 3 and February 5, 2020, the Huoshenshan Hospital and the Leishenshan Hospital were completed and put into use.

⑥ On the evening of February 7, 2020, Wuhan Fang Cang Hospital formally receive patients and gradually entered an institutionalized and standardized operation.

⑦ On February 13, 2020, the main leadership of Hubei Provincial Party Committeeed wuhan Municipal Party Committee were adjusted.

⑧ On February 16th, a three-day centralized investigation was conducted to concentrate on fighting the annihilation, grasp actual situation, and resolutely curb the spread of the epidemic. All villages, communities, estates, and residential areas in urban and rural areas implemented the most stringent closed-off management for 24 hours.