

COVID-19 pandemic in the African continent: forecasts of cumulative cases, new infections, and mortality

Short Title: Africa and the COVID-19 pandemic

Authors: Kinfu, Y (PHD, MPHIL)^{1,2*} ; Alam, U (PHD, MPH)³ ; Achoki, T (MD, PHD)^{4*}.

Tom Achoki: tachoki@sloan.mit.edu

Uzma Alam: uzma.alam@aya.yale.edu

Yohannes Kinfu: yohannes.kinfu@canberra.edu

Corresponding Authors (*):

1. Faculty of Health, University of Canberra, Australia
2. College of Medicine, Qatar University, Qatar
3. Africa Institute for Health Policy Foundation, Kenya
4. School of Health Systems and Public Health, University of Pretoria, South Africa

COVID-19 pandemic in the African continent: forecasts of cumulative cases, new infections, and mortality

ABSTRACT

Background: Africa is the last major region to capitulate to the SARS-CoV-2 (COVID-19) pandemic. The first confirmed COVID-19 case in the region was reported on February 14, but what lies ahead in terms of the course and magnitude of infection remains speculative. To the best of our knowledge, no study, using a robust methodology, provides the immediate and long-term trajectory of COVID-19 for the entire region or accounts for its local context. This paper is the first systematic attempt to provide estimates on how many people would contract the virus and how many would die in the coming few months across Africa.

Methods: The forecasts on caseloads and incidences are from a co-variate-based instrumental variable regression model. Fatality rates from Italy and China were further applied to generate mortality estimates after adjustments were made for differences in age-structure, health service quality, and living standards between each of the African countries and those of the reference population. We cover all countries that reported a confirmed case as of March 31, 2020.

Results: By the end of June, 16,283,085 people will contract COVID-19 (95% CI 718,403 to 98,358,799). With a cumulative caseload of 5,413,4517 (95% CI 1,332,953 to 8,489,940) and 906,625 (95% CI 173,821 to 4,742,917) Northern and Eastern Africa will respectively be the most and least affected sub-regions in the continent. Cumulative COVID-19 cases on June 30, 2020 are expected to reach 2,912,864 (95% CI 465,028 to 18,286,358) in Southern Africa, 2,787,913 (95% CI 517,489 to 15,056,314) in Western Africa, and 1,185,742 (95% CI 229,111 to 6,138,692) in Central Africa. New infections (incidence) for the month of April are expected to be the highest in Djibouti, 32.8 per 1000 (95% CI 6.25 to 171.77), while Morocco 1045 (95% CI 167 to 6,547) will register the highest number of deaths.

Conclusion: Our study shows that countries that are least urbanized and have a low level of socio-economic development, hence least connected to the outside world, are likely to register lower and slower transmissions, at least at the early stage of the epidemic. However, the same set of enabling factors that worked for their benefit are likely to go against them in implementing interventions that have lessened the impact of the disease elsewhere.

INTRODUCTION

The novel human coronavirus (SARS-CoV-2 or COVID-19) outbreak initially emerged in Wuhan, China, in late 2019. Since then, it has spread to 198 countries and territories around the world and has been declared a pandemic (1). As of March 31, 2020, there have been reported 750,874 covid-19 cases, 36,045 deaths, and 117,603 recoveries (1). The first confirmed COVID-19 case in Africa was reported in Egypt on February 14, 2020 and since then the number of confirmed infections in the region kept increasing and has surpassed over 5000 cases by March 31, 2020 (2, 3). Comoros, Lesotho, Malawi, and South Sudan are the only African countries that had not reported a confirmed case by March 31, 2020.

To date, no vaccine is available for COVID-19. Therefore, the ability to limit the devastating consequences of the disease on people's lives and livelihoods would have to rely on the implementation of effective preventative non-pharmaceutical interventions (NPIs). NPIs are multiple public health measures designed to reduce viral transmission rates in a population by reducing the reproduction number (R_0); the average number of secondary cases each case generates (4-6).

NPIs directly influence the course of the COVID-19 pandemic, including the rate of spread, and the expected duration of the pandemic. However, several characteristics of the virus such as incubation period (the time between infection and symptom onset), serial interval (the time between symptom onset of a primary and secondary case) the extent of asymptomatic cases, the possibility of pre-symptomatic infectiousness, and the case fatality rate (CFR), and also the possible role of weather in transmission, all remain ambiguous or mostly unknown (7, 8). Estimates for COVID-19 CFR range from 0.3-1% (9). Asymptomatic or mild presentation comprises the bulk of the reported cases, which is an estimated at 80% (10). Longitudinal viremia measurements from a small study (sample size of 16), suggests that there are high enough viral loads to trigger pre-symptomatic infectiousness for 1-2 days before the onset of symptoms (11). With the lack of clinical studies measuring viremia, the infectious period also remains largely unknown, with estimates ranging from few days to 10 days or more after the incubation period (11).

The complexity of the infection and recovery process, therefore, means that proper understanding of the epidemiological dynamics of COVID-19 within the local context is fundamental to combat the pandemic. Specifically, studies illustrating future trajectories of the disease can help nations and communities to develop early warning systems, avoid overwhelming healthcare services, minimize morbidity and mortality from the disease, evaluate the effects of interventions and the long-term consequences of the virus on peoples' livelihood. This is particularly true in Africa, where livelihoods are fragile, and previous epidemics, such as HIV/AIDS and, more recently, Ebola, has been known to exert enormous socioeconomic consequences (12-14). In addition, in the face of a new pandemic in the region, the already overstretched healthcare systems that are struggling to deliver essential healthcare services such as immunization and HIV/AIDS treatment would be in further jeopardy and risk losing the gains achieved so far in disease control efforts.

This underscores the need for decision-makers to be equipped with actionable information that brings visibility to contemporary and future epidemiological trends to guide interventions. This study, using a robust methodology, provides spatial and temporal trajectories of COVID-19 for the entire Africa region. To the best of our knowledge, this is the first such attempt and accounts for the local context and characteristics relevant to the epidemiology of the region.

The remaining part of this paper is divided into three sections. In section 2, we describe our covariate based predictive model and the input data that has been used in the modeling exercise. We also describe the methodology used for estimating deaths from COVID-19. Section 3 provides the results on the projected new infections, cumulated cases, and deaths due to COVID-19 across countries, covering the five sub regions. Finally, we discuss the key findings and implications of the study in section 4 and provide recommendations for action.

METHODS

The analysis is based on a linear instrumental variable (IV) regression framework. The basic model consists of two equations which can be expressed as follows (15-17):

$$Y = \delta + \beta_1 K + \beta_2 X + u_i$$

$$K = \theta + \alpha_1 T + z_i \dots \dots \dots \text{Equation (1)}$$

Here, Y is the outcome variable of interest, and in our case represents the number of confirmed cases of COVID-2019 as of March 31, 2020. K is the rate of infection between week 1 and week 2 and is an endogenous variable. T and X represent a set of instrumental and exogenous variables, respectively. u_i and z_i are error terms and both are assumed to have a zero mean and a nonzero-correlation (18-20).

Based on our understanding of the epidemiology of COVID-19 and the scanty literature, we have identified eleven regressors that are deemed associated with the trajectory of COVID-19 in the Africa region (4, 6). These are urbanicity, socio-demographic index, average household size, the age structure of the population, health care access, health quality index, adherence to international health regulations, and prevalence of HIV and asthma. In addition, we have controlled for time since the first reported case and the rate of expansion of the disease between week one and week two of the epidemic in the respective countries. This was modeled using air traffic as its instrument variable.

The data for analysis were assembled from various sources. The data on prevalence of HIV and asthma, as well as on socio-demographic index and the index of health care access and quality index, were from (21-23). The data on urbanization were obtained from the Population Reference Bureau (24), while average household size, age profile, and data on the total population used for the calculation of rates were all from the UN Population Division (25). The index of adherence to international health regulations was accessed from the WHO, while the number of confirmed cases was from the John Hopkins database of COVID-19 (26, 27).

The 11 variables selected for analyses were chosen out of an initial set of over 20 variables that included the proportion of households with at least one member aged 60 years or over, the proportion of households with at least one member aged 65 years or over, the World Bank's governance index, as well as the prevalence of TB and diabetes. STATA's Lasso software for model selection was used to narrow down the list of covariates (28). Furthermore, out of the 193 countries captured in our data, for a few countries, data were unavailable for selected variables, namely air traffic (25 countries), adherence to international health regulation (20 countries), and household size (24 countries). To

bridge the gap, we performed multiple imputation procedures, particularly by running 1000 imputations and 99 000 iterations using STATA's multiple imputation routine (29, 30). The resulting median values from the imputation exercise were subsequently used to estimate the prediction model specified earlier. We transformed all the regressors as well as the dependent variables to a log-scale before they were put into the model.

The goodness-of-fit for the predictive model was evaluated using k-fold cross-validation techniques (31, 32). The advantage of a k-fold cross-validation technique over other statistical validation procedures is that all observations are used for both training and validation, and each observation is used for validation exactly once (31).

Estimation of COVID-19 deaths was made using indirect standardization technique, with additional adjustments for differences in health system readiness and living standards. In the absence of age-specific infection data the proposed approach can be expressed in a general form as follows:

$$a_{yk} = [p_y \int w_{iy} c_{ik}] / b_{yk} \dots \dots \dots (Equation 2a)$$

Where,

$$b_{yk} = \sqrt[2]{S_y Q_y} / \sqrt[2]{S_k Q_k} \dots \dots \dots (Equation 2b)$$

Here, a_{yk} is the estimated number of deaths for country y ($y= 1, 2 \dots 47$) using country k's (China and Italy) COVID-19 fatality rate, general health system readiness and socio-economic standards as a reference. c_{ik} is the standard or reference age-specific fatality rate for age group i in standard country k, in our case obtained from two sources, China and Italy. w_{iy} is the population at age i in country y; p_y is population level COVID-19 prevalence rate and is an estimate we generated using the instrumental variable regression expressed in equation 1. The quantity in the square bracket represents age-standardized estimated number of deaths in country y, if age-specific fatality rates were to follow those of the standard populations. However, our study population differs from the two countries not just with respect to age structure but also living standard and health system readiness. Hence, to adjust the estimated deaths further we developed a correction factor that links the socio-economic status and health system readiness of each country with those two populations, which is represented by b_{yk} . In this case, b_{yk} is a geometric- mean of socio-demographic index (S) and health access and quality index (Q), and the relevant data for each country was sourced from IHME (21, 22). Analysis was performed using STATA version 16.0 (33).

RESULTS

Figure A1 in the Appendix shows a graphical comparison of observed and predicated COVID-19 confirmed cases (in log scale). Table A1 presents results from the cross-fold robustness test for the predictive model. The statistical diagnostic tests as well the patterns from the graphical analysis suggest that the predicative model sufficiently captures the observed data. As can be seen from the table, not only the RMSE, the MAE and the Pseudo- R^2 are consistent across samples, the error margins are also small. Hence, we have used the resulting model to forecast cumulative COVID-19 cases, estimate new infections and deaths from COVID-19 for each of the 47 African countries which have reported at least one case as of March 31, 2020.

Cumulative COVID-19 cases and point prevalence rates

Table 1 presents cumulative COVID-19 cases and prevalence rates for Africa and its five sub-regions for the period April 1 to June 30 2020. By the end of April cumulated infection across the region will reach 2,459,328 cases (CI 95% 424,155 to 14,380,080), and will further increase to 16, 283, 084 (CI 95% 2,718,402 to 98, 358, 798) by June 30, 2020. During the same period, cumulative cases in the hardest hit sub-region in the African continent, Northern Africa, is predicted to increase from 1.3 million at the end of April (CI 95% 207982 to 7,914,479) to over 7.5 million by June 30, 2020. This is followed by Southern, Western, and Central Africa sub-regions. Throughout the forecast period, Eastern Africa is predicated to experience the lowest case load and prevalence rate in the continent throughout the forecast period.

Table 1: Cumulative cases and prevalence rates for COVID-19, Africa: April 1 – June 30, 2020.

Panel A: Cumulated number of infected people at end of reported month; Forecast, April 1 – June 30

Sub-region	March 31st	Point estimate			Lower bound (95% C.I.)			Upper bound (95% C.I.)		
	Reported	April	May	June	April	May	June	April	May	June
Central	155	179089	599906	1185742	35748	117282	229111	897478	3069555	6138692
Eastern	375	136933	458692	906625	27122	88980	173821	693416	2371620	4742917
Northern	2454	1282284	4295345	8489940	207982	682342	1332953	7914479	27069102	54134517
Southern	1385	439947	1473716	2912864	72559	238049	465028	2673470	9143801	18286358
Western	1259	421075	1410499	2787913	80744	264904	517489	2201237	7528670	15056314
Africa	5628	2459328	8238158	16283084	424155	1391557	2718402	14380080	49182748	98358798

Panel B: cumulative infection rate per 1000 mid-population in each month: Forecast, April 1 – June 30, 2020

Sub-region	Point estimate			Lower bound (95% C.I.)			Upper bound (95% C.I.)		
	April	May	June	April	May	June	April	May	June
Central	0.147	0.491	0.968	0.029	0.095	0.185	0.744	2.54	5.068
Eastern	0.312	1.041	2.056	0.06	0.196	0.383	1.621	5.532	11.05
Northern	0.678	2.269	4.48	0.111	0.362	0.707	4.164	14.226	28.422
Southern	0.731	2.446	4.829	0.133	0.435	0.85	4.035	13.782	27.526
Western	0.114	0.383	0.755	0.022	0.073	0.143	0.586	1.999	3.99
Africa	0.306	1.025	2.023	0.056	0.183	0.357	1.694	5.784	11.553

Figure 1A shows spatial patterns of cumulative COVID-19 cases by June 30, 2020 (see also Appendix 2A for country estimates). In Northern Africa, the leading contributor to the burden of COVID-19 is Morocco. By the end of June 2020, Morocco will have 4,459,877 (CI 95% 689,547 to 28,845,720) cumulative cases of COVID-19 and this is almost double the estimated number for Algeria, a country with the next highest burden, 2,804,674 (CI 95% 438,629 to 17,935, 144) by the end of June 2020. In Southern Africa, South Africa and Swaziland are the leading contributors to the pandemic. By the end of June 2020, the countries are expected to have 2,581,366 (CI 95% 404,835 to 16,459, 651) and 254,403 (CI 95% 45,730 to 1,415,274) cumulative cases, respectively. In the Western Africa sub-region, cumulative cases of infection are dominated by Cote d'Ivoire and Ghana, despite Nigeria having a larger population than both countries combined.

To remove the effect of size, we also estimated prevalence rates shown in Panel B of Table 1 by combining the forecasts on cumulative cases with population estimates which we generated for each month from the UN annual population estimates. This suggests that population level prevalence of COVID-19 in Africa is expected to remain under 5 percent (CI 95% 0.6 to 15.9) throughout the prediction period, but there are wide inter-regional differences. In the Northern and Southern Africa sub-regions, cumulative infection rates are expected to be over 15 percent, with the rate in the respective subregions expected to hit as high as 34 percent (CI 95% 0.53 to 21.36) and 19 percent (CI 95% 0.34 to 10.13) by the end of June 30, 2020.




Figure 1A




Figure 1B




Figure 1C

Figure 1 (a) Cumulated COVID cases June 30, 2020; (b) COVID-19 incidence per 1000 population April 2020, and (c) Number of COVID-19 deaths April 2020

New infections and COVID-19 incidence rates across Africa

Across the months, the number of new COVID-19 infections are expected to increase from 2,453,700 (CI 95% 418,527 to 14,374,452) cases in April, to 5778830 (CI 95% 967402 to 34,802,668) cases in May, to 8,044,927 (CI 95% 1,326,846 to 49,176,051) cases by the end of July. This represents a 135% increase from April to May and a 39% increase from May to June, which suggests a possible slowing down of the pandemic with time.

Table 2: Number of new infections and incidence rates for COVID-19, Africa: April 1 – June 30, 2020.

Panel A: Number of new infection at end of reported month; Forecast, April 1 – June 30									
Sub-region	Point estimate			Lower bound (95% C.I)			Upper bound (95% C.I)		
	April	May	June	April	May	June	April	May	June
Central	178934	420817	585835	35593	81534	111829	897323	2172077	3069137
Eastern	136558	321759	447933	26747	61858	84842	693041	1678204	2371297
Northern	1279830	3013060	4194595	205528	474360	650611	7912025	19154623	27065415
Southern	438562	1033769	1439149	71174	165490	226979	2672085	6470331	9142557
Western	419816	989424	1377415	79485	184160	252585	2199978	5327433	7527645
Africa	2453700	5778830	8044927	418527	967402	1326846	14374452	34802668	49176051

Panel B: New infection rate per 1000 mid-population in each month: Forecast, April 1 – June 30, 2020									
Sub-region	Point estimate			Lower bound (95% C.I)			Upper bound (95% C.I)		
	April	May	June	April	May	June	April	May	June
Central	1.465	3.443	4.783	0.287	0.661	0.904	7.439	17.971	25.338
Eastern	3.112	7.306	10.159	0.596	1.364	1.869	16.203	39.143	55.246
Northern	6.768	15.916	22.135	1.093	2.518	3.45	41.624	100.668	142.099
Southern	7.305	17.16	23.857	1.322	3.027	4.147	40.342	97.523	137.619
Western	1.14	2.683	3.729	0.22	0.511	0.699	5.852	14.144	19.948
Africa	3.059	7.190	9.996	0.553	1.272	1.742	16.933	40.931	57.760

Sub-regional and country level differences are expected to deepen as the pandemic becomes more established in the region. In the coming three months incidence is expected to increase faster in Southern Africa, followed by Northern and Eastern sub-region. In each sub-region, we expect some countries to be affected more than others and become hotbeds of new infections (see Figure 2B). In places like Djibouti new infection rate is expected to reach as high as 32.81 (CI 95% 6.25 to 171.77) cases per 1000 population, followed by Swaziland 26.81 (CI 95% 4.97 to 144.39), Morocco 11.97 (CI 95% 1.98 to 71.59), Algeria 9.80 (CI 95% 1.57 to 60.72) and Cote d'Ivoire 6.65 (CI 95% 1.07 to 29.68) (see Appendix Table 3A and Figure 1B).

Estimated COVID-19 mortality

Deaths for COVID-19 are estimated for each country, using indirect standardisation techniques after making relevant adjustments to the fatality rates from China and Italy as described in the methods section. As can be seen in Figure 2, the adjustment shows relatively higher expected mortality using Italy as a standard compared to China, but generally are closer to each other. Hence, our final country level estimates presented in Appendix Table A4, Figure 1C and the regional summaries presented in

Table 3, were generated using the arithmetic mean of death numbers generated using the two standard populations. The estimates are for the month of April 2020.

Figure 2: Comparison of estimated number of deaths using China and Italy as standard, April 2020.




Table 3: Estimated number of COVID-19 deaths, April- June 2020

	Point estimate			Lower bound (95% C.I)			Upper bound (95% C.I)		
	April	May	June	April	May	June	April	May	June
Central	248	828	1632	49	162	316	1241	4234	8445
Eastern	141	473	933	28	92	180	712	2430	4851
Northern	1836	6144	12131	297	1032	2014	11354	38792	77497
Southern	468	1566	3092	77	262	512	2856	9757	19494
Western	445	1489	2936	86	278	543	2319	7915	15797
Africa	3138	10499	20723	537	1827	3564	18482	63128	126084

The first COVID-19 case in Africa was reported on February 14,2020 (2, 3). Our estimates suggest that at the end of April, after two months since the first COVID-19 cases were reported in the region, the number of deaths is expected to reach over 3 thousand deaths (CI 95% 537 to 18482). Most deaths in the region will come from the Northern Africa sub-region, which, as reported earlier, also had the largest number of COVID-19 cases and the highest prevalence rates of all sub-regions in the content. Southern and Western Africa are expected to experience comparable death toll, while deaths in the Eastern Africa sub-region are expected to be the lowest of the sub-regions in Africa. Some countries such as Algeria, Morocco and South Africa are expected to experience the highest number of casualties since the beginning of the pandemic, in excess of 400 deaths. Deaths in the three countries will comprise of almost two-thirds of all expected deaths in the region. In others, such as the Democratic Republic of Congo, Tunisia and Cote d'Ivoire death toll will reach between 150 and 300,

and contribute 20 % of the regional total, while half of the continent (about 29 countries) will experience less than 10 deaths each. The same trend continues in May and June, but with much higher number of expected deaths.

DISCUSSION

The COVID-19 pandemic has been spreading rapidly across different parts of the world and the African continent has not been spared. Given the global trajectory, what lies ahead in terms of the course and magnitude of infection in Africa in the days and months to come remains speculative. While there have been attempts elsewhere in capturing the trajectory of the COVID-19 pandemic using statistical models, this has not been the case for the African region as whole (4, 6, 8). To the best of our knowledge, no study, using a robust methodology, provides estimates on future trend of COVID-19 for the entire region or accounts for its local context.

Modeling and predicting the epidemiology and trajectory of a disease such as COVID-19 is a challenging exercise. Firstly, COVID-19 is a newly identified pathogen and knowledge about the characteristics of the virus, including how the disease is spread and the time from exposure to onset of symptoms are still evolving. Secondly, at the time of writing, less than a handful of countries had reached a plateau in their infection rate, and this makes modeling future trends an open-ended exercise (1). The fact that some people only experience mild symptoms and that even the best health system can only detect and treat those presenting to facilities also means that the available data on 'confirmed' cases represents only a fraction of the true picture of the pandemic.

Thirdly, literature on the association between disease prevalence and population level characteristics that are known to be the staple of social and descriptive epidemiology, and the backbone of predictive modeling exercises are yet to emerge for COVID-19 (4-6, 8). Further, at the individual level, knowledge on the determinants of COVID-19 morbidity and mortality is limited to selected characteristics such as age and pre-existing conditions, and even then, they are drawn from scanty data. The association with community and population level characteristics is even more limited, hence largely speculative. In this context, the task of developing a multi-country covariate-based predictive model for COVID-19 in any country or region of the world, let alone Africa, is likely to be guided by what is feasible, and to the extent, the available data permits it. It also means that there is a need for constant revision of predictive models as new data becomes available. In our case, the work is further complicated because in Africa data on key covariates are either lacking or when they exist tend to be biased or derived from other global covariate-based modeling exercises (21, 22).

Despite this, we have made efforts to mitigate these limitations, by curating data from various credible sources around the world and assessing them for consistence. In developing our model, we have used data sources from 193 countries globally and used statistical relationships to fill data gaps on covariates in the Africa region. Additionally, our model performance was assessed through rigorous out of sample calibration using k-fold cross-validation techniques and obtaining robust results. We have also restricted our forecast to the first few months (April-June) and refrained from long term exercise. This is because first we anticipate new and additional data that is more informative and will lead to better and improved estimates. Second, in the face of sparse data any long-term projection is more likely to be detached from reality and easily become a wild guess, and less useful to policy actions. Third, if the countries of the region will not have been able to put effective strategies to contain the disease before the end of June, the last month of our projection period, the extent and consequence of COVID-19 on the continent would be more than any predictive model can anticipate.

Within these caveats, our results show a heterogeneous picture of projected new infections, cumulative cases, and deaths across the African continent. Our study substantially contributes to the growing literature on understanding the trajectory of the COVID-19 pandemic, in addition to conveying an African perspective. This perspective is particularly important at a time of controlling a pandemic given that Africa is not only the second largest continent but also home to 17% of the total world population, with the current population at 1,332,805,306 (34).

Among our key findings are that African countries that are relatively more urbanized and have a higher socio-demographic index (SDI) are projected to experience a faster growth of the epidemic, at least during the initial periods. To illustrate this trend, Algeria, Morocco, and Tunisia, in the Northern African sub-region, which have a predominantly urban population, and are in close proximity to the high burden European countries, are projected to experience a higher incidence of COVID-19 infections in comparison to their neighbors. In the Southern Africa sub-region, South Africa, which has a proportionately higher urban population and greater international connectivity, is also projected to lead the rest of the pack in terms of new infections rates. In the Eastern Africa sub-region, Kenya, which is relatively more urbanized has a higher per capita incidence rate in comparison to other countries within the sub-region. However, it is to be noted that some of these countries also have higher living standards and better health system than most countries in the region. Hence the high prevalence rates and estimated deaths in these countries may in part reflect their greater ability to detect and confirm COVID-19 cases.

On the other hand, countries such as Angola, Botswana and Mozambique, with a lower international connectivity are projected to have lower rates of new infections at the initial periods. Similar trends are observed in Burundi and Tanzania in the Eastern Africa region. However, within a short time, as the epidemic grows many of these countries are projected to have an increased caseload of cumulative infections that are likely to overwhelm their health system capacity. Overall, these trends are consistent with other regions of the world, where since the onset of this pandemic, it has been clear that higher connectivity, particularly through air traffic, and higher population densities are key drivers in the introduction and growth of the epidemic (35-39).

The impact of population density is clearly illustrated by the projected new infections of COVID-19 in places like Djibouti and Rwanda. To compound this, close to 90% of urban settlements in Africa are comprised of informal settlements, most of which are overcrowded and lack basic amenities such as clean water and sanitation that are critical prevention measures for COVID-19 (40-42). Moreover, 60% of jobs in urban areas are in the informal economy (43), and if African countries were to implement stringent lockdowns these livelihoods would be largely affected and could lead to significant socioeconomic consequences. For instance, those that lose their jobs are likely to move back to their homes in rural areas, where many of the elderly population reside, putting them at risk. Similarly, as the economy slows down in several countries due to the COVID-19 pandemic, some sections of migrant workers from African countries might be forced to return to their home countries further complicating the dynamics of the pandemic.

Therefore, large scale measures that are aimed at limiting population movement, across countries as well as increasing the social distance among populations, are not enough, or at best may prove to be impracticable to address the pandemic in the African context. Comprehensive response measures should be contextualized and seek to address some of the underlying individual and structural factors that are likely to complicate the epidemic within these environments. It is crucial to balance interventions geared towards preventing the spread of the epidemic with the need to maintaining

livelihoods and social cohesion. Measures such as appropriate messaging, provision of adequate water and sanitation subsidies, food as well as targeted restriction of movement (e.g. from urban to rural) would go a long way to mitigate the spread.

Still, even with all these NPIs measures, a number of countries with weak healthcare delivery systems are expected to have a significant number of infections and fatalities. Therefore, the already overstretched healthcare systems in Africa need to anticipate and prepare to handle an increased number of patients as a consequence of the pandemic, on top of the other common disease conditions that are prevalent in the region. Unfortunately, this is a lesson Africa has already had to learn from the 2014 Ebola outbreak in West Africa. It is estimated there were an additional 11,00 deaths from malaria, HIV/AIDS and TB across Sierra-Leone, Guinea and Liberia (44), 60% decrease in the number of children treated for diarrhea and acute respiratory infections (45) and 38% increase in maternal mortality in Guinea and another 111% in Liberia (46, 47).

Understanding the factors that accelerate and those that mitigate the spread and mortality related to COVID-19, while accounting for local realities is fundamental for sound public health measures to tackle the pandemic. In fact, lessons from HIV/AIDS programming have taught us that highly effective health interventions fail if the local context is not recognized. Therefore, decision makers in Africa must realize that one size does not fit all and could lead to disastrous outcomes.

Effective detection and response to epidemics is premised on decision makers having the right epidemiological information. Unfortunately, many countries in Africa have weak health information systems that are not able to collect and process data rapidly to facilitate timely and targeted action. For example, Lesotho, despite being in the same region and with closer proximity to South Africa and Swaziland, countries that are at the epicentre of the COVID-19 pandemic in the Southern Africa sub-region, is yet to register a confirmed case. This is a clear example of the gap in diagnostic capacity and health information system than of the absence of infection in the country. Hence, the fight to stop the pandemic in Africa should be focused on both strengthening clinical capacity as well reporting systems with the health care system. Innovations around collecting data through digital platforms make it easier to combine different data streams, thereby helping decision-makers gain a fuller understanding of epidemics and the impact of the interventions (48-52). With close to 747 million SIM connections in Sub-Saharan Africa, representing 74% of the population on the continent (53), the response to COVID-19 pandemic provides an opportunity for countries to retool their data collection systems to meet their surveillance needs.

In addition, the mobile digital technologies could enable confidential self-reporting of symptoms to healthcare providers' catalyzing further investigation and epidemic control measures before spread beyond defined population subgroups. As an information and health communication tool, mobile platforms could also be leveraged to deliver appropriate health messages to the population relieving direct consultation pressure from health system providers

Efficient resource allocation and use is only feasible where there is improved visibility across the various components of the health system through high quality health information systems (54). Ideally, this should cover key areas such as human resources for health, infrastructure and equipment, medicines and other health commodities, and financial resources available. This is particularly needed in epidemic situations, where there are often huge distortions in demand for resources, that if not well addressed could lead to wastage and suboptimal population health outcomes.

Lastly, but equally important in the African context, the global health community and governments addressing the COVID-19 pandemic should be inclusive in their efforts because the efficacy of a response to any emergency is only as good as its weakest link. It is, therefore, important that governments include marginalized communities, such as refugees, migrants and internally displaced populations (IDPs) in their policies and actions. Across Africa, approximately 12.3 million people remain forcibly displaced, including 8.1 million internally displaced people and 4.2 million refugees (55). Moreover, Sub-Saharan Africa alone hosts more than 26% of the world's refugee population (56, 57). These populations are likely to face a different set of constraints that restricts their ability to prevent themselves from getting infected and accessing health care. The fact that they also live in under difficult circumstances means that governments across the region should maintain critical supply corridors for humanitarian assistance and address the needs of the most vulnerable within their borders. In the longer term this pandemic should also be a call for African governments to further strengthen engagement with international humanitarian actors in order to foster strong partnership building on local capacities.

CONCLUSIONS

It is true that some of the most successful responses to global health threats including recent ones like Ebola and HIV/AIDS have been characterized by multi-stakeholder partnerships. African health systems have been in the forefront in the implementation of such programs and would be best served to leverage that experience and resources in responding to COVID-19. Bringing all stakeholders together to ensure effective coordination, pooling of resources and delivery of evidence-based interventions is imperative for any sustainable response to such pandemic situations.

Within, the African context policy makers should also consider (a) impact of strategies that could potentially deepen health inequalities and (b) continuously use data driven approaches to identifying their unique most at risk/ vulnerable groups e.g. IDP, HIV positive, food insecure to support them in an equitable manner and bring them into the COVID-19 prevention framework. At a time of a pandemic no one community can be marginalized.

Finally, in as much as health systems are dealing with an emergence situation, measurement should not be relegated to a nice to have, but as a fundamental tool to guide strategic response actions. The measurement framework should be agile yet comprehensive, rather than being piecemeal. It should seek to combine data from multiple sources within the healthcare system into a joint assessment framework such that a consistent and informative narrative emerges to tackle the epidemic. The age-old call for accurate health metrics to address population health still remains valid. Our best estimates presented in the paper are forecast, and their ability to capture reality is as good as the underlying data. It is thus imperative to review the model and the results as new data becomes available, and update the estimates accordingly.

CONFLICTS OF INTEREST

Yohannes Kinfu (YK), Uzma Alam (UA) and Tom Achoki (TA) report no conflicts of interest.

AUTHORS CONTRIBUTIONS

YK conceptualised the study, designed methods and approach; undertook statistical analysis and drafted the manuscript. UA synthesised the literature, contributed to conceptualisation and drafted the manuscript. TA accessed data, contributed to methods and design, undertook the statistical analysis and drafted the manuscript. All authors contributed to the discussion, read and approved the final draft.

ACKNOWLEDGMENTS

This study was finalised when YK was in a government designated quarantine centre for COVID-19 at Sydney InterContinental Hotel, upon his return to Australia. YK would like to acknowledge the Commonwealth Government of Australia and the New South Wales Government for supporting his stay at the hotel free of charge and creating an environment that enabled him to continue with his research and teaching activities, while doing their part to combat COVID-19.


REFERENCES

1. WHO [WHO COVID-19 data](#) accessed April 3 2020
2. African CDC <https://africacdc.org/covid-19/> accessed April 2 2020
3. WHO EMRO <http://www.emro.who.int/index.html> accessed April 6 2020
4. Ferguson N.M, Laydon. D, Nedjati-Gilani. G, Imai.N, Ainslie.K, Baguelin.M, Bhatia.S, Boonyasiri.A et.al., Impact of non-pharmaceutical interventions (NPIs) to reduce COVID-19 mortality and healthcare demand. London: Imperial College COVID-19 Response Team, March 16 (2020). [Imperial College](#) accessed March 23 2020
5. Tian. H, Liu. Y, Wu.C, Chen. B, Kraemer. M et.al, Early evaluation of transmission control measures in response to the 2019 novel coronavirus outbreak in China. <https://www.medrxiv.org/content/10.1101/2020.01.30.20019844v4>
6. Murray. C. JL, IHME COVID-19 health service utilization forecasting team. Forecasting COVID-19 impact on hospital bed-days, ICU-days, ventilator-days and deaths by US state in the next 4 months medRxiv 2020.03.27.20043752; <https://doi.org/10.1101/2020.03.27.20043752>
7. WHO. Coronavirus disease (COVID2019) situation report-30. https://www.who.int/docs/defaultsource/coronaviruses/situation-reports/20200219sitrep30-covid19.pdf?sfvrsn=3346b04f_2 accessed March 31 2020
8. Martinez-Alvarez.M, Jarde.A, Usuf.E, Brotherton.H, Bittaye.M, Samateh Al, Antonio. M et.al., COVID-19 pandemic in West Africa Lancet Glob Health 2020 ^[L]_{SEP} April 1, 2020
9. Anderson RM, Heesterbeek H, Klinkenberg D, Hollingsworth TD. How will country- based mitigation measures influence the course of the COVID-19 epidemic? Lancet. Comment| Volume 395, ISSUE 10228, P931-934, March 21, 2020
10. European Centre for Disease Prevention and Control. Daily risk assessment on COVID19 <https://www.ecdc.europa.eu/en/cases-2019-ncov-eueea> accessed April 2 2020
11. Zou L Ruan F, Huang M, Liang L, Huang H, Hong Z, Yu J, et.al., SARSCoV2 viral load in upper respiratory specimens of infected patients. N Engl J Med 2020; 382:1177-1179
12. Kinfu. Y, Opiyo.C, & Wamukoya, M. (2014). Child Health and Mortality in sub-Saharan Africa: Trends, causes, and forecasts. In C. O. Odimegwu, & J. Kekovole (Eds.), *Continuity and change in sub-saharan African Demography* (Vol. 17, pp. 60-77). (Routledge African Studies; Vol. 17). New York, USA: Routledge <https://doi.org/10.4324/9781315879444>
13. Kastelic. K, Himelein. T, Mauro.T, Abubakarr. T,. The socio-economic impacts of Ebola in Sierra Leone : results from a high frequency cell phone survey (round three) (English). Washington, D.C: World Bank Group. <http://documents.worldbank.org/curated/en/873321467999676330/The-socio-economic-impacts-of-Ebola-in-Sierra-Leone-results-from-a-high-frequency-cell-phone-survey-round-three> accessed March 22 2020
14. Wooldridge. J 2010 Econometric Analysis of cross section and panel data. 2nd ed. Cambridge, MA: MIT press
15. Cameron A.C and Trivedi P.K. 2010 Micro econometrics using Stata. Rev ed College Station, TX: Stata Press
16. Davidson R and Mackinnon J.G. 1993 Estimation and inference in econometrics New York: Oxford University press
17. Martens. E.P, Pestman. W.R, De Boer. A, Belitser. S.V, & Klungel O.H. (2006). Instrumental Variables: Application and Limitations. Epidemiology, 17(3), 260-267

18. Cameron A.C and Trivedi P.K. 2005 Microeconometrics methods and applications New York: Cambridge University press
19. Greene. WH 2018 Econometric Analysis 8th ed. New York: Pearson
20. Global Burden of Disease Collaborative Network. Global Burden of Disease Study 2015 (GBD 2015) Socio-Demographic Index (SDI) 1980–2015. Seattle, United States: Institute for Health Metrics and Evaluation (IHME), 2016.
21. GBD 2016 Healthcare Access and Quality Collaborators. Measuring performance on the Healthcare Access and Quality Index for 195 countries and territories and selected subnational locations: a systematic analysis from the Global Burden of Disease Study 2016. *Lancet*. 2018 Jun 2;391(10136):2236-2271.
22. Global health data exchange <http://ghdx.healthdata.org/> accessed March 24 2020
23. Population reference bureau <https://www.prb.org/data/> accessed March 24 2020
24. UN Population division <https://www.un.org> accessed March 25 2020
25. WHO Health Statistics <https://www.who.int/healthinfo/statistics/en/> accessed March 22 2020
26. Johns Hopkins Coronavirus resource center <https://coronavirus.jhu.edu/data> accessed March 31 2020
27. Lasso Reference Manual, Release 16 College Station, TX: Stata Press.
28. Rubin. D.B Multiple imputation for non-response in surveys 1987. New York: Wiley
29. Schafer. J.L Analysis on incomplete multivariate data. 1997. Boca Raton FL. Chapman &Hall/ CRC
30. Hastie T, Tibshirani R, Friedman J. The elements of statistical learning: data mining, inference, and prediction. 2nd ed. New York, NY: Springer; 2009. 219-260
31. Rao C, Adair T, Kinfu Y. Using historical vital statistics to predict the distribution of under-five mortality by cause. *Clin Med Res*. 2011 Jun; 9(2):66-74.
32. Stata Corp. 2019. Stata Statistical Software: Release 16. College Station, TX: StataCorp LLC
33. UN Population <https://www.un.org/en/sections/issues-depth/population/index.html> accessed April 1 2020
34. Chinazzi. M, Davis. JT, Ajelli. M, Gioannini. C, Litvinova. M, Merler. S, pastore. PY, et.al., The effect of travel restrictions on the spread of the 2019 novel coronavirus (COVID-19) outbreak. *Science*. 2020 Mar 6
35. Wu.T, Leung. K, Leung.G.M, Now casting and forecasting the potential domestic and international spread of the 2019-nCoV outbreak originating in Wuhan, China: A modeling study. *Lancet* 395, 689–697 (2020).
36. Du. Z, Wang.L, Cauchemez. S, Xu. X, Wang. X, Cowling. B.J, Meyers. L.A, Risk for Transportation of 2019 Novel Coronavirus Disease from Wuhan to Other Cities in China. *Emerg Infect Dis*. 2020 May 17;26(5).
37. Kucharski AJ, Russell TW, Diamond C, Liu Y, Edmunds J, Funk S, Eggo RM et.al, Early dynamics of transmission and control of COVID-19: a mathematical modelling study. *Lancet Infect Dis*. 2020 Mar 11.
38. Gilbert.M, Pullano.G, Pinotti.F, Valdano. E, Poletto.C, Boëlle. P, D’Ortenzio.E et.al., Preparedness and vulnerability of African countries against importations of COVID-19: a modelling study *Lancet* 2020; 395: 871–77
39. Simiyu. S, Cairncross. S & Swilling. M. Understanding Living Conditions and Deprivation in Informal Settlements of Kisumu, Kenya. *Urban Forum* 30, 223–241 (2019).

40. Wakhungu. J, Huhhins. C.D, Lumumba. J 2010 Approaches to informal urban settlements in Africa: experiences from Kigali and Nairobi. Technical report. African Centre for Technology Studies (ACTS)
41. Güneralp. B, Lwasa. S, Masundire. S, Parnell. S, and Seto. K.c Environmental Research Letters, Volume 13, Number 1 Published 20 December 2017
42. AEO 2016 Africa Economic Outlook: Sustainable Cities and Structural Transformation 389 Report
43. Parpia. AS, Ndeffo-Mbah. ML, Wenzel. NS, Galvani. AP. Effects of Response to 2014–2015 Ebola Outbreak on Deaths from Malaria, HIV/AIDS, and Tuberculosis, West Africa. *Emerg Infect Dis.* 2016;22(3):433-441.
44. O’Fallon. B, Barry. MA, Brodish. P, Hazerjian. J. Rapid Assessment of Ebola-Related Implications for Reproductive, Maternal, Newborn and Child Health Service Delivery and Utilization in Guinea. *PLoS Curr.* 2015 Aug 4;7.
45. Mullan. Z The Cost of Ebola *Lancet Global Health.* Editorial Volume 3, Issue 8. Aug 2015.
46. Elston. JW, Cartwright. C, Ndumbi. P, Wright. J. The health impact of the 2014–15 Ebola outbreak. *Public Health* 2017; 143: 60–70.
47. Mekuria. LA, Rinke de Wit. TF, Spieker. N, Koech. R, Nyarango. R, et al., Analyzing data from the digital healthcare exchange platform for surveillance of antibiotic prescriptions in primary care in urban Kenya: A mixed-methods study. *Plos one* 14(11) 2019.
48. Guo. X, Chen. S, Zhang. X, Ju. X, Wang. X Exploring Patients' Intentions for Continuous Usage of mHealth Services: Elaboration-Likelihood Perspective Study *JMIR Mhealth Uhealth* 2020;8(4):e17258
49. Lee. Jm, Newman. MW, Lee. Jm, Gebremariam. A, Choi. P, Lewis. D, Nordgren. W, Costik. J, et.al., Real-World Use and Self-Reported Health Outcomes of a Patient-Designed Do-it Yourself Mobile Technology System for Diabetes: Lessons for Mobile Health. *Diabetes Technol Ther.* 2017 Apr;19(4):209-219.
50. Cahn. A, Akiroy. A, Raz. I Digital health technology and diabetes management. *J Diabetes.* 2018 Jan;10(1) 10-17
51. Achoki T. Health Technology in Emerging Global Epidemics: An African Policy Perspective. *Georgetown Journal of International Affairs.* Published 2016 July 22. Available at: <https://www.georgetownjournalofinternationalaffairs.org/online-edition/health-technology-in-emerging-global-epidemics-an-african-policy-perspective>
52. GMSA report <https://www.gsma.com/mobileeconomy/sub-saharan-africa/> accessed April 1 2020
53. Achoki. T, Hovels. A, Masiye. F, Lesego. A, Leufkens. H, Kinfu. Y. Technical and scale efficiency in the delivery of child health services in Zambia: results from data envelopment analysis. *BMJ Open.* 2017;7(1):e012321.
54. UNHCR Global trend <https://www.unhcr.org/globaltrends2018/> accessed March 31
55. World Economic Forum <https://www.weforum.org/agenda/2019/09/newly-released-data-show-refugee-numbers-at-record-levels/> accessed April 1 2020
56. OCHA Southern Africa Humanitarian Snapshot (April 2020)

Appendix Figure A1: Observed and predicted COVID-19 cases, March 31, 2020



Appendix Table A1: Predication model performance from K-fold cross-validation test

Robustness parameter	Sub-sample number								
	1	2	3	4	5	6	7	8	9
RMSE	1.476	1.457	1.483	1.449	1.483	1.484	1.485	1.487	1.485
MAE	1.114	1.141	1.128	1.137	1.159	1.139	1.146	1.134	1.143
Pseudo-R ²	0.718	0.721	0.717	0.719	0.730	0.719	0.728	0.728	0.726

Note : RMSE, Root Mean Square Error; MAE, Mean Absolute Error.

Appendix Table: A2, Forecast cumulative COVID-19 cases, June 30, 2020

Sub-region	country	Forecast	UI (95 %)	
			Lower	Upper
Central Africa	Angola	15719	3113	79367
Central Africa	Democratic Republic of the Congo	1072457	207209	5550741
Central Africa	Central African Republic	3124	802	12165
Central Africa	Congo	19568	3907	98020
Central Africa	Gabon	48074	8876	260386
Central Africa	Equatorial Guinea	26799	5204	138013
Eastern Africa	Burundi	107	28	409
Eastern Africa	Djibouti	216573	40078	1170401
Eastern Africa	Eritrea	10792	2375	49046
Eastern Africa	Ethiopia	37254	7804	177835
Eastern Africa	Kenya	210417	39944	1108419
Eastern Africa	Madagascar	18784	4128	85470
Eastern Africa	Mozambique	3569	799	15938
Eastern Africa	Rwanda	71937	14145	365853
Eastern Africa	Somalia	19459	4698	80596
Eastern Africa	Tanzania	13489	2756	66032
Eastern Africa	Uganda	11522	2424	54766
Eastern Africa	Zambia	292723	54642	1568153
Northern Africa	Algeria	2804674	438629	17935144
Northern Africa	Egypt	239030	43521	1312839
Northern Africa	Libya	7324	1376	38982
Northern Africa	Morocco	4459877	689547	28845720
Northern Africa	Sudan	35498	6813	184966
Northern Africa	Tunisia	943537	153067	5816867
Southern Africa	Botswana	899	174	4659
Southern Africa	Namibia	49889	9126	272724
Southern Africa	Swaziland	254403	45730	1415274
Southern Africa	South Africa	2581366	404835	16459651
Southern Africa	Zimbabwe	26308	5163	134051
Western Africa	Burkina Faso	182704	37583	888197
Western Africa	Benin	39620	8162	192316
Western Africa	Cote d'Ivoire	973685	179511	5281373
Western Africa	Cameroon	473097	86418	2589963
Western Africa	Cape Verde	10371	1999	53821
Western Africa	Ghana	554402	97775	3143553
Western Africa	The Gambia	9607	1983	46540
Western Africa	Guinea	59094	12044	289954
Western Africa	Guinea-Bissau	639	155	2645
Western Africa	Liberia	6242	1389	28056
Western Africa	Mali	1314	294	5864
Western Africa	Mauritania	17283	3675	81293
Western Africa	Niger	10013	2388	41979
Western Africa	Nigeria	177914	32202	982974
Western Africa	Sierra Leone	51	12	215
Western Africa	Senegal	244680	45907	1304131
Western Africa	Chad	11501	2583	51196
Western Africa	Togo	15696	3410	72245

Appendix Table: A3, Forecast of new infections COVID per 1000, April, 2020

Sub-region	country	Forecast	UI (95 %)	
			Lower	Upper
Central Africa	Angola	0.07	0.01	0.36
Central Africa	Democratic Republic of the Congo	1.82	0.36	9.14
Central Africa	Central African Republic	0.10	0.02	0.36
Central Africa	Congo	0.52	0.10	2.53
Central Africa	Gabon	3.39	0.64	17.78
Central Africa	Equatorial Guinea	2.89	0.57	14.47
Eastern Africa	Burundi	0.00	0.00	0.00
Eastern Africa	Djibouti	32.81	6.25	171.77
Eastern Africa	Eritrea	0.30	0.07	1.32
Eastern Africa	Ethiopia	0.05	0.01	0.23
Eastern Africa	Kenya	0.60	0.12	3.05
Eastern Africa	Madagascar	0.10	0.02	0.45
Eastern Africa	Mozambique	0.02	0.00	0.07
Eastern Africa	Rwanda	0.83	0.16	4.10
Eastern Africa	Somalia	0.18	0.05	0.74
Eastern Africa	Tanzania	0.03	0.01	0.15
Eastern Africa	Uganda	0.04	0.01	0.17
Eastern Africa	Zambia	2.38	0.46	12.36
Northern Africa	Algeria	9.80	1.57	60.72
Northern Africa	Egypt	0.35	0.06	1.87
Northern Africa	Libya	0.17	0.03	0.86
Northern Africa	Morocco	18.21	2.89	114.08
Northern Africa	Sudan	0.12	0.02	0.62
Northern Africa	Tunisia	11.97	1.98	71.59
Southern Africa	Botswana	0.05	0.01	0.28
Southern Africa	Namibia	2.80	0.53	14.86
Southern Africa	Swaziland	26.81	4.97	144.39
Southern Africa	South Africa	6.64	1.06	41.07
Southern Africa	Zimbabwe	0.23	0.05	1.11
Western Africa	Burkina Faso	1.32	0.27	6.24
Western Africa	Benin	0.50	0.11	2.33
Western Africa	Cote d'Ivoire	5.65	1.07	29.68
Western Africa	Cameroon	2.76	0.52	14.67
Western Africa	Cape Verde	2.76	0.54	13.92
Western Africa	Ghana	2.73	0.49	15.03
Western Africa	The Gambia	0.64	0.13	2.99
Western Africa	Guinea	0.65	0.14	3.10
Western Africa	Guinea-Bissau	0.04	0.01	0.19
Western Africa	Liberia	0.19	0.04	0.81
Western Africa	Mali	0.01	0.00	0.04
Western Africa	Mauritania	0.55	0.12	2.50
Western Africa	Niger	0.06	0.01	0.26
Western Africa	Nigeria	0.13	0.02	0.70
Western Africa	Sierra Leone	0.00	0.00	0.00
Western Africa	Senegal	2.15	0.41	11.15
Western Africa	Chad	0.11	0.02	0.46
Western Africa	Togo	0.28	0.06	1.26

Appendix Table: A4, Forecast number of deaths from COVID, April 30, 2020

Sub-region	country	Forecast	UI (95 %)	
			Lower	Upper
Central Africa	Angola	2	0	9
Central Africa	Democratic Republic of the Congo	234	47	1170
Central Africa	Central African Republic	1	0	3
Central Africa	Congo	2	1	12
Central Africa	Gabon	7	1	35
Central Africa	Equatorial Guinea	2	0	12
Eastern Africa	Burundi	0	0	0
Eastern Africa	Djibouti	46	9	240
Eastern Africa	Eritrea	2	0	8
Eastern Africa	Ethiopia	8	2	38
Eastern Africa	Kenya	26	5	131
Eastern Africa	Madagascar	3	1	15
Eastern Africa	Mozambique	1	0	3
Eastern Africa	Rwanda	11	2	55
Eastern Africa	Somalia	6	2	25
Eastern Africa	Tanzania	2	0	9
Eastern Africa	Uganda	1	0	6
Eastern Africa	Zambia	35	7	181
Northern Africa	Algeria	540	87	3343
Northern Africa	Egypt	37	7	197
Northern Africa	Libya	1	0	5
Northern Africa	Morocco	1046	167	6547
Northern Africa	Sudan	5	1	27
Northern Africa	Tunisia	207	35	1233
Southern Africa	Botswana	0	0	1
Southern Africa	Namibia	6	1	33
Southern Africa	Swaziland	29	5	155
Southern Africa	South Africa	430	70	2652
Southern Africa	Zimbabwe	3	1	15
Western Africa	Burkina Faso	33	7	157
Western Africa	Benin	7	2	35
Western Africa	Cote d'Ivoire	169	32	885
Western Africa	Cameroon	67	13	354
Western Africa	Cape Verde	2	0	9
Western Africa	Ghana	79	14	436
Western Africa	The Gambia	2	0	7
Western Africa	Guinea	11	2	54
Western Africa	Guinea-Bissau	0	0	1
Western Africa	Liberia	1	0	6
Western Africa	Mali	0	0	1
Western Africa	Mauritania	3	1	12
Western Africa	Niger	2	1	10
Western Africa	Nigeria	21	4	114
Western Africa	Sierra Leone	0	0	0
Western Africa	Senegal	43	8	220
Western Africa	Chad	2	0	9
Western Africa	Togo	3	1	11