

1 **Title: Characteristics of lymphocyte subsets and their predicting values for the**
2 **severity of COVID-19 patients**

3 Short title: Lymphocyte subsets in COVID-19 patients

4

5 Jingrong Wang^{1, #}, Xingqi Dong^{2, #}, Boting Zhang², Xinping Yang², Zhi Li¹, Xicheng
6 Wang^{2, *}, Shuguang Zuo^{3, *}

7

8 ¹Department of Critical Care, First People's Hospital of Yunnan Province, Kunming,
9 Yunnan Province, 650032, China

10 ²Department of Infectious Diseases, Yunnan Provincial Hospital of Infectious
11 Diseases, Kunming, Yunnan Province, 650301, China

12 ³Kaifeng Kangbien Medical Laboratory, Kaifeng, Henan Province, 475000, China

13

14 [#]These authors contributed equally to this article.

15 ^{*}Correspondences: wxch62597@foxmail.com(X.WANG);

16 zuosg@icloud.com(S.ZUO);

17 **Abstract**

18 Severe COVID-19 patients showed worse clinical outcomes compared to mild and
19 moderate patients. However, effective indicators are still lacking to predict the
20 severity of the disease. In the present study, we retrospectively analyzed the clinical
21 and laboratory data of 16 COVID-19 patients and found that the absolute counts of
22 three T-cells (CD3⁺, CD4⁺, and CD8⁺) were significantly lower in the moderate and
23 severe patients than those in mild patients and were significantly lower in severe
24 patients than in moderate patients on admission. With the recovery of the COVID-19,
25 serum levels of inflammatory biomarkers (CRP, PCT, and IL6) of moderate and
26 severe patients gradually decreased. In contrast, the counts of lymphocytes and their
27 subsets including CD3⁺, CD4⁺, and CD8⁺ T cells gradually increased in severe
28 patients, and eventually showed comparable levels with moderate patients. ROC
29 analysis showed that the counts of CD3⁺, CD4⁺, and CD8⁺ T-cells with AUC > 0.9
30 have potential values for predicting the severity of COVID-19 patients. In conclusion,
31 the reduction of CD3⁺, CD4⁺, and CD8⁺ T-cells is related to the severity of
32 COVID-19 and dynamic detection of these T-lymphocyte subsets may help predict
33 the outcome of the patients.

34 **Keywords:** SARS-CoV-2; COVID-19; Pneumonia; Lymphocytes

1 **Introduction**

2 In December 2019, a novel coronavirus (provisionally called 2019-nCoV) infected
3 pneumonia occurred in Wuhan, Hubei Province, China[1]. Phylogenic analysis
4 indicated that 2019-nCoV is a positive-chain RNA enveloped beta genus coronavirus,
5 and closely related to Severe Acute Respiratory Syndrome Coronavirus (SARS-CoV)
6 and Middle East respiratory syndrome Coronavirus (MERS-CoV)[2, 3]. The disease
7 caused by 2019-nCoV was recently named by the World Health Organization (WHO)
8 as Coronavirus Disease 2019 (COVID-19)[4]. Shortly after WHO announced the
9 official name of the disease, 2019-nCoV is officially named as Severe Acute
10 Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) by the International Committee
11 on Taxonomy of Viruses (ICTV)[5]. According to the daily report of the WHO, by
12 February 25,2020, the epidemic of SARS-CoV-2 has caused 80,239 confirmed cases,
13 of which 77,780 were confirmed in China, 2,666 were dead, and 2,459 were
14 confirmed in 33 countries outside China, 34 patients died[6]. COVID-19 caused by
15 SARS-CoV-2 has posed a serious threat to global public health.

16
17 SARS-CoV-2 can cause different clinical types (severity) of COVID-19. Mild and
18 some of the moderate patients can heal themselves, but other moderate patients will
19 develop into severe patients. The severe patients with underlying diseases often
20 develop organ dysfunction (such as shock, acute respiratory distress syndrome
21 (ARDS), acute heart injury, and acute renal failure), and some critical patients require
22 admission to the intensive care unit (ICU). Therefore, it is very imperative to find

1 indicators that can predict the severity of the disease.

2

3 COVID-19 patients are commonly accompanied by an increase of inflammatory
4 biomarkers [e.g., C-reactive protein (CRP), procalcitonin (PCT), and interleukin 6
5 (IL6)], leukopenia and lymphopenia[7, 8], which is more pronounced in severe
6 patients. However, the status of the lymphocyte subpopulations especially the T cell
7 subsets in these patients are not well understood. Furthermore, the dynamic changes
8 of the previously mentioned inflammatory biomarkers and the lymphocyte
9 subpopulations in COVID-19 patients remain unclear. With this in mind, we
10 retrospectively analyzed three commonly used inflammatory biomarkers (CRP, PCT,
11 and IL6), T-cell subsets in the peripheral blood as well as the clinical data of 16
12 SARS-CoV-2 infected patients to identify potential predictors regarding the severity
13 of the COVID-19 patients.

14

15 **Methods**

16 **Data collection**

17 The study was approved by the Ethics Committee of Yunnan Provincial Hospital of
18 Infectious Disease, AIDS Care Center. Oral consent was obtained from patients. The 16
19 laboratory-confirmed COVID-19 patients at Yunnan Provincial Hospital of Infectious
20 Disease from January 17 to February 8, 2020, were enrolled in this retrospective study.
21 All medical record information, including epidemiological, clinical, radiological,

1 laboratory and treatment data, was obtained through data collection forms of electronic
2 medical records. All data reviewed by a team of trained physicians.

3 **Laboratory examination**

4 Throat swab samples were collected from patients suspected of having SARS-CoV-2
5 infection and maintained in a collection tube with the virus-preservation solution. After
6 collection, total RNA was extracted within 2 hours using the virus RNA isolation kit
7 (DAAN Gene Co., Ltd. of Sun Yat-sen University, Guangzhou, China). The
8 quantitative reverse transcription PCR (RT-qPCR) assay was performed using a
9 SARS-CoV-2 nucleic acid detection kit according to the manufacturer's protocol
10 (Shanghai GeneDx Biotechnology Co., Ltd., and Shanghai BioGerm Medica
11 Biotechnology Co., Ltd., Shanghai, China). Lymphocyte subsets were detected by flow
12 cytometry using a 4-color BD Multitest™ CD3/CD8/CD45/CD4 kit (BD Biosciences,
13 USA) according to the manufacturer's protocol. Flow cytometry was performed on
14 FACSCalibur (BD Biosciences, USA).

15 **Clinical classification of the COVID-19 patients**

16 Clinical classification of the COVID-19 patients was performed according to the
17 "Guidelines of the Diagnosis and Treatment of COVID-19 Pneumonia (Version 5)"
18 published by the National Health Commission of China. Mild patients: the clinical
19 symptoms of patients are mild, and there is no pneumonia on CT imaging. Moderate
20 patients: patients with fever and respiratory symptoms, CT imaging show pneumonia.

1 Severe patients should meet at least one of the following conditions: shortness of breath
2 and the respiratory rate (RR) is greater than or equal to 30 times/min; Finger oxygen
3 saturation at resting is less than or equal to 93%; arterial oxygen partial pressure to
4 fractional inspired oxygen ($\text{PaO}_2/\text{FiO}_2$ ratio) is less than or equal to 300. Critical
5 patients should meet at least one of the following conditions: patients have respiratory
6 failure and need mechanical ventilation; Patients have shock; Patients with other organ
7 failure and require monitoring in the ICU.

8 **Statistical Analysis**

9 All statistical analyses were performed using R (version 3.5.1). Categorical variables
10 were described using frequency or percentage, and continuous variables were described
11 as mean, median, and quartile intervals. When the data were Gaussian distributed, the
12 mean of continuous variables was compared using ANOVA, and the differences
13 between the 3 groups were tested by Tukey's HSD test. When the data were not
14 Gaussian distributed, the median of continuous variables was compared using
15 non-parametric analysis and the significance is tested by the Kruskal Wallis test.
16 Significance between the two groups is tested by the Wilcoxon signed-rank test.
17 Correlation analysis was performed using Spearman's method. The ggplot2 package
18 (Version 3.2.1, <http://ggplot2.tidyverse.org>, <https://github.com/tidyverse/ggplot2>) was
19 used to visualize data with R. In all statistical analyses, $P < 0.05$ was considered
20 statistically significant. Receiver operating characteristic (ROC) curves were used to
21 assess the accuracy of variables predicting severely ill patients. A pROC package

1 (Version 1.16.1, <http://expasy.org/tools/pROC>) was used to visualize the ROC curve
2 and calculate the area under the curve (AUC).

3 **Results**

4 **Clinical characteristics of COVID-19 patients**

5 All admitted patients were confirmed by RT-qPCR measurement of SARS-CoV-2
6 RNA in throat swab samples. The 16 COVID-19 patients were divided into three
7 groups according to previously mentioned Guidelines, including 5 mild, 7 moderate,
8 and 4 severe cases [Oxygen saturation decreased ($\leq 90\%$)]. The median age was 49
9 years (3-69 years) in total patients, of which the median age of the severe group (61,
10 49-69 years) was older than that of the moderate (50, 32-57 years) and mild group (22,
11 3-40 years). The 16 patients included 9 males (3 in each of groups) and 7 females (2
12 were mild, 4 were moderate, and 1 was severe). Except for one patient who is a native
13 of Kunming (with a history of tourism in Wuhan), the remaining patients are from
14 Wuhan and have no history of contact with the seafood market. Four patients were
15 family clustered, of which three patients were mild and one patient was moderate. The
16 median time from the onset of symptoms to admission was 6 days (2 days for mild
17 patients, 7 days for moderate patients, and 7.5 days for severe patients). Three patients
18 had hypertension (one in the mild group, two in the severe group), and three patients
19 had diabetes (one in the moderate group, two in the severe group). Twelve patients
20 had a fever, of which the mild, moderate and severe groups included 3, 5, and 4
21 patients, respectively. Of the 16 patients, 7 had a cough, 6 had sputum production, 2

1 had chest tightness, 2 had shortness of breath, 6 had fatigue, 1 had a headache, 1 had a
2 sore throat, and 2 had body aches. CT imaging of 6 patients was normal, and the other
3 10 patients were abnormal. Patients with abnormal CT imaging showed abnormalities
4 in the subpleural, pulmonary base, and bronchoalveolar segments, which is consistent
5 with atypical upper respiratory symptoms. Although the CT scan showed abnormal
6 lesions close to the pleura, no pleural fluid was found in all patients, indicating that
7 the pleura was not affected, which is consistent with the absence of chest pain in all
8 patients. Three patients were treated with antibiotics (piracillin and sulbactam,
9 moxifloxacin, and meropenem). All patients received antiviral therapy with inhaled
10 IFN- α for 7 days. Fourteen patients received KALETRA [(lopinavir/ritonavir), BID,
11 500 mg/day], of which 12 patients completed a 14-day course, and one severe patient
12 was discontinued because of vomiting after 2 doses of medication, and another severe
13 patient was stopped due to hepatic dysfunction. Five patients received
14 immunoglobulin (400 mg/kg/day), of which one was a moderate patient and four were
15 severe patients. Of the four severe patients, one patient received a 4-day course of
16 glucocorticoids (prednisone, 80 mg/kg/day). As of February 24, 2020, 13 of the 16
17 patients had SARS-CoV-2 RNA converted to negative, and one patient in each group
18 had not converted to negative RNA. The median days from onset to virus eradication
19 was 21 days, 17 days for mild patients, 24 days for moderate patients, and 21 days for
20 severe patients.

21 **The Counts of lymphocyte subsets were associated with the severity of the**
22 **COVID-19 patients**

1 To find laboratory indicators associated with the severity of the disease at admission,
2 we compared the differences of CRP, PCT, IL6, WBC, neutrophil, lymphocyte (LC),
3 and T-lymphocyte subsets (CD3⁺, CD4⁺, and CD8⁺ T) cells among the 3 clinical types
4 of COVID-19 patients. As shown in **Figure 1**, Except for PCT, serum levels of CRP
5 and IL6 were significantly higher in the severe group than in the mild and moderate
6 group (all P<0.05). LC counts (P=0.02), CD3⁺ T-cell counts and ratio (P=0.019,
7 P=0.02), CD4⁺ T-cell counts (P=0.02), CD8⁺ T-cell counts and ratio (P=0.019,
8 P=0.02) were significantly lower in the severe group compared to the mild group. In
9 contrast, NLR (neutrophils to lymphocyte ratio) were significantly higher in the
10 moderate group than in the mild group (P=0.015). Of note, only CD3⁺ T-cell counts
11 (P=0.029), CD4⁺ T-cell counts (P=0.047) and CD8⁺ T-cell counts (P=0.047) were
12 significantly lower in the severe group than in the moderate group. Except for CD3⁺,
13 CD4⁺ and CD8⁺ T-cells counts, no differences in these indicators were observed
14 between the severe and the moderate group. These data suggest that the decreased
15 CD3⁺, CD4⁺ and CD8⁺ T-cell counts have the potential value to predict the severity of
16 COVID-19 patients.

17

18 **Dynamic analysis of the indicators in moderate and the severe COVID-19** 19 **patients**

20 We analyzed the dynamic changes of CRP, PCT, IL6, WBC, Neutrophil, LC in
21 moderate and severe groups. As shown in **Figure 2**, CRP levels gradually decreased
22 in both moderate and severe groups. PCT levels continued to be lower from 6 to 21

1 days in the moderate group, whereas it was initially higher and then gradually
2 decreased in the severe group. In the severe group, IL6 levels peaked at days 8-11 but
3 then continued to decline. In the moderate group, IL6 showed a downward trend in
4 fluctuations. Except for days 12-14, there was no significant difference in IL6 levels
5 between the two groups. At days 6-8 after disease onset, the LC counts in the
6 moderate group were higher than that of the severe group, but there was no statistical
7 difference. In the following days, the LC counts of the moderate group gradually
8 reached their highest levels and kept close to normal levels (1.5×10^9 cells/L). The
9 LC counts in the severe group gradually decreased, reached its lowest value on days
10 12-14, and then gradually increased. On days 18-20, LC counts in the severe group
11 reached a comparable level to the moderate group. However, there were statistically
12 significant differences in LC counts between the two groups over three consecutive
13 time periods ($P < 0.05$). There were no significant differences in WBC counts,
14 Neutrophil counts, and NLRs between the two groups throughout the course of the
15 disease.

16 Next, we analyzed the dynamic changes of CD3⁺, CD4⁺ and CD8⁺ T-cells in the
17 moderate and the severe COVID-19 patients. As shown in **Figure 2**, the CD3⁺, CD4⁺
18 and CD8⁺ T-cell counts in the moderate group were significantly higher than that of
19 the moderate group initially ($P < 0.05$). However, the counts of lymphocytes and their
20 subsets including CD3⁺, CD4⁺, and CD8⁺ T-cells in the severe group gradually
21 increased. With the disease recovery, all these cells reached a comparable level to the
22 moderate group. There were no significant differences in the percentage of CD3⁺,

1 CD4⁺ and CD8⁺ T-cells, CD4/CD8 ratio between the two groups throughout the
2 course of the disease.

3 To investigate the association between these indicators, we performed
4 Spearman's correlation analysis. As shown in **Figure 3**, LC counts were negatively
5 correlated with the majority of variables ($R < -0.3$, $P < 0.05$) except for WBC and
6 Neutrophil counts. In contrast, NLR was positively correlated with all the other
7 variables ($R > 0.3$, $P < 0.05$) except for LC ($R < -0.3$). Levels of CRP, PCT and IL6
8 displayed a positive correlation with each other ($R > 0.3$, $P < 0.05$). Of note, IL6 was
9 positively correlated with other variables ($R > 0.3$, $P < 0.05$), but negatively correlated
10 with LC counts ($R < -0.3$, $P < 0.05$). Moreover, counts of lymphocytes and their subsets
11 including CD3⁺, CD4⁺, and CD8⁺ T-cells displayed a positive correlation with each
12 other ($R > 0.9$, $P < 0.05$).

13 **Counts of T cells are potential predictors for severe COVID-19 patients**

14 To investigate the potential predictive value of these parameters, a receiver operating
15 characteristic (ROC) analysis was performed and the area under ROC curve (AUC)
16 was calculated. As shown in **Figure 4**, four variables (including the counts of
17 lymphocyte, CD3⁺, CD4⁺, and CD8⁺ T-cells) had an AUC of greater than 0.9, whereas
18 other variables had an AUC of lesser than 0.9. Furthermore, the cutoff values of the
19 four parameters with higher AUC were calculated from the ROC curves, with a value
20 of 0.855×10^9 cell/L for lymphocyte, 582 cells/ μ L for CD3⁺ T-cells, 347 cells/ μ L for
21 CD4⁺ T-cell and 171 cells/ μ L for CD8⁺ T-cells. Both CD3⁺ and CD4⁺ T-cells showed

1 the highest AUC which suggesting a more powerful prediction performance for
2 severe COVID-19 patients.

3 **Discussion**

4 Previous reports showed that the majority (63%-70%) of COVID-19 patients on
5 admission have lymphopenia[9, 10], and nonsurvivors develop more severe
6 lymphopenia throughout the course of the disease[10]. A case report also confirmed
7 this phenomenon, in which the nonsurvivor's lymphocytes continued to decrease[11].
8 These papers suggest that lymphopenia might be a common feature in the COVID-19
9 patients and the lymphocyte status might be a critical factor related to disease severity
10 and mortality. In addition to confirming the previously reported lymphopenia in the
11 COVID-19 patients, we found that the counts of lymphocytes as well as their subsets
12 CD3⁺, CD4⁺, and CD8⁺ T-cells are significantly decreased in severe patients compare
13 to mild and moderated patients on admission. We also found that the lymphocytes
14 especially the T-lymphocyte subsets gradually increased in severe patients with the
15 recovery of COVID-19 patients, suggesting that the recovery of T-lymphocytes is
16 associated with a favorable outcome in severe COVID-19 patients.

17

18 Previous studies have demonstrated a correlation between T-cell counts and the
19 severity of SARS-CoV and MERS-CoV infected patients[12], as well as explored the
20 possible mechanisms[13, 14]. Both SARS-CoV and MERS-CoV were found to be
21 able to directedly infect the T cells and induced the lymphocyte apoptosis[13, 14].

22 Other studies have attributed the activation of the hypothalamus-pituitary-adrenal

1 (HPA) axis under stress, which leads to excessive cortisol that causes apoptosis and
2 decreases of lymphocytes[15]. This phenomenon is also seen in other patients with
3 severe infections, such as sepsis[16]. Currently, there is no public evidence that
4 steroid therapy can improve the incidence or mortality of SARS[17], but it can further
5 aggravate the severity of lymphopenia[15]. Because lymphopenia is related to the
6 severity of COVID-19 patients, caution should be exercised in using glucocorticoids
7 in these patients. Even if low-dose glucocorticoids are needed to reduce pulmonary
8 edema in patients with COVID-19, cortisol concentrations and the status of
9 lymphocyte subsets in peripheral blood should be measured. It is unclear whether the
10 T-cell reduction is caused by SARS-CoV-2 direct infection of T-cells or by indirect
11 effects caused by systemic stress responses (such as HPA dysfunction). Further
12 research is needed to explore the corresponding mechanism in detail.

13

14 Effective predictors can help doctors provide appropriate supportive care for patients
15 with severe COVID-19. In this study, we found that four variables including
16 lymphocyte, CD3⁺, CD4⁺, and CD8⁺ T-cells showed the potential value to predict
17 severe patients. Both CD3⁺ and CD4⁺ T-cells showed the highest AUC (0.972) which
18 suggested a more powerful predictive performance. A recent study reported that the
19 NLR was a powerful prognostic factor for severe COVID-19 patients[18]. In their
20 study, patients aged 50 years or older and NLR equal to or greater than 3.13 may
21 serve as a high-risk factor for COVID-19 patients. In our study, we found that NLRs
22 for all severe patients all exceeded 5, and NLRs for moderate patients ranged from 3

1 to 4. Moreover, the AUC of NLR for predicting severe patients is less than 0.9,
2 indicating poor predictive performance. Another study showed that N8R (neutrophils
3 to CD8⁺ T-cell ratio) has better performance with a higher AUC value than NLR in
4 the ROC curve analysis, and may serve as a more powerful factor than NLR for
5 predicting the severity in COVID-19 patients[19]. However, neutrophils are
6 susceptible to many factors, and its ratio to CD8⁺ T cells will mask the actual
7 reduction of T cells. Therefore, we believe that using direct counting of T cells may
8 more accurately reflect the patient's immune status and thus have better predictive
9 performance.

10

11 Previous studies have shown that CD4⁺ T-cell counts are the main prognostic factor
12 for AIDS patients[20]. When the counts of CD4⁺ T cells are below 200 cells/ μ L,
13 patients are prone to secondary infections and tumors. HIV mainly infects CD4⁺ T
14 cells and causes them to decline. However, in the early stages of HIV infection, CD8⁺
15 T cells increase reactively[21]. In this study, we found that the CD4⁺ and CD8⁺ T
16 cells of COVID-19 patients decreased in a fixed ratio, which is different from
17 HIV-infected patients. In addition, we found that the cutoff values for CD4⁺ and CD8⁺
18 T cells are 347 and 171 cells/ μ L, respectively. Due to the small sample size of our
19 study, there may be some deviation in the cutoff values. A larger sample size study is
20 needed to determine the best cutoff values. Nonetheless, the cutoff values we obtained
21 are informative.

22

1 CRP, PCT, IL6 are commonly used inflammatory indicator for bacterial
2 infections[22]. CRP is an acute-phase protein that is an extremely sensitive but
3 non-specific biomarker in many kinds of systemic inflammatory processes, including
4 bacterial and viral infections[23]. PCT has been proposed to be a reliable marker for
5 distinguishing bacterial infections from other types of infections[24]. IL6 is always
6 used for monitoring early states of infectious because of its fast serum peak[25]. In
7 patients with COVID-19, about half of the patients had elevated concentrations of
8 CRP (>10 mg/L)[26, 27]. Levels of IL6 showed sustained increases in severe patients
9 compared to the mild patients[19]. In this study, we found that levels of CRP and IL6
10 were significantly higher in the moderate and severe patients than in mild patients on
11 admission. With the recovery of COVID-19 patients, the elevated CRP and IL6
12 gradually decreased to normal levels. However, there was no difference in CRP, PCT,
13 and IL6 levels between moderate and severe patients. ROC curve analysis also
14 indicated that all of these indicators could not predict the severity of the disease. In
15 SARS patients, the IL6 concentration was increased and positively correlated with the
16 severity of the disease[28]. Although we found that IL6 does not predict the severity
17 of COVID-19 patients, it does not mean that testing for IL6 is useless. It is well
18 known that IL6 plays a key role in cytokine release syndrome (CRS) in chimeric
19 antigen receptor (CAR) T-cell therapy of leukemia, as CRS patients can be found to
20 have high levels of IL-6[29]. IL6 receptor (IL6R) antibody (e.g. Tocilizumab) can
21 effectively control the CRS caused by CAR-T treatment[30]. Cytokine storms can
22 usually be observed in patients with severe COVID-19[9]. Of course, the mechanism

1 of cytokine storm caused by SAR-COV-2 may be different from CAR-T treatment.
2 However, given that both patients have higher levels of IL6, it is suggested that severe
3 COVID-19 patients with significantly increased IL6 may also benefit from IL6R
4 monoclonal antibody treatment. Most patients with COVID-19 had normal serum
5 PCT levels on admission, but their serum PCT levels increased when they had a
6 secondary infection[9]. Because patients with low lymphocytes are prone to bacterial
7 infections, dynamic testing for PCT may be helpful in diagnosing patients with
8 bacterial infections.

9

10 In conclusion, the counts of CD3⁺, CD4⁺, and CD8⁺ T-cells are significantly reduced
11 in severe patients compare to mild and moderated COVID-19 patients, and the
12 reduction of these T cells is related to the severity of the disease. Both the counts of
13 CD3⁺ and CD4⁺ T-cells can be used as predictive indicators for predicting the severity
14 of COVID-19 patients, and dynamic detection of the T-lymphocyte subsets may help
15 predict the outcome of the patients.

16 **Conflict of interest:**

17 The authors disclose no conflicts of interest.

18 **Data Availability:**

19 The data that support the findings of this study are available from the corresponding
20 author upon reasonable request.

1

2 **Acknowledgments:**

3 This study was supported financially by New Product Development Projects of Yunnan
4 Province (2016BC005), The “Thirteenth Five-Year” Science and Technology Major
5 Project for Prevention and Control of AIDS, Viral Hepatitis and Other Major Infectious
6 Diseases (2017ZX10202101-002-005) and the National Natural Science Foundation of
7 China (8136024).

8

9 **References**

- 10 1. Zhu N, Zhang D, Wang W, et al. A Novel Coronavirus from Patients with
11 Pneumonia in China, 2019. *N Engl J Med* **2020**; 382:727-33.
- 12 2. Sahin AR. 2019 Novel Coronavirus (COVID-19) Outbreak: A Review of the
13 Current Literature. *Eurasian Journal of Medical Investigation* **2020**.
- 14 3. Lu R, Zhao X, Li J, et al. Genomic characterisation and epidemiology of 2019
15 novel coronavirus: implications for virus origins and receptor binding. *Lancet* **2020**;
16 395:565-74.
- 17 4. Naming the coronavirus disease (COVID-2019) and the virus that causes it.
18 Available at:
19 [https://www.who.int/emergencies/diseases/novel-coronavirus-2019/technical-guidanc](https://www.who.int/emergencies/diseases/novel-coronavirus-2019/technical-guidance/naming-the-coronavirus-disease-(covid-2019)-and-the-virus-that-causes-it)
20 [e/naming-the-coronavirus-disease-\(covid-2019\)-and-the-virus-that-causes-it.](https://www.who.int/emergencies/diseases/novel-coronavirus-2019/technical-guidance/naming-the-coronavirus-disease-(covid-2019)-and-the-virus-that-causes-it)

- 1 5. Gorbalenya AE, Baker SC, Baric RS, et al. Severe acute respiratory
2 syndrome-related coronavirus: The species and its viruses – a statement of the
3 Coronavirus Study Group. *bioRxiv* **2020**:2020.02.07.937862.
- 4 6. Coronavirus disease 2019 (COVID-19) Situation Report – 36. Available at:
5 [https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200225-sitr
7 ep-36-covid-19.pdf?sfvrsn=2791b4e0_2](https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200225-sit
6 ep-36-covid-19.pdf?sfvrsn=2791b4e0_2).
- 8 7. Chang, Lin M, Wei L, et al. Epidemiologic and Clinical Characteristics of Novel
9 Coronavirus Infections Involving 13 Patients Outside Wuhan, China. *JAMA* **2020**.
- 10 8. Bai Y, Yao L, Wei T, et al. Presumed Asymptomatic Carrier Transmission of
11 COVID-19. *JAMA* **2020**.
- 12 9. Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel
13 coronavirus in Wuhan, China. *Lancet* **2020**; 395:497-506.
- 14 10. Wang D, Hu B, Hu C, et al. Clinical Characteristics of 138 Hospitalized Patients
15 With 2019 Novel Coronavirus-Infected Pneumonia in Wuhan, China. *JAMA* **2020**.
- 16 11. Xu Z, Shi L, Wang Y, et al. Pathological findings of COVID-19 associated with
17 acute respiratory distress syndrome. *Lancet Respir Med* **2020**.
- 18 12. Wong RS, Wu A, To KF, et al. Haematological manifestations in patients with
19 severe acute respiratory syndrome: retrospective analysis. *BMJ* **2003**; 326:1358-62.
- 20 13. Chu H, Zhou J, Wong BH, et al. Middle East Respiratory Syndrome Coronavirus
21 Efficiently Infects Human Primary T Lymphocytes and Activates the Extrinsic and
22 Intrinsic Apoptosis Pathways. *J Infect Dis* **2016**; 213:904-14.
14. Gu J, Gong E, Zhang B, et al. Multiple organ infection and the pathogenesis of

- 1 SARS. *J Exp Med* **2005**; 202:415-24.
- 2 15. Panesar NS, Lam CW, Chan MH, Wong CK, Sung JJ. Lymphopenia and
3 neutrophilia in SARS are related to the prevailing serum cortisol. *Eur J Clin Invest*
4 **2004**; 34:382-4.
- 5 16. Drewry AM, Samra N, Skrupky LP, Fuller BM, Compton SM, Hotchkiss RS.
6 Persistent lymphopenia after diagnosis of sepsis predicts mortality. *Shock* **2014**;
7 42:383-91.
- 8 17. Gomersall CD. Pro/con clinical debate: steroids are a key component in the
9 treatment of SARS. Pro: Yes, steroids are a key component of the treatment regimen
10 for SARS. *Crit Care* **2004**; 8:105-7.
- 11 18. Liu J, Liu Y, Xiang P, et al. Neutrophil-to-Lymphocyte Ratio Predicts Severe
12 Illness Patients with 2019 Novel Coronavirus in the Early Stage. *medRxiv*
13 **2020**:2020.02.10.20021584.
- 14 19. Liu J, Li S, Liu J, et al. Longitudinal characteristics of lymphocyte responses and
15 cytokine profiles in the peripheral blood of SARS-CoV-2 infected patients. *medRxiv*
16 **2020**:2020.02.16.20023671.
- 17 20. de Wolf F, Spijkerman I, Schellekens PT, et al. AIDS prognosis based on HIV-1
18 RNA, CD4+ T-cell count and function: markers with reciprocal predictive value over
19 time after seroconversion. *AIDS* **1997**; 11:1799-806.
- 20 21. Mudd JC, Lederman MM. CD8 T cell persistence in treated HIV infection. *Curr*
21 *Opin HIV AIDS* **2014**; 9:500-5.
- 22 22. Plesko M, Suvada J, Makohusova M, et al. The role of CRP, PCT, IL-6 and

- 1 presepsin in early diagnosis of bacterial infectious complications in paediatric
2 haemato-oncological patients. *Neoplasma* **2016**; 63:752-60.
- 3 23. Chew KS. What's new in Emergencies Trauma and Shock? C-reactive protein as a
4 potential clinical biomarker for influenza infection: More questions than answers. *J*
5 *Emerg Trauma Shock* **2012**; 5:115-7.
- 6 24. Schuetz P, Albrich W, Mueller B. Procalcitonin for diagnosis of infection and
7 guide to antibiotic decisions: past, present and future. *BMC Med* **2011**; 9:107.
- 8 25. Jawa RS, Anillo S, Huntoon K, Baumann H, Kulaylat M. Interleukin-6 in surgery,
9 trauma, and critical care part II: clinical implications. *J Intensive Care Med* **2011**;
10 26:73-87.
- 11 26. Chen H, Guo J, Wang C, et al. Clinical characteristics and intrauterine vertical
12 transmission potential of COVID-19 infection in nine pregnant women: a
13 retrospective review of medical records. *The Lancet* **2020**.
- 14 27. Chan JF, Yuan S, Kok KH, et al. A familial cluster of pneumonia associated with
15 the 2019 novel coronavirus indicating person-to-person transmission: a study of a
16 family cluster. *Lancet* **2020**; 395:514-23.
- 17 28. Zhang Y, Li J, Zhan Y, et al. Analysis of serum cytokines in patients with severe
18 acute respiratory syndrome. *Infect Immun* **2004**; 72:4410-5.
- 19 29. Shimabukuro-Vornhagen A, Godel P, Subklewe M, et al. Cytokine release
20 syndrome. *J Immunother Cancer* **2018**; 6:56.
- 21 30. Maude SL, Barrett D, Teachey DT, Grupp SA. Managing cytokine release
22 syndrome associated with novel T cell-engaging therapies. *Cancer J* **2014**; 20:119-22.

1 **Figure legends**

2 **Figure 1. Compare the indicators in the different type (severity) of COVID-19**
3 **patients on admission**

4 The lines in the middle of the boxes represent the median, boxes represent 25th and
5 75th percentiles and whiskers represent 5th and 95th percentiles. Significance between
6 different groups is tested by the Kruskal Wallis test. A p-value of less than 0.05 is
7 statistically significant.

8 **Figure 2. Dynamic analysis of indicators in moderate and the severe COVID-19**
9 **patients**

10 The lines represent mean, error bar represents SD. Significance between different
11 groups is tested by the Wilcoxon signed-rank test. Asterisk (*) indicates significant
12 ($P < 0.05$) differences between the two groups in various time points.

13 **Figure 3. Correlation analysis between indicators**

14 The correlation coefficient (R) was calculated using Spearman's correlation method. R
15 between 0.3 and 0.7 (-0.3 and -0.7) indicate a moderate positive (negative) correlation;
16 R between 0.7 and 1 (-0.7 and -1) indicate a strong positive (negative) correlation. Mark
17 "X" indicates no significant difference ($P > 0.05$).

18 **Figure 4. Receiver operator characteristic (ROC) analysis of the biomarker**
19 **performance**

- 1 The best cutoff values of the parameters with higher areas under the curve (AUC) were
- 2 calculated from the ROC curves.

Figure 1




Figure 2




Figure 4

