A structured open dataset of government interventions in response to COVID-19

Amelie Desvars-Larrive^{1,2}, Elma Dervic^{2,3}, Nils Haug^{2,3}, Thomas Niederkrotenthaler^{2,4}, Jiaying Chen^{2,3}, Anna Di Natale^{2,3}, Jana Lasser^{2,3}, Diana S. Gliga⁵, Alexandra Roux^{6,7}, Abhijit Chakraborty^{2,8}, Alexandr Ten⁹, Alija Dervic¹⁰, Andrea Pacheco¹¹, David Cserjan², Diana Lederhilger⁵, Dorontinë Berishaj⁵, Erwin Flores Tames², Huda Takriti², Jan Korbel^{2,3}, Jenny Reddish^{2,12}, Johannes Stangl⁵, Lamija Hadziavdic⁵, Laura Stoeger², Leana Gooriah¹¹, Lukas Geyrhofer², Marcia R. Ferreira², Rainer Vierlinger⁵, Samantha Holder⁵, Samuel Álvarez¹³, Simon Haberfellner⁵, Verena Ahne², Viktoria Reisch⁵, Vito D.P. Servedio², Xiao Chen⁵, Xochilt María Pocasangre-Orellana¹³, David Garcia^{2,3}, Stefan Thurner^{2,3,14}

1. Unit of Veterinary Public Health and Epidemiology, University of Veterinary Medicine Vienna, Veterinaerplatz 1, 1210 Vienna, Austria

2. Complexity Science Hub Vienna, Josefstaedter Strasse 39, 1080 Vienna, Austria

3. Section for Science of Complex Systems, Medical University of Vienna, Spitalgasse 23, 1090 Vienna, Austria

4. Unit Suicide Research & Mental Health Promotion, Center for Public Health, Department of Social and Preventive Medicine, Medical University of Vienna, Kinderspitalgasse 15, 1090 Vienna

5. Independent scholar

6. CERMES3, Ecole des Hautes Etudes en Sciences Sociales, 7 rue Guy Moquet, 94801 Villejuif, France

7. Gender, Sexuality, Health, CESP, INSERM, Paris-Saclay University, Paris-Sud University, UVSQ, Hôpital Paul Brousse, 16 avenue Paul Vaillant Couturier, 94807 Villejuif, France

8. Advanced Systems Analysis, International Institute for Applied Systems Analysis (IIASA), Schlossplatz 1, A-2361 Laxenburg, Austria

9. Flowers project-team, National Research Institute for Digital Sciences (INRIA), 200 avenue de la Vieille Tour, 33405 Talence cedex, France

10. Institute of Electrodynamics, Microwave and Circuit Engineering, Vienna University of Technology, Gusshausstrasse 25/354, 1040 Vienna, Austria

11. German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Deutscher Platz 5E, 04103 Leipzig, Germany

12. Seshat: The Global History Databank

13. Fundación Naturaleza El Salvador, Research Department, Colonia Escalón, San Salvador,

CP 1101, El Salvador

14. Santa Fe Institute, Santa Fe, NM 87501, USA

Corresponding author(s): Amélie Desvars-Larrive (amelie.desvars@vetmeduni.ac.at)

Abstract

In response to the COVID-19 pandemic, governments have implemented a wide range of nonpharmaceutical interventions (NPIs). Monitoring and documenting government strategies during the COVID-19 crisis is crucial to understand the progression of the epidemic. Following a content analysis strategy of existing public information sources, we developed a specific hierarchical coding scheme for NPIs. We generated a comprehensive structured dataset of government interventions and their respective timelines of implementation. To improve transparency and motivate collaborative validation process, information sources are shared via an open library. We also provide codes that enable users to visualise the dataset. Standardization and structure of the dataset facilitate inter-country comparison and the assessment of the impacts of different NPI categories on the epidemic parameters, population health indicators, the economy, and human rights, among others. This dataset provides an in-depth insight of the government strategies and can be a valuable tool for developing relevant preparedness plans for pandemic. We intend to further develop and update this dataset on a weekly basis until the end of December 2020.

Background & Summary

Non-pharmaceutical interventions (NPIs), also known as community mitigation strategies, aim to prevent the introduction of infectious diseases (preparedness and readiness measures), control their spread and reduce their burden on the health system (control measures). The general concept of containing the initial (exponential) spread of a disease is called "flattening the (epi-) curve" ¹. By reducing the growth rate of an epidemic, NPIs reduce the stress on the healthcare system and help gaining time to develop and produce vaccines and specific medications, which is of utmost importance in the case of emerging infectious diseases.

During the COVID-19 pandemic, governments have enforced a broad spectrum of intervention measures, under rapidly changing, unprecedented circumstances. Government responses to COVID-19 included the *laissez-faire* strategy, which implies doing little to nothing, the *herd immunity* strategy, which implies a few measures only or measures relying

on voluntary compliance, and more *aggressive* approaches based on the implementation of a wide range of stringent NPIs, sometimes even limiting civil rights and liberty ^{2,3}. Governments control policies have shown divergences in particular in the timeline of implementation and in the prioritization of the NPIs. In China for example, quarantine, social distancing, cordon sanitaire, and isolation of cases have been associated with improvements in the key epidemiological markers, including the number of infections and COVID-19-related deaths ⁴. In Hong Kong and Taiwan, which experienced severe acute respiratory syndrome (SARS) epidemics in 2002-2003 ^{5,6}, early government actions, strict social distancing measures, contact tracing, extensive and proactive testing, and high compliance of the population, have, to date, successfully mitigated the COVID-19 epidemic ^{7,8}. Following a *herd immunity* approach, similar to the one initially adopted by the UK government, the Swedish government did not introduce strict bans but formulated non-binding recommendations only ⁹. Predictive models, however, suggest that such a strategy might ultimately overwhelm the healthcare system due to shortages in critical care capacity ¹⁰.

Poor control policies have potentially dramatic repercussions on public health. Although there is an urgent need for data on government responses to COVID-19, there is currently a limited opportunity to capture information on country-based mitigation measures. Our project aims to generate a comprehensive and structured dataset of government interventions taken to control the spread of COVID-19, including the respective time schedules of their implementation, which can enable to assess the effectiveness of these interventions and facilitate an inter-country comparison.

During the COVID-19 crisis, other data collection initiatives have emerged on this topic; all of them offering public access to data. Nonetheless, most of them focus on a specific type of measures, e.g. the closure of educational institutions ^{11,12}, travel restrictions¹³, trade-related measures¹⁴, legal and economic responses¹⁵, or measures to ensure continuity of supply of personal protective equipment and critical medical products^{16,17}. Two datasets describe a larger range of interventions; however, the number of measures per country is limited ¹⁸, measures are not explicitly detailed, or sources are not provided¹⁹.

In the context of the current COVID-19 health crisis, open knowledge ²⁰ and data sharing are crucial to understand and help to mitigate the pandemic. In this article, we document and share the methodologies, tools and approaches used to produce the *Complexity Science Hub COVID-19 Control Strategies List* (CCCSL) dataset following the principles of open science. We provide a detailed description of the dataset and present examples of how the data can be used in order to provide insights into the global government responses to COVID-19. The dataset is readily usable for modelling and machine learning analyses.

By compiling the information on NPIs, the CCCSL dataset allows evaluating the government policies against COVID-19 and should further enable to estimate the impact of these measures on population health indicators, the economy, and human rights, among others. The CCCSL can also be used as a tool for designing emergency preparedness and response plans for epidemic respiratory diseases. Considering the imperative necessity for such data, version 1 of the dataset has already been released. The dataset is not complete and we continuously update it with new available records. Updates are planned until, at least, the end of December 2020.

Methods

We used a content analysis²¹⁻²³ strategy of existing information sources to develop a hierarchical coding scheme specific to NPIs implemented to mitigate the burden of COVID-19. First, based on a literature review on community mitigation strategies and expert knowledge, eight themes (thereafter called level 1 (L1) in the coding scheme) were identified and labelled: (i) Case identification, contact tracing and related measures, (ii) Environmental measures, (iii) Healthcare and public health capacity, (iv) Resource allocation, (v) Risk communication, (vi) Social distancing, (vii) Travel restriction, and (viii) Returning to normal life. A definition for each theme is provided in Table 1. At the start of our project, there were no previously published studies on NPIs against COVID-19 to be used as a reference for developing our labelling and coding scheme. Therefore, a list of NPIs that have been already implemented by different governments at this time (mid-March 2020) was compiled, that served as a preliminary template to generate a priori categories within a hierarchical coding scheme. Strategies that could provide assistance to the population (e.g., related to financial support or food supply) or that may encourage compliance with the measures (e.g. resource allocations, risk communication) were also included. Listed intervention measures were then assigned to one of the eight themes defined above. The specific details and descriptions of each NPI were coded into a priori categories (thereafter called level 2 (L2) in the coding scheme), and into subsequent *a priori* subcategories and codes whenever needed (thereafter called level 3 (L3) and level 4 (L4) in the coding scheme, respectively). Discrepancies in code assignments were discussed within the coding team and were resolved by consensus. The objective of this hierarchical coding scheme for NPIs was to standardize the data collection and obtain a structured dataset that uses a consistent taxonomy, and therefore, promotes common understanding.

On 19 March 2020, we set up a platform for students, researchers, and volunteers to collect data on the NPIs implemented by the governments for preventing and limiting the spread of COVID-19, including the time schedules for the implementation. Data collectors received clear instructions on the objective of the project and indications on how to proceed for data collection. The template of *a priori* themes, categories, subcategories, and codes described above was provided. Data collectors were asked to use the *a priori* coding system or to refer to the data curators if a measure could not be coded using the *a priori* coding system. Therefore, throughout the data collection process, new categories, subcategories, and codes emerged, derived directly from the text data sources. The emergent (inductive) categories and subcategories were openly coded by the data collectors or by the data curators. In a second step, inductive categories and subcategories were compared together and in relation to the entire dataset to detect co-occurrence (codes that partially or completely overlap) and redundancies. Codes with the same meaning were aggregated²⁴. The categories and subcategories were tightened up to the point that maximized mutual exclusivity and exhaustiveness²³. This resulted in a master list of coding categories (a list of all the codes that were developed and used in the study), including the curated *a priori* and inductive coding categories. The master list replaced the *a priori* template for categorisation of the measures during data collection. It was shared with the data collectors via a Google spreadsheet and updated daily.

A wide range of different public sources were used to populate, update and curate our dataset, including official government sources, peer-reviewed and non-peer-reviewed scientific papers, webpages of public health institutions (WHO, CDC, and ECDC), press releases, newspaper articles, and government communication through social media. We collected data on the following: (i) country, (ii) region (when measures were implemented at subnational-level), (iii) date of implementation of the measure, (iv) implemented measures coded following the four-level classification scheme described above (theme, category, subcategory and code), and (v) source. For each country, data were preferentially collected in the language of the country by native data collectors (i.e. Austria, Belgium, Bosnia and Herzegovina, Brazil, Canada, Croatia, Czech Republic, Ecuador, El Salvador, France, Germany, Honduras, Hong Kong, India, Italy, Kazakhstan, Kosovo, Kuwait, Mauritius, Mexico, Montenegro, North Macedonia, New Zealand, Portugal, Ireland, Romania, Serbia, Spain, Syria, Taiwan, and United Kingdom). If this was not possible, Google Translate was used to translate documents²⁵.

Data Records

A static copy of the dataset has been archived in figshare, including all NPIs recorded as of time of submission (4 May 2020), spanning the period 31 December 2019 to 25 April 2020 and covering 53 countries over all continents. A dynamic version of the dataset, which will be continually updated, can be accessed via Github: https://github.com/amel-github/covid19interventionmeasures) from Google Drive: or https://drive.google.com/open?id=1041U8iWPDSGI6KHIn9Dg7THkXIo3-gui, in CSV format. Each of the rows represents a single individual NPI and is identified by a unique ID. The master list of coding categories is also available from these repositories. We have also established a Github repository available at: https://github.com/amel-github/CCCSL-Codes and provide codes for importing and visualising the data into R statistical software²⁶. Furthermore, for purposes of transparency of data collection and to motivate collaborative validation process as well as a large use and development of our dataset, an open library is all collect available, that contains sources used to the data: https://www.zotero.org/groups/2488884/cccsl_covid_measure_project. In order to leverage on the potential of crowdsourcing for populating and curating the CCCSL dataset, we have launched a webpage dedicated to this project at: http://covid19-interventions.com/ where contributors can fill up a Google form at: https://bit.ly/2KsYOTn, if they wish to correct entries, add a measure, and/or provide a feedback.

The dataset contains the following fields:

ID – Unique identifier for the implemented measure. ID is also used in the Google Form to report erroneous entries.

Country – The country where the NPI measure was implemented.

ISO3 – Three-letter country code as published by the International Organization for Standardization.

Region – Subnational geographic area (e.g. region, department, municipality, city) where the NPI measure has been locally implemented (i.e. the measure was not implemented nationwide as of the mentioned date). The country name otherwise (i.e. measure implemented nationwide).

Date – Date of implementation of the NPI. Date of announcement was used when the date of implementation of the NPI could not be found and this was specified in the field *Comment*.

L1_Measure – Themes (L1 of the classification scheme). Eight themes were defined (see Table 1).

L2_Measure – Categories (L2 of the classification scheme). Table 1 provides the list of the categories for each theme.

L3_Measure – Subcategories (L3 of the classification scheme). Provides detailed information on the corresponding category (L2).

L4_Measure – Codes (L4 of the classification scheme). Corresponds to the finest level of description of the measures.

Comment – Provides the description of the measure as found in the text data source, translated into English. This field allows to judge the quality of the label for the different levels of the coding scheme and enables to re-assign the measure to the correct theme/category/subcategory/code in case of error or misinterpretation of the data collector ²⁷.

Source – Provides the reference for each entry. Enables to trace back potential changes in the meaning of the label during the translation ²⁷. Enables to access the description of the measures in the source language and/or to access to the information as it was dispatched originally.

At the time of submission, the CCCSL dataset included information for 3,274 intervention measures, from 53 countries, including 32 European countries, 12 Asian countries, five South American, one North American country, one Oceanian country, one African country, and the Diamond Princess cruise ship. Fig. 1, Table 2, and Table 3 summarize the dataset. A description of the measures grouped per themes (L1) for each country can be computed from the codes available in the Github (https://github.com/amel-github/CCCSL-Codes).

Technical Validation

After the initial data entry, the dataset was checked manually by the data curators. For each measure, concordance between L1, L2, L3, and L4 was checked. Moreover, the unique combinations of L1, L2, L3, and L4 were extracted and controlled for consistency. Typographical and coding errors were also minimized through a manual process. We also initiated a collaborative curation platform relying on internal and external collaborators who exchanged through Slack, Github, Skype, and via emails. This extended effort enabled us to correct typographical and coding errors, to remove line breaks, and to homogenize the dataset for universal use in different programming languages.

Beyond manual validations, we performed a technical validation step to detect possible duplicates. Using the *dplyr* package for the R Statistical Language²⁸, we identified any duplicate entries in the vector composed of country, region, date, and the codes from L1 to

L4. Those entries were flagged as possible duplicates and reviewed by hand by two curators, ensuring that the dataset does not contain duplicated entries. An R script to reproduce this step is provided at: <u>https://github.com/amel-github/CCCSL-Codes</u>.

While an important effort has been made for standardizing the records, the four-level-*a priori* coding scheme originally proposed showed limitations. First, the existing classification of NPI measures are discordant ²⁹⁻³¹. We proposed a classification scheme that best fitted our (emergency) needs and the specificity of the COVID-19 pandemic, but this scheme may be subjected to revisions in the future. Secondly, some NPIs have been uniquely implemented (e.g. the deportation of Chinese workers by the Kazakh government) which complicated the coding and categorisation process.

Access to the information from governmental or official source can be compromised if not performed timely. Indeed, several governments or national health agencies update their webpage to provide the latest information to the public. Therefore, access to previous content (i.e. previous restrictions and measures) from official sources was sometimes difficult and some links to online document sources may not exist in the future. Furthermore, while native speakers were recruited whenever possible for data collection, transliteration or translation errors may have occurred when extracting data from Google Translate translations.

We aim to maintain the quality level of the dataset with weekly updates on the countries currently described. Furthermore, we plan to increase the geographic coverage of the dataset, prioritizing large countries (e.g. China, USA, and Australia), those with a high number of reported cases (e.g. Vietnam, Iran, Turkey, Russia, Israel, Peru, Chile Poland, Pakistan, Philippines, Saudi Arabia), and those in which the epidemic is rising and which suffer from a data gap (i.e. African and South American countries). The same technical procedures and the classification scheme described above will be applied to any new information to be included in the dataset. Future versions will be subjected to extensive data validation processes. Four main factors will be considered in future updates: (1) subnational level data; (2) planned duration of measure at time of implementation; (3) new measures versus extension of a previous measure; and (4) measures related to returning to normal life (lifting of restrictions). We plan to stabilize our hierarchical coding scheme for NPIs implemented to contain COVID-19 within six months (measures related to the lifting of the restrictions and adaptive measures that accompany them will be included).

Usage Notes

The aim of this work is not only to improve the current knowledge on country-based intervention measures implemented to mitigate the burden of COVID-19, but also to characterize the political, public health, and economic strategies of the governments worldwide. Combined with publicly available data on the number of confirmed cases, recovered cases, and deaths ^{32,33}, the CCCSL dataset makes it possible to assess the effectiveness of the control policies on the COVID-19 epidemic, e.g. the epidemic growth rate or the daily reproduction numbers. The dataset can further benefit the risk assessment of lifting some restrictions and the development of exit strategies. It can also become an essential data source in the aftermath of the first wave of COVID-19, to guide government control policies anticipating a potential second wave of cases. We envision the CCCSL dataset to become a timely valuable and long-lasting data source for assessing the impact of the NPIs on global public health indicators, the economy, and human rights, among others. We provide below three examples of data usage that give an insight into the responsiveness and aggressiveness of the governments in their management of the COVID-19 crisis. In these examples, we excluded data about the Diamond Princess cruise ship, as it represents a special case, and NPIs related to "returning to normal life", as the collect of those measures is ongoing.

Identify worldwide decision patterns over time

We propose a visualisation of the global decision pattern over 52 countries using a network analysis of the NPIs recorded in the CCCSL at level 2 (categories) of the coding scheme (Fig. 2). The network analysis is based on the overlap in time of the implemented government interventions. Nodes represent NPIs; the size of each node is proportional to the frequency of the NPI (number of countries that implemented this NPI over all considered countries). The network links are directed; the source of a link corresponds to the measure that was implemented first. An interactive version of Fig. 2 is available online at: http://covid19-interventions.com/Network/interventionsNetwork.html. On the panel on the left side of the online Fig. 2, for a selected node (measure), the weight of each link (average overlap in time) is given.

Mapping the timeline of government interventions through the epidemic

We propose to visualise the time-series of the date of implementation of the NPIs recorded in the CCCSL at level 2 (categories) in the 52 countries (Diamond Princess excluded) using a heat map (Fig. 3). To highlight country-based differences in the timeline of implementation, we used the epidemic age instead of calendar time. For a given day, t, in a certain country, the epidemic age is defined as the time difference, t-t₀, measured in days, where t₀ is the first day when the number of confirmed cases was greater or equal to 10. The time-series data of the number of COVID-19 cases was retrieved from the COVID-19 Data Repository by the Center for Systems Science and Engineering (CSSE) at Johns Hopkins University ³³.

Country-cluster analysis of the government control strategies

In order to partition the countries based on the aggressiveness (number of NPIs) and responsiveness (timeline) of their control strategy, we applied a k-means clustering. We focused on mandatory government interventions (i.e. the theme "Risk communication" was not included) recorded in the CCCSL at level 2 (categories) that appeared in at least 15 countries, leading to a total number of 29 categories. The clustering algorithm uses the date of implementation of the measures in each country to build a feature vector based on the epidemic age (see above). We considered "anticipatory measures" as those implemented before day when 10 cases were reported; "early measures" as those implemented at the beginning of the epidemic, i.e. between the day when 10 cases were reported and the day when 200 cases were reported; and "late measures" as those implemented at a later stage of the epidemic, i.e. after the day when 200 cases were reported. The algorithm takes also into account the number of measures implemented at these different stages of the epidemic. The time-series data of the number of COVID-19 cases was retrieved from the COVID-19 Data Repository by the Center for Systems Science and Engineering (CSSE) at Johns Hopkins University³³. The optimal number of cluster, k, determined by optimizing the withincluster sums of squares, was seven, explaining 80.3% of the variance (Fig. 4). An interactive version of Fig. 4 is available online at: http://covid19interventions.com/CountryClusters.html.

Code Availability

A GitHub record of this project is accessible at <u>https://github.com/amel-github/covid19-</u> <u>interventionmeasures</u>. The codes used to describe the CCCSL dataset and the codes used to

explore the CCCSL dataset in the Usage Notes section are available at: https://github.com/amel-github/CCCSL-Codes.

Acknowledgements

D.G. and A.D.N. acknowledge funding from the Vienna Science and Technology Fund - WWTF (VRG16-005). The authors acknowledge William Schueller for his help in the recruitment of the team of data collectors and Petar Sekulic for his advises in the analysis of the dataset. We warmly thank Caspar Matzhold and Michaela Kaleta for checking and testing our code components.

Author contributions

A.D.L. managed and coordinated the production of the dataset and prepared it for publication, including: developing the coding scheme, collecting and curating the data, managing the team of data collectors, creating the library of sources, writing the data descriptor, and creating the tables. The following authors: E.D., N.H., T.N., E.C., A.D.N., J.L., D.S.G., and A.R. produced a substantial work to generate the dataset and prepare it for publication, including: developing the coding scheme, collecting the data, formatting the data for presentation of the published work, visualising the data, and creating the library of sources. D.G. created the webpage dedicated to this project. The following list of authors (listed alphabetically): A.C., A.T., A.D., A.P., D.C., D.L., D.B., E.F.T., H.T., J.A., J.K., J.R., J.S., L.H., L.S., L.G., L.G., M.R.F., R.V., S.H., S.Á., S.L., S.H., V.A., V.R., V.D.P.S., X.C. and X.M.P.O. includes the many individuals who collected and curated data through the dedicated platform and provided comments to facilitate the use of the dataset across different programming platforms. A.D.L. and D.G. jointly supervised this work. S.T. mentored the core team. All authors reviewed and approved the final manuscript.

Competing interests

The authors declare no competing interests.

Figure Legends

Fig. 1. Geographical coverage of the CCCSL and total number of recorded NPIs that were implemented in each country to control the spread of COVID-19. As of date of submission, the dataset includes 53 countries (including the Diamond Princess cruise ship) and dates of NPI implementation range from 31/12/2019 to 25/04/2020.

> **Fig. 2.** Network of global government decision patterns over time, built using NPIs recorded in the CCCSL at level 2 (categories). The network analysis includes 52 countries (Diamond Princess excluded from this analysis); dates of NPI implementation range from 31/12/2019 to 25/04/2020. The size of the nodes is proportional to the frequency of a given NPI (number of countries that implemented this NPI over all considered countries). The network is filtered such that only edges with weight greater than 0.5 are displayed.

> **Fig. 3.** Heat map of the date of implementation of the NPIs recorded in the CCCSL at level 2 (categories) in 52 countries. Time is in epidemic age with t_0 = day when 10 cases were reported (<u>http://covid19-interventions.com/CountryMeasuresHeatmap.svg</u>).

Fig. 4. Country-cluster analysis based on the number of mandatory government interventions and respective dates of implementation, as calculated using the epidemic age (t_0 = day when 10 cases were reported).

Tables

Table 1. Definition of the eight themes (L1) used to classify the NPIs and associated categories (L2).

Theme	Definition	Associated categories
		[1] Airport health check
		[2] Border health check
		[3] Enhance detection system
	Measures aiming at identifying cases (testing, surveillance), tracing the	[4] Health check at train stations
Case identification contact tracing and related	contacts (i.e. identification and follow-up of persons who may have	[5] Isolation of cases
	come into contact with an infected person 31), implementing isolation	[6] Quarantine
measures	for cases and quarantine for contacts and suspected cases (including	[7] Restriction in testing
	tracking).	[8] Scale up the system
		[9] Specific health channel for travellers
		[10] Surveillance
		[11] Tracing and monitoring
	Measures aiming at reducing the transmission of COVID-19 through	[1] Environmental cleaning and disinfection
Environmental measures	shared environment and surfaces. Those measures include the routine	
environmental measures	cleaning of frequently used surfaces, clothes and objects; minimizing	
	the sharing of objects; and ensuring appropriate ventilation 34 .	
Healthcare and nublic health canacity	Public health capacity under the International Health Regulations (IHR)	[1] Activate emergency response mechanisms
neartheare and public hearth capacity	(2005) is defined as the indispensable, fundamental actions that are	[2] Activate notification of cases

	the primary responsibility of each State Party for achieving the goal of	[3] Case management
	national health security, i.e. to prevent the spread of diseases and to	[4] Enhance emergency response mechanisms
	detect and investigate health risks in the community by efficient	[5] Enhance hand hygiene
	multisectoral action (e.g. laboratory services and national, regional	[6] Enhance healthcare capacity - Funds
	and global networks $^{ m 35}$. Healthcare capacity refers to patient capacity,	[7] Enhance laboratory testing
	healthcare workforce capacity, laboratory capacity, availability of	[8] Increase availability of PPE
	medical supplies and equipment.	[9] Increase healthcare workforce
		[10] Increase in medical supplies and equipment
		[11] Increase isolation and quarantine facilities
		[12] Increase patient capacity
		[13] Measures for blood donation
		[14] Precautions for handling and disposal of dead
		bodies
		[15] Reduce the burden on health system
		[16] Repurpose hospitals
		[17] Research
		[18] Scale up emergency response mechanisms
	Resource allocation covers the whole range of operations involved in	[1] Administrative procedures facilitated
	the allocation of budgets, deployment of resources, and distribution	[2] Ban export of necessity goods
Resource allocation	of goods for controlling a disease outbreak (these measures are most	[3] Crisis management plans
	often carried out by national, regional, or even local authorities). For	[4] Emergency laws

	example, under the IHR (2005), each State Party is required to	[5] Enforce lockdown
	develop, strengthen and maintain core public health capacities for	[6] Fixing maximum price for necessity goods
	surveillance and response by using existing national resources, such as	[7] Fixing price for specific protective products
	the national plans for pandemic preparedness 35,36 .	[8] Mobilize army and police to support application
		of laws related to covid-19 control
		[9] Provide international help
		[10] Receive international help
		[11] Setting price for specific protective products
		[12] The government provides food assistance
		[1] Lift quarantine
	Measures supporting the return to normal life, i.e. lifting of restriction	[2] Re-opening land border
leturning to normal life	measures (e.g. lift of quarantine, re-opening of schools, re-opening of	[3] Re-opening of shops
	shops) and adaptive measures.	[4] Reactivate the economy
Risk communication	All implemented methods for using communication strategically to	[1] Actively communicate with healthcare professionals
	includes health education, health literany health promotion, rick	[2] Actively communicate with stakeholders
	$\frac{1}{1000}$	[3] Call for return of nationals living abroad
	communication and social mobilization , we have included in this	[4] Educate and actively communicate with the
	category an voluntary, non-compulsory, recommended measures of	public
	any kinu.	[5] Travel alert

		[1] Closure of educational institutions
		[2] Mandatory use of masks
	Social distancing refers to methods for reducing frequency and	[3] Mass gathering cancellation
	closeness of contact between people in order to decrease the risk of	[4] Measures for special populations
	transmission of disease. Examples of social distancing include	[5] Public transport
Social distancing	cancellation of public events such as concerts, sports events, or	[6] Return operation of nationals living abroad
	movies, closure of office buildings, schools, and other public places,	[7] Return operation of nationals living in regions
	and restriction of access to public places such as shopping malls or	at risk
	other places where people gather ³⁶ .	[8] Small gathering cancellation
		[9] Special measures for certain establishments
		[10] Work safety protocols
		[1] Airport restriction
		[2] Airports closing for all international flights
Travel restriction		[3] Airports closing for domestic flights
		[4] Border restriction
	Travel restriction encompasses travel alert, entry and exit screening,	[5] Cancellation of international flights
	internal travel restrictions, border closure ³¹ , and cordon sanitaire.	[6] Commercial flights suspended
		[7] Cordon sanitaire
		[8] Denial entry of travellers from China
		[9] Denial entry to boats with more than 100
		passengers [10] Deportation of Chinese nationals

[11] Individual movement restrictions

[12] National lockdown

[13] Non-essential travels forbidden

[14] Outgoing domestic travel ban

[15] Outgoing oversea travel ban

[16] Partial cancellation of international flights

[17] Restriction of freight transport

[18] Stop public transports

[19] Train transport restrictions

[20] Travel restriction to high-risk areas

PPE: Personal Protective Equipment.

Table 2. Summary of the government interventions recorded in the CCCSL at level 1 (themes) of the coding scheme. As of date of submission, the dataset includes 53 countries (including the Diamond Princess cruise ship) and dates of NPI implementation range from 31/12/2019 to 25/04/2020.

Theme (L1)	Number of records	Frequency
Case identification, contact tracing and related	438	0.13
measures		
Environmental measures	32	0.01
Healthcare and public health capacity	477	0.15
Resource allocation	134	0.04
Returning to normal life	12	0.00
Risk communication	634	0.19
Social distancing	1,081	0.33
Travel restriction	466	0.14

Table 3. Top 20 most frequent categories (L2) of NPIs implemented to control the spread of COVID-19. As of date of submission, the dataset includes 53 countries (including the Diamond Princess cruise ship) and dates of NPI implementation range from 31/12/2019 to 25/04/2020.

Rank	Category (L2)	Number of records	Frequency	Examples of associated subcategories (L3)
1	Educate and actively communicate with the public	501	0.15	Communication via social media, TV, radio; Campaign for encouraging hand hygiene; Campaign for discouraging gathering; Campaign for respecting respiratory etiquette; Campaign for encouraging self- initiated quarantine; Campaign to promote the 2m distance.
2	Mass gathering cancellation	375	0.11	Cancellation of culture events; Cancellation of indoor activities (e.g. fitness gyms, swimming pools, music classes); Closure of place of worship; Sport events; Festivals, faith-based events; Requirement on the number of persons allowed.
3	Closure of educational institutions	236	0.07	Kindergartens; Schools; Colleges; Universities; Precise complete or partial closure.
4	Small gathering cancellation	205	0.06	Closure of restaurants, bars, cafes; Closure of non- essential shops; Mandatory home office; Family celebrations prohibited; Complete prohibition of gathering; Police and army control and sanctions; Precise the number of persons allowed.
5	Border restriction	140	0.04	Borders closed for people from China; Borders closed for people from high-risk areas other than China; Borders closed to all non-citizens; Borders closed for travellers but open for food supply; Borders closed to refugees; Border control.
6	Quarantine	125	0.04	Asymptomatic carriers; Contacts; Suspected cases; People who travelled to China or high-risk areas.

7	Measures for special populations	121	0.04	Measures to limit contact with the elderly; More exposed professionals (not healthcare), e.g. post officers, public transport drivers; Persons in prison; Vulnerable persons (e.g. homeless).
8	Crisis management plans	98	0.03	Develop all-of-society and business continuity plans; Implement repurpose community services plans; Implement repurpose government plans; Research funding.
9	Enhance emergency response mechanisms	93	0.03	Set up crisis unit (the mode is described for each government).
10	Enhance detection system	83	0.03	Broaden definition / special definition (L4: "of suspected cases"); Research and test of contacts; Research and test of suspected cases; Search for and/or monitor contacts; Test people who travelled in high-risk areas; Test vulnerable population.
11	Airport health check	80	0.02	Measure of the temperature; Test of all incoming people with fever or symptoms; Health screening; Health certificate requested.
12	Individual movement restrictions	75	0.02	Curfew; Non-essential movements forbidden.
13	Airport restriction	67	0.02	Denial entry of travellers from China; Denial entry of travellers from high-risk countries other than China; Specific health channel for travellers from high-risk areas; Denial entry of travellers from Wuhan.
14	Increase availability of PPE	65	0.02	Prohibition of export of PPE; Increase availability for healthcare professionals; L3 can precise the type of PPE (face masks, gloves, protective suits) while L4 can precise mean for increasing availability (e.g. import, increase production).
15	Increase patient capacity	59	0.02	Number of beds; Build emergency hospitals; Increase ICU capacity; Increase the number of isolation wards; Postponement of non-essential care and non-urgent operations in hospitals.

16	Return operation of nationals living in regions at risk	55	0.02	Precise country/region.
17	Enhance laboratory testing	51	0.02	Decrease time between test and result; Increase testing capacity (number tests).
18	Border health check	46	0.01	Measure of the temperature; Health screening; Health certificate requested; Health declaration; Test at the border.
19	Increase healthcare workforce	44	0.01	Mobilization of domestic resources for health; Exception to work law allowed; Train medical staff specially for covid-19; Travelling ban for healthcare professionals; Ease transport for health workers.
20	Travel alert	40	0.01	Level 1; Level 2; Level 3 (concerned country is specified)

References

- 1 Anderson, R. M., Heesterbeek, H., Klinkenberg, D. & Hollingsworth, T. D. How will country-based mitigation measures influence the course of the COVID-19 epidemic? *Lancet* **395**, 931-934, doi:10.1016/S0140-6736(20)30567-5 (2020).
- 2
 Ugarov, A. Inclusive Costs of NPI Measures for COVID-19 Pandemic: Three Approaches.
 medRxiv,
 2020.2003.2026.20044552,

 doi:10.1101/2020.03.26.20044552 (2020).
- 3 Studdert, D. M. & Hall, M. A. Disease control, civil liberties, and mass testing Calibrating restrictions during the Covid-19 pandemic. *N. Engl. J. Med.*, doi:10.1056/NEJMp2007637 (2020).
- 4 Pan, A. *et al.* Association of public health interventions with the epidemiology of the COVID-19 outbreak in Wuhan, China. *JAMA*, doi:10.1001/jama.2020.6130 (2020).
- 5 Chen, K.-T. *et al.* SARS in Taiwan: an overview and lessons learned. *Int. J. Infect. Dis.* **9**, 77-85, doi:<u>https://doi.org/10.1016/j.ijid.2004.04.015</u> (2005).
- 6 Hung, L. S. The SARS epidemic in Hong Kong: what lessons have we learned? J. R. Soc. Med. 96, 374-378, doi:10.1258/jrsm.96.8.374 (2003).
- 7 Cowling, B. J. *et al.* Impact assessment of non-pharmaceutical interventions against coronavirus disease 2019 and influenza in Hong Kong: an observational study. *Lancet Public Health*, doi:10.1016/S2468-2667(20)30090-6.
- 8 Wang, C. J., Ng, C. Y. & Brook, R. H. Response to COVID-19 in Taiwan: Big Data Analytics, New Technology, and Proactive Testing. *JAMA* **323**, 1341-1342, doi:10.1001/jama.2020.3151 (2020).
- 9 Folkhalsomyndighet. <<u>https://www.folkhalsomyndigheten.se/nyheter-och-press/</u>> (2020).
- 10 Rocklov, J. COVID-19 health care demand and mortality in Sweden in response to non-pharmaceutical (NPIs) mitigation and suppression scenarios. *medRxiv*, doi:10.1101/2020.03.20.20039594 (2020).
- 11Unesco.COVID-19EducationalDisruptionandResponse.(2020).<https://en.unesco.org/covid19/educationresponse>.
- 12 ISC Research. Coronavirus (COVID-19) update. The latest news about the international school closures. (2020). <<u>https://www.iscresearch.com/cornavirus-covid-19-update</u>>.
- 13IATA.AirTransport& COVID-19Coronavirus.(2020).<https://www.iata.org/en/programs/safety/health/diseases/>.
- 14 World Trade Organization. COVID-19: Trade and trade-related measures. (2020). <<u>https://www.wto.org/english/tratop_e/covid19_e/trade_related_goods_measure_e.htm</u>>.

- 15 PwC. Navigate Tax, Legal and Economic Measures in response to COVID-19. (2020). <<u>https://www.pwc.com/gx/en/issues/crisis-solutions/covid-19/response-</u> <u>navigator.html</u>>.
- 16 Hogan Lovells. The EU and various member states measures on Personal Protective Equipment in response to the COVID-19 outbreak. (2020). <<u>https://www.hoganlovells.com/en/publications/the-eu-and-various-memberstates-measures-on-personal-protective-equipment-in-response-to-the-covid-19outbreak>.</u>
- 17 World Customs Organization. List of national legislation* of countries that adopted temporary export restrictions on certain categories of critical medical supplies in response to COVID-19. (2020).
 <<u>http://www.wcoomd.org/en/topics/facilitation/activities-and-programmes/natural-disaster/list-of-countries-coronavirus.aspx</u>>.
- 18 U.S. Chamber of Commerce. Global Dashboard on COVID-19 Government Policies. (2020). <<u>https://www.uschamber.com/international-affairs-division/covid-dashboard</u>>.
- 19 Hale, T., Webster, S., Petherick, A., Phillips, T. & Kira, B. Oxford COVID-19 Government Response Tracker, Blavatnik School of Government. (2020). <<u>https://www.bsg.ox.ac.uk/research/research-projects/coronavirus-government-response-tracker</u>>.
- 20 Molloy, J. C. The open knowledge foundation: open data means better science. *PLoS Biol.* **9**, e1001195, doi:10.1371/journal.pbio.1001195 (2011).
- Vaismoradi, M., Turunen, H. & Bondas, T. Content analysis and thematic analysis: Implications for conducting a qualitative descriptive study. *Nursing & Health Sciences* 15, 398-405, doi:10.1111/nhs.12048 (2013).
- 22 Erlingsson, C. & Brysiewicz, P. A hands-on guide to doing content analysis. *Afr J Emerg Med* 7, 93-99, doi:<u>https://doi.org/10.1016/j.afjem.2017.08.001</u> (2017).
- 23 Weber, R. *Basic Content Analysis*. 2 edn, (1990).
- 24 Gläser, J. & Laudel, G. Life with and without coding: Two methods for early-stage data analysis in qualitative research aiming at causal explanations. 2013 14, doi:10.17169/fqs-14.2.1886 (2013).
- 25 Windsor, L. C., Cupit, J. G. & Windsor, A. J. Automated content analysis across six languages. *PLOS ONE* 14, e0224425, doi:10.1371/journal.pone.0224425 (2019).
- 26 R Core Team. R: A language and environment for statistical computing <<u>https://www.R-project.org/</u>> (R Foundation for Statistical Computing, Vienna, Austria, 2020).
- 27 Vaismoradi, M., Jones, J., Turunen, H. & Snelgrove, S. Theme development in qualitative content analysis and thematic analysis. *Journal of Nursing Education and Practice* **6**, 100-110, doi:10.5430/jnep.v6n5p100 (2016).

- 28 Wickham, H., François, R., Henry, L. & Müller, K. *dplyr*: A grammar of data manipulation <<u>https://CRAN.R-project.org/package=dplyr</u>> (2020).
- 29 Centers for Disease Control and Prevention. Interim pre-pandemic planning guidance : community strategy for pandemic influenza mitigation in the United States : early, targeted, layered use of nonpharmaceutical interventions. (Centers for Disease Control and Prevention, 2007).
- 30 European Centre for Disease Prevention and Control. Guide to revision of national pandemic influenza preparedness plans Lessons learned from the 2009 A(H1N1) pandemic. (ECDC, 2017).
- 31 World Health Organization. Non-pharmaceutical public health measures for mitigating the risk and impact of epidemic and pandemic influenza. (2019).
- 32 European Centre for Disease Prevention and Control. Download today's data on the geographic distribution of COVID-19 cases worldwide. (2020). <<u>https://www.ecdc.europa.eu/en/publications-data/download-todays-data-</u> geographic-distribution-covid-19-cases-worldwide>.
- 33 Johns Hopkins University Center for Systems Science and Engineering. 2019 Novel Coronavirus COVID-19 (2019-nCoV) Data Repository by Johns Hopkins CSSE. (2020). <<u>https://github.com/CSSEGISandData/COVID-19</u>>.
- 34 European Centre for Disease Prevention and Control. Technical report. Considerations relating to social distancing measures in response to COVID-19 – second update. (ECDC, 2020).
- 35 World Health Assembly. *International Health Regulations (2005)*. (World Health Organization, 2016).
- 36 Kinlaw, K. & Levine, R. J. Ethical guidelines in Pandemic Influenza— Recommendations of the Ethics Subcommittee of the Advisory Committee to the Director, Centers for Disease Control and Prevention. (Centers for Disease Control and Prevention, 2007).



Map based on Longitude (generated) and Latitude (generated). Color shows sum of Measure_L1. Details are shown for Group.1.





Country



