

Title: Characteristics of patients presenting, and not presenting, to the emergency department during the early days of COVID-19

Bjorn C. Westgard, MD, MA
Regions Hospital, HealthPartners Institute

Matthew W. Morgan, MD
Regions Hospital

Gabriela Vazquez-Benitez, PhD, MSC
HealthPartners Institute

Lauren O. Erickson, MS
HealthPartners Institute

Michael D. Zwank, MD
Regions Hospital

Abstract

Objective: Societal responses to the COVID-19 pandemic have had a substantial effect upon the number of patients seeking healthcare. An initial step in estimating the impact of these changes is characterizing the patients, visits, and diagnoses for whom care is being delayed or deferred.

Methods: We conducted an observational study, examining demographics and diagnoses for all patient visits to the ED of an urban Level-1 trauma center before and after the state declaration and compared them to visits from a similar period in 2019. We estimated the ratios of the before and after periods using Poisson regression, calculated the percent change with respect to 2019 for total ED visits, patient characteristics, and diagnoses, and then evaluated the interactions between each factor and the overall change in ED visits.

Results: There was a significant 35.2% drop in overall ED visits after the state declaration. Disproportionate declines were seen in visits by pediatric and older patients, women, and Medicare recipients as well as for presentations of syncope, cerebrovascular accidents, urolithiasis, abdominal and back pain. Significantly disproportionate increases were seen in ED visits for potential symptoms of COVID-19, including URIs, shortness of breath, and chest pain.

Conclusions: Patient concerns about health care settings and public health have significantly altered care-seeking during the COVID-19 pandemic. Overall and differential declines in ED visits for certain demographic groups and disease processes should prompt efforts to encourage care-seeking and research to monitor for the morbidity and mortality that is likely to result from delayed or deferred care.

Introduction

In response to the spread of the novel coronavirus 2019 (COVID-19) pandemic, state governments and health systems have enacted a range of mitigation strategies and operational changes to anticipate and address an increasing number of patients with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). At the same time, during the early days of the pandemic, health systems have also seen a decrease in the number of patients presenting for acute care unrelated to COVID-19¹. Given public health recommendations to mitigate the spread of COVID-19, including social distancing and stay-at-home orders, many patients are not seeking and receiving care for acute conditions which may put them at significant risk for preventable morbidity and mortality in the future. The characteristics of those patients who are and are not presenting to emergency departments as a result of stay at home orders during the early days of the SARS-CoV-2 pandemic have not yet been substantially examined in the medical literature. We report changes in the characteristics of patients and presentations to the emergency department (ED) of an urban Level-1 Trauma Center before and after the statewide announcement of “peacetime emergency” public health measures to respond to the pandemic on March 13, 2020².

Methods

We conducted an observational study of visits to the ED from February 15 to April 10, 2020, a period of 28 days before through 28 days after the state’s announcement. Accounting for days of the week, we then examined a cross-section of visits from a similar period in 2019 for historical comparison. We estimated the ratios of the before and after periods using Poisson regression and calculated the percent change with respect to 2019 for the total number of ED visits and for each

of the patient characteristics and diagnoses. We calculated type III p-values to evaluate the interactions between each of those factors and the overall change in ED visits. Data were collected as part of institutional operations and quality improvement and were therefore deemed by the institutional review board to be exempt from review.

Results

After the state declaration, the ED experienced a significant decline from an average of 250 visits daily visits for the 28 days before to an average of 167 daily visits for the 28 days after ($p<0.001$). This represents a decline of 35.2% in ED visits overall, or a daily decline of 4 visits, with respect to 2019. When we looked at billing, there was a 23% reduction in ED charges (95% CI 19-27%) and a 13% reduction in related hospital charges (9-17%) with respect to 2019 when adjusted for inflation.

We found significant changes in the distribution of patient demographics and visit characteristics after the state declaration, particularly in patient age, gender, race, insurance, arrival mode, and disposition. There were notable decreases in ED visits by patients under age 18 (-60.1%, 95% CI -68.0 to -50.2%, $p<0.001$) and over age 65 (-41.3%, -33.7 to -48.0%, $p<0.001$), women (-40.2%, -44.4 to -35.8%, $p<0.001$), African American and white patients (-37.0%, -42.6 to -30.7%; -37.0%, -42.6 to -30.7%, $p<0.001$), Medicare recipients (-40.8%, -47.8 to -32.7%, $p<0.001$) and those with other insurances (e.g. liability, no-fault, workman's compensation, -74.1%, -80.0 to -66.5%, $p<0.001$), ambulatory patients (-38.1%, -41.8 to -34.3%, $p=0.021$), and those who left prior to evaluation or discharge (-75.6%, -82.5 to -66.0%, $p=0.021$).

When we examined diagnoses, we noted significantly disproportionate decreases in patients presenting with syncope (-70.5%, -53.4 to -81.3%, $p<0.001$), cerebrovascular accidents (-58.3%, -73.3 to -34.9% $p=0.049$), abdominal pain (-43.3%, -50.8 to -34.8%, $p=0.045$), urolithiasis (-70.0%, -84.0 to -43.6%, $p=0.014$), and back pain (-50.7, -61.1 to -37.4%, $p=0.022$). We also saw significant increases in the proportion of patients presenting with upper respiratory infections (URIs, -10.0%, -29.4 to 14.8%, $p=0.007$), shortness of breath (25.1%, -6.4 to 67.3%, $p<0.001$), and chest pain (-13.1%, -27.2 to 3.7%, $p<0.001$). Of note, proportional increases were also seen for all of the mental health and substance-use diagnoses we examined, though these were not significantly disproportionate. For all other conditions, the declines in presentations were proportionate to the overall change in ED visits.

Discussion

Paralleling anecdotal reports, we noted a significant temporal association between our state's emergency declaration and an overall decline in daily ED visits. We found significant changes in ED patient demographics. The significant proportional decreases in visits by the overlapping populations of patients over 65 and Medicare recipients are likely a product of the extra prudence of higher risk individuals who are social distancing and staying at home. The significant proportional declines in visits by pediatric, ambulatory, and other patients likely reflect concerns about contracting COVID-19 in health care settings, overburdening the health care system with unrelated complaints, and adhering to public health recommendations. The decrease in number of patients who left prior to evaluation or discharge is likely due at least in part to the substantial operational changes made in response to the pandemic which have further streamlined patient triage, rooming, and evaluation.

The most alarming finding of this report, however, is simply the overall decline in care-seeking for acute and potentially life-threatening conditions unrelated to COVID-19¹. Societal responses to the COVID-19 pandemic have contributed to multiple phenomena, including diminished air pollution, traffic, and infectious disease transmission, which could all potentially contribute to improved population health over the long term¹. These phenomena may well influence exacerbations of respiratory conditions as well as COVID-19 infections and outcomes over the short term³, but most of the conditions seen and managed in the emergency department would not be expected to suddenly decrease in incidence. Variable disease severity could play a role, as patients with less acute conditions might be able to delay or defer seeking care. Though we did not examine low acuity presentations that might also be managed in other outpatient settings, we did find a disproportionate decrease in presentations of back pain and other non-specific pains. However, we also found unexpectedly disproportionate declines in visits for conditions of significantly higher acuity like syncope, cerebrovascular accidents, and urolithiasis, similar to those noted elsewhere for presentations of myocardial infarction^{4,5}. Additionally, we saw increases in ED visits for upper respiratory infections, chest pain, or shortness of breath, symptoms that might bring patients to the ED with concerns for COVID-19 infection. Similar trends in patient presentations have been seen around the world⁶, during prior disease outbreaks⁸, and in the outpatient setting⁸.

These are early days, and the timeline for the COVID-19 pandemic and resulting changes in patterns of ED utilization promise to be much longer than that covered in our preliminary study. This study is limited by its single center and cross-sectional nature, the short study time frame,

and lack of adjustment for multiple potentially confounding factors related to patients and their presentations to the ED, including disease severity, comorbidities, and COVID-19 risk factors. In spite of those limitations, we have found notable differential changes in the demographic factors, visits characteristics, and diagnoses of presentations to the ED unrelated to COVID-19. These will no doubt change along with the burdens of care as the pandemic progresses. Further efforts should and are being made to reassure and affirm the appropriateness of seeking emergency care⁹, particularly for the groups and disease processes that have been highlighted here and elsewhere^{4,5}. Further research will also be needed to examine these and other factors contributing to delayed or deferred care and to monitor for the morbidity and mortality that is likely to result^{5,8} and which may already be occurring¹⁰.

Acknowledgement: We would like to thank Hrafn P. Gudjonsson (“Patrick”) for his valuable assistance in querying and creating the data set used in this study.

References

1. Kolata, G. Amid the Coronavirus Crisis, Heart and Stroke Patients Go Missing. *New York Times*. Published online, April 25, 2020.
<https://www.nytimes.com/2020/04/25/health/coronavirus-heart-stroke.html>. Accessed April 27, 2020.
2. Bakst B. Walz advises no gatherings over 250 people as MN COVID-19 cases hit 14. *Minnesota Public Radio News*. Published online March 13, 2020.
<https://www.mprnews.org/story/2020/03/13/14-covid19-cases-now-in-mn-walz-to-unveil-new-control-measures>. Accessed April 20, 2020.
3. Exposure to air pollution and COVID-19 mortality in the United States. Xiao Wu, Rachel C. Nethery, Benjamin M. Sabath, Danielle Braun, Francesca Dominici. *medRxiv* 2020.04.05.20054502; doi: <https://doi.org/10.1101/2020.04.05.20054502>
4. Garcia S, Albaghdadi MS, Meraj PM, et al. Reduction in ST-Segment Elevation Cardiac Catheterization Laboratory Activations in the United States during COVID-19 Pandemic. *Journal of the American College of Cardiology*. Published online, April 7, 2020.
<https://doi.org/10.1016/j.jacc.2020.04.011>.
5. Tam DF, Cheung K, Lam S, et al. Impact of Coronavirus Disease 2019 (COVID-19) Outbreak on ST-Segment–Elevation Myocardial Infarction Care in Hong Kong, China. *Circ Cardiovasc Qual Outcomes*. 2020;13:e006631. Doi: [10.1161/circoutcomes.120.006631](https://doi.org/10.1161/circoutcomes.120.006631).

6. Ong CW, Goletti D. Impact of the global COVID-19 outbreak on the management of other communicable diseases. *International Journal of Tuberculosis and Lung Disease*. Published online, March 20, 2020. <http://dx.doi.org/10.5588/ijtld.20.0140>.
7. Huang YT, Lee YC, Hsiao CJ. Hospitalization for ambulatory-care-sensitive conditions in Taiwan following the SARS outbreak: a population-based interrupted time series study. *J Formos Med Assoc*. 2009;108(5):386–394. doi:10.1016/S0929-6646(09)60082-6.
8. Rosenbaum L. The Untold Toll — The Pandemic’s Effects on Patients without Covid-19. *New England Journal of Medicine*. Published online, April 17, 2020. doi:10.1056/NEJMms2009984.
9. American College of Emergency Physicians. Amidst COVID-19 Concerns, Emergency Physicians Urge Public Not to Delay Necessary Medical Care. Press release. Published online April 15, 2020. <https://www.emergencyphysicians.org/press-releases/2020/4-15-20-amidst-covid-19-concerns-emergency-physicians-urge-public-not-to-delay-necessary-medical-care>. Accessed April 20, 2020.
10. Gillum J, Song L, Kao J. There’s Been a Spike in People Dying at Home in Several Cities; That Suggests Coronavirus Deaths Are Higher Than Reported. *Propublica*. Published online April 14, 2020. <https://www.propublica.org/article/theres-been-a-spike-in-people-dying-at-home-in-several-cities-that-suggests-coronavirus-deaths-are-higher-than-reported>. Accessed April 20, 2020.




Figure 1. Trend of daily ED visits before and after state announcement

	Mar 14 – Apr 10, 2020	Feb 15 – Mar 13, 2020	Mar 16 – Apr 12, 2019	Feb 16 – Mar 15, 2019	% change 2020 with respect to 2019 (95% CI)	p-value ^a
Total No. of ED visits	4666	6993	6744	6547	-35.2 (-38.4 to -32.9)	
Demographics	N(%)	N(%)	N(%)	N(%)		
Age, median (IQR, range), y	42 (29-59, 0-104)	42 (29-60, 0-102)	44 (28-58, 0-100)	44 (28-59, 0-106)		
0-18 years	177 (3.8)	436 (6.2)	427 (6.3)	420 (6.4)	-60.1 (-68.0 to -50.2)	<0.001
18<65 years	3723 (79.8)	5246 (75.0)	5186 (76.9)	4990 (76.2)	-31.7 (-35.5 to -27.7)	
65≤ years	766 (16.4)	1311 (18.7)	1131 (16.8)	1137 (17.4)	-41.3 (-33.7 to -48.0)	
Sex						
Female	2171 (46.6)	3585 (51.3)	3386 (50.2)	3342 (51.0)	-40.2 (-44.4 to -35.8)	<0.001
Male	2492 (53.4)	3408 (48.7)	3358 (49.8)	3205 (49.0)	-30.2 (-35.0 to -25.1)	
Race						
Asian	316 (6.8)	475 (6.8)	416 (6.2)	374 (5.7)	-40.2 (-51.0 to -27.0)	0.035
African American	1304 (28.0)	2005 (28.7)	1962 (29.1)	1902 (29.1)	-37.0 (-42.6 to -30.7)	

White	2302 (49.3)	3503 (50.1)	3489 (51.2)	3305 (50.5)	-37.8 (-42.0 to -33.2)	
Hispanic or Latino	323 (6.9)	477 (6.8)	403 (6.0)	456 (7.0)	-23.4 (-36.9 to -6.9)	
Native American	62 (1.3)	99 (1.4)	95 (1.4)	114 (1.7)	-24.9 (-50.5 to 14.2)	
Native Hawaiian or OPI	13 (0.3)	16 (0.2)	19 (0.3)	21 (0.3)	-10.2 (-65.6 to 134.4)	
Other/multiracial	346 (7.4)	418 (6.0)	360 (5.3)	375 (5.7)	-13.8 (-29.6 to 5.6)	
Insurance						
Commercial	2322 (49.8)	3452 (49.4)	3261 (48.4)	3315 (50.6)	-31.6 (-36.3 to -26.6)	
Medicare	654 (14.0)	1128 (16.1)	1095 (16.2)	1119 (17.1)	-40.8 (-47.8 to -32.7)	
Medicaid	782 (16.8)	1175 (16.8)	1242 (18.4)	1193 (18.2)	-36.1 (-43.3 to -27.9)	<0.001
Self-pay	759 (16.3)	936 (13.4)	757 (11.2)	710 (10.8)	-23.9 (-33.9 to -12.5)	
Other	145 (3.1)	301 (4.7)	391 (5.7)	210 (3.2)	-74.1 (-80.0 to -66.5)	
Arrival						
Ambulatory	3151 (67.5)	4872 (69.7)	4783 (70.9)	4575 (69.9)	-38.1 (-41.8 to -34.3)	
EMS, helicopter, fire	1417 (30.4)	2000 (28.6)	1812 (26.9)	1823 (27.8)	-28.7 (-35.1 to -21.7)	0.021
Police	98 (2.1)	121 (1.7)	149 (2.2)	149 (2.3)	-19.0 (-42.9 to 14.9)	
ESI Acuity						
1 (high)	103 (2.2)	139 (2.0)	128 (1.9)	122 (1.9)	-29.4 (-50.5 to 0.8)	
2	1366 (29.3)	2198 (31.4)	2085 (30.9)	2096 (32)	-37.5 (-42.9 to -31.6)	
3	2252 (48.3)	3186 (45.6)	3271 (48.5)	2996 (45.8)	-35.3 (-39.8 to -30.3)	0.533
4	760 (16.3)	1216 (17.4)	1063 (15.8)	1109 (16.9)	-34.8 (-42.4 to -26.2)	
5 (low)	156 (3.3)	224 (3.2)	169 (2.5)	197 (3)	-18.8 (-39.2 to 8.5)	
Unknown	29 (0.6)	30 (0.4)	28 (0.4)	27 (0.4)		
Disposition						
Hospitalization	1230 (26.4)	1811 (25.9)	1787 (26.5)	1658 (25.3)	-37.0 (-42.9 to -30.5)	
Discharged	3360 (72.0)	4933 (70.5)	4712 (69.9)	4661 (71.2)	-32.6 (-36.5 to -28.5)	<0.001
AMA, eloped, LWBS	64 (1.4)	240 (3.4)	237 (3.5)	217 (3.3)	-75.6 (-82.5 to -66.0)	
Expired	12 (0.3)	9 (0.1)	8 (0.1)	11 (0.2)		
Diagnoses						
Dizziness (6) ^{bc}	103 (2.3)	178 (2.6)	190 (2.9)	166 (2.6)	-49.4 (-30.4 to -63.3)	0.123
Syncope (9) ^c	44 (1.0)	115 (1.7)	101 (1.5)	78 (1.2)	-70.5	0.001

					(-53.4 to -81.3)	
Headache (2) ^c	152 (3.3)	215 (3.1)	303 (4.5)	242 (3.7)	-43.5 (-56.8 to -26.2)	0.301
Cerebrovascular accident ^d	52 (1.1)	104 (1.5)	96 (1.4)	80 (1.2)	-58.3 (-73.3 to -34.9)	0.049
Chest pain (1) ^c	456 (9.8)	528 (7.6)	492 (7.3)	495 (7.6)	-13.1 (-27.2 to 3.7)	<0.001
STEMI/NSTEMI/Angina ^d	92 (2.0)	121 (1.7)	142 (2.1)	171 (2.6)	-8.4 (-35.5 to 30.0)	0.051
VTE disease ^c	24 (0.5)	38 (0.5)	40 (0.6)	43 (0.7)	-32.1 (-65.2 to 32.5)	0.890
Congestive heart failure ^c	115 (2.5)	144 (2.1)	134 (2.0)	145 (2.2)	-13.6 (-38.5 to 21.4)	0.093
Shortness of breath	261 (5.6)	198 (2.8)	157 (2.3)	149 (2.3)	25.1 (-6.4 to 67.3)	<0.001
URI/pharyngitis/sinusitis (7) ^d	247 (5.3)	282 (4.0)	253 (3.8)	260 (4.0)	-10.0 (-29.4 to 14.8)	0.007
Asthma/COPD ^d	179 (3.8)	264 (3.8)	253 (3.8)	261 (4.0)	-30.0 (-45.9 to -9.6)	0.550
Influenza/pneumonia ^d	152 (3.3)	358 (5.1)	153 (2.3)	215 (3.3)	-40.3 (-55.0 to -21.0)	0.531
COVID-19/coronavirus ^c	20 (0.4)	0 (0)	0 (0)	0 (0)		--
Abdominal pain (4) ^c	579 (12.4)	958 (13.7)	863 (12.8)	809 (12.4)	-43.3 (-50.8 to -34.8)	0.045
Appendicitis ^d	15 (0.3)	21 (0.3)	18 (0.3)	21 (0.3)	-16.7 (-66.6 to 107.85)	0.589
GB/biliary/pancreas ^d	49 (1.1)	66 (0.9)	86 (1.3)	75 (1.1)	-35.3 (-60.0 to 4.9)	0.998
Renal stone/colic ^d	22 (0.5)	52 (0.7)	62 (0.9)	44 (0.7)	-70.0 (-84.0 to -43.6)	0.014
Urinary tract infection ^d	156 (3.3)	240 (3.4)	243 (3.6)	214 (3.3)	-42.8 (-56.4 to -24.8)	0.365
Sepsis (3) ^c	148 (3.2)	183 (2.6)	146 (2.2)	154 (2.4)	-14.7 (-37.6 to 16.7)	0.081
Diabetes complications ^d	176 (3.9)	280 (4.2)	291 (4.5)	267 (4.3)	-42.3 (-55.1 to -25.85)	0.354
Diabetic ketoacidosis ^c	29 (0.6)	32 (0.5)	20 (0.3)	26 (0.4)	17.8 (-45.4 to 154.4)	0.126
Injuries ^d	607 (13.0)	1005 (14.4)	933 (13.8)	1080 (16.5)	-30.1 (-38.8 to -20.1)	0.212
Head injury (5) ^d	254 (5.8)	419 (6.4)	378 (5.9)	427 (7.0)	-31.5 (-44.4 to -15.7)	0.587
Fractures ^c	161 (3.5)	304 (4.3)	260 (3.9)	321 (4.9)	-34.6 (-49.2 to -15.9)	0.933
Dental complaints ^c	61 (1.3)	91 (1.3)	111 (1.6)	93 (1.4)	-43.8 (-63.3 to -14.1)	0.507
Back pain (8) ^d	165 (3.5)	342 (4.9)	349 (5.2)	357 (5.5)	-50.7 (-61.1 to -37.4)	0.022
Anxiety/stress ^d	195 (4.2)	251 (3.6)	292 (4.3)	249 (3.8)	-33.8 (-48.5 to -14.8)	0.860
Depression ^d	227 (5.1)	307 (4.6)	342 (5.3)	300 (4.8)	-35.1 (-48.5 to -18.3)	0.992
Suicidal ideation (10) ^c	179 (3.8)	201 (2.9)	225 (3.3)	196 (3.0)	-22.4 (-41.3 to 2.4)	0.196

Intoxication/Substance Use ^d	765 (19.6)	1040 (17.5)	994 (17.3)	945 (16.9)	-30.1 (-38.5 to -20.4)	0.203
Alcohol withdrawal ^c	60 (1.3)	60 (0.9)	69 (1.0)	53 (0.8)	-23.2 (-53.7 to 27.4)	0.507
Poisoning ^d	101 (2.2)	117 (1.7)	101 (1.5)	95 (1.5)	-18.8 (-44.8 to 19.5)	0.247

Table 1. Characteristics of ED patients and presentations before and after state announcement

Abbreviations: IQR, inter-quartile range; OPI, other Pacific Islander; EMS, Emergency Medical Services; ESI, Emergency Severity Index; AMA, against medical advice; LWBS, left without being seen; STEMI, ST-elevation myocardial infarction; NSTEMI, non-ST-elevation myocardial infarction; VTE, venous thromboembolic; URI, upper respiratory infection; COPD, chronic obstructive pulmonary disease; COVID-19, novel coronavirus disease 2019; and GB, gallbladder.

^a P-value represents the type III p-value for the interaction between the overall change in ED visits and the change in each category of patient and visit characteristics and diagnoses.

^b Numbers indicate historical top 10 diagnoses for visits to our ED.

^c Determined from a search of diagnosis names within the first 3 coder diagnoses.

^d Determined from a search of diagnostic related groups within the first 3 coder diagnoses.

