

A Noel Intervention Recurrent autoencoder for real time forecasting and non-pharmaceutical intervention selection to curb the spread of Covid-19 in the world

Qiyang Ge¹, Zixin Hu^{2,3}, Shudi Li⁴, Wei Lin¹, Li Jin^{2,3} and Momiao Xiong^{4,*}

¹The School of Mathematical Sciences, Fudan University, Shanghai, China.

²State Key Laboratory of Genetic Engineering and Innovation Center of Genetics and Development, School of Life Sciences, Fudan University, Shanghai, China.

²Human Phenome Institute, Fudan University, Shanghai, China.

⁴Department of Biostatistics and Data Science, School of Public Health, The University of Texas Health Science Center at Houston, Houston, TX 77030, USA.

Running Title: Forecasting and interventions of Covid-19 in the world

*Address for correspondence and reprints: Dr. Momiao Xiong, Department of Biostatistics and Data Science, School of Public Health, The University of Texas Health Science Center at Houston, P.O. Box 20186, Houston, Texas 77225, (Phone): 713-500-9894, (Fax): 713-500-0900, E-mail: Momiao.Xiong@uth.tmc.edu.

Highlights

- As the Covid-19 pandemic soars around the world, there is urgent need to forecast the number of cases worldwide at its peak, the length of the pandemic before receding and implement public health interventions to significantly stop the spread of Covid-19.
- Develop artificial intelligence (AI) and causal inference inspired methods for real-time forecasting and evaluation of interventions on the worldwide trajectory of the spread of Covid-19.
- We estimated the maximum number of cumulative cases under immediate additional intervention to be 2,271,648; under later additional intervention the number increased to 3,864,872 and the case ending time would be May 25, 2020.
- Without additional intervention, the spread of COVID-19 would be stopped on July 6, 2020.

ABSTRACT

Objective: Develop the AI and casual inference-inspired methods for forecasting and evaluating the effects of public health interventions on curbing the spread of Covid-19.

Methods: We developed recurrent neural network (RNN) for modeling the transmission dynamics of the epidemics and Counterfactual-RNN (CRNN) for evaluating and exploring public health intervention strategies to slow down the spread of Covid-19 worldwide. We applied the developed methods to real-time forecasting the confirmed cases of Covid-19 across the world. The data were collected from January 22 to April 18, 2020 by John Hopkins Coronavirus Resource Center (<https://coronavirus.jhu.edu/MAP.HTML>).

Results: The average errors of 1-step to 10-step forecasting were 2.9%. In the absence of a COVID-19 vaccine, we evaluated the potential effects of a number of public health measures. We found that the estimated peak number of new cases and cumulative cases, and the maximum number of cumulative cases worldwide with one week later additional intervention were reduced to 103,872, 2,104,800, and 2,271,648, respectively. The estimated total peak number of new cases and cumulative cases would be the same as the above and the maximum number of cumulative cases would be 3,864,872 in the world with 3 week later additional intervention. Duration time of the Covid-19 spread would be increased from 91 days to 123 days. Our estimation results showed that we were in the eve of stopping the spread of COVID-19 worldwide. However, we observed that transmission would quickly rebound if interventions were relaxed.

Conclusions: The accuracy of the AI-based methods for forecasting the trajectory of Covid-19 was high. The AI and causal inference-inspired methods are a powerful tool for helping public health planning and policymaking. We concluded that the spread of COVID-19 would be stopped very soon.

Keywords: Cov-19, artificial intelligence, recurrent neural networks, causal inference, forecasting, time series.

Introduction

As of April 28, 2020, global confirmed cases of Covid-19 passed 3,116,398 and has spread to 210 countries, causing fear globally (Anastassopoulou et al. 2020). The serious public health threat of Covid-19 has never been seen for more than one century. The government officers and people around the world are desperately trying to slow the spread of Covid-19 (Irfan, 2020). We must change our policies to deal with increased mobility of citizens and immediately implement the public health interventions to stop the spread of Covid-19 across the world. How computer modeling of Covid-19's transmission dynamics could help governments to quickly and strongly move slow down the spread of Covid-19?

Widely used statistical and computer methods for modeling of Covid-19 simulate the transmission dynamics of epidemics to understand their underlying mechanisms, forecast the trajectory of epidemics, and assess the potential impact of a number of public health measures on curbing the spread speed of Covid-19 (Li et al., 2020, Wu et al., 2020, Zhao et al., 2020, Kucharski et al., 2020, Tuite et al., 2020, Hellewell et al., 2020, Li et al. 2020). Although these epidemiological models are useful for estimating the dynamics of transmission, and evaluating the impact of intervention strategies, they have some serious limitations (Funk et al., 2018, Johansson et al., 2019). First, the epidemiological models consist of ordinary differential equations that have many unknown parameters, and depend on many assumptions. It is difficult to translate public interventions to these parameters. Most analyses used hypothesized parameters, which often lead to fitting data very poor. Health officers desperately want to track

the trajectory of epidemics and accurately estimate the peak time and number of cases, duration, and ending time and number of cases of Covid-19 for their health policy plan. However, the forecasting results of using the classical epidemiological models such as Susceptible-Exposed-Infectious-Removed (SEIR) models are highly unreliable. Second, the successful application of public health intervention planning highly depends on the model parameter identifiability. However, overall, the parameters in the complex compartmental dynamic models are unidentifiable (Roosa and Chowell, 2019, Roda et al., 2020). The values of parameters cannot be uniquely determined from the real data (Gábor et al., 2017). The variances of the estimators of these parameters are very high.

To overcome limitations of the epidemiological model approach, and assist public health planning and policy making, we formulated the real-time forecasting and evaluating multiple public health intervention problem into forecasting treatment response problem. We viewed the interventions as treatments where multiple interventions were administered at different time points. The number of new cases were taken as treatment responses. The ability to accurately estimate effects of public health interventions over time would allow health officers to determine what intervention strategies should be used and the optimal time at which to implement them (Lim et al., 2018). Recurrent Intervention Network (RIN) (Lim et al., 2018) where a recurrent neural network architecture for forecasting a nation's response (number of new cases) to a series of planned interventions were used to forecast and evaluate multiple public health interventions for Covid-19 worldwide. Potential outcomes of RIN were trajectory of the spread of Covid-19. Public health interventions including locking down residential buildings and compounds, strict self-quarantine for families, door-to-door inspection for suspected cases, matining social distancing, stopping mass gatherings, closure of schools and universities, vacating hotels and

university dormitories. To quantify comprehensive intervention strategies, an intervention variable was used as an input variable for each block of RIN. We cluster all the countries in the world into several groups. For each group, a value (weight) was assigned to each group such that the average prediction error of CRN was minimum. The RIN is taken as a general framework for investigating how Covid-19 evolves under different intervention plans, how individual nation responds to intervention over time, but also which are optimal timings for assigning interventions. Therefore, this approach will provide new tools to improve public health planning and policy making.

The RIN was applied to the surveillance data of confirmed and new Covid-19 cases in the world up to April 7, 2020. Data on the number of confirmed, new and death cases of Covid-19 from January 22, 2020 to April 7, 2020 were obtained from John Hopkins Coronavirus Resource Center (<https://coronavirus.jhu.edu/MAP.HTML>).

Methods

RIN as a Framework for modeling and forecasting the spread of Covid-19 over time with multiple interventions

The RIN uses sequence-to-sequence multi-input/output recurrent neural network (RNN) architectures to model health intervention plan and make multi-step prediction of the response trajectory of Covid-19 over time with multiple interventions (Lim et al., 2018). The RNN can learn the complex dynamics within the temporal ordering of input time series of Covid-19 and use an internal memory to remember. The health intervention plan has multiple intervention regimens. As shown in Figure 1, the RIN determines the intervention response (similar to counterfactual outputs) for a given set of planned interventions and evaluates the impact of

different intervention strategies and their implementation times on the curbing the spread of Covid-19 and provides timely selection of optimal sequence of intervention strategies.

The RIN is a RNN autoencoder. It consists of two RNNs: the encoder RNN (standard RNN or long short-term memory (LSTM) is used as encoder) and the decoder RNN (standard RNN or LSTM is used as the decoder). The RNN encoder models input time series (past history of the number of cases of Covid-19 over time) and predicts future response time series (number of cases of Covid-19 in the future with a planned sequence of interventions) (Srivastava et al., 2015). The latent state of the RNN encoder after reading in the entire input time series (past trajectory of Covid-19), is the representation (compressed latent features of the entire input time series) of the input trajectory of Covid-19. Unlike the standard decoder where the decoder reconstructs back the input time series from the latent representation, the RNN decoder uses the learned features of the dynamics of Covid-19 in the RNN encoder to forecast the potential response time series, given a sequence of planned public health interventions as an input to the RNN decoder. The feature vector learned in the RNN encoder is then provided as an input to the RNN decoder which initiate prediction of the future dynamics of Covid-19 under the future interventions (Figure 2).

Training the RIN

RIN training consisted of RNN encoder training and RNN decoder training. We first introduced RNN encoder training procedures. The basic RNN unit in the RIN consisted of input layer, hidden layer and output layer. The input variables in the RNN encoder included covariates X_t such as density of population, traffic flow, health facility resources, GDP, and social-economic status, intervention variable A_t and the numbers of cases (potential outcomes) Y_t at the time t . The state in the hidden layer at the time t was denoted by h_t . The output layer had the output

variable Y_{t+1} . A nonlinear activation function was exponential linear unit (ELU) (Clevert et al., 2015) which was defined as

$$f(x) = \begin{cases} x & x > 0 \\ \alpha(e^x - 1) & x \leq 0 \end{cases},$$

where $\alpha > 0$.

The input data were divided into several batches with length of 7 days. Each batch was used to train the RNN encoder which forecasted standard one-step-ahead intervention response \hat{Y}_{t+1} as close to the observed intervention response Y_{t+1} as possible via the nonlinear mapping (Supplementary A)

$$\hat{Y}_{t+1} = f(X_t, A_t, Y_t, h_{t-1}). \quad (1)$$

The mean-squared error was used as loss function for training the RNN encoder. The training was performed via the standard propagation algorithms (Supplementary A). After training was completed, the RNN encoder extracted the hidden state h_t that captured the internal features of the transmission dynamics of Covid-19 via performing a feed-forward pass over the training data on the RNN encoder (Supplementary A).

After the RNN encoder training was completed, we began to train the RNA decoder. A RNN unit in the RNN decoder consisted of input layer with intervention variable $A_{t+\tau}$, ($\tau = 1, 2, \dots$), hidden layer with hidden state $Z_{t+\tau-1}$ and output $Y_{t+\tau+1}$. For a given country, observations (intervention A_t and the number of cases Y_t were randomly divided into short batches of up to τ_b time steps. Each batch of short sequence starting at time t and ending at time $t + \tau_b - 1$ consisted of $\{h_t, (A_{t+1}, Y_{t+2}, \dots, A_{t+\tau_b-1}, Y_{t+\tau_b})\}$. The mean square errors were still used as the

RNN decoder loss function. The goal of RNN decoder training is to make its training loss function smallest (Supplementary A).

Forecasting Procedures

The trained RIN was used to forecast the future number of cumulative cases of Covid-19 for each country. The recursive multiple-step forecasting involved using a one-step model multiple times where the prediction for the preceding time step and intervention strategy were used as an input for making a prediction on the following time step. For example, for forecasting the number of new confirmed cases for the one more next day, the predicted number of new cases in one-step forecasting would be used as an observational input in order to predict day 2. Repeat the above process to obtain the two-step forecasting. The summation of the final forecasted number of new confirmed cases for each country was taken as the prediction of the total number of new confirmed cases of Covid-19 worldwide.

Data Collection

The analysis is based on surveillance data of confirmed cumulative and new COVID-19 cases worldwide as of April 7, 2020. Data on the number of cumulative and new cases and COVID-19-attributed deaths across 184 countries from January 22, 2020 to April 7 were obtained from John Hopkins Coronavirus Resource Center (<https://coronavirus.jhu.edu/MAP.HTML>).

Data Pre-processing

\log_2 was used to transform the original data: $\tilde{X} = \log_2(X + 1)$. A value of 1 was assigned to the intervention variable for China. A value of zero was assigned to the intervention variable if the country did not implement intervention. Evenly spaced numbers over the interval [0, 1] were assigned to the intervention variable for other countries. The values of intervention variable were

determined such that they minimized the prediction error via training process. The data were divided into batches. To train the RIN, the countries and starting day in each selected country were randomly selected. The length of each batch was 7. The values of each time series in the batch were divided by their mean values to normalize the data.

Results

Prediction accuracy of dynamics of Covid-19 using RIN

Accurate prediction of the spread of Covid-19 is important for health intervention plan for the future. To demonstrate that the RIN is an accurate forecasting method, the RIN was applied to confirmed accumulated cases of COVID-19 across 184 countries. Figure 3 plotted reported and one-step ahead predicted time-case curves of Covid-19 where blue dotted curve was the number of reported cumulative cases after the analysis completion. To further reliably evaluate the forecasting accuracy, we reported 10-step ahead forecasted numbers of cumulative cases and errors of Covid-19 of 8 countries in Table S1 starting with April 8, 2020. The forecasting errors were quite small.

Inputting a sequence of hypothesized public health intervention strategies, the outputs of the RIN decoder were counterfactual numbers of cases of Covid-19 to respond to the intervention strategies. Interventions were measured by number in the interval $[0, 1]$, where 1 indicated the strictest comprehensive public health intervention, 0 indicated no intervention and the values between 0 and 1 indicated the various less strict interventions. To intuitively illustrate the impact of the measure of intervention on the spread of Covid-19, we presented Figure S1. Figure S1 plotted counterfactual numbers of cases of Covid-19 over time worldwide to respond the interventions with values 0.3, 0.5, 0.7 and 1. We observed that if the measure of intervention was

1, the number of cases was dramatically decreased to zero. However, when the measure of intervention was 0.3, the number of cases exponentially increased. The measure of intervention had big effect on the spread of Covid-19.

The number of cases of Covid-19 was a function of the past history and the measure of intervention. Forecasting also depended on the measure of intervention. In Table S1, we also listed the measures of the interventions which provided information on the degrees of current interventions in the country. The measure of interventions in the most countries was 0.75. However, the current measure of interventions in UK was 0.5, the smallest in 8 countries. These results showed that the RIN for forecasting the trajectory of Covid-19 was accurate and reliable. Similar to causal inference, the RIN can be used to evaluate the impact of a sequence of multiple intervention strategies on the curbing the spread of Covid-19 if the interventions were viewed as treatments.

The number of cases of Covid-19 grows exponentially without additional intensive interventions

To investigate how Covid-19 pandemic surges around the world, we presented Figure 4 that showed the forecasted number of cumulative cases of Covid-19 worldwide over time, assuming that the current intention remains unchanged. We observed that the number of cumulative cases of Covid-19 exponentially grown and would reach extremely high number 199,554,596 on July 6, 2020 if none of additional comprehensive public health intervention was implemented. Similarly, Figure 5 and Figure S2 plotted time-case curves of Covid-19 of eight countries: Italy, Spain, Iran, Germany, USA, France, Belgium and UK, and worldwide with unchanged intervention strategies in the future, respectively. We also observed exponentially growth of the numbers of cases of Covid-19 for every country without additional intervention.

We are in the eve to successfully curb the spread of Covid-19

As Covid-19 Accelerates and exponentially grows, how to slow down the spread of Covid-19 is an urgent task for every country around world. To demonstrate that when the additional intervention was implemented, the number of new cases of Covid-19 would decrease, we presented Figures 6 and Figure 7. Figures 6 and 7 plotted the number of cumulative case and new case curves of Covid-19 over time for 12 countries: US, Italy, Spain, Germany, France, Iran, UK, Switzerland, Belgium, South Korea, Japan and Singapore under three invention scenarios, respectively. Scenario 1 started with the current interventions for one week, then transitioned to the intervention with a measure of 0.95 in two weeks. The scenario 2 started with the current intervention for the first week, and then transitioned to the intervention with a measure of 0.95 in three weeks. The scenario 3 started with the current intervention for one week, changed to 0.85 in three weeks and finally transitioned to the intervention with a measure of 0.95 in three weeks. Figures 6 and 7 showed that when the countries moved to intervention with measure of 0.95, the spread of Covid-19 in all 12 countries was curbed. Now the measures of interventions in the most of 12 countries were closer to 0.75 (Table S1). These countries were closing to stopping the spread of Covid-19 if additional interventions such as wearing face masks were implemented.

Next we investigated how various intervention strategies reduced the peak time and cumulative case numbers, and the final total number of cases. Table S2 showed the forecasted results of COVID-19 in 12 countries under three sequences of interventions (Scenarios 1-3). We can see that under all three scenarios, the peak times were before April 15, 2020 and the spread of COVID-19 would be sopped before the end of May, 2020.

Conclusion/Discussion

As an alternative to the epidemiologic transmission models, we formulated the real-time forecasting and evaluating multiple public health intervention problem into a novel causal inference problem. We viewed the interventions as treatments where multiple interventions were administered at different time points. The number of new cases were taken as treatment responses. The RIN uses sequence-to-sequence multi-input/output recurrent neural network as a tool for modeling the real-time trajectory of the transmission dynamics of Covid-19, health intervention planning and making multi-step prediction of the response trajectory of Covid-19 over time with multiple interventions. The RNN can learn the complex dynamics within the temporal ordering of input time series of Covid-19 and use an internal memory to remember the hidden features.

This AI and causal inference-inspired approach allows us to address three important questions. The first question is the prediction accuracy. Unlike other dynamic systems where the parameters in the systems and control variables are, in general, independent, the epidemic systems have intervention and system dependent parameters. We designed the intervention variable that quantified comprehensive intervention strategies and had close relationships with the parameters in the epidemic systems. Therefore, the RINs could take the parameters in the epidemic dynamic systems as input control variables that can be estimated in the RIN training. The RIN models were closer to real epidemic dynamic systems than the epidemiological models. Therefore, our results showed that the RIN substantially improved the accuracies of prediction and subsequently multiple-step forecasting.

The second question is how important is the intervention time. Since interventions are complicated and are difficult to quantify, we designed four intervention scenarios to represent the degrees and delays of interventions. Since the proposed methods combine the real data and

models, they allowed us to evaluate the consequences of multiple intervention strategies, while maintaining the analysis as close to the real data as possible. The RIN investigated the impact of multiple public intervention plans and intervention measures on the size, duration and time of the virus outbreak and recommended the appropriate intervention times.

We estimated the duration, peak time and ending time, peak number of new cases and cumulative cases, and maximum number of cumulative cases of COVID-19 under four intervention scenarios for 184 countries in the world. We observed that the number of cumulative cases of Covid-19 would exponentially grow and reach extremely high number 199,554,596 on July 6, 2020 if none of additional comprehensive public health intervention was implemented. However, we also found that top 12 countries with the largest number of the lab confirmed cumulative cases of COVID-19 were closing to stopping the spread of Covid-19 if additional interventions such as wearing face masks were implemented. We can see that under all three scenarios, the peak times were before April 15, 2020 and the spread of COVID-19 would be sopped before the end of May, 2020.

Conflict of interest

Non to declare

Funding

Dr. Wei Lin is supported by the National Key R&D Program of China (Grant no. 2018YFC0116600), by the National Natural Science Foundation of China (Grant nos. 11925103 & 61773125), and by the STCSM (Grant no. 18DZ1201000).

References

- Anastassopoulou C, Russo, L, Tsakris A, Siettos C. Data-based analysis, modelling and forecasting of the COVID-19 outbreak. 2020. doi: <https://doi.org/10.1101/2020.02.11.20022186>.
- Irfan U. The math behind why we need social distancing, starting right now. 2020. <https://www.vox.com/2020/3/15/21180342/coronavirus-covid-19-us-social-distancing>.
- Li Q, Guan X, Wu P, Wang X, Zhou L, Tong Y et al. Early transmission dynamics in Wuhan, China, of novel Coronavirus-infected pneumonia. *N Engl J Med*. 2020, Jan 29. doi: 10.1056/NEJMoa2001316. [Epub ahead of print].
- Wu JT, Leung K, Leung GM. Nowcasting and forecasting the potential domestic and international spread of the 2019-nCoV outbreak originating in Wuhan, China: a modelling study. *Lancet*. 2020; 395(10225):689-697.
- Zhao S, Musa SS, Lin Q et al. Estimating the unreported number of novel Coronavirus (2019-nCoV) cases in China in the first half of January 2020: A data-driven modelling analysis of the early outbreak. *J Clin Med*. 2020 Feb 1;9(2). pii: E388. doi: 10.3390/jcm9020388.
- Kucharski A, Russell T, Diamond C, Liu Y, CMMID nCoV working group, Edmunds J, Funk S, Eggo R. Analysis and projections of transmission dynamics of nCoV in Wuhan (2020) https://cmmid.github.io/ncov/wuhan_early_dynamics/index.html.
- Tuite AR, Fisman DN. Reporting, epidemic growth, and reproduction numbers for the 2019 novel coronavirus (2019-nCoV) epidemic. *Ann Intern Med*. 2020 Feb 5. doi: 10.7326/M20-0358. [Epub ahead of print].
- Hellewell J, Abbott S, Gimma A, et al. Centre for the Mathematical Modelling of Infectious Diseases COVID-19 Working Group, Funk S1, Eggo RM2. Feasibility of controlling COVID-19 outbreaks by isolation of cases and contacts. *Lancet Glob Health*. 2020 Feb 28. pii: S2214-109X(20)30074-7. doi: 10.1016/S2214-109X(20)30074-7. [Epub ahead of print].

Li R, Pei S, Chen B, Song Y, Zhang T, Yang W, Shaman J. Substantial undocumented infection facilitates the rapid dissemination of novel coronavirus (SARS-CoV2). *Science*. 2020 Mar 16. pii: eabb3221. doi: 10.1126/science.abb3221. [Epub ahead of print].

Roosa K, and Chowell G. Assessing parameter identifiability in compartmental dynamic models using a computational approach: application to infectious disease transmission models. *Theor Biol Med Model*. 2019; 16: 1.

Funk S, Camacho A, Kucharski AJ, Eggo RM, Edmunds WJ. Real-time forecasting of infectious disease dynamics with a stochastic semi-mechanistic model. *Epidemics*. 2018; 22:56-61.

Johansson MA, Apfeldorf KM, Dobson S et al. An open challenge to advance probabilistic forecasting for dengue epidemics. *Proc Natl Acad Sci U S A*. 2019;116(48):24268-24274.

Gábor A, Villaverde AF, Banga JR. Parameter identifiability analysis and visualization in large-scale kinetic models of biosystems. *BMC Syst Biol*. 2017;11(1):54.

Lim B, Alaa AM, and van der Schaar M. Forecasting treatment responses over time using recurrent marginal structural networks. In *Advances in Neural Information Processing Systems*, 7493–7503, 2018.

Clevert DA, Unterthiner T, Hochreiter S. Fast and accurate deep network learning by exponential linear units (ELUs). 2015, arXiv:1511.07289.

Bica I, Alaa AM, Jordon J, and van der Schaar M. (2020). Estimating Counterfactual Treatment Outcomes over Time Through Adversarially Balanced Representations. arXiv e-prints, page arXiv:2002.04083.

Figure Legend

Figure 1. Forecasting intervention response to curbing the spread of Covid-19 under a sequence of interventions.

Figure 2. Architecture of recurrent intervention network.

Figure 3. Reported and predicted time-case curves of Covid-19 worldwide where blue dotted curve was the number of reported cumulative cases after the analysis completion.

Figure 4. The numbers of cumulative and new cases of Covid-19 worldwide over time, assuming the current intention remains unchanged. The curves in blue color and red color represented the number of cumulative cases and the number of new cases, respectively.

Figure 5. Forecasted number of cumulative cases of Covid-19 of eight countries Italy (A), Spain (B), Iran (C), Germany (D), USA (E), France (F), UK (G) and Belgium (H) over time without additional interventions.

Figure 6. Number of cumulative case curves of Covid-19 over time for 12 countries under three invention scenarios. (A) Time-case plot for US, (B) Time-case plot for Italy, (C) Time-case plot

for Spain, (D) Time-case plot for Germany, (E) Time-case plot for France, (F) Time-case plot for Iran, (G) Time-case plot for UK, (H) Time-case plot for Switzerland, (I) Time-case plot for Belgium, (J) Time-case plot for South Korea, (K) Time-case plot for Japan, and (L) Time-case plot for Singapore.

Figure 7. Number of new case curves of Covid-19 over time for 12 countries under three invention scenarios. (A) Time-new case plot for US, (B) Time-new case plot for Italy, (C) Time-new case plot for Spain, (D) Time-new case plot for Germany, (E) Time-new case plot for France, (F) Time-new case plot for Iran, (G) Time-new case plot for UK, (H) Time-new case plot for Switzerland, (I) Time-new case plot for Belgium, (J) Time-new case plot for South Korea, (K) Time-new case plot for Japan, and (L) Time-new case plot for Singapore.




Figure 1. Forecasting intervention response to curbing the spread of Covid-19 under a sequence of interventions.




Figure 2. Architecture of recurrent intervention network.




Figure 3. Reported and predicted time-case curves of Covid-19 worldwide where blue dotted curve was the number of reported cumulative cases after the analysis completion.




Figure 4. The numbers of cumulative and new cases of Covid-19 worldwide over time, assuming the current intention remains unchanged. The curves in blue color and red color represented the number of cumulative cases and the number of new cases, respectively.




Figure 5. Forecasted number of cumulative cases of Covid-19 of eight countries Italy (A), Spain (B), Iran (C), Germany (D), USA (E), France (F), UK (G) and Belgium (H) over time without additional interventions.




Figure 6. Number of cumulative case curves of Covid-19 over time for 12 countries under three invention scenarios. (A) Time-case plot for US, (B) Time-case plot for Italy, (C) Time-case plot for Spain, (D) Time-case plot for Germany, (E) Time-case plot for France, (F) Time-case plot for Iran, (G) Time-case plot for UK, (H) Time-case plot for Switzerland, (I) Time-case plot for Belgium, (J) Time-case plot for South Korea, (K) Time-case plot for Japan, and (L) Time-case plot for Singapore.




Figure 7. Number of new case curves of Covid-19 over time for 12 countries under three invention scenarios. (A) Time-new case plot for US, (B) Time-new case plot for Italy, (C) Time-new case plot for Spain, (D) Time-new case plot for Germany, (E) Time-new case plot for France, (F) Time-new case plot for Iran, (G) Time-new case plot for UK, (H) Time-new case plot for Switzerland, (I) Time-new case plot for Belgium, (J) Time-new case plot for South Korea, (K) Time-new case plot for Japan, and (L) Time-new case plot for Singapore.

Supplementary Materials

Supplementary Note A

Supplementary Figure Legend

Figures S1. Counterfactual numbers of cases of Covid-19 over time worldwide to respond the interventions with values 0.3, 0.5, 0.7 and 1.

Figure S2. Time-case plot of Covid-19 worldwide under four invention scenarios

Table S1

Table S2

Supplementary Note A

Recurrent Intervention Network (RIN)

Simple RNN Unit

RIN consisted of an encoder RNN and a decoder RNN. Basic unit of RNN in both encoder and decoder had three layers: input, recurrent hidden and output layers. The input layer consisted of three types of variables: covariates $X_t = [X_t^1, \dots, X_t^k]^T$, a scalar intervention variable A_t and the numbers of cases (potential outcomes) Y_t at the time t . Define the input vector V_t as

$$V_t = \begin{bmatrix} X_t \\ A_t \\ Y_t \end{bmatrix}.$$

Let $h_t = [h_t^1, \dots, h_t^m]^T$ be a m dimensional hidden state vector. The data V_t is inputted into the input layer. The linear transformation $W_{vh} V_t$ of the data V_t is then sent to the hidden layer, where W_{vh} is a $m \times (k + 2)$ dimensional matrix. The hidden layer receives information from the input layer and hidden layer at the previous time point. The state is determined by the following nonlinear transformation of its received information:

$$h_t = f_h(W_{hh} h_{t-1} + W_{vh} X_t + b_h),$$

where W_{hh} is a $m \times m$ dimensional weight matrix that connect the previous state to the current state, and $b_h = [b_h^1, \dots, b_h^m]^T$ is a m dimensional bias vector that corrects the bias, and f_h is a element-wise nonlinear activation function and is often defined as the following “tanh” function:

$$\tanh(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}.$$

The neurons in hidden layer are connected to the output layer via a m dimensional weight vector W_{hy} . The output \hat{Y}_{t+1} is determined by

$$\hat{Y}_{t+1} = f_o(W_{hy}h_t + b_o),$$

where f_o is an activation function and b_o is the bias vector of the output neurons.

Long Short Term Memory Network Unit

Long Short Term Memory network (LSTM) is a special kind of RNN, capable of learning long-term dependencies (Hochreiter and Schmidhuber, 1997; Srivastava et al. 2015). Each LSTM unit has a cell that is a memory unit. The key to LSTMs is the cell state. The cell state is determined by four gates: forget gate, input gate, input modulation, and output gate. At each time step, the unit receives input from two external sources: the current input variables V_t and the previous hidden states h_{t-1} . The total input activates the gates through sigmoid function and the tanh nonlinear function. The forget gate layer decides what information we’re going to throw away from the cell state:

$$f_t = \sigma(W_{vf}V_t + W_{hf}h_{t-1}) + b_f.$$

The next step, the input gate layer decides which values we’ll update:

$$i_t = \sigma(W_{vi}V_t + W_{hi}h_{t-1} + b_i),$$

Next, input modulation gate creates a vector of new candidate values, g_t , that could be added to the state:

$$g_t = \tanh(W_{vc}V_t + W_{hc}h_{t-1} + b_c).$$

The old cell state is updated by

$$C_t = f_t \circ C_{t-1} + i_t \circ g_t,$$

where \circ denotes element-wise multiplication.

Output gate decides what parts of the cell state we're going to output:

$$O_t = \sigma(W_{vo}V_t + W_{ho}h_{t-1} + b_o),$$

$$h_t = o_t \circ \tanh(C_t),$$

$$Y_t = W_{hy}h_t + b_y.$$

Loss function

Since the output is continuous, the mean square errors are used as the loss function for the encoder:

$$L_{encoder} = \sum_{i=1}^n \sum_{t=1}^{T_i} \|Y_{t+1}^i - \hat{Y}_{t+1}^i\|^2. \quad (A1)$$

For the decoder, we assumed that observations were batched into shorter sequences of up to τ_b .

The loss function for the decoder is defined as

$$L_{decoder} = \sum_{i=1}^n \sum_{t=1}^{T_i} \sum_{\tau=2}^{\min\{T_i-t, \tau_b\}} \|Y_{t+\tau}^i - \hat{Y}_{t+\tau}^i\|^2. \quad (A2)$$

Algorithm for RIN training is summarized as follows (Lim et al., 2018).

Algorithm

Input: training and validation data $\{Y_1, \dots, Y_T, X_1, \dots, X_T, A_1, \dots, A_T\}$.

Output: Weights in the RNN encoder and RNN decoder.

Step 1 : Estimate parameters θ_E in encoder .

Fit the encoder and estimate θ_E by optimizing $L_{encoder}$ in equation (A1):

$$\theta_E \leftarrow \min_{\theta_E} L_{encoder}(\theta_E) .$$

Step 2: Estimate encoder states for initialization of decoder)

For $n = 1$ to N do

for $t = 1$ to T

$$h_t(i) = RNN(h_{t-1}(i), (i), A_t(i), Y_t(i); \theta_E)$$

end for

end for

Step 3: Estimate parameters in RNN decoder

Fit the decoder and estimate θ_D by optimizing $L_{decoder}$ in equation (A2):

$$\theta_D \leftarrow \min_{\theta_D} L_{decoder}(\theta_D) ..$$

Potential outcome framework for evaluation of public health interventions

Potential outcome framework is our basic model to evaluate the impact of the public health interventions on the spread of Covid-19. The potential outcome framework is often referred to the Neyman-Rubin model (Rubin 1974). Potential outcomes consist of actual (or observed) and

counterfactual (hypothesized) outcomes. We are interested in number of cases of Covid-19 under some specific intervention. We observed the number of cases of Covid-19 (actual observation) without intervention. However, we want to know what number of cases of Covid-19 (counterfactual, unobserved) would be if an intervention was implemented. To evaluate the effect of intervention, we should compare the difference between the observed actual number of cases of Covid-19 and the counterfactual number of cases of Covid-19. Our aim is to learn the counterfactual outcomes of Covid-19 under a sequence of public health intervention options and evaluate the impact of the intervention strategies on the spread of Covid-19.

Let X_t be a set of covariates and A_t be an intervention at time t . A_t can be a binary variable. $A_t = 1$ ($A_t = 0$) indicates that intervention is (not) implemented. A_t can also be continuous variable taking values in the interval $[0, 1]$. If A_t is a continuous variable, the value of A_t represents the intensity of intervention. $A_t = 1$ indicates that the intervention is the most strict and comprehensive public health intervention. Let $Y_{t+1} = Y(a_t)$ be the potential outcome under intervention a_t and be observed only when $A_t = a_t$. The potential outcome framework assumes the existence of the hypothetical outcome with some interventions which is not observed in the data. The hypothetical outcome under hypothetical intervention is called counterfactual outcome. The set $\{X_t, A_t, Y_{t+1}\}$ forms a potential framework for causal inference.

Assumptions

We introduce assumptions in the Neyman-Rubin model (Bica et al., 2020).

Assumption 1. (Consistency). If a nation receives an intervention $A_t = a_t$, then the potential outcome for the intervention a_t is equal to the observed (factual) outcome $Y_{t=1}(a_t) = Y_{t+1}$.

Assumption 2. (Overlap). For all $(a_1, x_1, \dots, a_{t-1}, x_{t-1})$, we have

$$0 < P(A_t = a_t | A_1 = a_1, X_1 = x_1, \dots, A_{t-1} = a_{t-1}, X_{t-1} = x_{t-1}) < 1 .$$

Assumption 3. Sequential strong ignorability. Conditional on $A_1 = a_1, X_1 = x_1, \dots, A_{t-1} = a_{t-1}, X_{t-1} = x_{t-1}$, the potential outcomes Y_{t+1} are independent of A_t ,

$$Y_{t+1}(a_t) \perp\!\!\!\perp A_t | A_1 = a_1, X_1 = x_1, \dots, A_{t-1} = a_{t-1}, X_{t-1} = x_{t-1} .$$

Assumption 3 implies that there is no confounders which affect both outcomes and interventions.

References


Hochreiter S, Schmidhuber J. Long short-term memory. *Neural computation*. 1997; 9(8): 1735-1780.

Srivastava N, Mansimov E, Salakhutdinov R. Unsupervised learning of video representations using LSTMs. 2015; arXiv:1502.04681.

Rubin DB. Estimating causal effects of treatments in randomized and non-randomized studies. *Journal of Educational Psychology*. 1974; 66: 688–701.

Lim B, Alaa AM, and van der Schaar M. Forecasting treatment responses over time using recurrent marginal structural networks. In *Advances in Neural Information Processing Systems*, 7493–7503, 2018.

Bica I, Alaa AM, Jordon J, and van der Schaar M. (2020). Estimating Counterfactual Treatment Outcomes over Time Through Adversarially Balanced Representations. arXiv e-prints, page arXiv:2002.04083.



Figures S1. Counterfactual numbers of cases of Covid-19 over time worldwide to respond the interventions with values 0.3, 0.5, 0.7 and 1.




Figure S2. Time-case plot of Covid-19 worldwide under four invention scenarios

Table S1. Forecasting errors of worldwide and eight countries. .

Country	Estimated	Reported	Error
Italy	0.75		
4/8/2020	138341	139422	-0.0078
4/9/2020	142657	143626	-0.0067
4/10/2020	147052	147577	-0.0036
4/11/2020	151508	152271	-0.005
4/12/2020	155822	156363	-0.0035
4/13/2020	160129	159516	0.00384
4/14/2020	164545	162488	0.01266
4/15/2020	169203	165155	0.02451
4/16/2020	174193	168941	0.03109
4/17/2020	179283	172434	0.03972
Spain	0.75		
4/8/2020	141942	148220	-0.0424
4/9/2020	145247	153222	-0.052
4/10/2020	150194	158273	-0.051
4/11/2020	155344	163027	-0.0471
4/12/2020	160104	166831	-0.0403
4/13/2020	164682	170099	-0.0318
4/14/2020	169139	172541	-0.0197
4/15/2020	173455	177644	-0.0236
4/16/2020	178560	184948	-0.0345
4/17/2020	183943	190839	-0.0361
Iran	0.75		
4/8/2020	62589	64586	-0.0309
4/9/2020	63903	66220	-0.035
4/10/2020	65892	68192	-0.0337
4/11/2020	67985	70029	-0.0292
4/12/2020	69973	71686	-0.0239
4/13/2020	71849	73303	-0.0198
4/14/2020	73643	74877	-0.0165
4/15/2020	75383	76389	-0.0132

4/16/2020	77438	77995	-0.0071
4/17/2020	79609	79494	0.00144
Germany	0.75		
4/8/2020	107663	113296	-0.0497
4/9/2020	110094	118181	-0.0684
4/10/2020	113823	122171	-0.0683
4/11/2020	117559	124908	-0.0588
4/12/2020	121054	127854	-0.0532
4/13/2020	124405	130072	-0.0436
4/14/2020	127755	131359	-0.0274
4/15/2020	130932	134753	-0.0284
4/16/2020	134696	137698	-0.0218
4/17/2020	138658	141397	-0.0194
US	0.60 ->0.65		
4/8/2020	397505	429052	-0.0735
4/9/2020	418314	462780	-0.0961
4/10/2020	446258	496535	-0.1013
4/11/2020	478409	526396	-0.0912
4/12/2020	509746	555313	-0.0821
4/13/2020	541540	580619	-0.0673
4/14/2020	572983	607670	-0.0571
4/15/2020	604249	636350	-0.0504
4/16/2020	642404	667801	-0.038
4/17/2020	683412	699706	-0.0233
France	0.6		
4/8/2020	86839	83057	0.04554
4/9/2020	92414	87366	0.05778
4/10/2020	99041	91738	0.0796
4/11/2020	105138	94863	0.10831
4/12/2020	111820	121712	-0.0813
4/13/2020	119355	125394	-0.0482
4/14/2020	127629	131361	-0.0284
4/15/2020	136860	134582	0.01693
4/16/2020	146225	147091	-0.0059
4/17/2020	156437	149130	0.04899
Belgium	0.6		
4/8/2020	22786	23403	-0.0264
4/9/2020	24285	24983	-0.0279
4/10/2020	25772	26667	-0.0335
4/11/2020	27465	28018	-0.0197
4/12/2020	29092	29647	-0.0187

4/13/2020	30819	30589	0.00751
4/14/2020	32630	31119	0.04856
4/15/2020	34521	33573	0.02825
4/16/2020	36689	34809	0.05401
4/17/2020	38952	36138	0.07786
UK	0.5		
4/8/2020	55949	61474	-0.0537
4/9/2020	59873	65872	-0.0478
4/10/2020	63963	74605	-0.0971
4/11/2020	68874	79874	-0.0875
4/12/2020	73829	85206	-0.0767
4/13/2020	78544	89570	-0.0564
4/14/2020	83393	94845	-0.0429
4/15/2020	88391	99483	-0.0202
4/16/2020	94533	104145	0.00944
4/17/2020	100968	109769	0.03214
Worldwide	0.7		
4/8/2020	1412237	1480202	-0.016
4/9/2020	1490688	1565278	-0.048
4/10/2020	1569708	1657526	-0.053
4/11/2020	1655960	1735650	-0.046
4/12/2020	1739545	1834721	-0.052
4/13/2020	1825078	1904838	-0.04
4/14/2020	1912021	1976191	-0.032
4/15/2020	2000928	2056054	-0.027
4/16/2020	2104800	2152646	-0.022
4/17/2020	2212119	2240190	-0.013

Table S2. COVID-19 dynamics in 12 countries and worldwide for three scenarios.

Scenario	Country	US	Italy	Spain	Germany	France	Iran
Scenario 1	Peak Time	4/15/2020	3/20/2020	3/24/2020	3/26/2020	4/15/2020	3/29/2020
	Peak Number (New)	57039	6557	9630	6933	9365	3186
	Peak Number (Cumulative)	755875	53578	49515	50871	146225	41495
	End Time	4/23/2020	4/23/2020	4/23/2020	4/23/2020	4/23/2020	4/23/2020
	End Cases	854416	182006	186792	140749	162640	80762
	Duration	91	84	83	88	91	65
Scenario 2	Peak Time	4/15/2020	3/20/2020	3/24/2020	3/26/2020	4/15/2020	3/29/2020
	Peak Number (New)	57039	6557	9630	6933	9365	3186
	Peak Number (Cumulative)	755875	53578	49515	50871	146225	41495
	End Time	5/5/2020	5/2/2020	5/2/2020	5/2/2020	5/5/2020	5/2/2020
	End Cases	1076779	197400	202656	152411	199269	87131
	Duration	103	93	92	97	103	74
Scenario 3	Peak Time	4/15/2020	3/20/2020	3/24/2020	3/26/2020	4/15/2020	3/29/2020
	Peak Number (New)	57039	6557	9630	6933	9365	3186
	Peak Number (Cumulative)	755875	53578	49515	50871	146225	41495
	End Time	5/25/2020	5/25/2020	5/25/2020	5/25/2020	5/25/2020	5/25/2020
	End Cases	1601040	267917	275228	205519	282349	115857
	Duration	123	116	115	120	123	97
Scenario	Country	Switzerland	Belgium	Korea South	Japan	Singapore	Worldwide
Scenario 1	Peak Time	3/22/2020	4/15/2020	4/7/2020	4/3/2020	4/4/2020	4/15/2020
	Peak Number (New)	1321	2168	1005	522	120	103872
	Peak Number (Cumulative)	8795	36689	11336	3139	1309	2104800
	End Time	4/23/2020	4/23/2020	4/22/2020	4/22/2020	4/23/2020	4/23/2020
	End Cases	32137	40346	15125	4954	1833	2271648
	Duration	59	80	90	88	91	91

Scenario 2	Peak Time	3/22/2020	4/15/2020	4/7/2020	4/3/2020	4/4/2020	4/15/2020
	Peak Number (New)	1321	2168	1005	522	120	103872
	Peak Number (Cumulative)	8795	36689	11336	3139	1309	2104800
	End Time	5/4/2020	5/5/2020	5/4/2020	5/4/2020	5/3/2020	5/4/2020
	End Cases	35563	48283	16612	5388	1973	2620765
	Duration	70	92	102	100	101	102
Scenario 3	Peak Time	3/22/2020	4/15/2020	4/7/2020	4/3/2020	4/4/2020	4/15/2020
	Peak Number (New)	1321	2168	1005	522	120	103872
	Peak Number (Cumulative)	8795	36689	11336	3139	1309	2104800
	End Time	5/25/2020	5/25/2020	5/25/2020	5/25/2020	5/24/2020	5/25/2020
	End Cases	46840	65700	21446	6753	2409	3864872
	Duration	91	112	123	121	122	123