

1 **Differential effects of socio-demographic factors on maternal haemoglobin**
2 **concentration in three sub-Saharan African Countries**

3

4 Dickson A. Amugsi^{1*}

5 damugsi2002@yahoo.com; damugsi@aphrc.org

6 Zacharie T. Dimbuene^{2, 3}

7 zacharie.tsala.dimbuene@gmail.com

8 Catherine Kyobutungi¹

9 ckyobutungi@aphrc.org

10

11 1. Maternal and Child Wellbeing Unit, African Population and Health Research Center,
12 APHRC Campus, P.O Box 10787-00100, Nairobi, Kenya

13 2. Department of Population Sciences and Development, University of Kinshasa,
14 Democratic Republic of the Congo

15 3. Social Analysis and Modeling Division, Statistics Canada, Ottawa, Canada K1A 0T6

16

17 * Corresponding author: Dr Dickson A. Amugsi

18 Email: damugsi2002@yahoo.com; damugsi@aphrc.org

19

20 Word count: 3493

21

22

23

24

25

26

27

28

29

30 **Abstract**

31 **Objective:** To investigate the effects of socio-demographic factors on maternal
32 haemoglobin (Hb) at different points of the conditional distribution of Hb concentration.

33 **Methods:** We analysed the Demographic and Health Surveys data from Ghana,
34 Democratic Republic of the Congo (DRC) and Mozambique, using Hb concentration of
35 mothers aged 15-49 years as an outcome of interest. We utilise quantile regression to
36 estimate the effects of the socio-demographic factors across specific points of the
37 maternal Hb concentration.

38 **Results:** The results showed crucial differences in the effects of socio-demographic
39 factors along the conditional distribution of Hb concentration. In Ghana, maternal
40 education had a positive effect on Hb concentration in the 5th and 10th quantiles. The
41 positive effect of education on maternal Hb concentration occurred across all quantiles
42 in Mozambique, with the largest effect at the lowest quantile (5th) and the smallest effect
43 at the highest quantile (90th). In contrast, maternal education had a negative effect on
44 the Hb concentration of mothers in the 50th, 75th and 90th quantiles in DRC. Maternal
45 body mass index (BMI) had a positive effect on Hb concentration of mothers in the 5th,
46 10th, 50th and 90th, and 5th to 50th quantiles in Ghana and Mozambique, respectively.
47 Breastfeeding had a significant positive effect on Hb concentration across all countries,
48 with the largest effect on Hb concentration of mothers in the lower quantiles. All the
49 household wealth indices had positive effects on maternal Hb concentration across
50 quantiles in Mozambique, with the largest effect among mothers in the upper quantiles.
51 However, in Ghana, living in a poor wealth index was inversely related with Hb
52 concentration of mothers in the 5th and 10th quantiles.

53 **Conclusions:** Our results showed that the effects of socio-demographic factors on
54 maternal Hb concentration vary along its distribution. Interventions to address maternal
55 anaemia should take these variations into account to identify the most vulnerable
56 groups.

57

58 **Keywords:** Haemoglobin concentration, socio-demographics, Africa, quantile
59 regression

60

61

62 **What this study adds**

- 63 • Quantile regression can be used effectively to analyse anaemia data
- 64 • Socio-demographic factors have differential effects on Hb at different points of its
65 distribution
- 66 • Interpreting results based on the mean effect (as in OLS) only provides a partial
67 picture
- 68 • Breastfeeding has positive effect on maternal Hb concentration
- 69 • The use of multicountry data revealed differences and commonalities between
70 countries

71

72

73

74

75

76

77

78

79

80

81

82

83

84

85

86

87

88

89 **Introduction**

90 Maternal anaemia or low haemoglobin (Hb) concentration, a condition in which the Hb is
91 lower than normal is a worldwide public health problem (1). It is caused by deficiencies
92 in iron, folate, copper, and other vital vitamins(2, 3). Also, infectious disease morbidity,
93 parasitic infections and genetic disorders among others could cause low Hb
94 concentration (3, 4). While the causes of low Hb concentration are multifaceted, the
95 evidence show that an estimated 50% of low Hb concentration cases reported
96 worldwide are due to iron deficiency (5-7). The available data suggest that anaemia
97 affects about 500 million women of reproductive age, globally (8, 9). The World Health
98 Organisation (WHO) estimates indicate that the global anaemia prevalence >30% (2, 6).
99 Consequently, the WHO included a target of reducing anaemia among women of
100 reproductive age by 50% by 2025 in its Global Nutrition Targets (GNT) (5). Similarly,
101 anaemia was recently added as an indicator to track the progress of sub-goal 2.2 of the
102 Sustainable Development Goals (SDGs) to end all forms of malnutrition by
103 2030(10). The problem of maternal low Hb concentration is particularly severe in sub-
104 Saharan Africa (SSA) where poverty is highly prevalent and nutritious food is not easily
105 accessible, coupled with high incidence of infectious diseases (11-13).

106

107 The consequences of low Hb concentration on the health of women include but are not
108 limited to increased risks of low birth weight, preterm birth, perinatal mortality, and
109 neonatal mortality (14). Low Hb concentration also places mothers at an elevated risk of
110 death during childbirth and postpartum (15). Additionally, the literature suggests that
111 deficient Hb concentration can result in cardiac decompensation (i.e. the failure of the
112 heart to maintain adequate blood circulation). It also elevates the risk of haemorrhage
113 and decreases the ability to tolerate blood loss, which can lead to circulatory shock and
114 death (16, 17). The consequences of low Hb concentration mentioned above calls for
115 an investigation that would examine the effects of putative factors across the conditional

116 distribution of the Hb concentration. Such an inquiry will provide entry points for
117 interventions to address maternal anaemia in developing countries. This study intends
118 to achieve this goal by using an analytical strategy that focuses on the effects of socio-
119 demographic factors at different stages of the conditional distribution of maternal Hb
120 concentration.

121

122 The existing literature has identified several factors that have both negative and positive
123 effects on maternal Hb concentration. Some of these factors include, maternal age,
124 education, parity, marital status, household size, socioeconomic status, place of
125 residence, body mass index (BMI) and breastfeeding (18-25). A study in Dhaka showed
126 a strong relationship between maternal age, education level, income level, and maternal
127 Hb concentration (18). Moreover, higher BMI, primiparity, and living in better-off
128 households were associated with higher levels of Hb (21, 22). On the contrary, low
129 family income and large family size are associated negatively with maternal Hb
130 concentration (24). Further, low maternal BMI, parity, living in poorest wealth quintile
131 and breastfeeding were inversely related to maternal Hb concentration (25). Other
132 studies have shown that being separated or widowed, using an intrauterine device and
133 place of residence increased the odds of low Hb concentration among women (22, 23).

134

135 Indeed, from the studies reviewed above, it appears the literature on maternal anaemia
136 is abundant. Nonetheless, there are shortfalls with the analytical strategies employed in
137 these studies. For example, almost all these studies used either Ordinary Least
138 Squares (OLS) or logistic regression to estimate the effects of socio-demographics on
139 Hb concentration. This may mask the differential effects that may occur throughout the
140 entire distribution of the Hb concentration--socio-demographic factors may influence
141 maternal Hb differently across the distribution of the Hb concentration. These
142 differences in the effects of socio-demographic factors on Hb concentration suggest the
143 need to undertake an analysis that has the potential to present a comprehensive and

144 complete picture of the effects of these factors on maternal Hb concentration. The
145 quantile regression analytical strategy utilised in this paper can determine the effects of
146 the socio-demographic factors at different points of the conditional distribution of
147 maternal Hb concentration. Therefore, the objective this study was to examine the
148 effects of sociodemographic factors on maternal Hb concentration using quantile
149 regression. This type of analysis is currently missing in the anaemia research arena.

150

151 **Methods**

152 *Data sources and study design*

153 This study involved a secondary analysis of the demographic and health survey (DHS)
154 data (26) from Ghana, Mozambique, and Democratic Republic of Congo (DRC). These
155 are nationally representative data collected every five years in low and middle income
156 countries (LMICs). We based the selection of the three countries on our previous
157 analysis (27), as well as availability of anaemia data. In designing the surveys, the DHS
158 ensured that the surveys are identical across all participating countries to facilitate
159 comparison between and among nations. The DHS utilised a two-stage sample design
160 in the selection of participating households in their surveys. The detail description of the
161 DHS design processes is published elsewhere (28-32).

162 *Study participants*

163 The study participants were women aged 15-49 years and who had complete anaemia
164 data. Information on study participants was obtained through a face-to-face interviews.
165 The DHS collected blood samples for anaemia testing from mothers who voluntarily
166 consented to be tested (30). Blood samples were drawn from a drop of blood taken from
167 a finger prick and collected in a microcuvette. Hb concentration analysis was
168 undertaken on-site using a battery-operated portable HemoCue analyser. Non-pregnant
169 women with a Hb concentration less than 7.0 g/dl, and pregnant women with a Hb
170 concentration lower than 9.0 g/dl were referred to a nearby health facility for immediate
171 treatment (30). The total samples per respective countries were as follows: Ghana (n=
172 2975), DRC (n= 9438) and Mozambique (n= 10961).

173

174

175

176 **Ethical statement**

177 The DHS study was undertaken based on high ethical standards (33). Data collectors
178 were trained to recognise and respect the rights of study participants. They also
179 informed participants of their rights to decide whether to participate in the study or not.
180 The risks and benefits of the study as well as steps taken to mitigate the potential risks
181 were adequately explained to study participants. The protocols of the study in each
182 country, including biomarker collection, were approved by the recognised ethics review
183 committees of each country, and the Institutional Review Board of ICF International,
184 USA. Written informed consent was obtained from each study participant before they
185 were allowed to participate in the study. The biomarker results were made available to
186 study participants (30). The DHS Program, USA, granted permission to the authors for
187 the use of the data.

188

189 **Measures**

190 *Outcome variable*

191 We used maternal Hb concentration (g/dl) as the outcome variable for this analysis. As
192 described in the preceding sections, DHS collected blood samples from eligible women
193 to test for anaemia using various strategies. The Hb concentration is captured first in the
194 DHS data as a continuous variable and then categorised into three levels of anaemia:
195 mild, moderate and severe. In this analysis, we used the Hb concentration as a
196 continuous variable.

197 *Predictor variables*

198 We grouped the predictor variables into three main categories: *maternal* (education,
199 age, BMI, employment status, parity, breastfeeding status, marital status and ANC

200 attendance); *household* (wealth index, sex of household head, household size, number
201 of children U5, decision making on large household purchases and husband/partner
202 education) and *community* (place of residence). The DHS created the household wealth
203 index using assets ownership and housing characteristics: type of roofing, and flooring
204 material, source of drinking water, sanitation facilities, ownership of television, bicycle,
205 motorcycle, automobile among others. The detail of the computation processes is
206 published elsewhere (30). The maternal BMI (Kg/m²) was obtained by dividing weight in
207 kilogrammes by height in meters squared.

208

209 **Data analysis**

210 *Outline of quantile regression model*

211 Koenker and Bassett (34) introduced the quantile regression (QR) as a *location*
212 *model* to extend Ordinary Least Squares (OLS). It is the case because OLS
213 summarises the distribution at the grand mean. However, the QR assesses more
214 general class of linear models in which, the conditional quantiles have a linear form to
215 fully account for the overall distribution of the response variable. To formalise the QR,
216 we consider a real-valued random variable Y characterized by the following distribution
217 function (34, 35);

$$218 F(y) = \Pr(Y \leq y) \tag{1}$$

219 Then for any $\alpha \in (0, 1)$, the α -th quantile of Y is defined as:

$$220 Q(\alpha) = \inf\{y: F(y) \geq \alpha\} \tag{2}$$

221 The common quantiles α from Equation (1) are $\alpha = .25$, $\alpha = .50$, and $\alpha = .75$ for the
222 first, quartile, the median and the third quartile. Therefore, unlike the OLS, which
223 minimises the squared differences around the mean, QR minimises the
224 weighted *absolute* difference between the observed value of y and the α -th quintile of Y .
225 The preceding discussion demonstrates that OLS is nested within QR (34, 35).

226

227 **Analytical approach utilised**

228 We used quantile regression (QR) (34) to examine the effects of the putative socio-
229 demographic factors on maternal Hb concentration. Using the QR, we were able to
230 investigate the effects of the predictor variables at different points of the conditional
231 distribution of the outcome variable (Hb concentration). This type of analysis cannot be
232 done with OLS, because standard OLS regression techniques summarise the average
233 relationship between a set of regressors and the outcome variable based on the
234 conditional mean function $E(y|x)$. Thus, it provides only a partial view of the
235 relationship, as we might be interested in describing the relationship at different points
236 in the conditional distribution of y . The QR, unlike OLS, provides a complete view of the
237 effects of the socio-demographic factors on maternal Hb concentration. Thus, making it
238 possible to identify the vulnerable groups that are in dire need of interventions. Further,
239 QR is more robust in handling non-normal errors and outliers compared with OLS (34).
240 Finally, QR provides a richer characterisation of the data, thereby illuminating the
241 effects of a covariate on the entire distribution of the outcome variable. In this analysis,
242 we also estimated OLS estimates for comparison purpose, and QR at the 5th, 10th, 25th,
243 50th, 75th and 90th quantiles (27, 36).

244

245 Furthermore, since we did not have a specific predictor variable of interest, all the socio-
246 demographic variables were included simultaneously in the models. They include,
247 maternal education, age, BMI, employment status, parity, breastfeeding status, marital
248 status, ANC attendance, household wealth index, sex of household head, household
249 size, number of children under five years, decision making on large household
250 purchases, husband/partner education and place of residence. The variables outlined
251 above were selected based on the literature, followed by bivariate analysis. Significant
252 variables in the bivariate analysis were included in the QR models.

253

254 **Results**

255 *Descriptive analysis of the characteristics of samples*

256 The descriptive results (Table 1) showed that maternal mean Hb concentration was
 257 relatively the same across all the three countries: Ghana (11.95±1.49), DRC
 258 (12.05±1.65) and Mozambique (11.64±1.73). Mothers in Ghana (5.27±4.92) spent a
 259 little more years in education than those in DRC (4.88±3.77) and Mozambique
 260 (3.46±3.45). The age of the study participants ranged from 28 years in Mozambique to
 261 31 years in Ghana. Moreover, mothers in Ghana tended to have higher mean BMI
 262 (24.34±4.96) relative to those in DRC (21.79±3.66) and Mozambique (22.53±3.70). In
 263 DRC, 68% of mothers indicated they were breastfeeding at the time of the study, while
 264 the number of breastfeeding mothers in Ghana and Mozambique stood at 58%,
 265 respectively.

266

267 **Table 1: Characteristics of the socio-demographic variables of the three countries**

Variables	Ghana (n=2975)		DRC (n=9438)		Mozambique (n=10961)	
	Mean/%	SD	Mean/%	SD	Mean/%	SD
Maternal-level variables						
Hb concentration (g/dl)	11.95	1.49	12.05	1.65	11.64	1.73
Women education (in years)	5.27	4.92	4.88	3.77	3.46	3.45
Age (in years)	30.65	6.96	29.13	6.94	28.38	7.18
Women's Body Mass Index (BMI)(Kg/m ²)	24.34	4.96	21.79	3.66	22.53	3.70
Working (yes)	79.3		76.0		37.5	
Parity	3.62	2.17	4.42	2.56	3.8	2.28
Breastfeeding (yes)	57.9		67.8		57.6	
Marital status						
Never married	6.4		4.2		5.6	
Married or Cohabiting	87.4		87.1		82.8	
Divorced/Widowed/Separated	6.2		8.7		11.6	
Household-level variables						
Wealth index						
Poorest	33.0		27.4		18.2	
Poor	21.1		22.9		18.9	
Middle	19.0		20.7		19.7	
Rich	15.1		17.1		22.1	
Richest	12.3		12.1		21.1	
Head of household (Female)	23.7		22.2		32.9	
Household size	5.80	2.82	6.74	2.86	6.16	2.79
Number of children under five years	1.73	0.93	2.16	1.04	1.86	0.98

Decision on household large purchases

Respondent alone	16.0		12.5		11.1	
Respondent and Husband/Partner	46.5		50.3		57.1	
Husband/Partner alone	24.1		36.7		30.7	
Someone else/Other	13.4		0.59		1.1	
Husband education (in years)	7.04	5.44	8.53	4.11	4.28	3.95
Community-level variables						
Place of Residence (% urban)	39.8		28.7		32.1	

SD: standard deviation; Hb: Haemoglobin;

268 **Notes:** SDs are reported only for continuous variables

269

270

271 *Quantile multivariable regression analysis of the effects of socio-demographic factors on*
 272 *maternal Hb concentration*

273 In Tables 2-4, we present the QR results of the effects of socio-demographic factors on
 274 maternal Hb concentration in Ghana, DRC and Mozambique. We also reported the OLS
 275 estimates for the purposes of comparison with the QR results. The results from the
 276 OLS analysis showed that maternal education had strong positive effects on Hb
 277 concentration in Ghana and Mozambique. Thus, in both countries, one year increase in
 278 education was associated with a unit increase maternal Hb concentration. A similar
 279 effects of BMI on Hb concentration were observed in all the three countries, so was
 280 breastfeeding practice.

281

282 However, in the QR analysis (Tables 2-4), the results revealed vital differences in
 283 effects at different points in the conditional distribution of maternal Hb concentration. For
 284 example, in Ghana, the effect of maternal years of education occurred at the first two
 285 lowest quantiles (5th and 10th), with the largest effect at the 5th quantile. Similarly, in
 286 Mozambique, one-year increase in maternal education was associated with a unit
 287 increase in Hb concentration across all quantiles, with the largest effect on mothers in
 288 the lowest quantile (5th) and the smallest effect at the highest quantile (90th).
 289 Interestingly, in DRC, maternal years of education had an inverse relationship with Hb
 290 concentration of mothers in the three upper quantiles. Thus, one-year increase in

291 maternal years of education was associated with 0.015, 0.020 and 0.023 units
292 decrease in maternal Hb concentration at the 5th, 75th and 90th quantiles, respectively.
293 In Ghana, maternal BMI had a significant positive effect on Hb concentration of mothers
294 in the 5th, 10th, 50th and 90th quantiles. The effect on the remaining two quantiles did not
295 reach statistical significance. The positive effects of BMI on maternal Hb concentration
296 was at the three lowest quantiles (5th, 10th and 25th) in DRC, while in Mozambique, a
297 unit increase in maternal BMI was associated with 0.031, 0.033, 0.029 and 0.018 units
298 increase in maternal Hb concentration at the 5th, 10th, 25th and 50th quantiles,
299 respectively. In each of the countries, the largest effect of maternal BMI on Hb
300 concentration occurred at the lower end of the Hb distribution (i.e. lower quantiles).

301
302 In Ghana, breastfeeding was positively and significantly associated with Hb
303 concentration of mothers in the 5th, 10th, 25th, and 50th quantiles, with the least effect
304 occurring at the 50th quantile. However, in DRC and Mozambique, breastfeeding had a
305 decreasing effect across all quantiles in the respective countries. The largest effect in
306 each country was at the lower end of the conditional distribution of the Hb
307 concentration, while the smallest effect was at the higher end of the distribution. In
308 Ghana, women participation in decision making regarding large household purchases
309 was associated with better Hb concentration among mothers in the 25th and 50th
310 quantiles, while the partner taking the decision alone was associated positively with the
311 Hb concentration at 5th, 25th and 50th quantiles. On the contrary, there was an inverse
312 effect of the partner alone, deciding on large household purchases on Hb concentration
313 of mothers in the 25th and 50th quantiles in DRC. The effects of female household
314 headship in DRC was mixed. It associated positively with Hb concentration of mothers
315 in the first three quantiles (5th, 10th and 25th), and negatively with the two upper
316 quantiles (75th and 90th). In Mozambique, the household wealth index had a significant
317 and increasing effect on maternal Hb concentration across almost all the quantiles. The
318 largest effect occurred at the highest end of the Hb distribution (90th quantile). In Ghana,
319 being in a lower wealth index was associated with low Hb concentration among mothers
320 in the 5th and 10th quantiles.

Table 2: Effects of socio-demographic factors on maternal Hb concentration in Ghana

Variables	OLS	Q5	Q10	Q25	Q 50	Q75	Q90
Maternal level variables							
Women education (in years)	0.015* (0.008)	0.063*** (0.016)	0.044** (0.016)	0.024 (0.013)	0.004 (0.010)	-0.004 (0.009)	-0.007 (0.015)
Age (in years)	0.009 (0.006)	0.013 (0.011)	0.024 (0.013)	0.012 (0.011)	0.006 (0.008)	-0.005 (0.008)	-0.001 (0.009)
Women's Body Mass Index (BMI) (Kg/m ²)	0.020** (0.006)	0.037** (0.012)	0.022* (0.010)	0.009 (0.009)	0.019* (0.008)	0.012 (0.008)	0.024* (0.010)
Mother working-No (reference)							
Mother working-Yes	0.121 (0.070)	0.008 (0.139)	0.003 (0.147)	0.129 (0.111)	0.142 (0.079)	0.095 (0.084)	0.159 (0.106)
Parity	-0.002 (0.021)	-0.009 (0.060)	-0.014 (0.064)	0.019 (0.035)	0.009 (0.031)	0.036 (0.030)	0.036 (0.029)
Breastfeeding-No (reference)							
Breastfeeding-Yes	0.216*** (0.060)	0.435*** (0.132)	0.424*** (0.128)	0.442*** (0.106)	0.252** (0.077)	-0.099 (0.085)	-0.070 (0.101)
Marital status							
Never married (reference)							
Married or Cohabiting	-0.180 (0.328)	-0.306 (0.650)	-0.416 (0.766)	-0.432 (0.523)	-0.435 (0.504)	0.141 (0.491)	-0.176 (0.573)
Divorced/Widowed/Separated	0.154 (0.159)	0.507 (0.406)	0.189 (0.330)	0.413 (0.214)	0.084 (0.176)	0.138 (0.193)	-0.384 (0.266)
Antenatal visits=0-3 (reference)							
Antenatal visits=4+	0.212* (0.093)	0.458 (0.247)	0.383 (0.214)	0.121 (0.167)	0.193 (0.121)	0.094 (0.131)	0.300* (0.138)
Household level variables							
Wealth index-poorest (reference)							
Wealth index-Poor	-0.199* (0.082)	-0.243 (0.177)	-0.411* (0.180)	-0.169 (0.141)	-0.188 (0.098)	-0.136 (0.096)	-0.071 (0.145)
Wealth index-Middle	-0.128 (0.096)	0.012 (0.198)	0.109 (0.191)	-0.076 (0.156)	-0.303** (0.117)	-0.079 (0.130)	0.100 (0.176)
Wealth index-Rich"	0.122 (0.119)	0.123 (0.266)	0.068 (0.257)	0.348 (0.195)	0.146 (0.128)	0.149 (0.135)	0.210 (0.228)
Wealth index-Richest	0.109 (0.141)	0.112 (0.302)	0.294 (0.300)	0.228 (0.245)	0.088 (0.149)	0.317 (0.176)	0.159 (0.229)
Head of household-Male (reference)							
Head of household-Female	0.067 (0.075)	0.016 (0.188)	0.149 (0.174)	0.102 (0.119)	0.005 (0.095)	-0.004 (0.091)	0.035 (0.109)
Household size	0.011 (0.013)	0.001 (0.045)	0.031 (0.033)	0.019 (0.017)	0.004 (0.014)	0.000 (0.017)	-0.003 (0.017)
Number of children under five	-0.060 (0.038)	-0.001 (0.108)	0.053 (0.094)	-0.095 (0.058)	-0.128** (0.045)	-0.087 (0.053)	0.006 (0.058)

Decision on large household purchases

Respondent alone (reference)							
Respondent and Husband/Partner	0.232** (0.080)	0.227 (0.237)	0.268 (0.190)	0.247* (0.124)	0.296** (0.106)	0.120 (0.096)	0.160 (0.116)
Husband/Partner alone	0.269** (0.088)	0.480* (0.236)	0.294 (0.203)	0.379** (0.121)	0.357** (0.118)	0.098 (0.101)	-0.033 (0.144)
Someone else/Other	0.008 (0.317)	-0.281 (0.617)	-0.153 (0.763)	-0.397 (0.501)	-0.158 (0.500)	0.227 (0.470)	0.252 (0.550)
Husband education (in years)	-0.002 (0.007)	-0.033* (0.014)	-0.025 (0.016)	-0.001 (0.011)	0.013 (0.009)	0.001 (0.008)	-0.006 (0.011)
Place of Residence-rural (reference)							
Place of Residence-Urban	-0.097 (0.077)	-0.128 (0.202)	-0.037 (0.175)	-0.041 (0.111)	-0.209** (0.081)	-0.181* (0.090)	-0.096 (0.140)
Observations	2975	2975	2975	2975	2975	2975	2975

Standard errors in parentheses; OLS: Ordinary least squares; Q: Quantile

* p<0.05; **p<0.01; ***p<0.001

Table 3: Effects of socio-demographic factors on maternal Hb concentration in DRC

Variables	OLS	Q5	Q10	Q25	Q 50	Q75	Q90
Maternal level variables							
Women education (in years)	-0.007 (0.006)	0.016 (0.018)	-0.007 (0.013)	0.006 (0.008)	-0.015** (0.006)	-0.020** (0.007)	-0.023* (0.011)
Age (in years)	0.004 (0.004)	-0.010 (0.012)	-0.007 (0.010)	0.006 (0.005)	0.012* (0.005)	0.005 (0.005)	0.003 (0.006)
Women's Body Mass Index (BMI)(Kg/m ²)	0.015** (0.005)	0.026*** (0.008)	0.021** (0.007)	0.014* (0.006)	0.007 (0.005)	0.009 (0.007)	0.001 (0.010)
Mother working-No (reference)							
Mother Working -Yes	-0.033 (0.041)	0.060 (0.107)	0.046 (0.082)	-0.010 (0.052)	-0.037 (0.044)	0.030 (0.051)	-0.038 (0.070)
Parity	0.016 (0.012)	0.052 (0.037)	0.041 (0.026)	0.017 (0.018)	0.003 (0.014)	0.014 (0.016)	0.023 (0.018)
Mother breastfeeding-No (reference)							
Mother breastfeeding-Yes	0.470*** (0.038)	0.745*** (0.117)	0.662*** (0.079)	0.686*** (0.057)	0.488*** (0.045)	0.355*** (0.057)	0.219*** (0.061)
Marital status							
Never married (reference)							
Married or Cohabiting"	-0.128 (0.092)	-0.219 (0.244)	-0.372* (0.169)	-0.186 (0.127)	-0.242** (0.092)	-0.145 (0.103)	0.108 (0.132)
Divorced/Widowed/Separated	0.071 (0.103)	-0.010 (0.307)	-0.060 (0.191)	-0.026 (0.141)	0.007 (0.106)	0.179 (0.126)	0.320* (0.149)
Antenatal visits=0-3 (reference)							
Antenatal visits=4+	0.025 (0.039)	0.011 (0.098)	0.027 (0.078)	0.030 (0.053)	0.014 (0.043)	0.016 (0.045)	0.095 (0.063)
Household level variables							
Wealth index-Poorest (reference)							
Wealth index-Poor	-0.030 (0.048)	0.024 (0.158)	0.091 (0.116)	-0.062 (0.074)	-0.056 (0.048)	-0.081 (0.061)	-0.048 (0.085)
Wealth index- Middle	0.104* (0.050)	0.219 (0.137)	0.282* (0.118)	0.106 (0.070)	0.058 (0.054)	-0.011 (0.068)	0.027 (0.087)
Wealth index- Rich	0.064 (0.058)	0.279 (0.148)	0.393** (0.137)	0.112 (0.082)	-0.004 (0.061)	-0.106 (0.073)	-0.020 (0.090)
Wealth index- Richest	-0.062 (0.079)	0.192 (0.241)	0.206 (0.168)	-0.072 (0.114)	-0.207* (0.095)	-0.039 (0.096)	-0.070 (0.125)
Head of household-Male (reference)							
Head of household-Female	0.019 (0.045)	0.278* (0.136)	0.249** (0.083)	0.211*** (0.057)	-0.033 (0.046)	-0.129* (0.057)	-0.179** (0.068)
Household size	0.001 (0.008)	0.035 (0.020)	0.001 (0.016)	-0.004 (0.012)	-0.002 (0.011)	0.002 (0.010)	-0.009 (0.011)
Number of children under five	-0.015 (0.021)	-0.107 (0.056)	-0.051 (0.053)	0.021 (0.031)	-0.037 (0.023)	-0.042 (0.026)	0.020 (0.035)

Table 4: Effects of socio-demographic factors on maternal Hb concentration in Mozambique

Variables	OLS	Q5	Q10	Q25	Q 50	Q75	Q90
Maternal level variables							
Women education (in years)	0.035*** (0.007)	0.055** (0.021)	0.043** (0.015)	0.030** (0.011)	0.037*** (0.008)	0.046*** (0.009)	0.027* (0.011)
Age (in years)	0.001 (0.004)	0.002 (0.009)	0.001 (0.007)	0.001 (0.006)	0.005 (0.004)	0.000 (0.005)	0.001 (0.005)
Women's Body Mass Index (BMI)(Kg/m ²)	0.019*** (0.005)	0.031* (0.015)	0.033** (0.011)	0.029*** (0.007)	0.018** (0.006)	0.008 (0.007)	0.008 (0.008)
Mother working-No (reference)							
Mother working-Yes	-0.049 (0.034)	0.004 (0.094)	-0.038 (0.071)	-0.092 (0.052)	0.001 (0.039)	-0.013 (0.042)	-0.080 (0.052)
Parity	0.025* (0.012)	0.069* (0.027)	0.061** (0.024)	0.024 (0.015)	0.003 (0.011)	0.023 (0.017)	0.004 (0.018)
Mother breastfeeding-No (reference)							
Mother breastfeeding-Yes	0.424*** (0.036)	0.732*** (0.100)	0.663*** (0.077)	0.520*** (0.055)	0.459*** (0.041)	0.310*** (0.051)	0.219*** (0.054)
Marital status							
Never married (reference)							
Married or Cohabiting	-0.067 (0.087)	-0.445* (0.210)	-0.275 (0.160)	-0.159 (0.136)	0.139 (0.106)	-0.055 (0.121)	0.075 (0.134)
Divorced/Widowed/Separated	-0.139 (0.092)	-0.495* (0.220)	-0.221 (0.205)	-0.144 (0.149)	-0.062 (0.113)	-0.138 (0.121)	-0.005 (0.141)
Antenatal visits=0-3 (reference)							
Antenatal visits = 4+	0.049 (0.036)	0.045 (0.086)	0.028 (0.067)	0.035 (0.048)	0.015 (0.038)	0.036 (0.041)	0.075 (0.051)
Household level variables							
Wealth index-Poorest (reference)							
Wealth index-Poor	0.255*** (0.054)	-0.114 (0.157)	0.198 (0.110)	0.174* (0.076)	0.285*** (0.063)	0.326*** (0.063)	0.429*** (0.074)
Wealth index- Middle	0.367*** (0.054)	0.246 (0.131)	0.397*** (0.109)	0.249*** (0.073)	0.335*** (0.055)	0.399*** (0.067)	0.552*** (0.075)
Wealth index-Rich	0.380*** (0.056)	0.242 (0.146)	0.406*** (0.118)	0.268*** (0.079)	0.337*** (0.059)	0.400*** (0.067)	0.594*** (0.085)
Wealth index-Richest	0.257*** (0.073)	-0.122 (0.191)	0.036 (0.150)	0.147 (0.109)	0.198** (0.074)	0.374*** (0.098)	0.474*** (0.111)
Head of household-Male (reference)							
Head of household-Female	-0.010 (0.039)	-0.061 (0.100)	-0.150 (0.083)	-0.040 (0.057)	0.075 (0.045)	-0.050 (0.047)	-0.074 (0.067)
Household size	0.015 (0.008)	0.014 (0.016)	0.007 (0.016)	0.004 (0.012)	0.028*** (0.008)	0.007 (0.011)	0.011 (0.017)
Number of children under five	-0.025 (0.022)	-0.077 (0.054)	-0.070 (0.051)	0.001 (0.035)	-0.053* (0.024)	0.015 (0.034)	0.003 (0.034)

Decision on large household purchases

Respondent alone (reference)

Respondent and Husband/Partner	0.010 (0.056)	-0.146 (0.143)	-0.092 (0.106)	-0.050 (0.084)	0.075 (0.064)	0.027 (0.062)	0.088 (0.093)
Husband/Partner alone	-0.032 (0.059)	-0.021 (0.149)	-0.061 (0.112)	-0.031 (0.088)	-0.050 (0.065)	-0.020 (0.069)	0.043 (0.099)
Someone else/Other	0.244 (0.168)	0.817 (0.605)	0.349 (0.230)	0.133 (0.267)	0.062 (0.227)	0.430* (0.172)	0.314 (0.260)
Husband education (in years)	0.011* (0.006)	-0.002 (0.015)	0.019 (0.012)	0.015 (0.008)	0.013 (0.006)	0.002 (0.007)	0.011 (0.010)
Place of Residence-Rural (reference)							
Place of Residence-Urban	-0.059 (0.046)	-0.278* (0.129)	-0.203* (0.088)	-0.184** (0.067)	0.019 (0.046)	-0.021 (0.058)	0.139 (0.072)
Observations	10961	10961	10961	10961	10961	10961	10961

Standard errors in parentheses; OLS: Ordinary least squares; Q: Quantile

* p<0.05; **p<0.01; ***p<0.001

336

337

338 Figures 1-3 are visual presentations of the effects of the various socio-demographic
339 factors on maternal Hb concentration in the three countries included in the analysis.

340

341 **Figure 1: Pictorial presentation of the effects of socio-demographic factors on**
342 **maternal Hb concentration in Ghana**

343

344 **Figure 2: Pictorial presentation of the effects of socio-demographic factors on**
345 **maternal Hb concentration in DRC**

346

347 **Figure 3: Pictorial presentation of the effects of socio-demographic factors on**
348 **maternal Hb concentration in Mozambique**

349

350

351

352

353 **Discussion**

354 This paper examined the effects of socio-demographic factors on maternal Hb
355 concentration in Ghana, DRC and Mozambique, using quantile regression
356 to understand the differential effects of putative socio-demographic factors at different
357 points of the conditional distribution of maternal Hb concentration. Our QR results
358 showed that in Ghana, one-year increase in maternal education was associated with an
359 improved Hb concentration of mothers in the 5th and 10th quantiles, while the effect on
360 the Hb of mothers in the other four quantiles did not reach statistical significance.
361 However, the OLS results show that one-year increase maternal education was
362 associated with a unit increase in Hb concentration of all mothers. This paints just a part
363 of the picture and therefore can be misleading. However, in Mozambique, maternal
364 education had a decreasing effect on Hb concentration across quantiles (from 5th to
365 90th). Our findings in the two countries imply a disproportionate positive effect of
366 maternal education accruing to mothers in the lower tail of the Hb distribution--improving
367 maternal education may be more impactful on the Hb concentration of mothers in the
368 lower than the upper quantiles. Conversely, a year increase in maternal years of
369 schooling was associated with a unit decrease in Hb concentration of mothers in the
370 three upper quantiles in DRC. These findings are puzzling as the literature suggests
371 that education consistently predict positive health outcomes in women(18, 25, 37,
372 38). For example, study using multicountry data concluded that women with higher
373 years of education were less likely to be anaemic relative to those with fewer years of
374 schooling (37). Further research is needed to elucidate the possible factors accounting
375 for the negative effect of maternal education on maternal health outcomes in DRC. Our
376 study together with the above studies, despite using different analytical strategies,
377 strongly suggest that education has positive effects on maternal health outcomes.

378

379 Our analysis also showed that maternal BMI has a significant positive effect on Hb
380 concentration in at least three quantiles in each country. The most significant effect of
381 BMI was among mothers in the lower quantiles in each country. This suggests that
382 interventions targeted at improving maternal BMI qualitatively are likely to be more

383 effective in increasing the Hb concentration of mothers in the lower tail of the Hb
384 distribution. It is worthy to note that the effects of BMI were not across all quantiles.
385 Hence, the OLS estimates, which suggested that a unit change in maternal BMI is
386 associated with a unit increase in Hb concentration among all mothers, may be
387 misleading. The QR findings, are, therefore, critical for identifying the groups that need
388 to be targeted in programme planning. The literature corroborated the findings of our
389 study. Several studies using either linear or logistic regression analytical strategies
390 suggested that women with higher BMI tend to have higher levels of Hb concentration
391 (22, 23, 37).

392

393 Similarly, breastfeeding had significant positive effect on maternal Hb concentration in
394 all the three countries. Mothers who were breastfeeding at the time of the survey tended
395 to have better Hb concentration compared with non-breastfeeding mothers. The largest
396 effects were observed among mothers in the lower quantiles, suggesting that
397 intervention to promote breastfeeding among lactating mothers may have more impact
398 on Hbs of mothers at the lower end of the Hb distribution. These findings may appear
399 puzzling because it is generally believed that lactating mothers tend to lose some iron to
400 their infants, which may have a bearing on their Hb concentration (39, 40).
401 Nevertheless, other evidence suggests that the iron contained in breast milk to children
402 is not significant enough to deplete the iron level of the mother, unless the mother is
403 already anaemic (41). The literature further suggests that mothers who are anaemic
404 during pregnancy and postpartum can recover through intake of iron rich diet and/or iron
405 supplement, and may not suffer low Hb concentration during lactation (41-43). The
406 preceding discussion suggests that breastfeeding may not necessarily deplete maternal
407 iron level, with the consequential negative effect on Hb concentration. Some available
408 evidence suggests a positive effect of breastfeeding on maternal Hb (39). Other studies
409 observed inverse relationship between breastfeeding and Hb concentration levels (40,
410 44). These mixed findings notwithstanding, the findings in the present study, together
411 with the literature suggest that breastfeeding can indeed have positive effects on

412 maternal Hb concentration levels, although the mechanism through which this happens
413 may be complicated.

414

415 Our findings in Mozambique suggest that household wealth index (HWI) had a positive
416 and increasing (from 5th to 90th) effect on Hb concentration across all quantiles. The
417 least effect was on mothers in the lower end of the Hb distribution, while the largest
418 effect was on mothers at the upper end of the distribution. Thus, improving HWI may be
419 more impactful on mothers at the upper quantiles relative to those at the lower
420 quantiles. The positive association between HWI and women health outcomes have
421 been substantially documented (39, 45, 46). The evidence is that mothers who live in
422 better-off households tend to have higher levels of Hb concentration (39). However, in
423 Ghana, mothers who live in poor households and are in the 5th and 10th quantiles
424 tended to have lower Hb concentration. The finding in Ghana is consistent with the
425 literature, which often identifies poverty as a risk factor of maternal health outcomes (25,
426 47).

427

428 An essential strength of this study is that the outcome variable was objectively
429 measured, thereby reducing the possible biases associated with subjective
430 measurements. The use of QR helped to examine the effects of the socio-demographic
431 factors at different points of the Hb concentration, and thus present a comprehensive
432 picture of the effects. Another necessary strength is the use of nationally representative
433 data, making it possible for the results to be generalised to all women of reproductive
434 age in the respective countries. We could not establish causality in this study due to the
435 cross-sectional nature of the data. Also, missing data is an essential limitation of
436 secondary data analysis. However, due to the robust measures put place by DHS to
437 ensure the completeness of their datasets, missing data was not an issue in our study.

438

439 **Conclusions**

440 We used quantile regression to examine the effects of socio-demographic factors on
441 maternal Hb concentration. Our analysis demonstrated substantially that the various
442 putative socio-demographic factors have differential effects on maternal Hb
443 concentration at different points of the Hb distribution in all countries. Interventions and
444 programmes to address maternal anaemia must take into account the different effects
445 of the various socio-demographic factors on Hb concentration throughout the different
446 percentiles of the Hb distribution. It may help identify suitable interventions for groups
447 most in need.

448

449 **Acknowledgements**

450 We wish to express our gratitude to The DHS Program, USA, for providing us access to
451 the data. We also want to acknowledge the institutions of respective countries that
452 played critical roles in the data collection processes.

453

454 **Competing Interest**

455 The authors have no competing interests to declare.

456 **Funding**

457 This study did not receive funding from any source.

458

459 **Data Sharing Statement**

460 This study was a re-analysis of existing data that are publicly available from The DHS
461 Program at <http://dhsprogram.com/publications/publication-fr221-dhs-final-reports.cfm>.
462 Data are accessible free of charge upon registration with the Demographic and Health
463 Survey program (The DHS Program). The registration is done on the DHS website
464 indicated above.

465 **Authors' Contribution**

466 DAA conceived and designed the study, interpreted the results, wrote the first draft of
467 the manuscript, and contributed to the revision of the manuscript. DAA and ZTD
468 analysed the data. ZTD and CK contributed to study design, data interpretation, and
469 critical revision of the manuscript. All authors take responsibility for any issues that
470 might arise from the publication of this manuscript.

471

472

473 References

- 474 1. World Health Organization. Anaemia 2018 [Available from: https://www.who.int/health-topics/anaemia#tab=tab_1.
- 475 2. World Health Organization. WHO. Nutritional anaemias: tools for effective prevention and
476 control. Nutrition [cited 2020 6.5]. Available from:
477 <http://www.who.int/nutrition/publications/micronutrients/anaemias-tools-prevention-control/en/>.
- 478 3. Chaparro CM, Suchdev PS. Anemia epidemiology, pathophysiology, and etiology in low- and
479 middle-income countries. *Annals of the New York Academy of Sciences*. 2019;1450(1):15-31.
- 480 4. Kassebaum NJ. The Global Burden of Anemia. *Hematology/oncology clinics of North America*.
481 2016;30(2):247-308.
- 482 5. World Health Organization. Global Targets 2025: to improve maternal, infant and young child
483 nutrition [cited 2020 6.5]. Available from: <https://www.who.int/nutrition/global-target-2025/en/>.
- 484 6. World Health Organization. Nutrition: Global Nutrition Targets 2025: Anaemia policy brief 2014
485 [cited 2020 6.5]. Available from:
486 https://www.who.int/nutrition/publications/globaltargets2025_policybrief_anaemia/en/.
- 487 7. Lopez A, Cacoub P, Macdougall IC, Peyrin-Biroulet L. Iron deficiency anaemia. *Lancet*.
488 2016;387(10021):907-16.
- 489 8. Young MF. Maternal anaemia and risk of mortality: a call for action. *Lancet Glob Health*.
490 2018;6(5):e479-e80.
- 491 9. Stevens GA, Finucane MM, De-Regil LM, Paciorek CJ, Flaxman SR, Branca F, et al. Global,
492 regional, and national trends in haemoglobin concentration and prevalence of total and severe anaemia
493 in children and pregnant and non-pregnant women for 1995-2011: a systematic analysis of population-
494 representative data. *Lancet Glob Health*. 2013;1(1):e16-25.
- 495 10. United Nations. Sustainable Development Goals 2015 [cited 2020 6.5]. Available from:
496 <http://www.un.org/sustainabledevelopment/sustainable-development-goals/>.
- 497 11. Ayoya MA, Bendeck MA, Zagr e NM, Tchibindat F. Maternal anaemia in West and Central Africa:
498 time for urgent action. *Public health nutrition*. 2012;15(5):916-27.
- 499 12. van den Broek N. Anaemia in pregnancy in sub-Saharan countries. *European Journal of*
500 *Obstetrics and Gynecology and Reproductive Biology*. 2001;96(1):4-6.
- 501 13. Tako EA, Zhou A, Lohoue J, Leke R, Taylor DW, Leke RFG. Risk factor for placental malaria and its
502 effect on pregnancy outcome in Yaounde, Cameroon. *Am J Trop Med Hyg*. 2005;72.
- 503 14. Rahman MM, Abe SK, Rahman MS, Kanda M, Narita S, Bilano V, et al. Maternal anemia and risk
504 of adverse birth and health outcomes in low- and middle-income countries: systematic review and
505 meta-analysis. *Am J Clin Nutr*. 2016;103(2):495-504.
- 506

- 507 15. Black RE, Victora CG, Walker SP, Bhutta ZA, Christian P, de Onis M, et al. Maternal and child
508 undernutrition and overweight in low-income and middle-income countries. *Lancet*.
509 2013;382(9890):427-51.
- 510 16. Kavle JA, Stoltzfus RJ, Witter F, Tielsch JM, Khalfan SS, Caulfield LE. Association between
511 anaemia during pregnancy and blood loss at and after delivery among women with vaginal births in
512 Pemba Island, Zanzibar, Tanzania. *Journal of health, population, and nutrition*. 2008;26(2):232-40.
- 513 17. Christian P, Mullany LC, Hurley KM, Katz J, Black RE. Nutrition and maternal, neonatal, and child
514 health. *Seminars in perinatology*. 2015;39(5):361-72.
- 515 18. Chowdhury HA, Ahmed KR, Jebunessa F, Akter J, Hossain S, Shahjahan M. Factors associated
516 with maternal anaemia among pregnant women in Dhaka city. *BMC women's health*. 2015;15:77.
- 517 19. Kofie P, Tarkang EE, Manu E, Amu H, Ayanore MA, Aku FY, et al. Prevalence and associated risk
518 factors of anaemia among women attending antenatal and post-natal clinics at a public health facility in
519 Ghana. *BMC Nutrition*. 2019;5(1):40.
- 520 20. Mockenhaupt FP, Rong B, Günther M, Beck S, Till H, Kohne E, et al. Anaemia in pregnant
521 Ghanaian women: importance of malaria, iron deficiency, and haemoglobinopathies. *Transactions of the*
522 *Royal Society of Tropical Medicine and Hygiene*. 2000;94(5):477-83.
- 523 21. Gaillard R, Eilers PH, Yassine S, Hofman A, Steegers EA, Jaddoe VW. Risk factors and
524 consequences of maternal anaemia and elevated haemoglobin levels during pregnancy: a population-
525 based prospective cohort study. *Paediatric and perinatal epidemiology*. 2014;28(3):213-26.
- 526 22. Hakizimana D, Nisingizwe MP, Logan J, Wong R. Identifying risk factors of anemia among women
527 of reproductive age in Rwanda - a cross-sectional study using secondary data from the Rwanda
528 demographic and health survey 2014/2015. *BMC public health*. 2019;19(1):1662-.
- 529 23. Bentley ME, Griffiths PL. The burden of anemia among women in India. *European journal of*
530 *clinical nutrition*. 2003;57(1):52-60.
- 531 24. Gobezie M, Mekonnen Z, Alem M, Enawgaw B. Prevalence and Predictors of Maternal Anemia
532 during Pregnancy in Gondar, Northwest Ethiopia: An Institutional Based Cross-Sectional Study. *Anemia*.
533 2014;2014:108593.
- 534 25. Harding KL, Aguayo VM, Namirembe G, Webb P. Determinants of anemia among women and
535 children in Nepal and Pakistan: An analysis of recent national survey data. *Maternal & child nutrition*.
536 2018;14 Suppl 4(Suppl 4):e12478.
- 537 26. The DHS Program. Demographic and Health Surveys [Available from:
538 <http://dhsprogram.com/data/available-datasets.cfm>.
- 539 27. Amugsi DA, Dimbuene ZT, Kimani-Murage EW, Mberu B, Ezeh AC. Differential effects of dietary
540 diversity and maternal characteristics on linear growth of children aged 6-59 months in sub-Saharan
541 Africa: a multi-country analysis. *Public health nutrition*. 2017;20(6):1029-45.
- 542 28. Ghana Statistical Service (GSS), Ghana Health Service (GHS), ICF Macro. Ghana Demographic and
543 Health Survey 2008 Accra, Ghana: GSS, GHS, and ICF Macro; 2009.
- 544 29. Ministry of Health and Social Services (MoHSS), Macro International Inc. Namibia Demographic
545 and Health Survey 2006-07 Windhoek, Namibia and Calverton, Maryland, USA: MoHSS and Macro
546 International Inc. ; 2008.
- 547 30. Ghana Statistical Service (GSS), Ghana Health Service (GHS), ICF International. Ghana
548 Demographic and Health Survey 2014. Rockville, Maryland, USA: GSS, GHS, and ICF International.; 2015.
- 549 31. National Population Commission (NPC) [Nigeria], ICF International. Nigeria Demographic and
550 Health Survey 2013. Abuja, Nigeria, and Rockville, Maryland, USA: NPC and ICF International. . 2014.
- 551 32. National Bureau of Statistics-Kenya, ICF International. 2014 KDHS Key Findings. Rockville,
552 Maryland, USA: KNBS and ICF International. 2015.
- 553 33. World Medical Association. Declaration of Helsinki: Ethical Principles for Medical Research
554 Involving Human Subjects. *Bulletin of the World Health Organization*. 2001;79(4).

- 555 34. Koenker R, Bassett G. Regression quantiles *Econometrica*. 1978;46(1):33-50.
- 556 35. Amugsi DA, Dimbuene ZT, Bakibinga P, Kimani-Murage EW, Haregu TN, Mberu B. Dietary
557 diversity, socioeconomic status and maternal body mass index (BMI): quantile regression analysis of
558 nationally representative data from Ghana, Namibia and Sao Tome and Principe. *BMJ Open*. 2016;6(9):
559 doi:10.1136/bmjopen-2016-012615.
- 560 36. Amugsi DA, Dimbuene ZT, Bakibinga P, Kimani-Murage EW, Haregu TN, Mberu B. Dietary
561 diversity, socioeconomic status and maternal body mass index (BMI): quantile regression analysis of
562 nationally representative data from Ghana, Namibia and Sao Tome and Principe. *BMJ Open*. 2016;6(9).
- 563 37. DHS. Anaemia among women and children [cited 2020 22.04]. Available from:
564 <https://www.dhsprogram.com/pubs/pdf/OD28/12Chapter12.pdf>.
- 565 38. Balarajan YS, Fawzi WW, Subramanian SV. Changing patterns of social inequalities in anaemia
566 among women in India: cross-sectional study using nationally representative data. *BMJ Open*. 2013;3(3).
- 567 39. Lakew Y, Biadgilign S, Haile D. Anaemia prevalence and associated factors among lactating
568 mothers in Ethiopia: evidence from the 2005 and 2011 demographic and health surveys. *BMJ open*.
569 2015;5(4):e006001-e.
- 570 40. Friel J, Qasem W, Cai C. Iron and the Breastfed Infant. *Antioxidants (Basel)*. 2018;7(4):54.
- 571 41. Petraro P, Duggan C, Urassa W, Msamanga G, Makubi A, Spiegelman D, et al. Determinants of
572 anemia in postpartum HIV-negative women in Dar es Salaam, Tanzania. *European journal of clinical*
573 *nutrition*. 2013;67(7):708-17.
- 574 42. Abioye AI, Aboud S, Premji Z, Etheredge AJ, Gunaratna NS, Sudfeld CR, et al. Iron
575 Supplementation Affects Hematologic Biomarker Concentrations and Pregnancy Outcomes among Iron-
576 Deficient Tanzanian Women. *J Nutr*. 2016;146(6):1162-71.
- 577 43. Jorgensen JM, Yang Z, Lönnerdal B, Chantry CJ, Dewey KG. Effect of iron supplementation during
578 lactation on maternal iron status and oxidative stress: A randomized controlled trial. *Maternal & child*
579 *nutrition*. 2017;13(4).
- 580 44. Abu-Ouf NM, Jan MM. The impact of maternal iron deficiency and iron deficiency anemia on
581 child's health. *Saudi Med J*. 2015;36(2):146-9.
- 582 45. Ejigu BA, Wencheke E, Berhane K. Spatial pattern and determinants of anaemia in Ethiopia. *PloS*
583 *one*. 2018;13(5):e0197171-e.
- 584 46. Gebremedhin S, Enquesslassie F, Umeta M. Prevalence and correlates of maternal anemia in
585 rural Sidama, Southern Ethiopia. *African journal of reproductive health*. 2014;18(1):44-53.
- 586 47. Derso T, Abera Z, Tariku A. Magnitude and associated factors of anemia among pregnant women
587 in Dera District: a cross-sectional study in northwest Ethiopia. *BMC Res Notes*. 2017;10(1):359.
- 588

