

1 **Estimation of the number of general anesthesia cases based on a series of nationwide**
2 **surveys on Twitter during COVID-19 in Japan: A statistical analysis**

3 Yosuke Fujii MD ^{1,3}, Hiroki Daijo MD, Ph.D. ^{2,3}, Kiichi Hirota MD, Ph.D. ^{1,*}

4 ¹ Department of Human Stress Response Science, Institute of Biomedical Science, Kansai
5 Medical University, Hirakata, Osaka 573-1010, Japan

6 ² Department of Anesthesia, Takarazuka Daiichi Hospital, Takakazuka, Hyogo 665-0832,
7 Japan

8 ³ These authors contributed equally to this work

9

10 *Corresponding author

11 Kiichi Hirota

12 Department of Human Stress Response Science, Institute of Biomedical Science, Kansai

13 Medical University, Hirakata, Osaka, Japan

14 phone: +81-72-804-2526

15 email: hif1@mac.com / khirota-kyt@umin.ac.jp

16

17 **Keywords:** Coronavirus disease 2019 (COVID-19); twitter; general anesthesia; quadratic

18 programming

19

20 **Word count: 2144**

21 **The number of tables: 2**

22 **The number of figures: 5**

23

24 **Abstract**

25 **Purpose**

26 The coronavirus disease (COVID-19), caused by severe acute respiratory syndrome
27 coronavirus 2 (SARS-CoV-2), has spread to more than 200 other countries. In light of this
28 situation, the Japanese Government declared a state of emergency in seven regions of Japan
29 on April 7, 2020, under the provisions of the law. The medical care delivery system has been
30 under pressure. Although various surgical societies have published guidelines on which to
31 base their surgical decisions, it is not clear how general anesthesia has been performed and
32 will be performed in Japan.

33 **Methods**

34 One of the services provided by Twitter is a voting function—Twitter Polls—through which
35 anonymous surveys can be conducted. We used quadratic programming to analyze the results
36 of a series of 17 surveys on Twitter, over the course of 22 weeks. The analysis focused on
37 solving the mathematical optimizing problem on the status of operating restrictions. Public
38 data provided by the Japanese Government were used to estimate the current changes in the
39 number of general anesthesia performed in Japan.

40 **Results**

41 The minimum number of general anesthesia per week was estimated at 66.1%, compared to

42 2015, on April 27, 2020. The time series trend was compatible with the results reported by the

43 Japanese Society of Anesthesiologists.

44 **Conclusion** The number of general anesthesia was reduced by up to two-thirds during the

45 COVID-19 pandemic in Japan, and was quantitatively estimated using the Twitter quick

46 questionnaire.

47

48 **Introduction**

49 Coronavirus (COVID-19), caused by severe acute respiratory syndrome coronavirus 2
50 (SARS-CoV-2), was first reported in Wuhan, Hubei, China, and has since spread to more than
51 200 other countries around the world at the time of writing [1]. In light of this situation, the
52 Japanese Government declared a state of emergency in seven regions of Japan on April 7,
53 2020, under the provisions of the law. This declaration was extended to the entire nation on
54 April 17, 2020. The incidence of pneumonia from COVID-19 is considerably higher than
55 from seasonal influenza, and the number of cases with no identifiable route of infection has
56 increased rapidly [2, 3]. The medical care delivery system has been under pressure [4].
57 Emphasis should be placed on maintaining medical care systems. Although various surgical
58 societies have published guidelines on which to base their surgical decisions
59 (<https://www.facs.org/covid-19/clinical-guidance/elective-case>;
60 [https://www.cms.gov/files/document/cms-non-emergent-elective-medical-](https://www.cms.gov/files/document/cms-non-emergent-elective-medical-recommendations.pdf)
61 [recommendations.pdf](https://www.cms.gov/files/document/cms-non-emergent-elective-medical-recommendations.pdf)), it is not clear how general anesthesia has been performed.
62 Social networks such as Twitter are becoming a part of our society, as various information is
63 being accumulated on the Web [5]. One of the services provided is a voting function called
64 Twitter Polls. Using this function, anonymous surveys can be conducted on Twitter in Japan
65 [6].

66 A series of surveys were conducted on Twitter on the status of operating restrictions. We
67 analyzed the results using quadratic programming to solve a mathematical optimizing
68 problem. Public data provided by the Japanese Government were used to estimate and
69 compare the current changes in the number of general anesthesia carried out in Japan. In
70 addition, the estimated time series trend was validated to be compatible with the results of the
71 surveys sponsored by the Japanese Society of Anesthesiologists (JSA).

72

73 **Methods**

74 **Twitter Surveys**

75 Using the Twitter account of @dajhiroki, a board-certified anesthesiologist with
76 approximately 1,300 followers, Twitter Polls was used to conduct seventeen 24-hour surveys,
77 which were held approximately one week apart from each other, from March 13 to August 14,
78 2020 (Supplementary Information S1). During the period, the spread of COVID-19 had
79 become a social problem in Japan.

80 The surveys used the same wording throughout the period, in the form of a choice of one of
81 the following four options:

- 82 1. No surgical restrictions
- 83 2. Partial restrictions (more than half of the usual)

84 3. Extensive restrictions (less than half of the usual)

85 4. No scheduled surgery

86

87 **Transition of responses to the survey**

88 Suppose there were $R = 4$ discrete categories of responses, transition between the responses

89 during the questionnaire period would be estimated (Fig. 1). P was a $R \times R$ transition matrix

90 whose elements p_{ij} show the probability transition from the i th category at time t to the j th

91 category at time $t + 1$. The transition matrix was

$$P = \begin{bmatrix} p_{11} & p_{12} & p_{13} & p_{14} \\ p_{21} & p_{22} & p_{23} & p_{24} \\ p_{31} & p_{32} & p_{33} & p_{34} \\ p_{41} & p_{42} & p_{43} & p_{44} \end{bmatrix}, \quad (1)$$

92 under the constraints of

$$\left\{ \begin{array}{l} p_{13}, p_{14}, p_{24}, p_{31}, p_{41}, p_{42} = 0 \\ 0 \leq p_{11}, p_{12}, p_{21}, p_{22}, p_{23}, p_{32}, p_{33}, p_{34}, p_{43}, p_{44} \leq 1 \\ p_{11} + p_{12} = 1 \\ p_{21} + p_{22} + p_{23} = 1 \\ p_{32} + p_{33} + p_{34} = 1 \\ p_{43} + p_{44} = 1 \end{array} \right. \quad (2)$$

93 The number of the responses to the web questionnaire at time t was aggregated as the

94 proportion $y_i(t)$, $i = \{1, 2, 3, 4\}$. When the individual transition was not available, it was not

95 possible to estimate the transition matrix from individual transition data using ordinary least

96 squares method. However, quadratic programming method could estimate the transition

97 matrix from proportional data [7]. The stochastic model of the relation between the actual

98 response counts and the estimated occurrence of $y_i(t)$ was described below with the error

99 term $u_j(t)$;

$$y_i(t+1) = \sum_i^R y_i(t)p_{ij} + u_j(t). \quad (3)$$

100 This equation could be written in linear algebraic form as follows:

$$y = Xp + u, \quad (4)$$

101 where

$$y = [y_1(1), y_1(2), \dots, y_1(t), y_2(1), y_2(2), \dots, y_2(t), \dots, y_R(1), y_R(2), \dots, y_R(t)]', \quad (5)$$

102 and

$$X_j = \begin{bmatrix} y_1(0) & y_2(0) & \dots & y_R(0) \\ y_1(1) & y_2(1) & \dots & y_R(1) \\ \vdots & \vdots & \ddots & \vdots \\ y_1(T-1) & y_2(T-1) & \dots & y_R(T-1) \end{bmatrix}, \quad (6)$$

103 for $j = \{1, 2, 3, 4\}$ so that

$$X = \begin{bmatrix} X_1 & 0 & \dots & 0 \\ 0 & X_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & X_R \end{bmatrix}, \quad (7)$$

$$p = [p_1(1), p_1(2), \dots, p_1(t), p_2(1), p_2(2), \dots, p_2(t), \dots, p_R(1), p_R(2), \dots, p_R(t)]', \quad (8)$$

104 and

$$u = [u_1(1), u_1(2), \dots, u_1(t), u_2(1), u_2(2), \dots, u_2(t), \dots, u_R(1), u_R(2), \dots, u_R(t)]'. \quad (9)$$

105 Estimation was performed minimizing the squared error term $|u|^2$ as

$$\begin{aligned} |u|^2 &= (\mathbf{y} - \mathbf{Xp})^2 \\ &= |\mathbf{y}|^2 - 2\mathbf{y}'\mathbf{Xp} + \mathbf{p}'\mathbf{X}'\mathbf{Xp}. \end{aligned} \quad (10)$$

106 The objective of quadratic programming was to find a vector p that would minimize

$$\frac{1}{2}\mathbf{p}'\mathbf{X}'\mathbf{Xp} - \mathbf{y}'\mathbf{Xp}, \quad (11)$$

107 subject to

$$\mathbf{Ap} \leq \mathbf{b}, \quad (12)$$

108 with a real matrix A and a real vector b as constraints for p denoted in equation (2).

109

110 **The number of general anesthesia under restriction**

111 The restriction rate of the number of general anesthesia was assumed for each survey

112 response: (1) cases of no surgical restrictions were almost the same as the past statistics, (2)

113 partial restrictions was about 70% restriction rate compared to past statistics, (3) extensive

114 restrictions was about 40% restriction rate compared to past statistics, and (4) no scheduled

115 surgery was about 10% restriction rate compared to the statistics (Fig. 2). Under such

116 assumption, the restriction rate for each hospital h at week t , $r_{t,i,h}$ was generated from the

117 probability distribution as follows:

118 (1) for no surgical restrictive hospitals, $r_{t,1,h}$ was generated from uniform distribution, U ,

119 ranging from 0.95 to 1.05.

$$r_{t,1,h} \sim U(0.95, 1.05), \quad (13)$$

120 (2–4) for restrictive management hospitals, $r_{t,i,h}$, $i = \{2, 3, 4\}$, was generated from beta
121 distribution, $Beta(\alpha, \beta)$. Their parameters $(\alpha_{t,i,h}, \beta_{t,i,h})$, $i = \{2, 3, 4\}$ were defined so that the
122 means were 0.7, 0.4, and 0.1, and the variance was 0.005, respectively (Fig. 2).

$$r_{t,i,h} \sim Beta(\alpha_{t,i,h}, \beta_{t,i,h}), \quad (14)$$

123 for $i = \{2, 3, 4\}$. The number of general anesthesia performed at hospital h in the t th week,
124 $R_{h,t}$, was generated as follows:

$$R_{h,t} \sim \begin{cases} \text{round}(N_h r_{t,1,h}) & \text{if } i = 1 \\ \text{Binomial}(r_{t,i,h}, N_h) & \text{otherwise} \end{cases}, \quad (15)$$

125 where N_h is the reference number of the general anesthesia performed at each hospital (data
126 available from e-Stat), *round* is a rounding function, and $\text{Binomial}(p, N)$ is a binomial
127 sampling function that performs N Bernoulli trials with probability p .

128 The survey response for each hospital at $t = 1$ was randomly allocated to the i th response
129 according to $y_i(t)$, and its status transitioned to the j th response by p_{ij} .

130 For sensitivity analysis, two other scenarios were considered. An optimistic scenario assumed
131 relatively high performance in operating rooms, with an average of 80% for “2. Partial
132 restrictions,” 50% for “3. Extensive restrictions,” and 25% for “4. No scheduled surgery.” A
133 pessimistic scenario assumed extremely low performance, with an average of 50% for “2.

134 Partial restrictions,” 30% for “3. Extensive restrictions,” and 3% for “4. No scheduled
135 surgery.”

136 Analysis and estimation were performed using the quadprog package (version 1.5.8) within
137 the R programming language (version 3.4.4). Point estimates and their statistical uncertainty
138 were presented as 95% confidential intervals (CI) with 1000 iteration. The data were obtained
139 from the Japan official statistics portal site e-Stat, a public database in Japan ([https://www.e-](https://www.e-stat.go.jp/stat-search/file-download?statInfId=000031565245&fileKind=0)
140 [stat.go.jp/stat-search/file-download?statInfId=000031565245&fileKind=0](https://www.e-stat.go.jp/stat-search/file-download?statInfId=000031565245&fileKind=0)), and were
141 preprocessed for analysis. The code is available online
142 (<https://github.com/yfujii08/covid19generalanesthesia>) and is available upon request from the
143 corresponding author.

144

145 **Nationwide report of the estimation of the number of general anesthesia from the**
146 **Japanese Society of Anesthesiologists**

147 The JSA determined the extent of the restriction of general anesthesia using questionnaires.
148 They sent questionnaires to all authorized institutes (n=1415) from April 23, 2020,
149 approximately every week. The proportion of the number of general anesthesia compared to
150 2015 was estimated and disclosed to only the members of JSA
151 (https://anesth.or.jp/img/upload/ckeditor/files/2004_07_08.pdf).

152

153 **Ethics**

154 This study is exempted from institutional review board approval because this study does not
155 contain human participants or research material derived from human participants.

156

157 **Results**

158 From March 13 to August 14, 2020, responses were solicited from 17 surveys almost weekly
159 via a Twitter account (@dajhiroki) in the Japanese language, with the number of responses
160 received per survey ranging from 47 to 288 (Supplementary Information S1). The results of
161 the surveys are presented in Table 1.

162 In the public database e-Stat, 3501 hospitals were registered as facilities that provide general
163 anesthesia. Of the 3311 hospitals that reported the number of general anesthesia, 1989
164 performed more than 100 general anesthesia.

165 From the survey conducted in the first week (March 13, 2020), 89.2% had no surgical
166 restrictions, but this proportion decreased to 24.7% in the survey conducted in the seventh
167 week (April 27, 2020) then slightly recovered to 38.9% (May 17, 2020). The proportion
168 increased to 87.2% (July 4, 2020) and was finally 80.0% at the end of the questionnaire
169 administration (July 31, 2020). The proportions of partial restrictions, extensive restrictions,

170 and no scheduled surgeries were 5.4%, 3.6%, and 1.8% (March 13, 2020); 37.8%, 29.5%, and
171 8.0% (April 27, 2020); 44.2%, 11.5%, and 5.3% (May 17, 2020); and 11%, 4%, and 5% (July
172 31, 2020), respectively (Fig. 3). The estimated proportion of each response and its transition
173 are shown in Fig. 3 and Table 1.

174 The number of general anesthesia performed decreased as the restrictions increased (Fig. 4
175 and Table 2). Before COVID-19, 4.45×10^4 cases of general anesthesia were performed per
176 week in Japan. The number of general anesthesia (10^4 per week unit) was estimated to be 4.21
177 [95% CI: 4.16, 4.26] in the first week (March 13, 2020), 2.95 [2.87, 3.02] in the seventh week
178 (April 27, 2020), and 4.10 [4.04, 4.16] in the last week survey (July 31, 2020) (Table 2).

179 Sensitivity analysis showed that the number of general anesthesia performed decreased
180 according to the intensity of the restrictions (Fig. 5). In the first week survey (March 13,
181 2020), the optimistic scenario had a median of 4.21×10^4 /week, while pessimistic scenario
182 had that of 4.14. In the seventh (April 27, 2020) and last week survey (July 31, 2020), the
183 optimistic scenario medians were 3.28 and 4.19, while those of the pessimistic scenario were
184 2.46 and 3.95, respectively. The maximum difference between optimistic and pessimistic
185 scenarios was 1.34 folds at May 2, 2020.

186 The comparison between the estimation of this survey and the nationwide reports from the
187 surveys of the JSA was shown in Fig. 5. The data before the end of April is missing because

188 the JSA questionnaire started from April 23, 2020. The JSA data showed a gradual decrease
189 in the number of general anesthesia compared to 2019: 87.3%, 79.2%, and 41.1% on April 23,
190 April 27, and May 4, respectively. After that, the number of the general anesthesia gradually
191 recovered parallel to the estimation of this survey.

192

193 **Discussion**

194 It is difficult to adequately control for the level of limitation of usual surgical care during the
195 spread of COVID-19. It is determined by adherence to official guidelines, restrictions on
196 medical supplies and medical staff, and the level of need for surgery. However, because
197 circumstances can change quickly, it is often determined with reference to the extent of
198 restrictions at a medical facility other than the home facility.

199 The staged approach—recommended by the Centers for Medicare and Medicaid Services
200 (CMS) and the American College of Surgeons (ACS)—is used as a guide to determine how to
201 perform surgery in situations where the preservation of ventilators and personal protective
202 equipment is necessary, and the ICU has been compromised—or is expected to be
203 compromised soon.

204 The number of respondents to our surveys increased over time, and it is thought that the
205 number of respondents was dependent on their knowledge of the status of operational

206 limitations. By using a series of surveys conducted through Twitter Polls, we were quickly
207 able to determine the extent of the restrictions on surgery at medical facilities across the
208 country about one month ahead of the survey by the JSA. By periodically soliciting responses
209 to the same survey on Twitter, we were able to estimate the extent to which operations would
210 be restricted nationally—over time.

211 Our methodology in this study is an estimation method that uses mathematical analysis. We
212 used public data from a database operated by the Japanese Government. The number of
213 hospitals accredited by the JSA is 1415. However, in this study, 1989 hospitals that perform
214 at least 100 surgeries per year under general anesthesia were used. The number of operations
215 performed at each hospital varies—mainly according to the size of the hospital. On the other
216 hand, since neither the size of the hospitals to which the survey respondents belonged nor the
217 changes in the number of operations were known, we used mathematical analysis to calculate
218 the number of operations.

219 There are several limitations in this study. First, the number and type of survey responses is a
220 potential limitation. It is possible that the number of anesthesiologists using Twitter was not
221 sufficient to obtain a definitive estimation, and there may be some bias in the responses from
222 those that do use it. Responses were low initially, with 47 in the early stages of the surveys,
223 but increased to 288 in the latter half. The growing penetration of the surveys and increasing

224 familiarity with the issue of operating room restrictions may have contributed to an increase
225 in participation, but this may have been biased. Second, we calculated the number of surgeries
226 as 95% to 105% for the criteria: "1. No surgical restrictions," 70% for "2. Partial restrictions,"
227 40% for "3. Extensive restrictions," and 10% for "4. No scheduled surgery." In the sensitivity
228 analysis, two scenarios were assumed, but the number of general anesthesia performed could
229 be approximately a 1.34-fold difference following the period of the surveys. Because it is
230 difficult to estimate, accurately, the extent of restrictions, intuitive values were adopted. It
231 needs to be compared with the descriptive statistics after the COVID-19 era.

232 In conclusion, we analyzed the results of a series of surveys on Twitter on the status of
233 operating restrictions using quadratic programming to solve a mathematical optimizing
234 problem. Public data provided by the Japanese Government were used to estimate and
235 compare the current changes in the number of general anesthesia carried out in Japan. The
236 number of general anesthesia was reduced by up to two-thirds during the COVID-19
237 pandemic in Japan, and was quantitatively estimated using the Twitter quick questionnaire.

238

239 **Acknowledgements**

240 We would like to thank Editage (www.editage.com) for English language editing.

241

242 **Availability of data and material**

243 All data generated or analyzed during this study are included in this published article. The
244 datasets used and/or analyzed during the current study are available from the corresponding
245 author on reasonable request.

246

247 **Ethics declarations**

248 This study is exempted from institutional review board approval because this study does
249 contain no human participants, or research material derived from human participants.

250

251 **Competing interests**

252 The authors declare no competing interests.

253

254 **Funding**

255 This work was supported by a research grant from the Kansai Medical University (KMU)
256 research consortium to K.H., the branding program as a world-leading research university on
257 intractable immune and allergic diseases from MEXT Japan.

258

259

260 **References**

- 261 1. Zhu N, Zhang D, Wang W, Li X, Yang B, Song J, et al. A Novel Coronavirus from
262 Patients with Pneumonia in China, 2019. *N Engl J Med.* 2020;382:727-33; doi:
263 10.1056/NEJMoa2001017.
- 264 2. Guan WJ, Ni ZY, Hu Y, Liang WH, Ou CQ, He JX, et al. Clinical Characteristics of
265 Coronavirus Disease 2019 in China. *N Engl J Med.* 2020;382:1708-20; doi:
266 10.1056/NEJMoa2002032.
- 267 3. Meselson M. Droplets and Aerosols in the Transmission of SARS-CoV-2. *N Engl J*
268 *Med.* 2020; 382: 2063 doi: 10.1056/NEJMc2009324.
- 269 4. Zarzaur BL, Stahl CC, Greenberg JA, Savage SA, Minter RM. Blueprint for
270 Restructuring a Department of Surgery in Concert With the Health Care System During a
271 Pandemic: The University of Wisconsin Experience. *JAMA Surg.* 2020; 155: 628-35 doi:
272 10.1001/jamasurg.2020.1386.
- 273 5. Aramaki E, Maskawa S, Morita M. Twitter Catches The Flu: Detecting Influenza
274 Epidemics using Twitter. *Proceedings of the Conference on Empirical Methods in Natural*
275 *Language Processing.* 2011:1568–76.
- 276 6. Sang E, Bos J. Predicting the 2011 Dutch Senate Election Results with Twitter.
277 *Proceedings of the Workshop on Semantic Analysis in Social Media.* 2012:53–60.

278 7. Matthew T. Jones. Estimating Markov Transition Matrices Using Proportions Data:

279 An Application to Credit Risk. International Monetary Fund; 2005. No. 5-219.

280

281

282 **Tables**

283 Table 1: Results of the Twitter surveys. The sum did not necessarily equal to 100% due to
 284 rounding.

Date of survey (2020)	3/13	3/20	3/27	4/6	4/10	4/17	4/27	5/2	5/9	5/17	5/22	6/19	7/4	7/24	7/31	8/7	8/14
Number of votes	112	47	73	160	221	232	288	184	93	113	54	157	86	142	100	77	107
1. No surgical restrictions	89%	77%	82%	64%	46%	27%	25%	29%	41%	39%	46%	78%	87%	80%	80%	87%	82%
2. Partially restricted	5%	17%	7%	23%	38%	38%	38%	40%	37%	44%	41%	19%	10%	16%	11%	10%	8%
3. Extensively restricted	4%	2%	4%	7%	13%	27%	30%	23%	19%	12%	9%	2%	2%	1%	4%	1%	1%
4. No scheduled surgery	2%	4%	7%	6%	3%	8%	7%	8%	3%	5%	4%	1%	0%	4%	5%	1%	8%

285

286 Table 2 : Estimated number of general anesthesia (10^4 cases per week) in Japan during the

2020	3/13	3/20	3/27	4/6	4/10	4/17	4/27	5/2	5/9	5/17	5/22	6/19	7/4	7/24	7/31	8/7	8/14
2.5%	4.16	4.03	4.02	3.97	3.71	3.22	2.87	2.88	3.11	3.27	3.38	3.75	4.14	4.15	4.04	4.11	4.16
50%	4.21	4.09	4.08	4.03	3.78	3.30	2.95	2.95	3.19	3.34	3.45	3.82	4.19	4.19	4.10	4.16	4.21
97.5%	4.26	4.14	4.14	4.08	3.84	3.36	3.02	3.03	3.26	3.42	3.52	3.88	4.23	4.24	4.16	4.21	4.25

287 period of the surveys.

288

289

290 **Figure Legends**

291 Figure 1: Transition scheme of the proportion of surveys. (1: green) no surgical restrictions,
292 (2: yellow) partial restrictions, (3: orange) extensive restrictions, and (4: red) no scheduled
293 surgery. p_{ij} is the transition probability from response i to response j .

294

295 Figure 2: Distribution of the degree of restriction for each response group. The mean
296 percentages of the restrictions were generated from probability distributions. In particular, no
297 surgical restrictions (green) were generated from the uniform distribution $U(0.95,1.05)$, while
298 partial restrictions (yellow), extensive restrictions (orange), and no scheduled surgery (red)
299 were generated from the beta distribution, $Beta(\alpha,\beta)$. The set of parameters for beta
300 distribution, (α,β) , were defined so that the respective means were 0.7, 0.4, and 0.1, and the
301 variances were 0.005.

302

303 Figure 3: Transition of the proportion of responses to the survey. The solid line was the
304 estimated transition and the dots were actual data.

305

306 Figure 4: Estimation of the number of general anesthesia in Japan from the survey results.

307 The violinplots show the distribution of the estimated number of general anesthesia

308 performed at 1,989 hospitals in Japan.

309

310 Figure 5: Estimation of the number of general anesthesia in Japan from the results of the

311 surveys for the three scenarios, as a sensitivity analysis and comparison between the

312 estimation of the number of general anesthesia and the report from the Japanese Society of

313 Anesthesiologists (JSA). The JSA reported the results of the estimated number of the general


314 anesthesia from 1415 authorized institutes from April 23, 2020.

315


316 **Additional information**


317 Supplementary Information S1

It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/) .



It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/) .





It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/) .

