

Mathematical model of COVID-19 spread in Turkey and South Africa

Abdon ATANGANA^{1,*}, Seda İĞRET ARAZ

¹AtanganaA@ufs.ac.za, Institute for Groundwater Studies,
Faculty of Natural and Agricultural Sciences, University of the Free State,
South Africa

*Department of Medical Research, China Medical University Hospital,
China Medical University, Taichung, Taiwan

²sedaaraz@siirt.edu.tr, Department of Mathematic Education,
Faculty of Education, Siirt University, Siirt 56100, Turkey

May 5, 2020

Abstract

A comprehensive study about the spread of COVID-19 cases in Turkey and South Africa has been presented in this paper. An exhaustive statistical analysis encompassing arithmetic, geometric, harmonic means, standard deviation, skewness, variance, Pearson and Spearman correlation was derived from the data collected from Turkey and South Africa within the period of 11 March 2020 to 3 May 2020 and 05 March and 3 of May respectively. It was observed that in the case of Turkey, a negative Spearman correlation for the number of infected class and a positive Spearman correlation for both the number of deaths and recoveries were obtained. This implied that the daily infections could decrease, while the daily deaths and number of recovered people could increase under current conditions. In the case of South Africa, a negative Spearman correlation for both daily deaths and daily infected people was obtained, indicating that these numbers may decrease if the current conditions are maintained. The utilization of a statistical technique predicted the daily number of infected, recovered and dead people for each country; and three results were obtained for Turkey, namely an upper boundary, a prediction from current situation and lower boundary. The prediction shows that Turkey may register in the near future approximately more than 6000 new infections in a day as worst case scenario; and less than 300 cases in the perfect scenario. However, the country could register in the near future a daily number of 27000 people recovered from COVID-19 in the perfect scenario; and less than 5000 people in a worst scenario. Moreover, Turkey in a worst-case scenario could record a high number of approximately 200 deaths per day; and less than 150 deaths in a perfect scenario. Similarly, in the case of South Africa, the prediction results show that in the near future the country could register about 500 new infected cases daily and more than 25 deaths in the worst scenario; while in a perfect scenario less than 50 new infected and zero death cases could be recorded. The histograms of the daily number of newly infected, recovered and death showed a sign of lognormal and normal distribution, which is presented using the Bell curving method parameters estimation. A new mathematical model COVID-19 comprised of nine classes was suggested; of which a formula of the reproductive number, well-posedness of the solutions and the stability analysis were presented in details. The suggested model was further extended to the scope of nonlocal operators for each case; whereby the Atangana-Seda numerical method was used to provide numerical solutions, and simulations were performed for different non-integer numbers. Additionally, sections devoted to control optimal and others dedicated to compare cases between Turkey and South Africa with the aim to comprehend why there are less numbers of deaths and infected people in South Africa than Turkey were presented in details.

Keywords: Statistical analysis, Bell curve, prediction, new COVID-19 model, non-local operators, optimal control, Turkey vs South Africa.

1 Introduction

It is a Thursday morning, 5 March in South Africa, everybody is busy with his daily routine, when the National Institute for Communicable Diseases confirmed the first positive case of COVID-19. A situation that was

known for other countries now has become true and real in South Africa. How did we get here? The outbreak of COVID-19 started in China, Wuhan City, around December 2019; but within a short period the spread crossed over to some countries in Europe like United Kingdom, Italy, Spain and France. The first confirmed patient was a 38-year old male who visited Italy and arrived back in South Africa on March 1, 2020. The patient, after noticing symptoms of fever, malaise, a sore throat, cough and headache consulted a private general practitioner on March 3. From the 5 March to 15 March 2020 the number of infected people increased significantly, as a result on 15 March 2020, a national state of disaster was declared by the President of South Africa to mitigate the spread of COVID-19. This announcement was followed by measures including immediate travel restrictions and closure of schools from 18 March 2020. On 23 March, the South African government announced a country lockdown with effect on 26 March 2020. By the end of April, South Africa officially had 5647 confirmed cases. To date, South Africa is officially confirmed an African country with more confirmed cases, with 3471 active confirmed cases, 2073 recovered and 103 deaths due to COVID-19. In the case of Turkey, the first case of COVID-19 was confirmed and recorded on the 11 March 2020. Four days later, Turkey registered its first death caused by COVID-19. COVID-19 spread like wildfire in Turkey; in which by the 21 st of April the country had confirmed approximately 95591 cases of infected people, with 14918 number of recovered people and 2259 deaths recorded. The rapid spreading of COVID-19, has raised the total number of confirmed cases to 120200, of which 48900 have recovered and 3200 have died by the end of April 2020. In comparison to other European countries such as Iran, it is recorded that the total number of confirmed cases in Turkey surpassed it exceedingly; resulting in Turkey to be categorized as the most affected country in terms of numbers of confirmed cases within the settlement of the Middle East. Furthermore, Turkey's total number of confirmed cases by the 20 April was also recorded to exceed that of China; even though there were some raised concerns that the total confirmed cases in China could have been underestimated. The consideration of these statistics prompted researchers from Turkey and South Africa to undertake research in different fields of science, technology and engineering in the last 3 months, since their future is left uncertain. As the virologists are focusing their attention in developing a vaccine that could be used to prevent the spread of the deadly virus; mathematicians rely on modelling techniques to produce multi-scenarios models that could be utilized to foresee the future [1-6]. Therefore, as mathematicians our role is to use and apply mathematical tools, particularly mathematical models, on suggested scenarios that could be helpful in predicting the future. In this paper, we present a detailed analysis of spread in both countries and structured the paper as follows: Section 2 presents a detailed statistical analysis of COVID-19 spread in Turkey. Then we present a detailed statistical analysis of COVID-19 spread in South Africa. Also after using the inverse problem approach and the Bell curving approach we present the parameter's estimation, we present a comparative analysis between Turkey and South Africa. In Section 3, we suggest a new mathematical model of COVID-19 that takes into account nine classes, including, susceptible, infected with 5 sub-classes, recovered class, death and vaccinated. Then presents the positivity of solutions of the model as well as the reproduction number; and also deals with local and global asymptotic stability of disease free equilibrium and endemic points. In Section 4 we present an analysis of the suggested model with non-local operators. In section 5, we present numerical the suggested mathematical model for COVID-19 using Atangana-Seda scheme for fractional and fractal-fractional operators. In section 6, we present the optimal control of the disease. Finally we present a discussion, recommendations and conclusion respectively.

2 Statistical analysis of COVID-19 spread in Turkey and South Africa

To understand the impact of COVID-19, collection of numbers of daily new infected, recovered and deaths are performed all over the globe, such process follows a discrete approach. Thus, to understand and predict the impact of the Covid-19 on humans, statistics is associated with such collection, analysis, interpretation, organization and presentation. We shall recall that, this mathematical branch is wider applicable in numerous academic fields for example, natural and social science, business and government. Some important and useful statistical formula are means, variance, skewness, correlation, linear regression, Pearson's correlation coefficient, Spearman's rank correlation test and many order. In this section, we present some formulas that will be used in this work for interpretation and prediction purposes. We define a data set whose values can be chosen as x_1, x_2, \dots, x_n . We start with the arithmetic mean, \bar{x} , which states the mean of the x_1, x_2, \dots, x_n . The arithmetic

mean can be computed as

$$\text{Arithmetic mean} = \frac{1}{n} \sum_{i=1}^n x_i. \quad (2.1)$$

The formula of the geometric mean is

$$\text{Geometric mean} = \left(\prod_{i=1}^n x_i \right)^{1/n}. \quad (2.2)$$

The formula of the harmonic mean is

$$\text{Harmonic mean} = \left(\frac{\sum_{i=1}^n \frac{1}{x_i}}{n} \right)^{-1}. \quad (2.3)$$

The formula of the standard deviation is

$$\text{Standard deviation} = \left(\frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2 \right)^{1/2}. \quad (2.4)$$

The formula of the skewness is

$$\text{Skewness} = \frac{\frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^3}{\left(\frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2 \right)^{3/2}}. \quad (2.5)$$

The formula of the variance is

$$\text{Variance} = \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^2. \quad (2.6)$$

The formula of the covariance is

$$\text{Covariance} = \sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y}). \quad (2.7)$$

The formula of the Pearson correlation is

$$\text{Pearson correlation} = \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{\bar{xy}}. \quad (2.8)$$

The formula of the Spearman correlation is

$$\text{Spearman correlation} = 1 - \frac{6 \sum_{i=1}^n (\text{rank}x_i - \text{rank}y_i)^2}{n(n^2 - 1)} \quad (2.9)$$

where rank enables to compared a numeric value with other values in the same list.

2.1 Statistical analysis for Turkey

In this section, we aim to provide a detailed statistical analysis of the collected data from Turkey. These data include, daily number of new infected, daily numbers of deaths, daily numbers of recovered and finally daily numbers of tested individuals. The collected data are from 11 March 2020 to 3 May 2020. The main aim of this section is to predict what could possibly happen in the near future using the reliability level method, additionally, to find which distribution each class follows. With the collected data, we will first present histogram, pie chart and nonlinear graphs for each class. The histograms will help identify the density of probability associated to each set of collected data. Additionally, we provide a polynomial fitting against collected. The results are presented in Figure 1 to 16. For each case, we present arithmetic, geometric, harmonic means respectively, skewness, variance, covariance, Pearson correlation and Spearman correlation and their results are presented in Table 1.

In Figure 1, 2 and 3, we present some statistical simulation about number of infected people due to COVID-19 in Turkey from 11 March 2020 to 3 May 2020.




Figure 1. Number of infected people in Turkey from 11 March 2020 to 3 May 2020.




Figure 2. Number of infected people in Turkey from 11 March 2020 to 3 May 2020.




Figure 3. Number of infected people in Turkey from 11 March 2020 to 3 May 2020.

In Figure 4,5 and 6, we present some statistical simulation about number of recovered people due to COVID-19 in Turkey from 11 March 2020 to 3 May 2020.




Figure 4. Number of recovered people in Turkey from 11 March 2020 to 3 May 2020.

NUMBER OF RECOVERED




Figure 5. Number of recovered people in Turkey from 11 March 2020 to 3 May 2020.




Figure 6. Number of recovered people in Turkey from 11 March 2020 to 3 May 2020.

In Figure 7, 8 and 9, we present some statistical simulation about number of died people due to COVID-19 in Turkey from 11 March 2020 to 3 May 2020.




Figure 7. Number of death people in Turkey from 11 March 2020 to 3 May 2020.




Figure 8. Number of death people in Turkey from 11 March 2020 to 3 May 2020.




Figure 9. Number of death people in Turkey from 11 March 2020 to 3 May 2020.

In Figure 10, 11 and 12, we present some statistical simulation about number of tested people due to COVID-19 in Turkey from 11 March 2020 to 3 May 2020.




Figure 10. Number of tested people in Turkey from 11 March 2020 to 3 May 2020.




Figure 11. Number of tested people in Turkey from 11 March 2020 to 3 May 2020.




Figure 12. Number of tested people in Turkey from 11 March 2020 to 3 May 2020.

2.1.1 Regression analysis

Regression analysis which is also used in epidemiologic research enables us to examine amongst a set of variables. Here aim is to estimate outcomes benefitting from this set of variables. To do this, we find a prediction model in which we obtain model that fits best to the considered data and explains the response variable. We can utilize all possible independent variables, interactions, transformations with these models. To evaluate goodness of fit

for the obtained model, we can utilize *R square* ability which is one of the different techniques used in regression diagnostics.

Linear regression models are given by

$$y = \beta_0 + \beta_i x_i + e_i \quad (2.10)$$

where β_0, β_i are the unknown constants, x_i are the independent variables, y is the dependent variable and e_i are the error terms in given data. If the value of *R square* is close to zero, this means that the significance of fit for model is unsuitable to predict outcomes. In other words, the obtained model used is not suitable for the given data and the obtained model should be left aside and another model should be found.

If the value of *R square* is close to one, this means that the significance of fit for model is suitable to predict outcomes. In this case, it can be passed to the other step of control analysis.

We firstly present a predictive analysis for infected people. According to the results obtained, we obtain a linear regression which is calculated as

$$y = -29772.4 + 2786.833x. \quad (2.11)$$

F - test was calculated as 1.94×10^{-32} . *R square* was calculated as 0.93445. We can conclude from these data that the significance of fit for the obtained model is suitable for the considered data. Also we present polynomial regression which is calculated as

$$y = -0.0551x^4 + 9682.6x^3 - 0.6 \times 10^{-8}x^2 + 0.2 \times 10^{-13}x - 0.2 \times 10^{-17}. \quad (2.12)$$

For this polynomial, *R square* was calculated as 0.9993. We present polynomial fitting data for infected people from 11 March to 3 May 2020.




Figure 13. Polynomial fitting data for infected people in Turkey from 11 March 2020 to 3 May 2020.

We present a predictive analysis for recovered people. According to the results obtained, we get a linear regression which can be calculated as

$$y = -13029.8 + 845.9233x. \quad (2.13)$$

F - test was calculated as 1.39×10^{-12} . *R square* was calculated as 0.622381. We can say that the significance of fit for model is not enough suitable for the considered data. To overcome this case, we can suggest another regression model

$$y = 0.028x^4 - 4922x^3 + 0.3 \times 10^{-8}x^2 - 0.9 \times 10^{-12}x + 0.1 \times 10^{-17} \quad (2.14)$$

which is polynomial. For this polynomial, *R square* was calculated as 0.9987. We present a simulation about polynomial fitting data for recovered people from 11 March to 3 May 2020.




Figure 14. Polynomial fitting data for recovered people in Turkey from 11 March 2020 to 3 May 2020.

We present a predictive analysis for died people. According to the results obtained, we get a linear regression which can be calculated as

$$y = -822.246 + 70.72746x. \quad (2.15)$$

F – test was calculated as 1.75×10^{-28} . *R square* was calculated as 0.907007. We can say that the significance of fit for model is suitable for the considered data. Also, we can present regression model

$$y = -0.0266x^3 + 3512.1x^2 - 0.2 \times 10^{-8}x + 0.2 \times 10^{-12} \quad (2.16)$$

which is polynomial of third order. For this polynomial, *R square* was calculated as 0.9971. We present a simulation about polynomial fitting data for died people from 11 March to 3 May 2020.




Figure 15. Polynomial fitting data for died people in Turkey from 11 March 2020 to 3 May 2020.

We firstly present a predictive analysis for tested people. According to the results obtained, we get a linear regression which can be calculated as

$$y = -257388 + 22572.98x. \quad (2.17)$$

F - test was calculated as 5.17×10^{-28} . R square was calculated as 0.903051. We can say that the significance of fit for model is suitable for the considered data. Also, we give polynomial regression model

$$y = 520.26x^2 - 0.5 \times 10^{-7}x + 0.1 \times 10^{-12} \quad (2.18)$$

which is polynomial of second order. For this polynomial, R square was calculated as 0.9962. We present a simulation about polynomial fitting data for tested people from 11 March to 3 May 2020.




Figure 16. Polynomial fitting data for tested people in Turkey from 11 March 2020 to 3 May 2020.

We present some statistical data about corona in Turkey in Table 1.

Table 1. Some data about corona in Turkey.

	Infected	Recovered	Death
Arithmetic mean	2334.166667	1169.462963	56.63636364
Geometric mean	1029.876948	592.2822821	41.57620243
Harmonic mean	21.75203938	142.4039454	8.182426471
Standard deviation	1641.461139	1701.397957	48.05766445
Skewness	-0.07519891	1.416616957	0.100241163
Variance	2644498.472	2841148.434	2045.624143
Covariance	17324.10185	21927.12037	607.7037037
Pearson Correlation	0.683518717	0.834653045	0.862085432
Spearman Correlation	-17.9276729	0.188679245	-8.39622641

We now present lognormal distribution for all cases in Turkey from 11 March to 03 May 2020.




Figure 17. Lognormal distribution for all cases in Turkey from 11 March 2020 to 3 May 2020.

2.1.2 Prediction about corona data in Turkey

In this section, we aim at performing prediction using existing data and reliability level method. The collected data will be considered from 11 March 2020 to 3 May 2020. The future prediction will start from 3 May 2020 to 15 of June 2020. This will help us give a prediction on numbers of new daily infected class, recovered, daily numbers of deaths in Turkey within this period. The prediction will consist of three different graphs comprising upper boundaries, middle lines and low boundaries. The upper boundaries represent the worse cases scenario, of course a scenario that is not needed for deaths class and infected but an ideal one for recovered class, and the lower boundaries representing perfect scenarios (A scenario that is needed) for Turkey to get rid of the infection. These results of prediction for future daily new infected, recovered and deaths are represented graphically in Figure 18, 19 and 20 respectively.




Figure 18. Prediction of daily number of infected in Turkey using Forecast Sheet.

(1)




Figure 19. Prediction of daily number of recovered in Turkey using Forecast Sheet.




Figure 20. Prediction of daily number of deaths in Turkey using Forecast Sheet.

2.2 Detailed Statistical Analysis for South Africa

In this section, we aim to provide a detailed statistical analysis of the collected data representing the evolution COVID-19 spread within the republic of South Africa. These data include, daily number of new infected and daily numbers of deaths. The collected data are from 5 March 2020 to 3 May 2020 [7]. The main aim of this section is to predict what could possibly happen in the near future using the reliability level method, additionally, to find which distribution each class follows. With the collected data, we will first present histogram, pie chart

and nonlinear graphs for each class. The histograms will help identify the density of probability associated to each set of collected data. Additionally, we provide a polynomial fitting against collected. The results are presented in Figure 21 to 30. For each case, we present arithmetic, geometric, harmonic means respectively, skewness, variance, covariance, Pearson correlation and Spearman correlation and their results are presented in Table 2.

In Figure 21, 22 and 23, we present some statistical simulation about number of infected people due to COVID-19 in South Africa from 5 March 2020 to 3 May 2020.




Figure 21. Number of infected people in South Africa from 5 March 2020 to 3 May 2020.




Figure 22. Number of infected people in South Africa from 5 March 2020 to 3 May 2020.




Figure 23. Number of infected people in South Africa from 5 March 2020 to 3 May 2020.

In Figure 24, 25 and 26, we present some statistical simulation about number of died people due to COVID-19 in South Africa from 15 March 2020 to 3 May 2020.




Figure 24. Number of death in South Africa from 15 March 2020 to 3 May 2020.




Figure 25. Number of death in South Africa from 15 March 2020 to 3 May 2020.




Figure 26. Number of death in South Africa from 15 March 2020 to 3 May 2020.

Now we present regression analysis about COVID-19 in South Africa from 5 March 2020 to 3 May 2020. We firstly present a predictive analysis for infected people. According to the results obtained, we get a linear regression which can be calculated as

$$y = -4488415 + 102.2293x. \tag{2.19}$$

F – test was calculated as 4.84×10^{-31} . *R square* was calculated as 0.902781. We can say that the significance

of fit for model is suitable for the considered data. We can give another regression model

$$y = -0.4 \times 10^{-6}x^6 + 0.9253x^5 - 101611x^4 + 0.6 \times 10^{-9}x^3 - 0.2 \times 10^{-14}x^2 + 0.3 \times 10^{-18}x - 0.3 \times 10^{-22} \quad (2.20)$$

which is polynomial of sixth order. For this polynomial, *R square* was calculated as 0.9978. We present a simulation about polynomial fitting data for infected people from 5 March to 3 May 2020.




Figure 27. Polynomial fitting data for infected in South Africa from 5 March 2020 to 3 May 2020.

Now we present a predictive analysis for died people. According to the results obtained, we get a linear regression which can be calculated as

$$y = -29.2547 + 2.51587x. \quad (2.21)$$

F – test was calculated as 5.36×10^{-22} . *R square* was calculated as 0.858225. We can say that the significance of fit for model is for the considered data. We can suggest another regression model

$$y = -0.2 \times 10^{-5}x^4 + 3.7609x^3 - 247847x^2 + 0.7 \times 10^{-9}x - 0.8 \times 10^{-13} \quad (2.22)$$

which is polynomial of fourth order. For this polynomial, *R square* was calculated as 0.9958. We present a simulation about polynomial fitting data for died people from 15 March to 3 May 2020.




Figure 28. Polynomial fitting data for death in South Africa from 15 March 2020 to 3 May 2020.

2.2.1 Prediction about corona data in South Africa

In this section, we aim at performing prediction using existing data collected representing daily numbers of new infected, deaths and reliability level method. The collected data will be considered from 5 March 2020 corresponding to the first day of confirmed case of COVID-19 in South Africa to 3 May 2020. The future prediction will start from 3 May 2020 to 15 of June 2020. This will help us give a prediction on numbers of new daily infected class, recovered, daily numbers of deaths in South Africa within this period. The prediction will consist of three different graphs comprising upper boundaries, middle lines and low boundaries. The upper boundaries represent the worse cases scenario, of course a scenario that is not needed for deaths class and infected but an ideal one for recovered class, and the lower boundaries representing perfect scenarios (A scenario that is needed) for South Africa to get rid of the infection. These results of prediction for future daily new infected, recovered and deaths are represented graphically in Figure 29 and 30 respectively. The prediction of daily new infected in the case of South Africa seems to follow the upper boundaries and low boundary for daily number of deaths.




Figure 29. Prediction of daily number of infected in South Africa using Forecast Sheet.




Figure 30. Prediction of daily number of deaths in South Africa using Forecast Sheet.

We now give some data about corona in South Africa in Table 2.

Table 2. Some data about corona in South Africa.

	Infected	Death
Arithmetic mean	113,05000	2,62000
Geometric mean	57,35444	3,184267
Harmonic mean	11,569120	2,260659
Standard deviation	109,71834	3,613410
Skewness	1,1222456	1,594677
Variance	11837,4808	12,79560
Covariance	1492,5916	31,91000
Pearson Correlation	0,7921566	0,618164
Spearman Correlation	-5,389406	-6,168775

We present lognormal distribution for infected and death cases about COVID-19 in Figure 31.




Figure 31. Lognormal distribution for all cases in South Africa.

2.3 Parameters estimation using the Bell curving approach

In the previous section, we presented the graph of a day-to-day evolution of COVID-19 spread including infected, recovery and death for South Africa and Turkey. To be honest, one cannot for sure tell if those curves follow the normal distribution or lognormal distribution. Therefore in this section, two cases are considered. In the first case, we assume a lognormal curve and second we assume normal distribution curve.

Case I : We consider the lognormal density of probability

$$L_p(x) = \frac{1}{x} \frac{1}{\sigma\sqrt{2\pi}} \exp\left[-\frac{1}{2} \frac{(\ln x - \mu)^2}{\sigma^2}\right]. \quad (2.23)$$

We now define a function β that captures daily occurrences

$$\beta = O_0 \exp\left[-\frac{1}{2} \frac{(\ln x - \mu)^2}{\sigma^2}\right]. \quad (2.24)$$

We aim to estimate O_0 , σ and μ . To achieve this, we consider first four different days

$$d_1 < d_2 < d_3 < d_4 \text{ where } d_i = \ln x_i. \quad (2.25)$$

We first start by estimating μ , by assuming a proportion

$$\frac{\beta(d_2)}{\beta(d_1)} = \exp \left[-\frac{1}{2\sigma^2} \left\{ (d_2 - \mu)^2 - (d_1 - \mu)^2 \right\} \right] \quad (2.26)$$

and

$$\frac{\beta(d_4)}{\beta(d_3)} = \exp \left[-\frac{1}{2\sigma^2} \left\{ (d_4 - \mu)^2 - (d_3 - \mu)^2 \right\} \right]. \quad (2.27)$$

To proceed, we apply on both sides the ln function

$$\ln \left[\frac{\beta(d_2)}{\beta(d_1)} \right] = -\frac{1}{2\sigma^2} \left\{ (d_2 - \mu)^2 - (d_1 - \mu)^2 \right\} \Rightarrow \sigma^2 = -\frac{1}{2 \ln \left[\frac{\beta(d_2)}{\beta(d_1)} \right]} \left\{ (d_2 - \mu)^2 - (d_1 - \mu)^2 \right\} \quad (2.28)$$

$$\ln \left[\frac{\beta(d_4)}{\beta(d_3)} \right] = -\frac{1}{2\sigma^2} \left\{ (d_4 - \mu)^2 - (d_3 - \mu)^2 \right\} \Rightarrow \sigma^2 = -\frac{1}{2 \ln \left[\frac{\beta(d_4)}{\beta(d_3)} \right]} \left\{ (d_4 - \mu)^2 - (d_3 - \mu)^2 \right\}.$$

Due to the equality, we can now have

$$\frac{1}{\ln \left[\frac{\beta(d_2)}{\beta(d_1)} \right]} \left\{ (d_2 - \mu)^2 - (d_1 - \mu)^2 \right\} = \frac{1}{\ln \left[\frac{\beta(d_4)}{\beta(d_3)} \right]} \left\{ (d_4 - \mu)^2 - (d_3 - \mu)^2 \right\}. \quad (2.29)$$

Thus

$$\frac{\ln \left[\frac{\beta(d_4)}{\beta(d_3)} \right]}{\ln \left[\frac{\beta(d_2)}{\beta(d_1)} \right]} = \frac{(d_4 + d_3 - 2\mu) - (d_4 - d_3)}{(d_2 + d_1 - 2\mu) - (d_2 - d_1)}. \quad (2.30)$$

The solution is the above is

$$\mu = \frac{\frac{1}{2} (d_4 + d_3) (d_4 - d_3) - \frac{\ln \left[\frac{\beta(d_4)}{\beta(d_3)} \right]}{\ln \left[\frac{\beta(d_2)}{\beta(d_1)} \right]} (d_2 + d_1) (d_2 - d_1)}{\frac{1}{2} (d_4 - d_3) - \frac{\ln \left[\frac{\beta(d_4)}{\beta(d_3)} \right]}{\ln \left[\frac{\beta(d_2)}{\beta(d_1)} \right]} (d_2 - d_1)}. \quad (2.31)$$

Having μ , we can determine

$$\sigma = \sqrt{\frac{1}{2 \ln \left[\frac{\beta(d_2)}{\beta(d_1)} \right]} \left\{ -(d_2 - \mu)^2 + (d_1 - \mu)^2 \right\}}. \quad (2.32)$$

Alternatively, we consider 8 days to capture more facts x_i , $i = 1, 2, 3, 4, 5, 6, 7, 8$, we put $d_i = \ln x_i$. We assume a proportionality λ of $\{d_1, d_2, d_3, d_4\}$ and $\{d_5, d_6, d_7, d_8\}$. Therefore

$$P_1 = \frac{(d_4 + d_3 - 2\mu) - (d_4 - d_3)}{(d_2 + d_1 - 2\mu) - (d_2 - d_1)} \quad (2.33)$$

and

$$P_2 = \frac{(d_8 + d_7 - 2\mu) - (d_8 - d_7)}{(d_6 + d_5 - 2\mu) - (d_6 - d_5)}. \quad (2.34)$$

We now assume that P_1 is proportional to P_2 then

$$\frac{(d_4 + d_3 - 2\mu) - (d_4 - d_3)}{(d_2 + d_1 - 2\mu) - (d_2 - d_1)} = \lambda \frac{(d_8 + d_7 - 2\mu) - (d_8 - d_7)}{(d_6 + d_5 - 2\mu) - (d_6 - d_5)}. \quad (2.35)$$

For simplicity, we put $d_i + d_j = A_{ij} = A_{ji}$

$$d_{14} = \frac{d_4 - d_3}{d_2 - d_1}, d_{58} = \frac{d_8 - d_7}{d_6 - d_5}. \quad (2.36)$$

Therefore, the above can be reformulated as

$$\frac{A_{43} - 2\mu}{A_{21} - 2\mu} d_{14} = \lambda \frac{A_{87} - 2\mu}{A_{65} - 2\mu} d_{58}. \quad (2.37)$$

Also we write

$$(A_{43}A_{65} - 2\mu(A_{43} + A_{65}) + 4\mu^2) \frac{d_{14}}{\lambda d_{58}} = (A_{21}A_{87} - 2\mu(A_{21} + A_{87}) + 4\mu^2) \quad (2.38)$$

and

$$4\mu^2 \left\{ \frac{d_{14}}{\lambda d_{58}} - 1 \right\} - 2\mu \left\{ \frac{d_{14}}{\lambda d_{58}} (A_{43} + A_{65}) + (A_{21} + A_{87}) \right\} + A_{43}A_{65} \frac{d_{14}}{\lambda d_{58}} - A_{21}A_{87} = 0. \quad (2.39)$$

Thus we have

$$\mu_{1,2} = \frac{\left(\frac{d_{14}}{\lambda d_{58}} (A_{43} + A_{65}) + (A_{21} + A_{87}) \right) \pm \sqrt{\left(\frac{d_{14}}{\lambda d_{58}} (A_{43} + A_{65}) + (A_{21} + A_{87}) \right)^2 - 4 \left\{ \frac{d_{14}}{\lambda d_{58}} - 1 \right\} (A_{43}A_{65} \frac{d_{14}}{\lambda d_{58}} - A_{21}A_{87})}}{4 \left\{ \frac{d_{14}}{\lambda d_{58}} - 1 \right\}}. \quad (2.40)$$

Thus for Case I, we get

$$O_0 = \frac{\sum_{j=1}^4 \beta(\ln x_j)}{\sum_{j=1}^4 \frac{1}{x_j} \exp \left[-\frac{1}{2} \frac{(\ln x_j - \mu)^2}{\sigma^2} \right]}. \quad (2.41)$$

Case II, we get

$$O_0 = \frac{\sum_{j=1}^8 \beta(\ln x_j)}{\sum_{j=1}^8 \frac{1}{x_j} \exp \left[-\frac{1}{2} \frac{(\ln x_j - \mu)^2}{\sigma^2} \right]} \quad (2.42)$$

$$\lambda = \frac{\ln \left[\frac{\beta(d_4)}{\beta(d_3)} \right]}{\ln \left[\frac{\beta(d_2)}{\beta(d_1)} \right]} \times \frac{\ln \left[\frac{\beta(d_6)}{\beta(d_5)} \right]}{\ln \left[\frac{\beta(d_8)}{\beta(d_7)} \right]}.$$

For each case, the cumulative distribution function can be calculated by

$$\Phi(x) = \frac{1}{2} \left[1 + \operatorname{erf} \left(\frac{(\ln x_j - \mu)}{\sigma \sqrt{2}} \right) \right]. \quad (2.43)$$

Case II: We assume that the curve follows the normal distribution, thus

$$\Phi(x) = \frac{1}{\sigma \sqrt{2\pi}} \exp \left[-\frac{1}{2} \left(\frac{x - \mu}{\sigma} \right)^2 \right]. \quad (2.44)$$

However, we consider the following function

$$\lambda(x) = \lambda_0 \exp \left[-\frac{1}{2} \left(\frac{x - \mu}{\sigma} \right)^2 \right]. \quad (2.45)$$

We aim to determine λ_0, σ and μ . Here we choose three points d_1, d_2, d_3 such that $\lambda(d_2)$ corresponds to the maximum point. Following the procedure presented earlier, we have

$$\frac{\ln \left[\frac{\lambda(d_3)}{\lambda(d_2)} \right]}{\ln \left[\frac{\lambda(d_2)}{\lambda(d_1)} \right]} = \frac{(d_3 + d_2 - 2\mu) - (d_3 - d_2)}{(d_2 + d_1 - 2\mu) - (d_2 - d_1)}. \quad (2.46)$$

Thus

$$\mu = \frac{\frac{1}{2} (d_3 + d_2) (d_3 - d_2) - \frac{\ln \left[\frac{\lambda(d_3)}{\lambda(d_2)} \right]}{\ln \left[\frac{\lambda(d_2)}{\lambda(d_1)} \right]} (d_2 + d_1) (d_2 - d_1)}{\frac{1}{2} (d_3 + d_2) - \frac{\ln \left[\frac{\lambda(d_3)}{\lambda(d_2)} \right]}{\ln \left[\frac{\lambda(d_2)}{\lambda(d_1)} \right]} (d_2 - d_1)}. \quad (2.47)$$

With μ in hand, we determine

$$\sigma = \sqrt{\frac{1}{2 \ln \left[\frac{\lambda(d_2)}{\lambda(d_1)} \right]} \left\{ (d_1 - \mu)^2 - (d_2 - \mu)^2 \right\}} \quad (2.48)$$

and

$$\lambda_0 = \frac{\sum_{j=1}^3 \lambda(d_j)}{\sum_{j=1}^3 \exp \left[-\frac{1}{2} \frac{(d_j - \mu)^2}{\sigma^2} \right]}. \quad (2.49)$$

Particular case if we consider the case where $d_2 - d_1 = d_3 - d_2$ that is $\lambda(d_1) = \lambda(d_3)$ due to symmetry of normal distribution, then we get

$$\mu = \frac{\frac{1}{2} (d_3 + d_2) (d_3 - d_2) + (d_2 + d_1) (d_2 - d_1)}{\frac{1}{2} (d_3 + d_2) + (d_2 - d_1)}. \quad (2.50)$$

2.4 Comparison: Turkey vs South Africa

In this subsection, we present a comparison between Turkey and South Africa about COVID-19.

Table 3: Comparison between Turkey and South Africa updated on COVID-19

Country	South Africa:5 March to 02 May 2020	Turkey:11 March to 02 May 2020
Number of infected	6336	124375
Number of death	123	3336
Number of recovered	2549	58259
Total tested	230686	1,033,617
Lockdown date	26 March 2020 (Total lockdown)	11April 2020, (Partial lockdown: Less than 20 and older than 65)
Donations	NA	† 1,865,799,782 up to 28 April 2020

The analysis presented in this section does not aim at praising nor criticizing any country; but just to assess the effect of lockdown and its regulations, and to perceive if this concept can help save humans before the vaccine. The fundamental question to answer here is to know why South Africa has less number of deaths and infected people than Turkey; if it recorded its first confirmed case six days earlier before Turkey recorded its first. The answer may rely on two fundamental facts which include the period lockdown was implemented and the type of lockdown put in place. The South African government publicized on 23 March 2020 a 21-day of national lockdown which started effectively from midnight 27 March. This was announced 22 days after the first confirmed case was recorded in the country. The lockdown came with strict measures encompassing, immediate deployment of South African National Defence force to ensure that all people living within the territory of South Africa obey the lockdown rules. Only workers considered necessary to operative response to the pandemic were exempted, namely: health caregivers, security service providers, essential service providers that are fundamental to the rudimentary functioning of economy as well as other workers in industries that cannot be economically shut down. This implies that the mentioned categories were permitted to go to their places of work during the lockdown. On the other hand, the numbers of people at gatherings apart from funerals were limited to 50 people; while restaurants, taverns, bottle stores and shops that are not selling indispensable goods were forced to close. Thus, a large population was not allowed to leave their houses except for essential needs. Consequently, the movement between provinces, metropolitan and districts were also restricted unless for essential reasons that cannot be catered for within provincial boundaries. The South African government further closed all of its

national borders and only allowed the transportation for indispensable reasons. Likewise, all international and domestic passenger flights were prohibited, except those assigned to evacuate citizens from foreign countries and certain repatriations due to COVID-19. However, the measures taken by Turkey were not implemented swiftly upon the confirmation of its first COVID-19 case. It is recorded that the Turkish government announced a partial lockdown on 11 April 2020, a month after the country registered its first confirmed case of COVID-19. Prior to the announcement of the partial lockdown, mosques, cafes, night clubs, and all universities within the country were already closed on 11 March 2020. The restriction measures applied only on people younger than 20 years old and older than 65 years old, who were not allowed to leave their homes except for indispensable reasons. In addition, the government ordered a ban on movements between 30 major cities with metropolitan status as well as Zonguldak; whereby the lockdown is applied every weekend since 11 April 2020 and also 21-23 April 2020. Punishment (money) is applied for people who go out. Very importantly the government has totally banned the sale of masks, but provided free masks to its people to be compulsorily utilized in public places. However, the newly placed order exempted health care assistance, funerals, military and passenger transports from the ban; provided that certain conditions are met. Although both countries have put severe measures to protect their citizens from the deadly disease; there are still records of rising number of infected and dead people in both Turkey and South Africa. Does that mean that the lockdown regulations are worthless? Absolutely not! It is only that several citizens in respective countries are not adhering to the rules and regulations put in place by their respective government authorities. This results from the concept of social distancing being largely misunderstood, as it is not clearly defined to mean whether persons should stay one meter away from one another or only from any infected humans, contaminated air and other objects because of the nature in which COVID-19 can be spread. Nevertheless, due to a long incubation period of COVID-19, approximately 14 days maximum; which renders ordinary citizens not to differentiate an infected person from others; then it is crucial that stringent measures be implemented, which will prohibit people from leaving their homes; and in case they had gone out, they should maintain the one metre distance away from each other and frequently wash their hands upon touching any object.

3 Mathematical model of COVID-19 in South Africa and Turkey

Mathematical models of infected diseases are deemed not that useful by some people who feel that they cannot be utilized to develop a vaccine or cure of any given disease. However, it is important to note that the principal aim of these mathematical models is to describe a system using mathematical tools, concepts and language. Hence, throughout the history of human beings, researchers working within the field of mathematics have developed more accurate and efficient mathematical models. For instance, history has made reference to one of the well-known Newtonian laws which described very accurately many problems in our daily lives; although they are coupled with some limits. In instances where these laws failed, two other well-known concepts namely; theory of relativity and quantum mechanics using mathematical formulas can be utilized instead. Generally, these concepts are of great importance in all fields of science such as in natural sciences including chemistry, biology, physics, and earth science, in engineering such as computer science, and electrical engineering, as well as in social science where their applicability to economics, sociology, psychology and political science can be relevant. In other words, mathematical models can help to provide a clear explanation of a system and investigate the effect of several components, and later make accurate predictions based on the observed facts. In the current situation under study, due to the magnitude of fear imposed by COVID-19 on humans, it is therefore paramount for mathematicians to provide conceptual models, using mathematical tools called differential and integral operators, to suggest well-constructed mathematical models that will be used to understand and predict the spread of COVID-19.

In this section, a mathematical model that takes into account nine classes (susceptible, infected which has 5 sub-classes, recovered, death and vaccinated classes), the dynamic is presented and explained with the subsequent diagrams; but the death class is omitted because it can produce a complex model. The created model incorporates the lockdown effect, represented by a coefficient that takes into account the social distancing and a contact coefficient.

$$\begin{aligned}
 \dot{S} &= \Lambda - (\alpha(x) + \gamma_1 + \mu_1) S \\
 \dot{I} &= \alpha(x) S - (\varepsilon + \xi + \lambda + \mu_1) I \\
 \dot{I}_A &= \xi I - (\theta + \mu + \chi + \mu_1) I_A \\
 \dot{I}_D &= \varepsilon I - (\eta + \varphi + \mu_1) I_D \\
 \dot{I}_R &= \eta I_D + \theta I_A - (v + \xi + \mu_1) I_R \\
 \dot{I}_T &= \mu I_A + v I_R - (\sigma + \tau + \mu_1) I_T \\
 \dot{R} &= \lambda I + \varphi I_D + \chi I_A + \xi I_R + \sigma I_T - (\Phi + \mu_1) R \\
 \dot{D} &= \tau I_T - \mu_1 D \\
 \dot{V} &= \gamma_1 S + \Phi R - \mu_1 V - \tau I_T
 \end{aligned} \tag{3.1}$$

where

$$\begin{aligned}
 \dot{N} &= \Lambda - \mu_1 N - \tau I_T \\
 \alpha(x) &= \frac{k_1 p e^{-x}}{N} (I + w (\beta I_D + \gamma I_A + \delta_1 I_R)).
 \end{aligned} \tag{3.2}$$

$S(t)$ is the class of individuals that are susceptible to contact COVID-19 at time t . $I(t)$ is the class of individuals that are susceptible to contacted COVID-19, but have no symptoms and have not been tested. $I_A(t)$ is the class of individuals that have some symptoms but not tested yet. $I_D(t)$ is the class of individuals that have contacted COVID-19, have been tested positive, but no symptoms. $I_R(t)$ is the class of individuals that have contacted COVID-19, have been tested positive and have symptoms. $I_T(t)$ is the class of individuals that have contacted COVID-19 and one is critical condition. $R(t)$ is the class of recovered individuals at time t . $D(t)$ is the number of death at time t . $V(t)$ is the class of individuals that have been vaccinated.

Table 4. Parameters of the suggested COVID-19 model

Φ	: The rate at which recovered and vaccinated
μ_1	: Turkish natural mortal rate
$\alpha(x)$: The infection force
β	: Transmission rate of $I_D(t)$ class
γ	: Transmission rate of $I_A(t)$ class
δ_1	: Transmission rate of $I_R(t)$ class
ε	: Proportion rate of detection relative to asymptomatic class
θ	: Proportion rate of detection relative to symptomatic class
ξ	: Rate that infected class are not aware of their status
η	: Rate that infected class are aware of their status
μ	: Rate that non-tested join class $I_T(t)$
v	: Rate that tested join class $I_T(t)$
τ	: The mortal rate due to COVID-19
λ	: Recovery rate of class $I(t)$

- κ : Recovery rate of class $I_A(t)$
- ξ : Recovery rate of class $I_R(t)$
- φ : Recovery rate of class $I_D(t)$
- σ : Recovery rate of class $I_T(t)$
- k_1 : Contact rate
- p : Proportion that a contact is sufficient enough to lead to transmission
- w : Transmission coefficient for the infected classes
- Λ : Recruitment rate into class $S(t)$
- γ_1 : Rate of vaccination

The initial conditions are given as

$$\begin{aligned} N(0) &= N_0, S(0) = S_0, I(0) = I_0, I_A(0) = I_A^0, I_D(0) = I_D^0, \\ I_R(0) &= I_R^0, I_T(0) = I_T^0, R(0) = R_0, D(0) = D_0, V(0) = V_0. \end{aligned} \quad (3.3)$$

We present a diagram which summarizes COVID-19 model which is described by the system (3.1) in Figure 21.




Figure 32. Diagram summarizing COVID-19 model given by the system (3.1).

3.1 Boundness and positivity of the solutions

In this section, we show that $\forall t \geq 0$, the system solution is positive, that the model is well-posed and biologically feasible. We define the norm

$$\|f\| = \sup_{t \in D_f} |f(t)|. \quad (3.4)$$

We assume that all the class

$$S(\alpha + \gamma_1) > 0, \quad \forall t \geq 0 \quad (3.5)$$

due to the model under this assumption. We write

$$\begin{aligned}\dot{I}(t) &= \alpha S - (\varepsilon + \xi + \lambda + \mu_1) I, \quad \forall t \geq 0 \\ &\geq -(\varepsilon + \xi + \lambda + \mu_1) I(t), \quad \forall t \geq 0 \\ &\geq I^0 e^{-(\varepsilon + \xi + \lambda + \mu_1)t}, \quad \forall t \geq 0\end{aligned}\tag{3.6}$$

Since $I(t) \geq 0, \forall t \geq 0$, then

$$\begin{aligned}\dot{I}_A(t) &= \xi I(t) - (\theta + \mu + \chi + \mu_1) I_A(t), \quad \forall t \geq 0 \\ &\geq -(\theta + \mu + \chi + \mu_1) I_A(t), \quad \forall t \geq 0\end{aligned}\tag{3.7}$$

then

$$I_A(t) \geq I_A^0 e^{-(\theta + \mu + \chi + \mu_1)t}, \quad \forall t \geq 0.\tag{3.8}$$

The same with $I_D(t)$ class

$$I_D(t) \geq I_D^0 e^{-(\eta + \varphi + \mu_1)t}, \quad \forall t \geq 0.\tag{3.9}$$

$I_A(t)$ and $I_D(t)$ are positive $\forall t \geq 0$ and $\eta, \theta \geq 0$ then

$$\begin{aligned}I_R(t) &\geq I_R^0 e^{-(v + \xi + \mu_1)t}, \quad \forall t \geq 0, \\ I_T(t) &\geq I_T^0 e^{-(\sigma + \tau + \mu_1)t}, \quad \forall t \geq 0, \\ R(t) &\geq R_0 e^{-(\Phi + \mu_1)t}, \quad \forall t \geq 0, \\ D(t) &\geq D_0 e^{-\mu t}, \quad \forall t \geq 0.\end{aligned}\tag{3.10}$$

Also

$$\begin{aligned}\dot{V}(t) &\geq -\mu_1 V(t) - \tau I_T(t), \quad \forall t \geq 0 \\ V(t) &\geq V_0 e^{-\mu_1 t} - \tau \int_0^t e^{-\mu_1(t-\tau)} I_T(\tau) d\tau\end{aligned}\tag{3.11}$$

$V(t) \geq 0$ if $V_0 e^{-\mu_1 t} > \tau \int_0^t e^{-\mu_1(t-\tau)} I_T(\tau) d\tau, \forall t \geq 0$. With $S(t)$, we have to assume that

$$\|\alpha\|_\infty < \infty \Rightarrow \|I\|_\infty + w \|I_D\|_\infty + \gamma \|I_A\|_\infty + \delta_1 \|I_R\|_\infty < \infty\tag{3.12}$$

such that

$$\begin{aligned}\dot{S}(t) &= \Lambda - (\alpha(x) + \gamma_1 + \mu_1) S, \quad \forall t \geq 0 \\ &\geq -(\alpha(x) + \gamma_1 + \mu_1) S, \quad \forall t \geq 0 \\ &\geq -(|\alpha(x)| + \gamma_1 + \mu_1) S, \quad \forall t \geq 0, \\ &\geq -\left(\sup_{x \in D_\alpha} |\alpha(x)| + \gamma_1 + \mu_1\right) S, \quad \forall t \geq 0 \\ &\geq -(\|\alpha\|_\infty + \gamma_1 + \mu_1) S, \quad \forall t \geq 0.\end{aligned}\tag{3.13}$$

This implies that

$$S(t) \geq S_0 e^{-(\|\alpha\|_\infty + \gamma_1 + \mu_1)t}, \quad \forall t \geq 0.\tag{3.14}$$

Therefore $\forall t \geq 0$, if the initial conditions are positive, all the solutions are positive with extra condition on $V(t)$ class that

$$V_0 e^{-\mu_1 t} > \tau \int_0^t e^{-\mu_1(t-\tau)} I_T(\tau) d\tau, \forall t \geq 0.\tag{3.15}$$

Now in the absence of the COVID-19, we have

$$N(t) \leq \frac{\Lambda}{\mu_1}. \quad (3.16)$$

The above inequality is called threshold population level. This obtains because we assume that the total number of population must be increased or be constant

$$\frac{dN(t)}{dt} \geq 0 \Rightarrow \Lambda - \mu_1 N \geq 0 \quad (3.17)$$

therefore $N(t) \leq \frac{\Lambda}{\mu_1}$. It is therefore biologically feasible that

$$\Omega = \left\{ (S, I, I_A, I_D, I_R, I_T, R, D, V) \in \mathbb{R} : 0 \leq S + I + I_A + I_D + I_R + I_T + R + D + V = N \leq \frac{\Lambda}{\mu_1} \right\}. \quad (3.18)$$

The disease free equilibrium point is

$$\left(\frac{\Lambda}{\gamma_1 - \mu_1}, 0, 0, 0, 0, 0, 0, 0, \frac{\Lambda \gamma_1}{\mu_1 (\gamma_1 - \mu_1)} \right). \quad (3.19)$$

We now derive the reproduction number using the next generation operator technique [9]. We have 3 infected classes $I(t)$, $I_A(t)$ and $I_D(t)$. The matrix F and V will be obtained from

$$\begin{aligned} \dot{I} &= \alpha(x)S - (\varepsilon + \xi + \lambda + \mu_1)I \\ \dot{I}_A &= \xi I - (\theta + \mu + \chi + \mu_1)I_A \\ \dot{I}_D &= \varepsilon I - (\eta + \varphi + \mu_1)I_D. \end{aligned} \quad (3.20)$$

We obtain the following matrices

$$F = \begin{bmatrix} \delta(x) & \delta(x)w\gamma & \delta(x)w\beta \\ \xi & 0 & 0 \\ \varepsilon & 0 & 0 \end{bmatrix} \quad (3.21)$$

and

$$V = \begin{bmatrix} (\varepsilon + \xi + \lambda + \mu_1) & 0 & 0 \\ 0 & (\theta + \mu + \chi + \mu_1) & 0 \\ 0 & 0 & (\eta + \varphi + \mu_1) \end{bmatrix}. \quad (3.22)$$

For simplicity

$$V = \begin{bmatrix} l_1 & 0 & 0 \\ 0 & l_2 & 0 \\ 0 & 0 & l_3 \end{bmatrix} \quad (3.23)$$

where

$$\begin{aligned} l_1 &= \varepsilon + \xi + \lambda + \mu_1 \\ l_2 &= \theta + \mu + \chi + \mu_1 \\ l_3 &= \eta + \varphi + \mu_1. \end{aligned} \quad (3.24)$$

Then we have

$$V^{-1} = \begin{bmatrix} \frac{1}{l_1} & 0 & 0 \\ 0 & \frac{1}{l_2} & 0 \\ 0 & 0 & \frac{1}{l_3} \end{bmatrix}. \quad (3.25)$$

So we write the following

$$FV^{-1} = \begin{bmatrix} \delta(x) & \delta(x)w\gamma & \delta(x)w\beta \\ \xi & 0 & 0 \\ \varepsilon & 0 & 0 \end{bmatrix} \begin{bmatrix} \frac{1}{l_1} & 0 & 0 \\ 0 & \frac{1}{l_2} & 0 \\ 0 & 0 & \frac{1}{l_3} \end{bmatrix} \quad (3.26)$$

and we have

$$FV^{-1} = \begin{bmatrix} \frac{\delta(x)}{l_1} & \frac{\delta(x)w\gamma}{l_2} & \frac{\delta(x)w\beta}{l_3} \\ \frac{\xi}{l_1} & 0 & 0 \\ \frac{\varepsilon}{l_1} & 0 & 0 \end{bmatrix}. \quad (3.27)$$

We now present disease equilibrium points. We achieve this by solving

$$\begin{aligned} \Lambda - (\alpha(x) + \gamma_1 + \mu_1)S &= 0 \\ \alpha(x)S - (\varepsilon + \xi + \lambda + \mu_1)I &= 0 \\ \xi I - (\theta + \mu + \chi + \mu_1)I_A &= 0 \\ \varepsilon I - (\eta + \varphi + \mu_1)I_D &= 0 \\ \eta I_D + \theta I_A - (v + \xi + \mu_1)I_R &= 0 \\ \mu I_A + v I_R - (\sigma + \tau + \mu_1)I_T &= 0 \\ \lambda I + \varphi I_D + \chi I_A + \xi I_R + \sigma I_T - (\Phi + \mu_1)R &= 0 \\ \tau I_T - \mu_1 D &= 0 \\ \gamma_1 S + \Phi R - \mu_1 V - \tau I_T &= 0 \end{aligned} \quad (3.28)$$

This implies that

$$\begin{aligned} D &= \frac{\tau}{\mu_1} I_T \\ I_A &= \frac{\xi}{\theta + \mu + \chi + \mu_1} I \\ I_D &= \frac{\varepsilon}{\eta + \varphi + \mu_1} I \\ I_R &= \left(\frac{\eta \varepsilon}{\eta + \varphi + \mu_1} + \frac{\theta \xi}{\theta + \mu + \chi + \mu_1} \right) \frac{I}{v + \xi + \mu_1} \\ I_T &= \left(\frac{\mu \xi}{\theta + \mu + \chi + \mu_1} + \frac{\chi \eta \varepsilon}{\eta + \varphi + \mu_1} + \frac{v \theta \xi}{\theta + \mu + \chi + \mu_1} \right) \frac{I}{\sigma + \tau + \mu_1}. \end{aligned} \quad (3.29)$$

Thus

$$\begin{aligned} S^* &= \frac{\delta(x)}{N} \left(1 + \frac{\beta \varepsilon w}{\eta + \varphi + \mu_1} + \frac{\gamma \xi w}{\theta + \mu + \chi + \mu_1} + \frac{\delta_1 \eta \varepsilon w}{(\eta + \varphi + \mu_1)(v + \xi + \mu_1)} \right. \\ &\quad \left. + \frac{\delta_1 \theta \xi w}{(\theta + \mu + \chi + \mu_1)(v + \xi + \mu_1)} \right), \\ \Lambda - (\delta(x)(I^* + w\beta I_D^* + \gamma w I_A^* + w\delta_1 I_R^*) + (\gamma_1 + \mu_1)) S^* &= 0 \end{aligned} \quad (3.30)$$

and

$$\frac{\Lambda + (\gamma_1 + \mu_1) S^*}{A} = I^* \quad (3.31)$$

where

$$A = \frac{\delta(x)}{N} \left(1 + \frac{\beta \varepsilon w}{\eta + \varphi + \mu_1} + \frac{\gamma \xi w}{\theta + \mu + \chi + \mu_1} + \frac{\delta_1 \eta \varepsilon w}{(\eta + \varphi + \mu_1)(v + \xi + \mu_1)} \right. \\ \left. + \frac{\delta_1 \theta \xi w}{(\theta + \mu + \chi + \mu_1)(v + \xi + \mu_1)} \right). \quad (3.32)$$

That is

$$\begin{aligned}
 I^* &= \frac{\Lambda A + (\gamma_1 + \mu_1)(\xi + \varepsilon + \lambda + \mu_1)}{A^2}, \\
 I_A^* &= \frac{\xi}{\theta + \mu + \chi + \mu_1} \left(\frac{\Lambda A + (\gamma_1 + \mu_1)(\xi + \varepsilon + \lambda + \mu_1)}{A^2} \right), \\
 I_D^* &= \frac{\varepsilon}{\eta + \varphi + \mu_1} \left(\frac{\Lambda A + (\gamma_1 + \mu_1)(\xi + \varepsilon + \lambda + \mu_1)}{A^2} \right), \\
 I_R^* &= \left(\frac{\eta\varepsilon}{\eta + \varphi + \mu_1} + \frac{\theta\xi}{\theta + \mu + \chi + \mu_1} \right) \left(\frac{\Lambda A + (\gamma_1 + \mu_1)(\xi + \varepsilon + \lambda + \mu_1)}{A^2} \right), \\
 I_T^* &= \frac{\Lambda A + (\gamma_1 + \mu_1)(\xi + \varepsilon + \lambda + \mu_1)}{A^2(\sigma + \tau + \mu_1)} \left(\frac{\mu\xi}{\theta + \mu + \chi + \mu_1} + \frac{\chi\eta\varepsilon}{\eta + \varphi + \mu_1} + \frac{v\theta\xi}{\theta + \mu + \chi + \mu_1} \right) \\
 D^* &= \frac{\tau}{\mu_1} \left(\frac{\mu\xi}{\theta + \mu + \chi + \mu_1} + \frac{\chi\eta\varepsilon}{\eta + \varphi + \mu_1} + \frac{v\theta\xi}{\theta + \mu + \chi + \mu_1} \right) \frac{\Lambda A + (\gamma_1 + \mu_1)(\xi + \varepsilon + \lambda + \mu_1)}{A^2(\sigma + \tau + \mu_1)}.
 \end{aligned} \tag{3.33}$$

Also we get

$$R^* = \frac{\Lambda A + (\gamma_1 + \mu_1)(\xi + \varepsilon + \lambda + \mu_1)}{A^2(\Phi + \mu_1)} \left\{ \begin{aligned} &\lambda + \frac{\eta\varepsilon}{\eta + \varphi + \mu_1} + \frac{\chi\xi}{\theta + \mu + \chi + \mu_1} \\ &+ \frac{\xi}{v + \xi + \mu_1} \left(\frac{\eta\varepsilon}{\eta + \varphi + \mu_1} + \frac{\theta\xi}{\theta + \mu + \chi + \mu_1} \right) + \frac{\sigma\mu\xi}{(\theta + \mu + \chi + \mu_1)(\sigma + \tau + \mu_1)} \\ &+ \frac{v\eta\varepsilon\sigma}{(\eta + \varphi + \mu_1)(\sigma + \tau + \mu_1)} + \frac{\sigma v\theta\xi}{(\theta + \mu + \chi + \mu_1)(\sigma + \tau + \mu_1)} \end{aligned} \right\} \tag{3.34}$$

and

$$\begin{aligned}
 V^* &= \frac{\gamma_1(\xi + \varepsilon + \lambda + \mu_1)}{\mu_1 A} + \frac{\Phi\Lambda A + (\gamma_1 + \mu_1)\Phi(\xi + \varepsilon + \lambda + \mu_1)}{A^2(\Phi + \mu_1)} \\
 &\times \left\{ \begin{aligned} &\lambda + \frac{\eta\varepsilon}{\eta + \varphi + \mu_1} + \frac{\chi\xi}{\theta + \mu + \chi + \mu_1} \\ &+ \frac{\xi}{v + \xi + \mu_1} \left(\frac{\eta\varepsilon}{\eta + \varphi + \mu_1} + \frac{\theta\xi}{\theta + \mu + \chi + \mu_1} \right) + \frac{\sigma\mu\xi}{(\theta + \mu + \chi + \mu_1)(\sigma + \tau + \mu_1)} \\ &+ \frac{v\eta\varepsilon\sigma}{(\eta + \varphi + \mu_1)(\sigma + \tau + \mu_1)} + \frac{\sigma v\theta\xi}{(\theta + \mu + \chi + \mu_1)(\sigma + \tau + \mu_1)} \end{aligned} \right\}.
 \end{aligned} \tag{3.35}$$

For the COVID-19 endemic with this model, we need to have

$$\dot{I}(t) > 0, \dot{I}_A(t) > 0 \text{ and } \dot{I}_D(t) > 0. \tag{3.36}$$

This implies

$$\begin{aligned}
 \alpha(x)S - (\varepsilon + \xi + \lambda + \mu_1)I &> 0, \quad \forall t \geq 0 \\
 \xi I - (\theta + \mu + \chi + \mu_1)I_A &> 0, \quad \forall t \geq 0 \\
 \varepsilon I - (\eta + \varphi + \mu_1)I_D &> 0, \quad \forall t \geq 0.
 \end{aligned} \tag{3.37}$$

Thus

$$\begin{aligned}
 I_A &< \frac{\xi}{\theta + \mu + \chi + \mu_1} I \\
 I_D &< \frac{\varepsilon}{\eta + \varphi + \mu_1} I \\
 \frac{\alpha(x)S}{\varepsilon + \xi + \lambda + \mu_1} &> I.
 \end{aligned} \tag{3.38}$$

We use the fact that $\frac{S}{N} < 1$

$$\frac{\delta(x)}{\varepsilon + \xi + \lambda + \mu_1} (I + w\beta I_D + \gamma w I_A + w\delta_1 I_R) > I \tag{3.39}$$

noting that

$$\begin{aligned}
 I_A &< \frac{\xi}{\theta + \mu + \chi + \mu_1} I \\
 I_D &< \frac{\varepsilon}{\eta + \varphi + \mu_1} I \\
 I_R &< \frac{\eta}{v + \xi + \mu_1} I_D + \frac{\theta}{v + \xi + \mu_1} I_A \\
 &< \frac{\eta\varepsilon}{(v + \xi + \mu_1)(\eta + \varphi + \mu_1)} I + \frac{\xi\theta}{(\theta + \mu + \chi + \mu_1)(v + \xi + \mu_1)} I.
 \end{aligned} \tag{3.40}$$

Also

$$\begin{aligned}
 w\beta I_D &< \frac{w\beta\varepsilon I}{\eta + \varphi + \mu_1} \\
 w\gamma_1 I_A &< \frac{w\gamma_1 \xi I}{\theta + \mu + \chi + \mu_1}.
 \end{aligned} \tag{3.41}$$

Therefore we have the following inequality in terms of I

$$\frac{\delta(x)}{\varepsilon + \xi + \lambda + \mu_1} \left(\frac{I + \frac{w\beta\varepsilon}{\eta + \varphi + \mu_1} I + \frac{w\gamma_1 \xi}{\theta + \mu + \chi + \mu_1} I}{+ \frac{w\eta\varepsilon\delta_1}{(v + \xi + \mu_1)(\eta + \varphi + \mu_1)} I + \frac{w\delta_1 \xi \theta}{(\theta + \mu + \chi + \mu_1)(v + \xi + \mu_1)} I} \right) > I \tag{3.42}$$

and

$$\frac{\delta(x)}{\varepsilon + \xi + \lambda + \mu_1} \left(\frac{1 + \frac{w\beta\varepsilon}{\eta + \varphi + \mu_1} + \frac{w\gamma_1 \xi}{\theta + \mu + \chi + \mu_1}}{+ \frac{w\eta\varepsilon\delta_1}{(v + \xi + \mu_1)(\eta + \varphi + \mu_1)} + \frac{w\delta_1 \xi \theta}{(\theta + \mu + \chi + \mu_1)(v + \xi + \mu_1)}} \right) > 1. \tag{3.43}$$

Therefore

$$R_0 > 1 \tag{3.44}$$

where

$$R_0 = \frac{\delta(x)}{\varepsilon + \xi + \lambda + \mu_1} \left(\frac{1 + \frac{w\beta\varepsilon}{\eta + \varphi + \mu_1} + \frac{w\gamma_1 \xi}{\theta + \mu + \chi + \mu_1}}{+ \frac{w\eta\varepsilon\delta_1}{(v + \xi + \mu_1)(\eta + \varphi + \mu_1)} + \frac{w\delta_1 \xi \theta}{(\theta + \mu + \chi + \mu_1)(v + \xi + \mu_1)}} \right). \tag{3.45}$$

This shows that we have a unique endemic equilibrium when $R_0 > 1$.

3.2 Local and global stability of the Disease-free equilibrium

Lemma 1. The disease-free equilibrium E_0 of the COVID-19 system is locally asymptotically stable when $R_0 < 1$ and unstable when $R_0 > 1$. The Jacobian matrix for COVID-19 system is given by

$$\begin{bmatrix}
 -(\gamma_1 + \mu_1) & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
 0 & -(\varepsilon + \xi + \lambda + \mu_1) & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
 0 & \xi & -(\theta + \mu + \chi + \mu_1) & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
 0 & \varepsilon & 0 & -(\eta + \varphi + \mu_1) & 0 & 0 & 0 & 0 & 0 & 0 \\
 0 & 0 & \theta & \eta & -(v + \xi + \mu_1) & 0 & 0 & 0 & 0 & 0 \\
 0 & 0 & \mu & \eta & v & -(\sigma + \tau + \mu_1) & 0 & 0 & 0 & 0 \\
 0 & \lambda & \chi & \varphi & \xi & 0 & -(\Phi + \mu_1) & 0 & 0 & 0 \\
 0 & 0 & 0 & 0 & 0 & \sigma & \tau & -\mu_1 & 0 & 0 \\
 \gamma_1 & 0 & 0 & 0 & 0 & -\tau & \Phi & 0 & -\mu_1 & -\mu_1
 \end{bmatrix}$$

It is known that the disease-free equilibrium E_0 asymptotically stable if and only if the $tr(J(E_0)) < 0$ and the $\det(J(E_0)) > 0$. For the suggested COVID-19 the trace of $J(E_0)$ is

$$tr(J(E_0)) = -(\gamma_1 + 9\mu_1 + 2\xi + \varepsilon + \lambda + \theta + \Phi + \tau + \mu + \varphi + \eta + v + \chi) < 0. \tag{3.46}$$

The determinant of $J(E_0)$ is

$$\begin{aligned}
 \det(J(E_0)) &= (\gamma_1 + \mu_1)(\varepsilon + \xi + \lambda + \mu_1)(\theta + \mu + \chi + \mu_1)(\eta + \varphi + \mu_1) \\
 &\quad \times (v + \xi + \mu_1)(\sigma + \tau + \mu_1)(\Phi + \mu_1)\mu_1^2 > 0.
 \end{aligned} \tag{3.47}$$

In this case, we can conclude that the disease-free equilibrium of the suggested model for COVID-19 under vaccination and treatment is locally asymptotically stable.

Theorem 2. The COVID-19 model disease-free equilibrium is globally asymptotically stable within the feasible interval if $R_0 < 1$ and unstable if $R_0 > 1$.

Proof. We use the Lyapunov function defined by

$$L = \frac{1}{l_1}I + \frac{1}{l_2}I_A + \frac{1}{l_3}I_D. \quad (3.48)$$

Therefore its derivative along the solutions of the COVID-19 model

$$\frac{dL}{dt} = \frac{1}{l_1} \frac{dI}{dt} + \frac{1}{l_2} \frac{dI_A}{dt} + \frac{1}{l_3} \frac{dI_D}{dt} \quad (3.49)$$

where

$$\begin{aligned} l_1 &= \varepsilon + \xi + \lambda + \mu_1 \\ l_2 &= \theta + \mu + \chi + \mu_1 \\ l_3 &= \eta + \varphi + \mu_1. \end{aligned} \quad (3.50)$$

Then we write

$$\begin{aligned} \frac{dL}{dt} &= \frac{1}{l_1} (\alpha(x)S - l_1I) + \frac{1}{l_2} (\xi I - l_2I_A) + \frac{1}{l_3} (\varepsilon I - l_3I_D) \\ &= \frac{1}{l_1 l_2 l_3} (\alpha(x) S l_2 l_3 - l_1 l_2 l_3 I + l_1 l_3 \xi I - l_1 l_2 l_3 I_A + l_1 l_2 \varepsilon I - l_1 l_2 l_3 I_D). \end{aligned} \quad (3.51)$$

We have on the other hand that

$$\alpha(x) = \frac{\delta(x)}{N} (I + w(\beta I_D + \gamma I_A + \delta_1 I_R)) \quad (3.52)$$

and we get

$$\frac{dL}{dt} = \frac{1}{l_1 l_2 l_3} \left(\frac{\delta(x)}{N} (I + w(\beta I_D + \gamma I_A + \delta_1 I_R)) S l_2 l_3 - l_1 l_2 l_3 I + l_1 l_3 \xi I - l_1 l_2 l_3 I_A + l_1 l_2 \varepsilon I - l_1 l_2 l_3 I_D \right). \quad (3.53)$$

Thus

$$\begin{aligned} \frac{dL}{dt} &\leq \frac{S}{N} (I + I_D + I_A + I_R) \left[\frac{\delta(x)}{\varepsilon + \xi + \lambda + \mu_1} \left(1 + \frac{w\beta\varepsilon}{\eta + \varphi + \mu_1} + \frac{w\delta_1\xi}{\theta + \mu + \chi + \mu_1} + \frac{w\delta_1\eta\varepsilon}{(\xi + \varphi + \mu_1)(v + \xi + \mu_1)} + \frac{w\delta_1\xi\theta}{(v + \xi + \mu_1)(\theta + \mu + \chi + \mu_1)} \right) \right] \\ &\leq \frac{S}{N} (I + I_D + I_A + I_R) (R_0 - 1) \leq 0 \text{ if } R_0 \leq 1. \end{aligned} \quad (3.54)$$

Therefore the COVID-19 will be eliminated if and only if $R_0 < 1$. In particular, all parameters in the COVID-19 model are positive thus the Lyapunov decreases if and only if $R_0 \leq 1$ constant if and only if

$$I = I_D = I_A = I_R = 0. \quad (3.55)$$

Therefore L is a Lyapunov function within the feasible biological interval and the bigger compact invariant set in $\{S, I, I_D, I_A, I_R, I_T, R, D, V \in \Omega : \frac{dL}{dt} \leq 0\}$ is the point E_0 . By the well-known LaSalle's invariance concept [8] each solution of the COVID-19 model suggested in this work with initial condition in Ω leads to E_0 when $t \rightarrow \infty$ only if $R_0 \leq 1$. Conclusion, the disease-free equilibrium E_0 of the COVID-19 model suggested here which includes treatment and vaccination is globally asymptotically stable.

3.3 Local and global stability of the endemic equilibrium

We compute first the Jacobian matrix of the COVID-19 model for endemic equilibrium case

$$JE_* = \begin{bmatrix} \alpha^*(x) - (\gamma_1 + \mu_1) & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ \alpha^*(x) & -l_1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & \xi & -l_2 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & \varepsilon & 0 & -l_3 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \theta & \eta & -l_4 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \mu & 0 & v & -l_5 & 0 & 0 & 0 & 0 \\ 0 & \lambda & \chi & \varphi & \xi & \sigma & -l_6 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \tau & -\mu_1 & 0 & 0 \\ \gamma_1 & 0 & 0 & 0 & 0 & -\tau & \Phi & 0 & -\mu_1 & 0 \end{bmatrix}. \quad (3.56)$$

We now construct a characteristic equation

$$P = \det |-JE_* + I_M \lambda| = 0 \quad (3.57)$$

where I_M is the 9×9 unit matrix. Then we have

$$\det \begin{bmatrix} \lambda + l & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ \alpha^* & \lambda + l_1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & \xi & \lambda + l_2 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & \varepsilon & 0 & \lambda + l_3 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \theta & \eta & \lambda + l_4 & 0 & 0 & 0 & 0 \\ 0 & 0 & \mu & 0 & v & \lambda + l_5 & 0 & 0 & 0 \\ 0 & \lambda & \chi & \varphi & \xi & \sigma & \lambda + l_6 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \tau & \lambda + \mu_1 & 0 \\ \gamma_1 & 0 & 0 & 0 & 0 & -\tau & \Phi & 0 & \lambda + \mu_1 \end{bmatrix}. \quad (3.58)$$

From the above, we obtain the

$$P(\lambda) = \lambda^9 + a_1 \lambda^8 + a_2 \lambda^7 + a_3 \lambda^6 + a_4 \lambda^5 + a_5 \lambda^4 + a_6 \lambda^3 + a_7 \lambda^2 + a_8 \lambda + a_9. \quad (3.59)$$

The square Hurwitz matrix associate to $P(\lambda)$ is given as

$$H = \begin{bmatrix} a_1 & a_3 & a_5 & a_7 & a_9 & 0 & 0 & 0 & 0 \\ 1 & a_2 & a_4 & a_6 & a_8 & 0 & 0 & 0 & 0 \\ 0 & a_1 & a_3 & a_5 & a_7 & a_9 & 0 & 0 & 0 \\ 0 & 1 & a_2 & a_4 & a_6 & a_8 & 0 & 0 & 0 \\ 0 & 0 & a_1 & a_3 & a_5 & a_7 & a_9 & 0 & 0 \\ 0 & 0 & 1 & a_2 & a_4 & a_6 & a_8 & 0 & 0 \\ 0 & 0 & 0 & a_1 & a_3 & a_5 & a_7 & a_9 & 0 \\ 0 & 0 & 0 & 1 & a_2 & a_4 & a_6 & a_8 & 0 \\ 0 & 0 & 0 & 0 & a_1 & a_3 & a_5 & a_7 & a_9 \end{bmatrix}. \quad (3.60)$$

Then we have

$$H_1 = a_1 > 0$$

$$H_2 = a_1 a_2 - a_3 > 0$$

$$H_3 = -a_1^2 a_4 + a_1 a_2 a_3 + a_1 a_5 - a_3^2 > 0$$

$$H_4 = a_1^2 a_2 a_6 - a_1^2 a_4^2 - a_1 a_2^2 a_5 + a_1 a_2 a_3 a_4 - a_1 a_2 a_7 - a_1 a_3 a_6 + 2a_1 a_4 a_5 + a_2 a_3 a_5 - a_3^2 a_4 + a_3 a_7 - a_5^2 > 0$$

$$H_5 = a_1^3 a_4 a_8 - a_1^3 a_6^2 - a_1^2 a_2 a_3 a_8 - a_1^2 a_2 a_4 a_7 + 2a_1^2 a_2 a_5 a_6 + a_1^2 a_3 a_4 a_6 - a_1^2 a_4^2 a_5 + a_1 a_2^2 a_3 a_7 - a_1 a_2^2 a_5^2 - a_3^3 + a_1 a_2 a_3 a_4 a_5 - a_1^2 a_4 a_9 - a_1^2 a_5 a_8 + 2a_1^2 a_6 a_7 + a_1 a_2 a_3 a_9 - a_1 a_2 a_5 a_7 + a_1 a_1^3 a_8 - 3a_1 a_3 a_5 a_6 - a_1 a_2 a_7 + 2a_1 a_4 a_5^2 - a_2 a_3^2 a_7 + a_2 a_3 a_5^2$$

(3.61)

$$+ a_3^3 a_6 - a_3^2 a_4 a_5 + a_1 a_5 a_9 - a_1 a_7^2 - a_9 a_3^2 + 2a_3 a_5 a_7 > 0$$

$$H_6 = -a_1^3 a_8^2 a_2 + 2a_1^3 a_4 a_6 a_8 - a_1^3 a_6^3 - a_1^2 a_2^2 a_6 a_9 + 2a_1^2 a_2^2 a_7 a_8 - a_1^2 a_2 a_3 a_6 a_8 + a_1^2 a_4^2 a_2 a_9 - 3a_1^2 a_2 a_6 a_4 a_7 + 2a_1^2 a_2 a_5 a_6^2 - a_1^2 a_4^2 a_3 a_8 + a_1^2 a_6^2 a_3 a_4 + a_1^2 a_4^3 a_7 - a_1^2 a_4^2 a_5 a_6 + a_2^3 a_1 a_5 a_9 - a_1 a_1^2 a_7^2 - a_1 a_2^2 a_3 a_4 a_9 - a_1 a_2^2 a_3 a_8 a_5 + 2a_1 a_2^2 a_3 a_6 a_7 + a_1 a_2^2 a_4 a_5 a_7 - a_1 a_2^2 a_5^2 a_6 + a_1 a_2 a_3^2 a_4 a_8 - a_1 a_2 a_3^2 a_6^2 - a_1 a_2 a_3 a_7 a_4^2 + a_1 a_2 a_3 a_4 a_5 a_6 + 2a_1^2 a_2 a_8 a_9 + a_1^2 a_3 a_8^2 - 2a_1^2 a_4 a_6 a_9 - 2a_1^2 a_4 a_7 a_8 - 2a_1^2 a_5 a_6 a_8 + 3a_1^2 a_6^2 a_7 - a_1 a_2^2 a_7 a_9 + 3a_1 a_2 a_3 a_6 a_9 - 3a_1 a_2 a_3 a_7 a_8 - 2a_1 a_2 a_4 a_5 a_9 + 3a_1 a_2 a_4 a_7^2 - a_1 a_2 a_5 a_7 a_6 + a_1 a_3^2 a_6 a_8 + 2a_1 a_4 a_7 a_9 + 2a_1 a_4 a_3 a_5 a_8 + a_1 a_4 a_3 a_6 a_7 - 3a_1 a_3 a_5 a_6^2 - 2a_1 a_4^2 a_5 a_7 + 2a_1 a_4 a_5^2 a_6 + 2a_1 a_5 a_6 a_9 - a_2^2 a_3 a_5 a_9 + a_2^2 a_3 a_7^2 + a_2 a_3^2 a_4 a_9 + a_2 a_3^2 a_5 a_8 - 2a_2 a_3^2 a_6 a_7 - a_2 a_3 a_4 a_5 a_7 + a_2 a_3 a_6 a_5^2 - a_3^3 a_4 a_8 + a_3^3 a_6^2 + a_3^2 a_4^2 a_7 - a_3^2 a_4 a_5 a_6 - a_1 a_2 a_9^2 - 2a_1 a_3 a_8 a_9 + a_2 a_3 a_7 a_9 + 2a_1 a_7 a_5 a_8 + a_2 a_5^2 a_9 - a_2 a_7^2 a_5 + a_3^2 a_7 a_8 - 2a_4 a_3 a_7^2 + 3a_5 a_3 a_6 a_7 - 3a_1 a_6 a_7^2 - 2a_9 a_6 a_3^2 - a_3 a_8 a_5^2 + a_4 a_5^2 a_7 - a_5^3 a_6 + a_3 a_9^2 - 2a_7 a_5 a_9 + a_7^3 > 0$$

$$\begin{aligned}
 H_7 = & a_1^4 a_8^3 + 2a_1^3 a_2 a_6 a_8 a_9 - 3a_1^3 a_2 a_7 a_8^2 - a_1^3 a_3 a_6 a_8^2 + a_1^3 a_4^2 a_8 a_9 - 2a_1^3 a_8^2 a_4 a_5 - a_1^3 a_4 a_6^2 a_9 + 3a_1^3 a_4 a_6 a_8 a_7 \\
 & + a_1^3 a_5 a_6^2 a_8 - a_1^3 a_6^3 a_7 + a_1^2 a_2^2 a_9^2 a_4 - 2a_1^2 a_2^2 a_8 a_9 a_5 + 3a_1^2 a_2 a_3 a_5 a_8^2 - 3a_1^2 a_2 a_4 a_6 a_7^2 - 2a_1^2 a_4^2 a_3 a_7 a_8 \\
 & - 2a_1^2 a_2^2 a_7 a_9 a_6 + 3a_1^2 a_2^2 a_7^2 a_8 - 3a_1^2 a_2 a_3 a_4 a_8 a_9 + 2a_1^2 a_2 a_5 a_6^2 a_7 + a_1^2 a_4 a_3^2 a_8^2 + a_1^2 a_4^2 a_3 a_6 a_9 + a_1^2 a_4^2 a_5^2 a_8 \\
 & - a_1^2 a_2 a_3 a_6 a_7 a_8 + a_1^2 a_2 a_4^2 a_7 a_9 + 2a_1^2 a_2 a_5 a_4 a_6 a_9 + a_1^2 a_2 a_5 a_4 a_7 a_8 - 2a_1^2 a_2 a_8 a_6 a_5^2 + a_1^2 a_4^3 a_7^2 - a_1 a_2^3 a_7^3 \\
 & - a_1^2 a_3 a_4 a_5 a_6 a_8 + a_1^2 a_6^2 a_3 a_4 a_7 - a_1^2 a_4^3 a_9 a_5 - a_1^2 a_4^2 a_5 a_6 a_7 - a_1 a_2^3 a_3 a_9^2 + 2a_1 a_2^2 a_5 a_7 a_9 + 2a_1 a_2^2 a_3^2 a_8 a_9 \\
 & - a_1 a_2^2 a_3 a_4 a_7 a_9 - 3a_1 a_2^2 a_3 a_5 a_7 a_8 + 2a_1 a_2^2 a_3 a_6 a_7^2 - a_1 a_2^2 a_4 a_9 a_5^2 + a_1 a_2^2 a_4 a_5 a_7^2 + a_1 a_2^2 a_5^3 a_8 - a_1 a_2^3 a_9^3 \\
 & - a_1 a_2^2 a_5^2 a_6 a_7 - a_1 a_2 a_2^3 a_8^2 - a_1 a_2 a_2^3 a_4 a_6 a_9 + 2a_1 a_2 a_2^3 a_4 a_7 a_8 + a_1 a_2 a_2^3 a_5 a_6 a_8 - a_1 a_2 a_2^3 a_6^2 a_7 \\
 & + a_1 a_2 a_3 a_4^2 a_5 a_9 - a_1 a_2 a_3 a_4^2 a_7^2 - a_1 a_2 a_3 a_4 a_5^2 a_8 + a_1 a_2 a_3 a_4 a_5 a_6 a_7 - 3a_1^3 a_8^2 a_9 - 2a_1^2 a_2 a_6 a_9^2 + a_5^2 a_9^2 \\
 & + 4a_1^2 a_2 a_7 a_8 a_9 + 4a_1^2 a_3 a_7 a_8^2 - a_1^2 a_4^2 a_9^2 + 2a_1^2 a_4 a_5 a_9 a_8 - a_1^2 a_4 a_6 a_9 a_7 - 3a_1^2 a_4 a_7^2 a_8 + 2a_1^2 a_5^2 a_8^2 \\
 & - 5a_1^2 a_5 a_6 a_7 a_8 + 3a_1^2 a_6^2 a_7^2 + a_1 a_2^2 a_5 a_9^2 - a_1 a_2^2 a_9 a_7^2 + a_1 a_2 a_3 a_4 a_9^2 + a_1 a_2 a_3 a_5 a_8 a_9 + 5a_1 a_2 a_3 a_6 a_7 a_9 \\
 & - 5a_1 a_2 a_3 a_7^2 a_8 - 5a_1 a_2 a_4 a_5 a_7 a_9 + 3a_1 a_2 a_4 + 3a_1 a_2 a_4 a_7^3 + a_1 a_2 a_5^2 a_7 a_8 - a_1 a_2 a_5 a_6 a_7^2 + a_3^4 a_8^2 \\
 & + a_1 a_2^3 a_4 a_8 a_9 - 4a_1 a_2^3 a_5 a_8^2 + a_1 a_2^3 a_6 a_7 a_8 - 3a_1 a_3 a_4 a_5 a_6 a_9 + 4a_1 a_3 a_4 a_5 a_7 a_8 + a_1 a_3 a_4 a_6 a_7^2 \\
 & + 3a_1 a_3 a_5^2 a_6 a_8 - 3a_1 a_3 a_5 a_6^2 a_7 + 2a_1 a_4^2 a_5^2 a_9 - 2a_1 a_4^2 a_5 a_7^2 - 2a_1 a_4 a_5^3 a_8 + 2a_1 a_4 a_5^2 a_6 a_7 - a_5^3 a_6 a_7 \\
 & + a_2^2 a_3^2 a_9^2 - 2a_2^2 a_3 a_5 a_7 a_9 + a_2^2 a_3 a_7^3 - 2a_2 a_3^3 a_8 a_9 + a_2 a_3^2 a_4 a_7 a_9 + 3a_2 a_3^2 a_5 a_7 a_8 - 2a_2 a_3^2 a_6 a_7^2 \\
 & + a_2 a_3 a_4 a_5^2 a_9 - a_2 a_3 a_4 a_5 a_7^2 - a_2 a_3 a_5^3 a_8 + a_2 a_3 a_5^2 a_6 a_7 + a_3^3 a_4 a_6 a_9 - 2a_3^3 a_4 a_7 a_8 - a_3^3 a_5 a_6 a_8 \\
 & + a_3^3 a_6^2 a_7 - a_3^2 a_4^2 a_5 a_9 + a_3^2 a_4^2 a_7^2 + a_3^2 a_5^2 a_4 a_8 - a_3^2 a_4 a_5 a_6 a_7 + 3a_1^2 a_8 a_9^2 + a_1 a_3 a_6 a_9^2 - 6a_1 a_3 a_7 a_8 a_9 \\
 & + 2a_1 a_4 a_7^2 a_9 - 3a_1 a_5^2 a_8 a_9 + 3a_5 a_6 a_7 a_9 + 4a_1 a_5 a_7^2 a_8 - 3a_1 a_6 a_7^3 - 2a_2 a_3 a_5 a_9^2 + a_2 a_3 a_7^2 a_9 + a_5^4 a_8 \\
 & + 2a_2 a_5^2 a_7 a_9 - a_2 a_5 a_7^3 - a_3^2 a_4 a_9^2 + 3a_3^2 a_5 a_8 a_9 - 3a_3^2 a_6 a_7 a_9 + 2a_3^2 a_7^2 a_8 + 2a_3 a_4 a_5 a_7 a_9 - 2a_3 a_4 a_7^3 \\
 & - 4a_3 a_5^2 a_7 a_8 + 3a_3 a_5 a_6 a_7^2 - a_4 a_5^3 a_9 + a_4 a_5^2 a_7^2 + 2a_3 a_7 a_9^2 - 3a_5 a_7^2 a_9 a_7^4 > 0.
 \end{aligned}$$

Theorem 3. If $R_0 \geq 1$, the endemic equilibrium point E_* of the COVID-19 system is globally asymptotically stable.

Proof. We prove this using the Lyapunov function

$$\begin{aligned}
 L(S^*, I^*, I_A^*, I_D^*, I_R^*, I_T^*, R^*, D^*, V^*) = & \left(S - S^* - S^* \log \frac{S^*}{S} \right) + \left(I - I^* - I^* \log \frac{I^*}{I} \right) \\
 & + \left(I_A - I_A^* - I_A^* \log \frac{I_A^*}{I_A} \right) + \left(I_D - I_D^* - I_D^* \log \frac{I_D^*}{I_D} \right) \\
 & + \left(I_R - I_R^* - I_R^* \log \frac{I_R^*}{I_R} \right) + \left(I_T - I_T^* - I_T^* \log \frac{I_T^*}{I_T} \right) \quad (3.62) \\
 & + \left(R - R^* - R^* \log \frac{R^*}{R} \right) + \left(D - D^* - D^* \log \frac{D^*}{D} \right) \\
 & + \left(V - V^* - V^* \log \frac{V^*}{V} \right).
 \end{aligned}$$

Therefore taking the derivative respect to t on both sides gives

$$\begin{aligned}
 \frac{dL}{dt} = \dot{L} = & \left(\frac{S - S^*}{S} \right) \dot{S} + \left(\frac{I - I^*}{I} \right) \dot{I} + \left(\frac{I_A - I_A^*}{I_A} \right) \dot{I}_A + \left(\frac{I_D - I_D^*}{I_D} \right) \dot{I}_D \\
 & + \left(\frac{I_R - I_R^*}{I_R} \right) \dot{I}_R + \left(\frac{I_T - I_T^*}{I_T} \right) \dot{I}_T + \left(\frac{R - R^*}{R} \right) \dot{R} + \left(\frac{D - D^*}{D} \right) \dot{D} \quad (3.63) \\
 & + \left(\frac{V - V^*}{V} \right) \dot{V}
 \end{aligned}$$

replacing $\dot{S}, \dot{I}, \dot{I}_A, \dot{I}_D, \dot{I}_R, \dot{I}_T, \dot{R}, \dot{D}$ and \dot{V} by their values, we obtain

$$\begin{aligned}
 \frac{dL}{dt} = & \left(\frac{S - S^*}{S} \right) (\Lambda - (\alpha(x) + \gamma_1 + \mu_1) S) + \left(\frac{I - I^*}{I} \right) (\alpha(x) S - l_1 I) \\
 & + \left(\frac{I_A - I_A^*}{I_A} \right) (\xi I - l_2 I_A) + \left(\frac{I_D - I_D^*}{I_D} \right) (\varepsilon I - l_3 I_D) \\
 & + \left(\frac{I_R - I_R^*}{I_R} \right) (\eta I_D + \theta I_A - l_4 I_R) + \left(\frac{I_T - I_T^*}{I_T} \right) (\mu I_A + v I_R - l_5 I_T) \\
 & + \left(\frac{R - R^*}{R} \right) (\lambda I + \varphi I_D + \chi I_A + \xi I_R + \sigma I_T - l_6 R) + \left(\frac{D - D^*}{D} \right) (\tau I_T - \mu_1 D) \\
 & + \left(\frac{V - V^*}{V} \right) (\gamma_1 S + \Phi R - \mu_1 V - \tau I_T).
 \end{aligned} \tag{3.64}$$

Then we have

$$\begin{aligned}
 \frac{dL}{dt} = & \left(\frac{S - S^*}{S} \right) (\Lambda - (\alpha(x)(S - S^*) + \gamma_1(S - S^*) + \mu_1(S - S^*))) \\
 & + \left(\frac{I - I^*}{I} \right) (\alpha(x)(S - S^*) - l_1(I - I^*)) \\
 & + \left(\frac{I_A - I_A^*}{I_A} \right) (\xi(I - I^*) - l_2(I_A - I_A^*)) + \left(\frac{I_D - I_D^*}{I_D} \right) (\varepsilon(I - I^*) - l_3(I_D - I_D^*)) \\
 & + \left(\frac{I_R - I_R^*}{I_R} \right) (\eta(I_D - I_D^*) + \theta(I_A - I_A^*) - l_4(I_R - I_R^*)) \\
 & + \left(\frac{I_T - I_T^*}{I_T} \right) (\mu(I_A - I_A^*) + v(I_R - I_R^*) - l_5(I_T - I_T^*)) \\
 & + \left(\frac{R - R^*}{R} \right) \left(\begin{array}{c} \lambda(I - I^*) + \varphi(I_D - I_D^*) + \chi(I_A - I_A^*) + \xi(I_R - I_R^*) \\ + \sigma(I_T - I_T^*) - l_6(R - R^*) \end{array} \right) \\
 & + \left(\frac{D - D^*}{D} \right) (\tau(I_T - I_T^*) - \mu_1(D - D^*)) \\
 & + \left(\frac{V - V^*}{V} \right) (\gamma_1(S - S^*) + \Phi(R - R^*) - \mu_1(V - V^*) - \tau(I_T - I_T^*)).
 \end{aligned} \tag{3.65}$$

They can be separated in two part as follows

$$\begin{aligned}
 \frac{dL}{dt} = & \frac{(S - S^*)^2}{S} (-\alpha(x) - \gamma_1 - \mu_1) + \Lambda - \frac{S^*}{S} \Lambda - l_1 \frac{(I - I^*)^2}{I} + \alpha(x) S - \alpha(x) S^* \\
 & - \alpha(x) \frac{I^*}{I} S + \alpha(x) \frac{I^*}{I} S^* - l_2 \frac{(I_A - I_A^*)^2}{I_A} + \xi I - \xi I^* - \xi \frac{I_A^*}{I_A} I + \xi \frac{I_A^*}{I_A} I^* \\
 & - l_3 \frac{(I_D - I_D^*)^2}{I_D} + \varepsilon I - \varepsilon I^* - \varepsilon \frac{I_D^*}{I_D} I + \varepsilon \frac{I_D^*}{I_D} I^* - l_4 \frac{(I_R - I_R^*)^2}{I_R} + \eta I_D - \eta I_D^* \\
 & - \eta I_D \frac{I_R^*}{I_R} + \eta \frac{I_R^*}{I_R} + \theta I_A - \theta I_A^* - \theta I_A \frac{I_R^*}{I_R} + \theta I_A^* \frac{I_R^*}{I_R} - l_5 \frac{(I_T - I_T^*)^2}{I_T} + \mu I_A - \mu I_A^* \\
 & - \mu I_A \frac{I_T^*}{I_T} + \mu I_A^* \frac{I_T^*}{I_T} + v I_R - v I_R^* - v I_R \frac{I_T^*}{I_T} + v I_R^* \frac{I_T^*}{I_T} - l_6 \frac{(R - R^*)^2}{R} + \lambda I - \lambda I^* \\
 & - \lambda I \frac{R^*}{R} + \lambda I^* \frac{R^*}{R} + \varphi I_D - \varphi I_D^* - \varphi I_D \frac{R^*}{R} + \varphi I_D^* \frac{R^*}{R} + \chi I_A - \chi I_A^* \\
 & - \chi I_A \frac{R^*}{R} + \chi I_A^* \frac{R^*}{R} + \xi I_R - \xi I_R^* - \xi I_R \frac{R^*}{R} + \xi I_R^* \frac{R^*}{R} + \sigma I_T - \sigma I_T^* \\
 & - \sigma I_T \frac{R^*}{R} + \sigma I_T^* \frac{R^*}{R} - \mu_1 \frac{(D - D^*)^2}{D} + \tau I_T - \tau I_T^* - \tau I_T \frac{D^*}{D} + \tau I_T^* \frac{D^*}{D} \\
 & - \mu_1 \frac{(V - V^*)^2}{V} - \gamma_1 S - \gamma_1 S^* - \gamma_1 S \frac{V^*}{V} + \gamma_1 S^* \frac{V^*}{V} + \Phi R - \Phi R^* \\
 & - \Phi R \frac{V^*}{V} + \Phi R^* \frac{V^*}{V} - \tau I_T + \tau I_T^* + \tau I_T \frac{V^*}{V} - \tau I_T^* \frac{V^*}{V}.
 \end{aligned} \tag{3.66}$$

This can be simplified as

$$\frac{dL}{dt} = \Pi - \Gamma \tag{3.67}$$

where

$$\begin{aligned}
 \Pi = & \Lambda + \alpha(x) S + \alpha(x) \frac{I^*}{I} S^* + \xi I + \xi \frac{I_A^*}{I_A} I^* + \varepsilon I + \varepsilon \frac{I_D^*}{I_D} I^* + \eta I_D \\
 & + \eta \frac{I_R^*}{I_R} + \theta I_A + \theta I_A^* \frac{I_R^*}{I_R} + \mu I_A + \mu I_A^* \frac{I_T^*}{I_T} + v I_R + v I_R^* \frac{I_T^*}{I_T} + \lambda I \\
 & + \lambda I^* \frac{R^*}{R} + \varphi I_D + \varphi I_D^* \frac{R^*}{R} + \chi I_A + \chi I_A^* \frac{R^*}{R} + \xi I_R + \xi I_R^* \frac{R^*}{R} \\
 & + \sigma I_T + \sigma I_T^* \frac{R^*}{R} + \tau I_T + \tau I_T^* \frac{D^*}{D} + \gamma_1 S + \gamma_1 S^* \frac{V^*}{V} + \Phi R^* \frac{V^*}{V} \\
 & + \Phi R + \tau I_T^* + \tau I_T \frac{V^*}{V}
 \end{aligned} \tag{3.68}$$

and

$$\begin{aligned}
 \Gamma = & \frac{(S - S^*)^2}{S} (\alpha(x) + \gamma_1 + \mu_1) + \frac{S^*}{S} \Lambda + l_1 \frac{(I - I^*)^2}{I} + \alpha(x) S^* \\
 & + \alpha(x) \frac{I^*}{I} S + l_2 \frac{(I_A - I_A^*)^2}{I_A} + \xi I^* + \xi \frac{I_A^*}{I_A} I + l_3 \frac{(I_D - I_D^*)^2}{I_D} + \varepsilon I^* \\
 & + \varepsilon \frac{I_D^*}{I_D} I + l_4 \frac{(I_R - I_R^*)^2}{I_R} + \eta I_D^* + \eta I_D \frac{I_R^*}{I_R} + \theta I_A^* + \theta I_A \frac{I_R^*}{I_R} + \mu I_A^* \\
 & + \mu I_A \frac{I_T^*}{I_T} + v I_R^* + v I_R \frac{I_T^*}{I_T} + l_6 \frac{(R - R^*)^2}{R} + \lambda I^* + l_5 \frac{(I_T - I_T^*)^2}{I_T} \\
 & + \lambda I \frac{R^*}{R} + \varphi I_D^* + \varphi I_D \frac{R^*}{R} + \chi I_A^* + \chi I_A \frac{R^*}{R} + \xi I_R^* + \xi I_R \frac{R^*}{R} + \sigma I_T^* \\
 & + \sigma I_T \frac{R^*}{R} + \mu_1 \frac{(D - D^*)^2}{D} + \tau I_T^* + \tau I_T \frac{D^*}{D} + \mu_1 \frac{(V - V^*)^2}{V} \\
 & + \gamma_1 S^* + \gamma_1 S \frac{V^*}{V} + \Phi R^* + \Phi R \frac{V^*}{V} + \tau I_T^* + \tau I_T \frac{V^*}{V}.
 \end{aligned} \tag{3.69}$$

Therefore having $\Pi < \Gamma$, this implies $\frac{dL}{dt} < 0$, however

$$0 = \Pi - \Gamma \quad \Rightarrow \quad \frac{dL}{dt} = 0 \quad (3.70)$$

if

$$S = S^*, I = I^*, I_A = I_A^*, I_D = I_D^*, I_R = I_R^*, I_T = I_T^*, R = R^*, D = D^* \quad \text{and} \quad V = V^*. \quad (3.71)$$

We can now conclude that the largest compact invariant set for COVID-19 model in

$$\left\{ (S^*, I^*, I_A^*, I_D^*, I_R^*, I_T^*, R^*, D^*, V^*) \in \Omega : \frac{dL}{dt} = 0 \right\} \quad (3.72)$$

is the point $\{E_*\}$ the endemic equilibrium of the COVID-19 model. Therefore using the Lasalle's invariance principle, we conclude that E_* is globally asymptotically stable in Ω if $\Pi < \Gamma$.

4 Modeling with non-local operators

Due to complexities around the spread of COVID-19, it is really hard to produce predictions. Especially, when multi-scenarios are requested. Indeed it has been reported that local operators including can not provide non-local processes for example change in processes. In this section, we present an analysis of COVID-19 model with local operators including Caputo-Caputo-Fabrizio, Atangana-Baleanu and the new introduced fractal-fractional operators. We first present the definition of each operator. We start with the definition of Caputo fractional derivative

$${}_0^C D_t^\alpha f(t) = \frac{1}{\Gamma(1-\alpha)} \int_0^t \frac{d}{d\tau} f(\tau) (t-\tau)^{-\alpha} d\tau. \quad (4.1)$$

Caputo-Fabrizio fractional derivative

$${}_0^{CF} D_t^\alpha f(t) = \frac{M(\alpha)}{1-\alpha} \int_0^t \frac{d}{d\tau} f(\tau) \exp\left[-\frac{\alpha}{1-\alpha}(t-\tau)\right] d\tau. \quad (4.2)$$

Atangana-Baleanu fractional derivative

$${}_0^{ABC} D_t^\alpha f(t) = \frac{AB(\alpha)}{1-\alpha} \int_0^t \frac{d}{d\tau} f(\tau) E_\alpha\left[-\frac{\alpha}{1-\alpha}(t-\tau)^\alpha\right] d\tau. \quad (4.3)$$

The fractal-fractional derivative with power-law kernel

$${}_0^{FFP} D_t^{\alpha,\beta} f(t) = \frac{1}{\Gamma(1-\alpha)} \frac{AGd}{dt^\beta} \int_0^t f(\tau) (t-\tau)^{-\alpha} d\tau \quad (4.4)$$

where

$$\frac{df(t)}{dt^\beta} = \lim_{t \rightarrow t_1} \frac{f(t) - f(t_1)}{t^{2-\beta} - t_1^{2-\beta}} (2-\beta). \quad (4.5)$$

The fractal-fractional derivative with exponential decay kernel

$${}_0^{FFE} D_t^{\alpha,\beta} f(t) = \frac{M(\alpha)}{1-\alpha} \frac{AGd}{dt^\beta} \int_0^t f(\tau) \exp\left[-\frac{\alpha}{1-\alpha}(t-\tau)\right] d\tau. \quad (4.6)$$

The fractal-fractional derivative with Mittag-Leffler kernel

$${}_0^{FFM} D_t^{\alpha,\beta} f(t) = \frac{AB(\alpha)}{1-\alpha} \frac{AGd}{dt^\beta} \int_0^t f(\tau) E_\alpha\left[-\frac{\alpha}{1-\alpha}(t-\tau)^\alpha\right] d\tau. \quad (4.7)$$

The associated integral operators of the last three operators are given as

$$\begin{aligned} {}_0^{FFP} J_t^{\alpha,\beta} f(t) &= \frac{1}{\Gamma(\alpha)} \int_0^t (t-\tau)^{\alpha-1} \tau^{1-\beta} f(\tau) d\tau, \\ {}_0^{FFE} J_t^{\alpha,\beta} f(t) &= \frac{1-\alpha}{M(\alpha)} t^{1-\beta} f(t) + \frac{\alpha}{M(\alpha)} \int_0^t \tau^{1-\beta} f(\tau) d\tau, \\ {}_0^{FFM} J_t^{\alpha,\beta} f(t) &= \frac{1-\alpha}{AB(\alpha)} t^{1-\beta} f(t) + \frac{\alpha}{AB(\alpha)\Gamma(\alpha)} \int_0^t (t-\tau)^{\alpha-1} \tau^{1-\beta} f(\tau) d\tau. \end{aligned} \quad (4.8)$$

4.1 Positive solutions with non-local operators

In this subsection, we present a detailed analysis of positiveness of the solutions for COVID-19 model with non-local operators. We start with ABC derivative case

$$\begin{aligned}
 {}_0^{ABC}D_t^\alpha S &= \Lambda - (\alpha(x) + \gamma_1 + \mu_1) S \\
 {}_0^{ABC}D_t^\alpha I &= \alpha(x) S - (\varepsilon + \xi + \lambda + \mu_1) I \\
 {}_0^{ABC}D_t^\alpha I_A &= \xi I - (\theta + \mu + \chi + \mu_1) I_A \\
 {}_0^{ABC}D_t^\alpha I_D &= \varepsilon I - (\eta + \varphi + \mu_1) I_D \\
 {}_0^{ABC}D_t^\alpha I_R &= \eta I_D + \theta I_A - (v + \xi + \mu_1) I_R \\
 {}_0^{ABC}D_t^\alpha I_T &= \mu I_A + v I_R - (\sigma + \tau + \mu_1) I_T \\
 {}_0^{ABC}D_t^\alpha R &= \lambda I + \varphi I_D + \chi I_A + \xi I_R + \sigma I_T - (\Phi + \mu_1) R \\
 {}_0^{ABC}D_t^\alpha D &= \tau I_T - \mu_1 D \\
 {}_0^{ABC}D_t^\alpha V &= \gamma_1 S + \Phi R - \mu_1 V - \tau I_T.
 \end{aligned} \tag{4.9}$$

The norm and all hypothesis of the classical are valid here also

$$\begin{aligned}
 {}_0^{ABC}D_t^\alpha I &= \alpha(x) S - (\varepsilon + \xi + \lambda + \mu_1) I \\
 &\geq -(\varepsilon + \xi + \lambda + \mu_1) I.
 \end{aligned} \tag{4.10}$$

This produces

$$\begin{aligned}
 I(t) &\geq I(0) E_\alpha \left[-\frac{\alpha(\varepsilon + \xi + \lambda + \mu_1) t^\alpha}{AB(\alpha) - (1 - \alpha)(\varepsilon + \xi + \lambda + \mu_1)} \right], \forall t \geq 0 \\
 S(t) &\geq S(0) E_\alpha \left[-\frac{\alpha(\|\alpha(x)\|_\infty + \gamma_1 + \mu_1) t^\alpha}{AB(\alpha) - (1 - \alpha)(\|\alpha(x)\|_\infty + \gamma_1 + \mu_1)} \right], \forall t \geq 0 \\
 I_A(t) &\geq I_A(0) E_\alpha \left[-\frac{\alpha(\theta + \mu + \chi + \mu_1) t^\alpha}{AB(\alpha) - (1 - \alpha)(\theta + \mu + \chi + \mu_1)} \right], \forall t \geq 0 \\
 I_D(t) &\geq I_D(0) E_\alpha \left[-\frac{\alpha(\eta + \varphi + \mu_1) t^\alpha}{AB(\alpha) - (1 - \alpha)(\eta + \varphi + \mu_1)} \right], \forall t \geq 0 \\
 I_R(t) &\geq I_R(0) E_\alpha \left[-\frac{\alpha(v + \xi + \mu_1) t^\alpha}{AB(\alpha) - (1 - \alpha)(v + \xi + \mu_1)} \right], \forall t \geq 0 \\
 I_T(t) &\geq I_T(0) E_\alpha \left[-\frac{\alpha(\sigma + \tau + \mu_1) t^\alpha}{AB(\alpha) - (1 - \alpha)(\sigma + \tau + \mu_1)} \right], \forall t \geq 0 \\
 R(t) &\geq R(0) E_\alpha \left[-\frac{\alpha(\Phi + \mu_1) t^\alpha}{AB(\alpha) - (1 - \alpha)(\Phi + \mu_1)} \right], \forall t \geq 0 \\
 D(t) &\geq D(0) E_\alpha \left[-\frac{\alpha\mu_1 t^\alpha}{AB(\alpha) - (1 - \alpha)\mu_1} \right], \forall t \geq 0 \\
 V(t) &\geq V(0) E_\alpha \left[-\frac{\alpha\mu_1 t^\alpha}{AB(\alpha) - (1 - \alpha)\mu_1} \right], \forall t \geq 0.
 \end{aligned} \tag{4.11}$$

This shows that if all the initial conditions are positive then all solutions are positive with the Atangana-Baleanu derivative. With Caputo-Fabrizio, we have

$$\begin{aligned}
 I(t) &\geq I(0) \exp \left[-\frac{\alpha(\varepsilon + \xi + \lambda + \mu_1)t}{M(\alpha) - (1-\alpha)(\varepsilon + \xi + \lambda + \mu_1)} \right], \forall t \geq 0 \\
 S(t) &\geq S(0) \exp \left[-\frac{\alpha(\|\alpha(x)\|_\infty + \gamma_1 + \mu_1)t}{M(\alpha) - (1-\alpha)(\|\alpha(x)\|_\infty + \gamma_1 + \mu_1)} \right], \forall t \geq 0 \\
 I_A(t) &\geq I_A(0) \exp \left[-\frac{\alpha(\theta + \mu + \chi + \mu_1)t}{M(\alpha) - (1-\alpha)(\theta + \mu + \chi + \mu_1)} \right], \forall t \geq 0 \\
 I_D(t) &\geq I_D(0) \exp \left[-\frac{\alpha(\eta + \varphi + \mu_1)t}{M(\alpha) - (1-\alpha)(\eta + \varphi + \mu_1)} \right], \forall t \geq 0 \\
 I_R(t) &\geq I_R(0) \exp \left[-\frac{\alpha(v + \xi + \mu_1)t}{M(\alpha) - (1-\alpha)(v + \xi + \mu_1)} \right], \forall t \geq 0 \\
 I_T(t) &\geq I_T(0) \exp \left[-\frac{\alpha(\sigma + \tau + \mu_1)t}{M(\alpha) - (1-\alpha)(\sigma + \tau + \mu_1)} \right], \forall t \geq 0 \\
 R(t) &\geq R(0) \exp \left[-\frac{\alpha(\Phi + \mu_1)t}{M(\alpha) - (1-\alpha)(\Phi + \mu_1)} \right], \forall t \geq 0 \\
 D(t) &\geq D(0) \exp \left[-\frac{\alpha\mu_1 t}{M(\alpha) - (1-\alpha)\mu_1} \right], \forall t \geq 0 \\
 V(t) &\geq V(0) \exp \left[-\frac{\alpha\mu_1 t}{M(\alpha) - (1-\alpha)\mu_1} \right], \forall t \geq 0.
 \end{aligned} \tag{4.12}$$

This shows that all solutions are positive if all the initial conditions are positive with Caputo-Fabrizio. With Caputo derivative, we have

$$\begin{aligned}
 I(t) &\geq I(0) E_\alpha [-(\varepsilon + \xi + \lambda + \mu_1)t^\alpha], \forall t \geq 0 \\
 S(t) &\geq S(0) E_\alpha [-(\|\alpha(x)\|_\infty + \gamma_1 + \mu_1)t^\alpha], \forall t \geq 0 \\
 I_A(t) &\geq I_A(0) E_\alpha [-(\theta + \mu + \chi + \mu_1)t^\alpha], \forall t \geq 0 \\
 I_D(t) &\geq I_D(0) E_\alpha [-(\eta + \varphi + \mu_1)t^\alpha], \forall t \geq 0 \\
 I_R(t) &\geq I_R(0) E_\alpha [-(v + \xi + \mu_1)t^\alpha], \forall t \geq 0 \\
 I_T(t) &\geq I_T(0) E_\alpha [-(\sigma + \tau + \mu_1)t^\alpha], \forall t \geq 0 \\
 R(t) &\geq R(0) E_\alpha [-(\Phi + \mu_1)t^\alpha], \forall t \geq 0 \\
 D(t) &\geq D(0) E_\alpha [-\mu_1 t^\alpha], \forall t \geq 0 \\
 V(t) &\geq V(0) E_\alpha [-\mu_1 t^\alpha], \forall t \geq 0.
 \end{aligned} \tag{4.13}$$

This shows that all solutions are positive if all the initial conditions are positive with Caputo.

For fractal-fractional case, without loss of generality, we present the proof for I class and the rest can be deduced similarly. We start with power-law case

$$\begin{aligned}
 {}_0^{FFP}D_t^{\alpha,\beta} I &= \alpha(x) S - (\varepsilon + \xi + \lambda + \mu_1) I, \forall t \geq 0 \\
 &\geq -(\varepsilon + \xi + \lambda + \mu_1) I, \forall t \geq 0.
 \end{aligned} \tag{4.14}$$

and

$$\begin{aligned}
 {}_0^{RL}D_t^{\alpha,\beta} I &\geq -t^{1-\beta} (\varepsilon + \xi + \lambda + \mu_1) I, \forall t \geq 0 \\
 &\geq -b^{1-\beta} (\varepsilon + \xi + \lambda + \mu_1) I, \forall t \geq 0.
 \end{aligned} \tag{4.15}$$

Thus, we have

$$\begin{aligned}
 I(t) &\geq I(0) E_\alpha [-b^{1-\beta} (\varepsilon + \xi + \lambda + \mu_1) t^\alpha], \forall t \geq 0 \\
 S(t) &\geq S(0) E_\alpha [-b^{1-\beta} (\|\alpha(x)\|_\infty + \gamma_1 + \mu_1) t^\alpha], \forall t \geq 0 \\
 I_A(t) &\geq I_A(0) E_\alpha [-b^{1-\beta} (\theta + \mu + \chi + \mu_1) t^\alpha], \forall t \geq 0 \\
 I_D(t) &\geq I_D(0) E_\alpha [-b^{1-\beta} (\eta + \varphi + \mu_1) t^\alpha], \forall t \geq 0 \\
 I_R(t) &\geq I_R(0) E_\alpha [-b^{1-\beta} (v + \xi + \mu_1) t^\alpha], \forall t \geq 0 \\
 I_T(t) &\geq I_T(0) E_\alpha [-b^{1-\beta} (\sigma + \tau + \mu_1) t^\alpha], \forall t \geq 0 \\
 R(t) &\geq R(0) E_\alpha [-b^{1-\beta} (\Phi + \mu_1) t^\alpha], \forall t \geq 0 \\
 D(t) &\geq D(0) E_\alpha [-b^{1-\beta} \mu_1 t^\alpha], \forall t \geq 0 \\
 V(t) &\geq V(0) E_\alpha [-b^{1-\beta} \mu_1 t^\alpha], \forall t \geq 0.
 \end{aligned}
 \tag{4.16}$$

With exponential kernel, we have

$$\begin{aligned}
 I(t) &\geq I(0) \exp \left[-\frac{b^{1-\beta} \alpha (\varepsilon + \xi + \lambda + \mu_1) t}{M(\alpha) - (1-\alpha) (\varepsilon + \xi + \lambda + \mu_1)} \right], \forall t \geq 0 \\
 S(t) &\geq S(0) \exp \left[-\frac{b^{1-\beta} \alpha (\|\alpha(x)\|_\infty + \gamma_1 + \mu_1) t}{M(\alpha) - (1-\alpha) (\|\alpha(x)\|_\infty + \gamma_1 + \mu_1)} \right], \forall t \geq 0 \\
 I_A(t) &\geq I_A(0) \exp \left[-\frac{b^{1-\beta} \alpha (\theta + \mu + \chi + \mu_1) t}{M(\alpha) - (1-\alpha) (\theta + \mu + \chi + \mu_1)} \right], \forall t \geq 0 \\
 I_D(t) &\geq I_D(0) \exp \left[-\frac{b^{1-\beta} \alpha (\eta + \varphi + \mu_1) t}{M(\alpha) - (1-\alpha) (\eta + \varphi + \mu_1)} \right], \forall t \geq 0 \\
 I_R(t) &\geq I_R(0) \exp \left[-\frac{b^{1-\beta} \alpha (v + \xi + \mu_1) t}{M(\alpha) - (1-\alpha) (v + \xi + \mu_1)} \right], \forall t \geq 0 \\
 I_T(t) &\geq I_T(0) \exp \left[-\frac{b^{1-\beta} \alpha (\sigma + \tau + \mu_1) t}{M(\alpha) - (1-\alpha) (\sigma + \tau + \mu_1)} \right], \forall t \geq 0 \\
 R(t) &\geq R(0) \exp \left[-\frac{b^{1-\beta} \alpha (\Phi + \mu_1) t}{M(\alpha) - (1-\alpha) (\Phi + \mu_1)} \right], \forall t \geq 0 \\
 D(t) &\geq D(0) \exp \left[-\frac{b^{1-\beta} \alpha \mu_1 t}{M(\alpha) - (1-\alpha) \mu_1} \right], \forall t \geq 0 \\
 V(t) &\geq V(0) \exp \left[-\frac{b^{1-\beta} \alpha \mu_1 t}{M(\alpha) - (1-\alpha) \mu_1} \right], \forall t \geq 0.
 \end{aligned}
 \tag{4.17}$$

With Mittag-Leffler kernel, we obtain

$$\begin{aligned}
 I(t) &\geq I(0) E_\alpha \left[-\frac{b^{1-\beta} \alpha (\varepsilon + \xi + \lambda + \mu_1) t^\alpha}{AB(\alpha) - (1-\alpha) (\varepsilon + \xi + \lambda + \mu_1)} \right], \forall t \geq 0 \\
 S(t) &\geq S(0) E_\alpha \left[-\frac{b^{1-\beta} \alpha (\|\alpha(x)\|_\infty + \gamma_1 + \mu_1) t^\alpha}{AB(\alpha) - (1-\alpha) (\|\alpha(x)\|_\infty + \gamma_1 + \mu_1)} \right], \forall t \geq 0 \\
 I_A(t) &\geq I_A(0) E_\alpha \left[-\frac{b^{1-\beta} \alpha (\theta + \mu + \chi + \mu_1) t^\alpha}{AB(\alpha) - (1-\alpha) (\theta + \mu + \chi + \mu_1)} \right], \forall t \geq 0 \\
 I_D(t) &\geq I_D(0) E_\alpha \left[-\frac{b^{1-\beta} \alpha (\eta + \varphi + \mu_1) t^\alpha}{AB(\alpha) - (1-\alpha) (\eta + \varphi + \mu_1)} \right], \forall t \geq 0 \\
 I_R(t) &\geq I_R(0) E_\alpha \left[-\frac{b^{1-\beta} \alpha (v + \xi + \mu_1) t^\alpha}{AB(\alpha) - (1-\alpha) (v + \xi + \mu_1)} \right], \forall t \geq 0 \\
 I_T(t) &\geq I_T(0) E_\alpha \left[-\frac{b^{1-\beta} \alpha (\sigma + \tau + \mu_1) t^\alpha}{AB(\alpha) - (1-\alpha) (\sigma + \tau + \mu_1)} \right], \forall t \geq 0 \\
 R(t) &\geq R(0) E_\alpha \left[-\frac{b^{1-\beta} \alpha (\Phi + \mu_1) t^\alpha}{AB(\alpha) - (1-\alpha) (\Phi + \mu_1)} \right], \forall t \geq 0 \\
 D(t) &\geq D(0) E_\alpha \left[-\frac{b^{1-\beta} \alpha \mu_1 t^\alpha}{AB(\alpha) - (1-\alpha) \mu_1} \right], \forall t \geq 0 \\
 V(t) &\geq V(0) E_\alpha \left[-\frac{b^{1-\beta} \alpha \mu_1 t^\alpha}{AB(\alpha) - (1-\alpha) \mu_1} \right], \forall t \geq 0.
 \end{aligned} \tag{4.18}$$

5 Numerical analysis of COVID-19 models from classical to nonlocal operators: Application of Atangana-Seda numerical scheme

While analytical methods are adequate to provide exact solution of a giving equation, or systems of equations, it is important to note that when dealing with nonlinear equations, analytical methods cannot be used. In particular, the model of COVID-19 suggested in this work either with classical or non-local operators contains nonlinear components therefore analytical methods is ineffective. Very recently, Atangana and Seda [10] made use of Newton polynomial to introduce an alternative numerical scheme that can be used to solving nonlinear equations arising in many field of science, technology and engineering, the method has been recognized to be very efficient and accurate. In this section, we will make use of the Atangana-Seda scheme to solve the suggested mathematical model for COVID-19 for different differential operators.

We start with classical case for numerical solution of COVID-19 model

$$\begin{aligned}
 \dot{S} &= \Lambda - (\alpha(x) + \gamma_1 + \mu_1) S \\
 \dot{I} &= \alpha(x) S - (\varepsilon + \xi + \lambda + \mu_1) I \\
 \dot{I}_A &= \xi I - (\theta + \mu + \chi + \mu_1) I_A \\
 \dot{I}_D &= \varepsilon I - (\eta + \varphi + \mu_1) I_D \\
 \dot{I}_R &= \eta I_D + \theta I_A - (v + \xi + \mu_1) I_R \\
 \dot{I}_T &= \mu I_A + v I_R - (\sigma + \tau + \mu_1) I_T \\
 \dot{R} &= \lambda I + \varphi I_D + \chi I_A + \xi I_R + \sigma I_T - (\Phi + \mu_1) R \\
 \dot{D} &= \tau I_T - \mu_1 D \\
 \dot{V} &= \gamma_1 S + \Phi R - \mu_1 V - \tau I_T.
 \end{aligned} \tag{5.1}$$

For simplicity, we write above equation as follows;

$$\begin{aligned}
 \dot{S} &= \tilde{S}(t, S, I, I_A, I_D, I_R, I_T, R, D, V) \\
 \dot{I} &= \tilde{I}(t, S, I, I_A, I_D, I_R, I_T, R, D, V) \\
 \dot{I}_A &= \tilde{I}_A(t, S, I, I_A, I_D, I_R, I_T, R, D, V) \\
 \dot{I}_D &= \tilde{I}_D(t, S, I, I_A, I_D, I_R, I_T, R, D, V) \\
 \dot{I}_R &= \tilde{I}_R(t, S, I, I_A, I_D, I_R, I_T, R, D, V) \\
 \dot{I}_T &= \tilde{I}_T(t, S, I, I_A, I_D, I_R, I_T, R, D, V) \\
 \dot{R} &= \tilde{R}(t, S, I, I_A, I_D, I_R, I_T, R, D, V) \\
 \dot{D} &= \tilde{D}(t, S, I, I_A, I_D, I_R, I_T, R, D, V) \\
 \dot{V} &= \tilde{V}(t, S, I, I_A, I_D, I_R, I_T, R, D, V).
 \end{aligned} \tag{5.2}$$

where

$$\begin{aligned}
 \tilde{S}(t, S, I, I_A, I_D, I_R, I_T, R, D, V) &= \Lambda - (\alpha(x) + \gamma_1 + \mu_1) S \\
 \tilde{I}(t, S, I, I_A, I_D, I_R, I_T, R, D, V) &= \alpha(x) S - (\varepsilon + \xi + \lambda + \mu_1) I \\
 \tilde{I}_A(t, S, I, I_A, I_D, I_R, I_T, R, D, V) &= \xi I - (\theta + \mu + \chi + \mu_1) I_A \\
 \tilde{I}_D(t, S, I, I_A, I_D, I_R, I_T, R, D, V) &= \varepsilon I - (\eta + \varphi + \mu_1) I_D \\
 \tilde{I}_R(t, S, I, I_A, I_D, I_R, I_T, R, D, V) &= \eta I_D + \theta I_A - (v + \xi + \mu_1) I_R \\
 \tilde{I}_T(t, S, I, I_A, I_D, I_R, I_T, R, D, V) &= \mu I_A + v I_R - (\sigma + \tau + \mu_1) I_T \\
 \tilde{R}(t, S, I, I_A, I_D, I_R, I_T, R, D, V) &= \lambda I + \varphi I_D + \chi I_A + \xi I_R + \sigma I_T - (\Phi + \mu_1) R \\
 \tilde{D}(t, S, I, I_A, I_D, I_R, I_T, R, D, V) &= \tau I_T - \mu_1 D \\
 \tilde{V}(t, S, I, I_A, I_D, I_R, I_T, R, D, V) &= \gamma_1 S + \Phi R - \mu_1 V - \tau I_T.
 \end{aligned} \tag{5.3}$$

After applying fractal-fractional integral with exponential kernel, we have the following

$$\begin{aligned}
 S(t_{p+1}) &= S(t_p) + \left[\begin{aligned} &\tilde{S}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\ &-\tilde{S}(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \end{aligned} \right] \\
 &\quad + \int_{t_p}^{t_{p+1}} \tilde{S}(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) d\tau \\
 I(t_{p+1}) &= I(t_p) + \left[\begin{aligned} &\tilde{I}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\ &-\tilde{I}(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \end{aligned} \right] \\
 &\quad + \int_{t_p}^{t_{p+1}} \tilde{I}(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) d\tau \\
 I_A(t_{p+1}) &= I_A(t_p) + \left[\begin{aligned} &\tilde{I}_A(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\ &-\tilde{I}_A(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \end{aligned} \right] \\
 &\quad + \int_{t_p}^{t_{p+1}} \tilde{I}_A(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) d\tau \\
 I_D(t_{p+1}) &= I_D(t_p) + \left[\begin{aligned} &\tilde{I}_D(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\ &-\tilde{I}_D(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \end{aligned} \right] \\
 &\quad + \int_{t_p}^{t_{p+1}} \tilde{I}_D(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) d\tau
 \end{aligned}$$

$$\begin{aligned}
 I_R(t_{p+1}) &= I_R(t_p) + \left[\begin{array}{c} \widetilde{I}_R(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\ -\widetilde{I}_R(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \end{array} \right] \quad (5.4) \\
 &\quad + \int_{t_p}^{t_{p+1}} \widetilde{I}_R(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) d\tau \\
 I_T(t_{p+1}) &= I_T(t_p) + \left[\begin{array}{c} \widetilde{I}_T(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\ -\widetilde{I}_T(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \end{array} \right] \\
 &\quad + \int_{t_p}^{t_{p+1}} \widetilde{I}_T(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) d\tau \\
 R(t_{p+1}) &= R(t_p) + \left[\begin{array}{c} \widetilde{R}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\ -\widetilde{R}(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \end{array} \right] \\
 &\quad + \int_{t_p}^{t_{p+1}} \widetilde{R}(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) d\tau \\
 D(t_{p+1}) &= D(t_p) + \left[\begin{array}{c} \widetilde{D}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\ -\widetilde{D}(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \end{array} \right] \\
 &\quad + \int_{t_p}^{t_{p+1}} \widetilde{D}(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) d\tau \\
 V(t_{p+1}) &= V(t_p) + \left[\begin{array}{c} \widetilde{V}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\ -\widetilde{V}(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \end{array} \right] \\
 &\quad + \int_{t_p}^{t_{p+1}} \widetilde{V}(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) d\tau
 \end{aligned}$$

We can have the following scheme for this model

$$\begin{aligned}
 S^{p+1} &= S^p + \frac{1-\alpha}{M(\alpha)} \left[\begin{array}{c} \widetilde{S}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\ -\widetilde{S}(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \end{array} \right] \\
 &\quad + \frac{\alpha}{M(\alpha)} \left\{ \begin{array}{l} \frac{23}{12} \widetilde{S}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \Delta t \\ -\frac{4}{3} \widetilde{S}(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \Delta t \\ +\frac{5}{12} \widetilde{S}(t_{p-2}, S^{p-2}, I^{p-2}, I_A^{p-2}, I_D^{p-2}, I_R^{p-2}, I_T^{p-2}, R^{p-2}, D^{p-2}, V^{p-2}) \Delta t \end{array} \right\} \\
 I^{p+1} &= I^p + \frac{1-\alpha}{M(\alpha)} \left[\begin{array}{c} \widetilde{I}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\ -\widetilde{I}(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \end{array} \right] \\
 &\quad + \frac{\alpha}{M(\alpha)} \left\{ \begin{array}{l} \frac{23}{12} \widetilde{I}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \Delta t \\ -\frac{4}{3} \widetilde{I}(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \Delta t \\ +\frac{5}{12} \widetilde{I}(t_{p-2}, S^{p-2}, I^{p-2}, I_A^{p-2}, I_D^{p-2}, I_R^{p-2}, I_T^{p-2}, R^{p-2}, D^{p-2}, V^{p-2}) \Delta t \end{array} \right\} \\
 I_A^{p+1} &= I_A^p + \frac{1-\alpha}{M(\alpha)} \left[\begin{array}{c} \widetilde{I}_A(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\ -\widetilde{I}_A(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \end{array} \right] \\
 &\quad + \frac{\alpha}{M(\alpha)} \left\{ \begin{array}{l} \frac{23}{12} \widetilde{I}_A(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \Delta t \\ -\frac{4}{3} \widetilde{I}_A(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \Delta t \\ +\frac{5}{12} \widetilde{I}_A(t_{p-2}, S^{p-2}, I^{p-2}, I_A^{p-2}, I_D^{p-2}, I_R^{p-2}, I_T^{p-2}, R^{p-2}, D^{p-2}, V^{p-2}) \Delta t \end{array} \right\}
 \end{aligned}$$

$$\begin{aligned}
 I_D^{p+1} &= I_D^p + \frac{1-\alpha}{M(\alpha)} \left[\begin{aligned} &\widetilde{I}_D(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\ &-\widetilde{I}_D(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \end{aligned} \right] \\
 &+ \frac{\alpha}{M(\alpha)} \left\{ \begin{aligned} &\frac{23}{12} \widetilde{I}_D(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \Delta t \\ &-\frac{4}{3} \widetilde{I}_D(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \Delta t \\ &+\frac{5}{12} \widetilde{I}_D(t_{p-2}, S^{p-2}, I^{p-2}, I_A^{p-2}, I_D^{p-2}, I_R^{p-2}, I_T^{p-2}, R^{p-2}, D^{p-2}, V^{p-2}) \Delta t \end{aligned} \right\} \\
 I_R^{p+1} &= I_R^p + \frac{1-\alpha}{M(\alpha)} \left[\begin{aligned} &\widetilde{I}_R(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\ &-\widetilde{I}_R(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \end{aligned} \right] \quad (5.5) \\
 &+ \frac{\alpha}{M(\alpha)} \left\{ \begin{aligned} &\frac{23}{12} \widetilde{I}_R(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \Delta t \\ &-\frac{4}{3} \widetilde{I}_R(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \Delta t \\ &+\frac{5}{12} \widetilde{I}_R(t_{p-2}, S^{p-2}, I^{p-2}, I_A^{p-2}, I_D^{p-2}, I_R^{p-2}, I_T^{p-2}, R^{p-2}, D^{p-2}, V^{p-2}) \Delta t \end{aligned} \right\} \\
 I_T^{p+1} &= I_T^p + \frac{1-\alpha}{M(\alpha)} \left[\begin{aligned} &\widetilde{I}_T(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\ &-\widetilde{I}_T(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \end{aligned} \right] \\
 &+ \frac{\alpha}{M(\alpha)} \left\{ \begin{aligned} &\frac{23}{12} \widetilde{I}_T(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \Delta t \\ &-\frac{4}{3} \widetilde{I}_T(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \Delta t \\ &+\frac{5}{12} \widetilde{I}_T(t_{p-2}, S^{p-2}, I^{p-2}, I_A^{p-2}, I_D^{p-2}, I_R^{p-2}, I_T^{p-2}, R^{p-2}, D^{p-2}, V^{p-2}) \Delta t \end{aligned} \right\} \\
 R^{p+1} &= R^p + \frac{1-\alpha}{M(\alpha)} \left[\begin{aligned} &\widetilde{R}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\ &-\widetilde{R}(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \end{aligned} \right] \\
 &+ \frac{\alpha}{M(\alpha)} \left\{ \begin{aligned} &\frac{23}{12} \widetilde{R}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \Delta t \\ &-\frac{4}{3} \widetilde{R}(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \Delta t \\ &+\frac{5}{12} \widetilde{R}(t_{p-2}, S^{p-2}, I^{p-2}, I_A^{p-2}, I_D^{p-2}, I_R^{p-2}, I_T^{p-2}, R^{p-2}, D^{p-2}, V^{p-2}) \Delta t \end{aligned} \right\} \\
 D^{p+1} &= D^p + \frac{1-\alpha}{M(\alpha)} \left[\begin{aligned} &\widetilde{D}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\ &-\widetilde{D}(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \end{aligned} \right] \\
 &+ \frac{\alpha}{M(\alpha)} \left\{ \begin{aligned} &\frac{23}{12} \widetilde{D}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \Delta t \\ &-\frac{4}{3} \widetilde{D}(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \Delta t \\ &+\frac{5}{12} \widetilde{D}(t_{p-2}, S^{p-2}, I^{p-2}, I_A^{p-2}, I_D^{p-2}, I_R^{p-2}, I_T^{p-2}, R^{p-2}, D^{p-2}, V^{p-2}) \Delta t \end{aligned} \right\} \\
 V^{p+1} &= V^p + \frac{1-\alpha}{M(\alpha)} \left[\begin{aligned} &\widetilde{V}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\ &-\widetilde{V}(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \end{aligned} \right] \\
 &+ \frac{\alpha}{M(\alpha)} \left\{ \begin{aligned} &\frac{23}{12} \widetilde{V}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \Delta t \\ &-\frac{4}{3} \widetilde{V}(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \Delta t \\ &+\frac{5}{12} \widetilde{V}(t_{p-2}, S^{p-2}, I^{p-2}, I_A^{p-2}, I_D^{p-2}, I_R^{p-2}, I_T^{p-2}, R^{p-2}, D^{p-2}, V^{p-2}) \Delta t \end{aligned} \right\}
 \end{aligned}$$

Now, we handle the following model with classical derivative

$$\begin{aligned}
 \dot{S} &= \Lambda - (\alpha(x) + \gamma_1 + \mu_1) S \\
 \dot{I} &= \alpha(x) S - (\varepsilon + \xi + \lambda + \mu_1) I \\
 \dot{I}_A &= \xi I - (\theta + \mu + \chi + \mu_1) I_A \\
 \dot{I}_D &= \varepsilon I - (\eta + \varphi + \mu_1) I_D \\
 \dot{I}_R &= \eta I_D + \theta I_A - (v + \xi + \mu_1) I_R \\
 \dot{I}_T &= \mu I_A + v I_R - (\sigma + \tau + \mu_1) I_T \\
 \dot{R} &= \lambda I + \varphi I_D + \chi I_A + \xi I_R + \sigma I_T - (\Phi + \mu_1) R \\
 \dot{D} &= \tau I_T - \mu_1 D \\
 \dot{V} &= \gamma_1 S + \Phi R - \mu_1 V - \tau I_T
 \end{aligned} \tag{5.6}$$

where initial conditions are

$$\begin{aligned}
 S(0) &= 57780000, I(0) = 1, I_A(0) = 1, I_D(0) = 1, I_R(0) = 1, \\
 I_T(0) &= 1, R(0) = 0.1, D(0) = 0.2, V(0) = 0.1.
 \end{aligned} \tag{5.7}$$

Also the parameters are chosen as

$$\begin{aligned}
 \Lambda &= 57000000, k = 3, p = 0.5, \eta = 0.12, \chi = 0.015, v = 0.027, x = 0.4, \\
 \theta &= 0.301, \gamma = 0.09, \beta = 0.013, \lambda = 0.0345, \varphi = 0.0345, \delta_1 = 0.01, \\
 \gamma_1 &= 0.4, \mu_1 = 0.3, \varepsilon = 0.161, \xi = 0.015, \sigma = 0.015, \tau = 0.0199, \Phi = 0.2.
 \end{aligned} \tag{5.8}$$

We present numerical simulation for COVID-19 model in Figure 33 and 34.




Figure 33. Numerical visualization for COVID-19 model in South Africa.




Figure 34. Numerical visualization for COVID-19 model in South Africa.

For same model, initial conditions are chosen

$$\begin{aligned} S(0) &= 81000000, I(0) = 1, I_A(0) = 1, I_D(0) = 1, I_R(0) = 1, \\ I_T(0) &= 1, R(0) = 0.1, D(0) = 0.2, V(0) = 0.1. \end{aligned} \quad (5.9)$$

Also the parameters are

$$\begin{aligned} \Lambda &= 80000000, k = 2, p = 0.5, \eta = 0.12, \chi = 0.015, v = 0.027, x = 0.4, \theta = 0.301, \\ \gamma &= 0.09, \beta = 0.013, \gamma_1 = 0.4, \mu_1 = 0.3, \varepsilon = 0.161, \xi = 0.015, \sigma = 0.015, \\ \tau &= 0.0199, \Phi = 0.2, \lambda = 0.0345, \varphi = 0.0345, \delta_1 = 0.01. \end{aligned} \quad (5.10)$$

We present numerical simulation for COVID-19 model in Figure 35 and 36.




Figure 35. Numerical visualization for COVID-19 model in Turkey.




Figure 36. Numerical visualization for COVID-19 model in Turkey.

Now, we replace the classical differential operator will be replaced by the operator with power-law, exponential decay and Mittag-Leffler kernels. We start with exponential decay kernel

$$\begin{aligned}
 {}_0^C F D_t^\alpha S &= \Lambda - (\alpha(x) + \gamma_1 + \mu_1) S \\
 {}_0^C F D_t^\alpha I &= \alpha(x) S - (\varepsilon + \xi + \lambda + \mu_1) I \\
 {}_0^C F D_t^\alpha I_A &= \xi I - (\theta + \mu + \chi + \mu_1) I_A \\
 {}_0^C F D_t^\alpha I_D &= \varepsilon I - (\eta + \varphi + \mu_1) I_D \\
 {}_0^C F D_t^\alpha I_R &= \eta I_D + \theta I_A - (v + \xi + \mu_1) I_R \\
 {}_0^C F D_t^\alpha I_T &= \mu I_A + v I_R - (\sigma + \tau + \mu_1) I_T \\
 {}_0^C F D_t^\alpha R &= \lambda I + \varphi I_D + \chi I_A + \xi I_R + \sigma I_T - (\Phi + \mu_1) R \\
 {}_0^C F D_t^\alpha D &= \tau I_T - \mu_1 D \\
 {}_0^C F D_t^\alpha V &= \gamma_1 S + \Phi R - \mu_1 V - \tau I_T.
 \end{aligned} \tag{5.11}$$

For simplicity, we write above equation as follows;

$$\begin{aligned}
 {}_0^C F D_t^\alpha S &= \tilde{S}(t, S, I, I_A, I_D, I_R, I_T, R, D, V) \\
 {}_0^C F D_t^\alpha I &= \tilde{I}(t, S, I, I_A, I_D, I_R, I_T, R, D, V) \\
 {}_0^C F D_t^\alpha I_A &= \tilde{I}_A(t, S, I, I_A, I_D, I_R, I_T, R, D, V) \\
 {}_0^C F D_t^\alpha I_D &= \tilde{I}_D(t, S, I, I_A, I_D, I_R, I_T, R, D, V) \\
 {}_0^C F D_t^\alpha I_R &= \tilde{I}_R(t, S, I, I_A, I_D, I_R, I_T, R, D, V) \\
 {}_0^C F D_t^\alpha I_T &= \tilde{I}_T(t, S, I, I_A, I_D, I_R, I_T, R, D, V) \\
 {}_0^C F D_t^\alpha R &= \tilde{R}(t, S, I, I_A, I_D, I_R, I_T, R, D, V) \\
 {}_0^C F D_t^\alpha D &= \tilde{D}(t, S, I, I_A, I_D, I_R, I_T, R, D, V) \\
 {}_0^C F D_t^\alpha V &= \tilde{V}(t, S, I, I_A, I_D, I_R, I_T, R, D, V).
 \end{aligned} \tag{5.12}$$

After applying fractal-fractional integral with exponential kernel, we have the following

$$\begin{aligned}
 S(t_{p+1}) &= S(t_p) + \frac{1-\alpha}{M(\alpha)} \left[\tilde{S}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \right. \\
 &\quad \left. - \tilde{S}(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \right] \\
 &\quad + \frac{\alpha}{M(\alpha)} \int_{t_p}^{t_{p+1}} \tilde{S}(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) d\tau \\
 I(t_{p+1}) &= I(t_p) + \frac{1-\alpha}{M(\alpha)} \left[\tilde{I}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \right. \\
 &\quad \left. - \tilde{I}(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \right] \\
 &\quad + \frac{\alpha}{M(\alpha)} \int_{t_p}^{t_{p+1}} \tilde{I}(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) d\tau \\
 I_A(t_{p+1}) &= I_A(t_p) + \frac{1-\alpha}{M(\alpha)} \left[\tilde{I}_A(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \right. \\
 &\quad \left. - \tilde{I}_A(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \right] \\
 &\quad + \frac{\alpha}{M(\alpha)} \int_{t_p}^{t_{p+1}} \tilde{I}_A(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) d\tau \\
 I_D(t_{p+1}) &= I_D(t_p) + \frac{1-\alpha}{M(\alpha)} \left[\tilde{I}_D(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \right. \\
 &\quad \left. - \tilde{I}_D(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \right] \\
 &\quad + \frac{\alpha}{M(\alpha)} \int_{t_p}^{t_{p+1}} \tilde{I}_D(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) d\tau
 \end{aligned}$$

$$\begin{aligned}
 I_R(t_{p+1}) &= I_R(t_p) + \frac{1-\alpha}{M(\alpha)} \left[\begin{aligned} &\widetilde{I}_R(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\ &- \widetilde{I}_R(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \end{aligned} \right] \quad (6.13) \\
 &\quad + \frac{\alpha}{M(\alpha)} \int_{t_p}^{t_{p+1}} \widetilde{I}_R(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) d\tau \\
 I_T(t_{p+1}) &= I_T(t_p) + \frac{1-\alpha}{M(\alpha)} \left[\begin{aligned} &\widetilde{I}_T(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\ &- \widetilde{I}_T(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \end{aligned} \right] \\
 &\quad + \frac{\alpha}{M(\alpha)} \int_{t_p}^{t_{p+1}} \widetilde{I}_T(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) d\tau \\
 R(t_{p+1}) &= R(t_p) + \frac{1-\alpha}{M(\alpha)} \left[\begin{aligned} &\widetilde{R}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\ &- \widetilde{R}(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \end{aligned} \right] \\
 &\quad + \frac{\alpha}{M(\alpha)} \int_{t_p}^{t_{p+1}} \widetilde{R}(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) d\tau \\
 D(t_{p+1}) &= D(t_p) + \frac{1-\alpha}{M(\alpha)} \left[\begin{aligned} &\widetilde{D}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\ &- \widetilde{D}(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \end{aligned} \right] \\
 &\quad + \frac{\alpha}{M(\alpha)} \int_{t_p}^{t_{p+1}} \widetilde{D}(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) d\tau \\
 V(t_{p+1}) &= V(t_p) + \frac{1-\alpha}{M(\alpha)} \left[\begin{aligned} &\widetilde{V}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\ &- \widetilde{V}(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \end{aligned} \right] \\
 &\quad + \frac{\alpha}{M(\alpha)} \int_{t_p}^{t_{p+1}} \widetilde{V}(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) d\tau
 \end{aligned}$$

We can have the following scheme for this model

$$\begin{aligned}
 S^{p+1} &= S^p + \frac{1-\alpha}{M(\alpha)} \left[\begin{aligned} &\widetilde{S}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\ &- \widetilde{S}(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \end{aligned} \right] \\
 &\quad + \frac{\alpha}{M(\alpha)} \left\{ \begin{aligned} &\frac{23}{12} \widetilde{S}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \Delta t \\ &- \frac{4}{3} \widetilde{S}(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \Delta t \\ &+ \frac{5}{12} \widetilde{S}(t_{p-2}, S^{p-2}, I^{p-2}, I_A^{p-2}, I_D^{p-2}, I_R^{p-2}, I_T^{p-2}, R^{p-2}, D^{p-2}, V^{p-2}) \Delta t \end{aligned} \right\} \\
 I^{p+1} &= I^p + \frac{1-\alpha}{M(\alpha)} \left[\begin{aligned} &\widetilde{I}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\ &- \widetilde{I}(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \end{aligned} \right] \\
 &\quad + \frac{\alpha}{M(\alpha)} \left\{ \begin{aligned} &\frac{23}{12} \widetilde{I}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \Delta t \\ &- \frac{4}{3} \widetilde{I}(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \Delta t \\ &+ \frac{5}{12} \widetilde{I}(t_{p-2}, S^{p-2}, I^{p-2}, I_A^{p-2}, I_D^{p-2}, I_R^{p-2}, I_T^{p-2}, R^{p-2}, D^{p-2}, V^{p-2}) \Delta t \end{aligned} \right\} \\
 I_A^{p+1} &= I_A^p + \frac{1-\alpha}{M(\alpha)} \left[\begin{aligned} &\widetilde{I}_A(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\ &- \widetilde{I}_A(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \end{aligned} \right] \\
 &\quad + \frac{\alpha}{M(\alpha)} \left\{ \begin{aligned} &\frac{23}{12} \widetilde{I}_A(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \Delta t \\ &- \frac{4}{3} \widetilde{I}_A(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \Delta t \\ &+ \frac{5}{12} \widetilde{I}_A(t_{p-2}, S^{p-2}, I^{p-2}, I_A^{p-2}, I_D^{p-2}, I_R^{p-2}, I_T^{p-2}, R^{p-2}, D^{p-2}, V^{p-2}) \Delta t \end{aligned} \right\}
 \end{aligned}$$

$$\begin{aligned}
 I_D^{p+1} &= I_D^p + \frac{1-\alpha}{M(\alpha)} \left[\begin{aligned} &\widetilde{I}_D(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\ &- \widetilde{I}_D(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \end{aligned} \right] \\
 &+ \frac{\alpha}{M(\alpha)} \left\{ \begin{aligned} &\frac{23}{12} \widetilde{I}_D(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \Delta t \\ &- \frac{4}{3} \widetilde{I}_D(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \Delta t \\ &+ \frac{5}{12} \widetilde{I}_D(t_{p-2}, S^{p-2}, I^{p-2}, I_A^{p-2}, I_D^{p-2}, I_R^{p-2}, I_T^{p-2}, R^{p-2}, D^{p-2}, V^{p-2}) \Delta t \end{aligned} \right\} \\
 I_R^{p+1} &= I_R^p + \frac{1-\alpha}{M(\alpha)} \left[\begin{aligned} &\widetilde{I}_R(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\ &- \widetilde{I}_R(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \end{aligned} \right] \quad (5.14) \\
 &+ \frac{\alpha}{M(\alpha)} \left\{ \begin{aligned} &\frac{23}{12} \widetilde{I}_R(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \Delta t \\ &- \frac{4}{3} \widetilde{I}_R(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \Delta t \\ &+ \frac{5}{12} \widetilde{I}_R(t_{p-2}, S^{p-2}, I^{p-2}, I_A^{p-2}, I_D^{p-2}, I_R^{p-2}, I_T^{p-2}, R^{p-2}, D^{p-2}, V^{p-2}) \Delta t \end{aligned} \right\} \\
 I_T^{p+1} &= I_T^p + \frac{1-\alpha}{M(\alpha)} \left[\begin{aligned} &\widetilde{I}_T(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\ &- \widetilde{I}_T(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \end{aligned} \right] \\
 &+ \frac{\alpha}{M(\alpha)} \left\{ \begin{aligned} &\frac{23}{12} \widetilde{I}_T(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \Delta t \\ &- \frac{4}{3} \widetilde{I}_T(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \Delta t \\ &+ \frac{5}{12} \widetilde{I}_T(t_{p-2}, S^{p-2}, I^{p-2}, I_A^{p-2}, I_D^{p-2}, I_R^{p-2}, I_T^{p-2}, R^{p-2}, D^{p-2}, V^{p-2}) \Delta t \end{aligned} \right\} \\
 R^{p+1} &= R^p + \frac{1-\alpha}{M(\alpha)} \left[\begin{aligned} &\widetilde{R}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\ &- \widetilde{R}(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \end{aligned} \right] \\
 &+ \frac{\alpha}{M(\alpha)} \left\{ \begin{aligned} &\frac{23}{12} \widetilde{R}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \Delta t \\ &- \frac{4}{3} \widetilde{R}(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \Delta t \\ &+ \frac{5}{12} \widetilde{R}(t_{p-2}, S^{p-2}, I^{p-2}, I_A^{p-2}, I_D^{p-2}, I_R^{p-2}, I_T^{p-2}, R^{p-2}, D^{p-2}, V^{p-2}) \Delta t \end{aligned} \right\} \\
 D^{p+1} &= D^p + \frac{1-\alpha}{M(\alpha)} \left[\begin{aligned} &\widetilde{D}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\ &- \widetilde{D}(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \end{aligned} \right] \\
 &+ \frac{\alpha}{M(\alpha)} \left\{ \begin{aligned} &\frac{23}{12} \widetilde{D}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \Delta t \\ &- \frac{4}{3} \widetilde{D}(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \Delta t \\ &+ \frac{5}{12} \widetilde{D}(t_{p-2}, S^{p-2}, I^{p-2}, I_A^{p-2}, I_D^{p-2}, I_R^{p-2}, I_T^{p-2}, R^{p-2}, D^{p-2}, V^{p-2}) \Delta t \end{aligned} \right\} \\
 V^{p+1} &= V^p + \frac{1-\alpha}{M(\alpha)} \left[\begin{aligned} &\widetilde{V}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\ &- \widetilde{V}(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \end{aligned} \right] \\
 &+ \frac{\alpha}{M(\alpha)} \left\{ \begin{aligned} &\frac{23}{12} \widetilde{V}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \Delta t \\ &- \frac{4}{3} \widetilde{V}(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \Delta t \\ &+ \frac{5}{12} \widetilde{V}(t_{p-2}, S^{p-2}, I^{p-2}, I_A^{p-2}, I_D^{p-2}, I_R^{p-2}, I_T^{p-2}, R^{p-2}, D^{p-2}, V^{p-2}) \Delta t \end{aligned} \right\}
 \end{aligned}$$

For Mittag-Leffler kernel, we can have the following

$$\begin{aligned}
 S^{p+1} &= S^p + \frac{1-\alpha}{AB(\alpha)} \widetilde{S}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\
 &\quad + \frac{\alpha}{AB(\alpha)\Gamma(\alpha)} \sum_{r=2}^p \int_{t_r}^{t_{r+1}} \widetilde{S}(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V)(t_{p+1} - \tau)^{\alpha-1} d\tau
 \end{aligned}$$

$$\begin{aligned}
 I^{p+1} &= I^p + \frac{1-\alpha}{AB(\alpha)} \tilde{I}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\
 &\quad \frac{\alpha}{AB(\alpha)\Gamma(\alpha)} \sum_{r=2}^p \int_{t_r}^{t_{r+1}} \tilde{I}(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) (t_{p+1} - \tau)^{\alpha-1} d\tau \\
 I_A^{p+1} &= I_A^p + \frac{1-\alpha}{AB(\alpha)} \tilde{I}_A(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\
 &\quad \frac{\alpha}{AB(\alpha)\Gamma(\alpha)} \sum_{r=2}^p \int_{t_r}^{t_{r+1}} \tilde{I}_A(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) (t_{p+1} - \tau)^{\alpha-1} d\tau \\
 I_D^{p+1} &= I_D^p + \frac{1-\alpha}{AB(\alpha)} \tilde{I}_D(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\
 &\quad \frac{\alpha}{AB(\alpha)\Gamma(\alpha)} \sum_{r=2}^p \int_{t_r}^{t_{r+1}} \tilde{I}_D(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) (t_{p+1} - \tau)^{\alpha-1} d\tau \\
 I_R^{p+1} &= I_R^p + \frac{1-\alpha}{AB(\alpha)} \tilde{I}_R(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\
 &\quad \frac{\alpha}{AB(\alpha)\Gamma(\alpha)} \sum_{r=2}^p \int_{t_r}^{t_{r+1}} \tilde{I}_R(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) (t_{p+1} - \tau)^{\alpha-1} d\tau \\
 I_T^{p+1} &= I_T^p + \frac{1-\alpha}{AB(\alpha)} \tilde{I}_T(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\
 &\quad \frac{\alpha}{AB(\alpha)\Gamma(\alpha)} \sum_{r=2}^p \int_{t_r}^{t_{r+1}} \tilde{I}_T(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) (t_{p+1} - \tau)^{\alpha-1} d\tau \\
 R^{p+1} &= R^p + \frac{1-\alpha}{AB(\alpha)} \tilde{R}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\
 &\quad \frac{\alpha}{AB(\alpha)\Gamma(\alpha)} \sum_{r=2}^p \int_{t_r}^{t_{r+1}} \tilde{R}(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) (t_{p+1} - \tau)^{\alpha-1} d\tau \\
 D^{p+1} &= D^p + \frac{1-\alpha}{AB(\alpha)} \tilde{D}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\
 &\quad \frac{\alpha}{AB(\alpha)\Gamma(\alpha)} \sum_{r=2}^p \int_{t_r}^{t_{r+1}} \tilde{D}(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) (t_{p+1} - \tau)^{\alpha-1} d\tau \\
 V^{p+1} &= V^p + \frac{1-\alpha}{AB(\alpha)} \tilde{V}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\
 &\quad \frac{\alpha}{AB(\alpha)\Gamma(\alpha)} \sum_{r=2}^p \int_{t_r}^{t_{r+1}} \tilde{V}(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) (t_{p+1} - \tau)^{\alpha-1} d\tau
 \end{aligned} \tag{5.15}$$

We can get the following numerical scheme

$$\begin{aligned}
 S^{p+1} &= \frac{1-\alpha}{AB(\alpha)} \tilde{S}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\
 &+ \frac{\alpha(\Delta t)^\alpha}{AB(\alpha)\Gamma(\alpha+1)} \sum_{r=2}^p \tilde{S}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \\
 &\times [(p-r+1)^\alpha - (p-r)^\alpha] \\
 &+ \frac{\alpha(\Delta t)^\alpha}{AB(\alpha)\Gamma(\alpha+2)} \sum_{r=2}^p \left[\begin{array}{l} \tilde{S}(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ -\tilde{S}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha (p-r+3+2\alpha) \\ -(p-r)^\alpha (p-r+3+3\alpha) \end{array} \right] \\
 &+ \frac{\alpha(\Delta t)^\alpha}{2AB(\alpha)\Gamma(\alpha+3)} \sum_{r=2}^p \left[\begin{array}{l} \tilde{S}(t_r, S^r, I^r, I_A^r, I_D^r, I_R^r, I_T^r, R^r, D^r, V^r) \\ -2\tilde{S}(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ +\tilde{S}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (3\alpha+10)(p-r) \\ +2\alpha^2 + 9\alpha + 12 \end{array} \right] \\ -(p-r)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (5\alpha+10)(p-r) \\ +6\alpha^2 + 18\alpha + 12 \end{array} \right] \end{array} \right]
 \end{aligned} \tag{5.16}$$

$$\begin{aligned}
 I^{p+1} &= \frac{1-\alpha}{AB(\alpha)} \tilde{I}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\
 &+ \frac{\alpha(\Delta t)^\alpha}{AB(\alpha)\Gamma(\alpha+1)} \sum_{r=2}^p \tilde{I}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \\
 &\times [(p-r+1)^\alpha - (p-r)^\alpha] \\
 &+ \frac{\alpha(\Delta t)^\alpha}{AB(\alpha)\Gamma(\alpha+2)} \sum_{r=2}^p \left[\begin{array}{l} \tilde{I}(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ -\tilde{I}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha (p-r+3+2\alpha) \\ -(p-r)^\alpha (p-r+3+3\alpha) \end{array} \right] \\
 &+ \frac{\alpha(\Delta t)^\alpha}{2AB(\alpha)\Gamma(\alpha+3)} \sum_{r=2}^p \left[\begin{array}{l} \tilde{I}(t_r, S^r, I^r, I_A^r, I_D^r, I_R^r, I_T^r, R^r, D^r, V^r) \\ -2\tilde{I}(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ +\tilde{I}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (3\alpha+10)(p-r) \\ +2\alpha^2 + 9\alpha + 12 \end{array} \right] \\ -(p-r)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (5\alpha+10)(p-r) \\ +6\alpha^2 + 18\alpha + 12 \end{array} \right] \end{array} \right]
 \end{aligned}$$

$$\begin{aligned}
 I_A^{p+1} &= \frac{1-\alpha}{AB(\alpha)} \widetilde{I}_A(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\
 &+ \frac{\alpha(\Delta t)^\alpha}{AB(\alpha)\Gamma(\alpha+1)} \sum_{r=2}^p \widetilde{I}_A(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \\
 &\times [(p-r+1)^\alpha - (p-r)^\alpha] \\
 &+ \frac{\alpha(\Delta t)^\alpha}{AB(\alpha)\Gamma(\alpha+2)} \sum_{r=2}^p \left[\begin{array}{l} \widetilde{I}_A(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ -\widetilde{I}_A(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha (p-r+3+2\alpha) \\ -(p-r)^\alpha (p-r+3+3\alpha) \end{array} \right] \\
 &+ \frac{\alpha(\Delta t)^\alpha}{2AB(\alpha)\Gamma(\alpha+3)} \sum_{r=2}^p \left[\begin{array}{l} \widetilde{I}_A(t_r, S^r, I^r, I_A^r, I_D^r, I_R^r, I_T^r, R^r, D^r, V^r) \\ -2\widetilde{I}_A(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ +\widetilde{I}_A(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (3\alpha+10)(p-r) \\ +2\alpha^2 + 9\alpha + 12 \end{array} \right] \\ -(p-r)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (5\alpha+10)(p-r) \\ +6\alpha^2 + 18\alpha + 12 \end{array} \right] \end{array} \right]
 \end{aligned}$$

$$\begin{aligned}
 I_D^{p+1} &= \frac{1-\alpha}{AB(\alpha)} \widetilde{I}_D(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\
 &+ \frac{\alpha(\Delta t)^\alpha}{AB(\alpha)\Gamma(\alpha+1)} \sum_{r=2}^p \widetilde{I}_D(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \\
 &\times [(p-r+1)^\alpha - (p-r)^\alpha] \\
 &+ \frac{\alpha(\Delta t)^\alpha}{AB(\alpha)\Gamma(\alpha+2)} \sum_{r=2}^p \left[\begin{array}{l} \widetilde{I}_D(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ -\widetilde{I}_D(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha (p-r+3+2\alpha) \\ -(p-r)^\alpha (p-r+3+3\alpha) \end{array} \right] \\
 &+ \frac{\alpha(\Delta t)^\alpha}{2AB(\alpha)\Gamma(\alpha+3)} \sum_{r=2}^p \left[\begin{array}{l} \widetilde{I}_D(t_r, S^r, I^r, I_A^r, I_D^r, I_R^r, I_T^r, R^r, D^r, V^r) \\ -2\widetilde{I}_D(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ +\widetilde{I}_D(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (3\alpha+10)(p-r) \\ +2\alpha^2 + 9\alpha + 12 \end{array} \right] \\ -(p-r)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (5\alpha+10)(p-r) \\ +6\alpha^2 + 18\alpha + 12 \end{array} \right] \end{array} \right]
 \end{aligned}$$

$$\begin{aligned}
 I_R^{p+1} &= \frac{1-\alpha}{AB(\alpha)} \widetilde{I}_R(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\
 &+ \frac{\alpha(\Delta t)^\alpha}{AB(\alpha)\Gamma(\alpha+1)} \sum_{r=2}^p \widetilde{I}_R(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \\
 &\times [(p-r+1)^\alpha - (p-r)^\alpha] \\
 &+ \frac{\alpha(\Delta t)^\alpha}{AB(\alpha)\Gamma(\alpha+2)} \sum_{r=2}^p \left[\begin{array}{l} \widetilde{I}_R(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ -\widetilde{I}_R(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha (p-r+3+2\alpha) \\ -(p-r)^\alpha (p-r+3+3\alpha) \end{array} \right] \\
 &+ \frac{\alpha(\Delta t)^\alpha}{2AB(\alpha)\Gamma(\alpha+3)} \sum_{r=2}^p \left[\begin{array}{l} \widetilde{I}_R(t_r, S^r, I^r, I_A^r, I_D^r, I_R^r, I_T^r, R^r, D^r, V^r) \\ -2\widetilde{I}_R(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ +\widetilde{I}_R(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (3\alpha+10)(p-r) \\ +2\alpha^2 + 9\alpha + 12 \end{array} \right] \\ -(p-r)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (5\alpha+10)(p-r) \\ +6\alpha^2 + 18\alpha + 12 \end{array} \right] \end{array} \right]
 \end{aligned}$$

$$\begin{aligned}
 I_T^{p+1} &= \frac{1-\alpha}{AB(\alpha)} \widetilde{I}_T(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\
 &+ \frac{\alpha(\Delta t)^\alpha}{AB(\alpha)\Gamma(\alpha+1)} \sum_{r=2}^p \widetilde{I}_T(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \\
 &\times [(p-r+1)^\alpha - (p-r)^\alpha] \\
 &+ \frac{\alpha(\Delta t)^\alpha}{AB(\alpha)\Gamma(\alpha+2)} \sum_{r=2}^p \left[\begin{array}{l} \widetilde{I}_T(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ -\widetilde{I}_T(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha (p-r+3+2\alpha) \\ -(p-r)^\alpha (p-r+3+3\alpha) \end{array} \right] \\
 &+ \frac{\alpha(\Delta t)^\alpha}{2AB(\alpha)\Gamma(\alpha+3)} \sum_{r=2}^p \left[\begin{array}{l} \widetilde{I}_T(t_r, S^r, I^r, I_A^r, I_D^r, I_R^r, I_T^r, R^r, D^r, V^r) \\ -2\widetilde{I}_T(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ +\widetilde{I}_T(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (3\alpha+10)(p-r) \\ +2\alpha^2 + 9\alpha + 12 \end{array} \right] \\ -(p-r)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (5\alpha+10)(p-r) \\ +6\alpha^2 + 18\alpha + 12 \end{array} \right] \end{array} \right]
 \end{aligned}$$

$$\begin{aligned}
 R^{p+1} &= \frac{1-\alpha}{AB(\alpha)} \tilde{R}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\
 &+ \frac{\alpha(\Delta t)^\alpha}{AB(\alpha)\Gamma(\alpha+1)} \sum_{r=2}^p \tilde{R}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \\
 &\times [(p-r+1)^\alpha - (p-r)^\alpha] \\
 &+ \frac{\alpha(\Delta t)^\alpha}{AB(\alpha)\Gamma(\alpha+2)} \sum_{r=2}^p \left[\begin{array}{l} \tilde{R}(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ -\tilde{R}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha (p-r+3+2\alpha) \\ -(p-r)^\alpha (p-r+3+3\alpha) \end{array} \right] \\
 &+ \frac{\alpha(\Delta t)^\alpha}{2AB(\alpha)\Gamma(\alpha+3)} \sum_{r=2}^p \left[\begin{array}{l} \tilde{R}(t_r, S^r, I^r, I_A^r, I_D^r, I_R^r, I_T^r, R^r, D^r, V^r) \\ -2\tilde{R}(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ +\tilde{R}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (3\alpha+10)(p-r) \\ +2\alpha^2 + 9\alpha + 12 \end{array} \right] \\ -(p-r)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (5\alpha+10)(p-r) \\ +6\alpha^2 + 18\alpha + 12 \end{array} \right] \end{array} \right]
 \end{aligned}$$

$$\begin{aligned}
 D^{p+1} &= \frac{1-\alpha}{AB(\alpha)} \tilde{D}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\
 &+ \frac{\alpha(\Delta t)^\alpha}{AB(\alpha)\Gamma(\alpha+1)} \sum_{r=2}^p \tilde{D}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \\
 &\times [(p-r+1)^\alpha - (p-r)^\alpha] \\
 &+ \frac{\alpha(\Delta t)^\alpha}{AB(\alpha)\Gamma(\alpha+2)} \sum_{r=2}^p \left[\begin{array}{l} \tilde{D}(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ -\tilde{D}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha (p-r+3+2\alpha) \\ -(p-r)^\alpha (p-r+3+3\alpha) \end{array} \right] \\
 &+ \frac{\alpha(\Delta t)^\alpha}{2AB(\alpha)\Gamma(\alpha+3)} \sum_{r=2}^p \left[\begin{array}{l} \tilde{D}(t_r, S^r, I^r, I_A^r, I_D^r, I_R^r, I_T^r, R^r, D^r, V^r) \\ -2\tilde{D}(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ +\tilde{D}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (3\alpha+10)(p-r) \\ +2\alpha^2 + 9\alpha + 12 \end{array} \right] \\ -(p-r)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (5\alpha+10)(p-r) \\ +6\alpha^2 + 18\alpha + 12 \end{array} \right] \end{array} \right]
 \end{aligned}$$

$$\begin{aligned}
 V^{p+1} &= \frac{1-\alpha}{AB(\alpha)} \tilde{V}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\
 &+ \frac{\alpha(\Delta t)^\alpha}{AB(\alpha)\Gamma(\alpha+1)} \sum_{r=2}^p \tilde{V}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \\
 &\times [(p-r+1)^\alpha - (p-r)^\alpha] \\
 &+ \frac{\alpha(\Delta t)^\alpha}{AB(\alpha)\Gamma(\alpha+2)} \sum_{r=2}^p \left[\begin{array}{l} \tilde{V}(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ -\tilde{V}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha (p-r+3+2\alpha) \\ -(p-r)^\alpha (p-r+3+3\alpha) \end{array} \right] \\
 &+ \frac{\alpha(\Delta t)^\alpha}{2AB(\alpha)\Gamma(\alpha+3)} \sum_{r=2}^p \left[\begin{array}{l} \tilde{V}(t_r, S^r, I^r, I_A^r, I_D^r, I_R^r, I_T^r, R^r, D^r, V^r) \\ -2\tilde{V}(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ +\tilde{V}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (3\alpha+10)(p-r) \\ +2\alpha^2 + 9\alpha + 12 \end{array} \right] \\ -(p-r)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (5\alpha+10)(p-r) \\ +6\alpha^2 + 18\alpha + 12 \end{array} \right] \end{array} \right]
 \end{aligned}$$

For power-law kernel, we can have the following

$$\begin{aligned}
 S^{p+1} &= \frac{1}{\Gamma(\alpha)} \sum_{r=2}^p \int_{t_r}^{t_{r+1}} \tilde{S}(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) (t_{p+1} - \tau)^{\alpha-1} d\tau \\
 I^{p+1} &= \frac{1}{\Gamma(\alpha)} \sum_{r=2}^p \int_{t_r}^{t_{r+1}} \tilde{I}(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) (t_{p+1} - \tau)^{\alpha-1} d\tau \\
 I_A^{p+1} &= \frac{1}{\Gamma(\alpha)} \sum_{r=2}^p \int_{t_r}^{t_{r+1}} \tilde{I}_A(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) (t_{p+1} - \tau)^{\alpha-1} d\tau \\
 I_D^{p+1} &= \frac{1}{\Gamma(\alpha)} \sum_{r=2}^p \int_{t_r}^{t_{r+1}} \tilde{I}_D(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) (t_{p+1} - \tau)^{\alpha-1} d\tau \\
 I_R^{p+1} &= \frac{1}{\Gamma(\alpha)} \sum_{r=2}^p \int_{t_r}^{t_{r+1}} \tilde{I}_R(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) (t_{p+1} - \tau)^{\alpha-1} d\tau \\
 I_T^{p+1} &= \frac{1}{\Gamma(\alpha)} \sum_{r=2}^p \int_{t_r}^{t_{r+1}} \tilde{I}_T(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) (t_{p+1} - \tau)^{\alpha-1} d\tau \\
 R^{p+1} &= \frac{1}{\Gamma(\alpha)} \sum_{r=2}^p \int_{t_r}^{t_{r+1}} \tilde{R}(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) (t_{p+1} - \tau)^{\alpha-1} d\tau \\
 D^{p+1} &= \frac{1}{\Gamma(\alpha)} \sum_{r=2}^p \int_{t_r}^{t_{r+1}} \tilde{D}(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) (t_{p+1} - \tau)^{\alpha-1} d\tau \\
 V^{p+1} &= \frac{1}{\Gamma(\alpha)} \sum_{r=2}^p \int_{t_r}^{t_{r+1}} \tilde{V}(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) (t_{p+1} - \tau)^{\alpha-1} d\tau
 \end{aligned} \tag{5.17}$$

We can get the following numerical scheme

$$\begin{aligned}
 S^{p+1} &= \frac{(\Delta t)^\alpha}{\Gamma(\alpha+1)} \sum_{r=2}^p \tilde{S}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \\
 &\quad \times [(p-r+1)^\alpha - (p-r)^\alpha] \\
 &\quad + \frac{(\Delta t)^\alpha}{\Gamma(\alpha+2)} \sum_{r=2}^p \left[\begin{array}{l} \tilde{S}(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ -\tilde{S}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\quad \times \left[\begin{array}{l} (p-r+1)^\alpha (p-r+3+2\alpha) \\ -(p-r)^\alpha (p-r+3+3\alpha) \end{array} \right] \\
 &\quad + \frac{(\Delta t)^\alpha}{2\Gamma(\alpha+3)} \sum_{r=2}^p \left[\begin{array}{l} \tilde{S}(t_r, S^r, I^r, I_A^r, I_D^r, I_R^r, I_T^r, R^r, D^r, V^r) \\ -2\tilde{S}(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ +\tilde{S}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\quad \times \left[\begin{array}{l} (p-r+1)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (3\alpha+10)(p-r) \\ +2\alpha^2 + 9\alpha + 12 \end{array} \right] \\ -(p-r)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (5\alpha+10)(p-r) \\ +6\alpha^2 + 18\alpha + 12 \end{array} \right] \end{array} \right] \\
 I^{p+1} &= \frac{(\Delta t)^\alpha}{\Gamma(\alpha+1)} \sum_{r=2}^p \tilde{I}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \\
 &\quad \times [(p-r+1)^\alpha - (p-r)^\alpha] \\
 &\quad + \frac{(\Delta t)^\alpha}{\Gamma(\alpha+2)} \sum_{r=2}^p \left[\begin{array}{l} \tilde{I}(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ -\tilde{I}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\quad \times \left[\begin{array}{l} (p-r+1)^\alpha (p-r+3+2\alpha) \\ -(p-r)^\alpha (p-r+3+3\alpha) \end{array} \right] \\
 &\quad + \frac{(\Delta t)^\alpha}{2\Gamma(\alpha+3)} \sum_{r=2}^p \left[\begin{array}{l} \tilde{I}(t_r, S^r, I^r, I_A^r, I_D^r, I_R^r, I_T^r, R^r, D^r, V^r) \\ -2\tilde{I}(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ +\tilde{I}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\quad \times \left[\begin{array}{l} (p-r+1)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (3\alpha+10)(p-r) \\ +2\alpha^2 + 9\alpha + 12 \end{array} \right] \\ -(p-r)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (5\alpha+10)(p-r) \\ +6\alpha^2 + 18\alpha + 12 \end{array} \right] \end{array} \right] \\
 I_A^{p+1} &= \frac{(\Delta t)^\alpha}{\Gamma(\alpha+1)} \sum_{r=2}^p \tilde{I}_A(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \\
 &\quad \times [(p-r+1)^\alpha - (p-r)^\alpha] \\
 &\quad + \frac{(\Delta t)^\alpha}{\Gamma(\alpha+2)} \sum_{r=2}^p \left[\begin{array}{l} \tilde{I}_A(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ -\tilde{I}_A(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\quad \times \left[\begin{array}{l} (p-r+1)^\alpha (p-r+3+2\alpha) \\ -(p-r)^\alpha (p-r+3+3\alpha) \end{array} \right] \\
 &\quad + \frac{(\Delta t)^\alpha}{2\Gamma(\alpha+3)} \sum_{r=2}^p \left[\begin{array}{l} \tilde{I}_A(t_r, S^r, I^r, I_A^r, I_D^r, I_R^r, I_T^r, R^r, D^r, V^r) \\ -2\tilde{I}_A(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ +\tilde{I}_A(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\quad \times \left[\begin{array}{l} (p-r+1)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (3\alpha+10)(p-r) \\ +2\alpha^2 + 9\alpha + 12 \end{array} \right] \\ -(p-r)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (5\alpha+10)(p-r) \\ +6\alpha^2 + 18\alpha + 12 \end{array} \right] \end{array} \right]
 \end{aligned}
 \tag{5.18}$$

$$\begin{aligned}
 I_D^{p+1} &= \frac{(\Delta t)^\alpha}{\Gamma(\alpha+1)} \sum_{r=2}^p \widetilde{I}_D(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \\
 &\quad \times [(p-r+1)^\alpha - (p-r)^\alpha] \\
 &\quad + \frac{(\Delta t)^\alpha}{\Gamma(\alpha+2)} \sum_{r=2}^p \left[\begin{aligned} &\widetilde{I}_D(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ &- \widetilde{I}_D(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{aligned} \right] \\
 &\quad \times \left[\begin{aligned} &(p-r+1)^\alpha (p-r+3+2\alpha) \\ &-(p-r)^\alpha (p-r+3+3\alpha) \end{aligned} \right] \\
 &\quad + \frac{(\Delta t)^\alpha}{2\Gamma(\alpha+3)} \sum_{r=2}^p \left[\begin{aligned} &\widetilde{I}_D(t_r, S^r, I^r, I_A^r, I_D^r, I_R^r, I_T^r, R^r, D^r, V^r) \\ &- 2\widetilde{I}_D(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ &+ \widetilde{I}_D(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{aligned} \right] \\
 &\quad \times \left[\begin{aligned} &(p-r+1)^\alpha \left[\begin{aligned} &2(p-r)^2 + (3\alpha+10)(p-r) \\ &+ 2\alpha^2 + 9\alpha + 12 \end{aligned} \right] \\ &-(p-r)^\alpha \left[\begin{aligned} &2(p-r)^2 + (5\alpha+10)(p-r) \\ &+ 6\alpha^2 + 18\alpha + 12 \end{aligned} \right] \end{aligned} \right]
 \end{aligned}$$

$$\begin{aligned}
 I_R^{p+1} &= \frac{(\Delta t)^\alpha}{\Gamma(\alpha+1)} \sum_{r=2}^p \widetilde{I}_R(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \\
 &\quad \times [(p-r+1)^\alpha - (p-r)^\alpha] \\
 &\quad + \frac{(\Delta t)^\alpha}{\Gamma(\alpha+2)} \sum_{r=2}^p \left[\begin{aligned} &\widetilde{I}_R(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ &- \widetilde{I}_R(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{aligned} \right] \\
 &\quad \times \left[\begin{aligned} &(p-r+1)^\alpha (p-r+3+2\alpha) \\ &-(p-r)^\alpha (p-r+3+3\alpha) \end{aligned} \right] \\
 &\quad + \frac{(\Delta t)^\alpha}{2\Gamma(\alpha+3)} \sum_{r=2}^p \left[\begin{aligned} &\widetilde{I}_R(t_r, S^r, I^r, I_A^r, I_D^r, I_R^r, I_T^r, R^r, D^r, V^r) \\ &- 2\widetilde{I}_R(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ &+ \widetilde{I}_R(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{aligned} \right] \\
 &\quad \times \left[\begin{aligned} &(p-r+1)^\alpha \left[\begin{aligned} &2(p-r)^2 + (3\alpha+10)(p-r) \\ &+ 2\alpha^2 + 9\alpha + 12 \end{aligned} \right] \\ &-(p-r)^\alpha \left[\begin{aligned} &2(p-r)^2 + (5\alpha+10)(p-r) \\ &+ 6\alpha^2 + 18\alpha + 12 \end{aligned} \right] \end{aligned} \right]
 \end{aligned}$$

$$\begin{aligned}
 I_T^{p+1} &= \frac{\alpha(\Delta t)^\alpha}{\Gamma(\alpha+1)} \sum_{r=2}^p \widetilde{I}_T(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \\
 &\quad \times [(p-r+1)^\alpha - (p-r)^\alpha] \\
 &\quad + \frac{(\Delta t)^\alpha}{\Gamma(\alpha+2)} \sum_{r=2}^p \left[\begin{aligned} &\widetilde{I}_T(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ &- \widetilde{I}_T(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{aligned} \right] \\
 &\quad \times \left[\begin{aligned} &(p-r+1)^\alpha (p-r+3+2\alpha) \\ &-(p-r)^\alpha (p-r+3+3\alpha) \end{aligned} \right] \\
 &\quad + \frac{(\Delta t)^\alpha}{2\Gamma(\alpha+3)} \sum_{r=2}^p \left[\begin{aligned} &\widetilde{I}_T(t_r, S^r, I^r, I_A^r, I_D^r, I_R^r, I_T^r, R^r, D^r, V^r) \\ &- 2\widetilde{I}_T(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ &+ \widetilde{I}_T(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{aligned} \right] \\
 &\quad \times \left[\begin{aligned} &(p-r+1)^\alpha \left[\begin{aligned} &2(p-r)^2 + (3\alpha+10)(p-r) \\ &+ 2\alpha^2 + 9\alpha + 12 \end{aligned} \right] \\ &-(p-r)^\alpha \left[\begin{aligned} &2(p-r)^2 + (5\alpha+10)(p-r) \\ &+ 6\alpha^2 + 18\alpha + 12 \end{aligned} \right] \end{aligned} \right]
 \end{aligned}$$

$$\begin{aligned}
 R^{p+1} &= \frac{(\Delta t)^\alpha}{\Gamma(\alpha+1)} \sum_{r=2}^p \tilde{R}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \\
 &\times [(p-r+1)^\alpha - (p-r)^\alpha] \\
 &+ \frac{(\Delta t)^\alpha}{\Gamma(\alpha+2)} \sum_{r=2}^p \left[\begin{array}{l} \tilde{R}(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ -\tilde{R}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha (p-r+3+2\alpha) \\ -(p-r)^\alpha (p-r+3+3\alpha) \end{array} \right] \\
 &+ \frac{(\Delta t)^\alpha}{2\Gamma(\alpha+3)} \sum_{r=2}^p \left[\begin{array}{l} \tilde{R}(t_r, S^r, I^r, I_A^r, I_D^r, I_R^r, I_T^r, R^r, D^r, V^r) \\ -2\tilde{R}(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ +\tilde{R}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (3\alpha+10)(p-r) \\ +2\alpha^2 + 9\alpha + 12 \end{array} \right] \\ -(p-r)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (5\alpha+10)(p-r) \\ +6\alpha^2 + 18\alpha + 12 \end{array} \right] \end{array} \right]
 \end{aligned}$$

$$\begin{aligned}
 D^{p+1} &= \frac{(\Delta t)^\alpha}{\Gamma(\alpha+1)} \sum_{r=2}^p \tilde{D}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \\
 &\times [(p-r+1)^\alpha - (p-r)^\alpha] \\
 &+ \frac{(\Delta t)^\alpha}{\Gamma(\alpha+2)} \sum_{r=2}^p \left[\begin{array}{l} \tilde{D}(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ -\tilde{D}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha (p-r+3+2\alpha) \\ -(p-r)^\alpha (p-r+3+3\alpha) \end{array} \right] \\
 &+ \frac{(\Delta t)^\alpha}{2\Gamma(\alpha+3)} \sum_{r=2}^p \left[\begin{array}{l} \tilde{D}(t_r, S^r, I^r, I_A^r, I_D^r, I_R^r, I_T^r, R^r, D^r, V^r) \\ -2\tilde{D}(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ +\tilde{D}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (3\alpha+10)(p-r) \\ +2\alpha^2 + 9\alpha + 12 \end{array} \right] \\ -(p-r)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (5\alpha+10)(p-r) \\ +6\alpha^2 + 18\alpha + 12 \end{array} \right] \end{array} \right]
 \end{aligned}$$

$$\begin{aligned}
 V^{p+1} &= \frac{(\Delta t)^\alpha}{\Gamma(\alpha+1)} \sum_{r=2}^p \tilde{V}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \\
 &\times [(p-r+1)^\alpha - (p-r)^\alpha] \\
 &+ \frac{(\Delta t)^\alpha}{\Gamma(\alpha+2)} \sum_{r=2}^p \left[\begin{array}{l} \tilde{V}(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ -\tilde{V}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha (p-r+3+2\alpha) \\ -(p-r)^\alpha (p-r+3+3\alpha) \end{array} \right] \\
 &+ \frac{(\Delta t)^\alpha}{2\Gamma(\alpha+3)} \sum_{r=2}^p \left[\begin{array}{l} \tilde{V}(t_r, S^r, I^r, I_A^r, I_D^r, I_R^r, I_T^r, R^r, D^r, V^r) \\ -2\tilde{V}(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ +\tilde{V}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (3\alpha+10)(p-r) \\ +2\alpha^2 + 9\alpha + 12 \end{array} \right] \\ -(p-r)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (5\alpha+10)(p-r) \\ +6\alpha^2 + 18\alpha + 12 \end{array} \right] \end{array} \right]
 \end{aligned}$$

Now, we handle the following model

$$\begin{aligned}
 {}_0^{ABC}D_t^\alpha S &= \Lambda - (\alpha(x) + \gamma_1 + \mu_1) S \\
 {}_0^{ABC}D_t^\alpha I &= \alpha(x) S - (\varepsilon + \xi + \lambda + \mu_1) I \\
 {}_0^{ABC}D_t^\alpha I_A &= \xi I - (\theta + \mu + \chi + \mu_1) I_A \\
 {}_0^{ABC}D_t^\alpha I_D &= \varepsilon I - (\eta + \varphi + \mu_1) I_D \\
 {}_0^{ABC}D_t^\alpha I_R &= \eta I_D + \theta I_A - (v + \xi + \mu_1) I_R \\
 {}_0^{ABC}D_t^\alpha I_T &= \mu I_A + v I_R - (\sigma + \tau + \mu_1) I_T \\
 {}_0^{ABC}D_t^\alpha R &= \lambda I + \varphi I_D + \chi I_A + \xi I_R + \sigma I_T - (\Phi + \mu_1) R \\
 {}_0^{ABC}D_t^\alpha D &= \tau I_T - \mu_1 D \\
 {}_0^{ABC}D_t^\alpha V &= \gamma_1 S + \Phi R - \mu_1 V - \tau I_T
 \end{aligned} \tag{5.19}$$

where initial conditions are

$$\begin{aligned}
 S(0) &= 57780000, I(0) = 1, I_A(0) = 1, I_D(0) = 1, I_R(0) = 1, \\
 I_T(0) &= 1, S(0) = 0.1, S(0) = 0.2, S(0) = 0.1.
 \end{aligned} \tag{5.20}$$

Also the parameters are chosen as

$$\begin{aligned}
 \Lambda &= 57000000, k = 3, p = 0.5, \eta = 0.12, \chi = 0.015, v = 0.027, x = 0.4, \theta = 0.301, \\
 \gamma &= 0.09, \beta = 0.013, \gamma_1 = 0.4, \mu_1 = 0.3, \varepsilon = 0.161, \xi = 0.015, \sigma = 0.015, \\
 \tau &= 0.0199, \Phi = 0.2, \lambda = 0.0345, \varphi = 0.0345, \delta_1 = 0.01.
 \end{aligned} \tag{5.21}$$

We present numerical simulation for COVID-19 model in Figure 37 and 38.




Figure 37. Numerical visualization for COVID-19 model in South Africa for $\alpha = 0.75$.




Figure 38. Numerical visualization for COVID-19 model in South Africa for $\alpha = 0.75$.

For same model, initial conditions are chosen

$$\begin{aligned} S(0) &= 81000000, I(0) = 1, I_A(0) = 1, I_D(0) = 1, I_R(0) = 1, \\ I_T(0) &= 1, R(0) = 0.1, D(0) = 0.2, V(0) = 0.1. \end{aligned} \quad (5.22)$$

Also the parameters are

$$\begin{aligned} \Lambda &= 80000000, k = 2, p = 0.5, \eta = 0.12, \chi = 0.015, v = 0.027, x = 0.4, \theta = 0.301, \\ \gamma &= 0.09, \beta = 0.013, \gamma_1 = 0.4, \mu_1 = 0.3, \varepsilon = 0.161, \xi = 0.015, \sigma = 0.015, \\ \tau &= 0.0199, \Phi = 0.2, \lambda = 0.0345, \varphi = 0.0345, \delta_1 = 0.01. \end{aligned} \quad (5.23)$$

We present numerical simulation for COVID-19 model in Figure 39 and 40.




Figure 39. Numerical visualization for COVID-19 model in Turkey for $\alpha = 0.8$.




Figure 40. Numerical visualization for COVID-19 model in Turkey for $\alpha = 0.8$.

Now, we replace the classical differential operator will be replaced by the operator with power-law, exponential decay and Mittag-Leffler kernels. We start with exponential decay kernel

$$\begin{aligned}
 {}_0^{FFE}D_t^{\alpha,\beta}S &= \Lambda - (\alpha(x) + \gamma_1 + \mu_1)S \\
 {}_0^{FFE}D_t^{\alpha,\beta}I &= \alpha(x)S - (\varepsilon + \xi + \lambda + \mu_1)I \\
 {}_0^{FFE}D_t^{\alpha,\beta}I_A &= \xi I - (\theta + \mu + \chi + \mu_1)I_A \\
 {}_0^{FFE}D_t^{\alpha,\beta}I_D &= \varepsilon I - (\eta + \varphi + \mu_1)I_D \\
 {}_0^{FFE}D_t^{\alpha,\beta}I_R &= \eta I_D + \theta I_A - (v + \xi + \mu_1)I_R \\
 {}_0^{FFE}D_t^{\alpha,\beta}I_T &= \mu I_A + v I_R - (\sigma + \tau + \mu_1)I_T \\
 {}_0^{FFE}D_t^{\alpha,\beta}R &= \lambda I + \varphi I_D + \chi I_A + \xi I_R + \sigma I_T - (\Phi + \mu_1)R \\
 {}_0^{FFE}D_t^{\alpha,\beta}D &= \tau I_T - \mu_1 D \\
 {}_0^{FFE}D_t^{\alpha,\beta}V &= \gamma_1 S + \Phi R - \mu_1 V - \tau I_T.
 \end{aligned} \tag{5.24}$$

For simplicity, we write above equation as follows;

$$\begin{aligned}
 {}_0^{FFE}D_t^{\alpha,\beta}S &= \tilde{S}(t, S, I, I_A, I_D, I_R, I_T, R, D, V) \\
 {}_0^{FFE}D_t^{\alpha,\beta}I &= \tilde{I}(t, S, I, I_A, I_D, I_R, I_T, R, D, V) \\
 {}_0^{FFE}D_t^{\alpha,\beta}I_A &= \tilde{I}_A(t, S, I, I_A, I_D, I_R, I_T, R, D, V) \\
 {}_0^{FFE}D_t^{\alpha,\beta}I_D &= \tilde{I}_D(t, S, I, I_A, I_D, I_R, I_T, R, D, V) \\
 {}_0^{FFE}D_t^{\alpha,\beta}I_R &= \tilde{I}_R(t, S, I, I_A, I_D, I_R, I_T, R, D, V) \\
 {}_0^{FFE}D_t^{\alpha,\beta}I_T &= \tilde{I}_T(t, S, I, I_A, I_D, I_R, I_T, R, D, V) \\
 {}_0^{FFE}D_t^{\alpha,\beta}R &= \tilde{R}(t, S, I, I_A, I_D, I_R, I_T, R, D, V) \\
 {}_0^{FFE}D_t^{\alpha,\beta}D &= \tilde{D}(t, S, I, I_A, I_D, I_R, I_T, R, D, V) \\
 {}_0^{FFE}D_t^{\alpha,\beta}V &= \tilde{V}(t, S, I, I_A, I_D, I_R, I_T, R, D, V).
 \end{aligned} \tag{5.25}$$

After applying fractal-fractional integral with exponential kernel, we have the following

$$\begin{aligned}
 S(t_{p+1}) &= S(t_p) + \frac{1-\alpha}{M(\alpha)} \left[t_p^{1-\beta} \tilde{S}(\tau, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \right. \\
 &\quad \left. - t_{p-1}^{1-\beta} \tilde{S}(\tau, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \right] \\
 &\quad + \frac{\alpha}{M(\alpha)} \int_{t_p}^{t_{p+1}} \tilde{S}(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) \tau^{1-\beta} d\tau \\
 I(t_{p+1}) &= I(t_p) + \frac{1-\alpha}{M(\alpha)} \left[t_p^{1-\beta} \tilde{I}(\tau, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \right. \\
 &\quad \left. - t_{p-1}^{1-\beta} \tilde{I}(\tau, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \right] \\
 &\quad + \frac{\alpha}{M(\alpha)} \int_{t_p}^{t_{p+1}} \tilde{I}(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) \tau^{1-\beta} d\tau \\
 I_A(t_{p+1}) &= I_A(t_p) + \frac{1-\alpha}{M(\alpha)} \left[t_p^{1-\beta} \tilde{I}_A(\tau, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \right. \\
 &\quad \left. - t_{p-1}^{1-\beta} \tilde{I}_A(\tau, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \right] \\
 &\quad + \frac{\alpha}{M(\alpha)} \int_{t_p}^{t_{p+1}} \tilde{I}_A(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) \tau^{1-\beta} d\tau \\
 I_D(t_{p+1}) &= I_D(t_p) + \frac{1-\alpha}{M(\alpha)} \left[t_p^{1-\beta} \tilde{I}_D(\tau, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \right. \\
 &\quad \left. - t_{p-1}^{1-\beta} \tilde{I}_D(\tau, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \right] \\
 &\quad + \frac{\alpha}{M(\alpha)} \int_{t_p}^{t_{p+1}} \tilde{I}_D(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) \tau^{1-\beta} d\tau
 \end{aligned}$$

$$\begin{aligned}
 I_R(t_{p+1}) &= I_R(t_p) + \frac{1-\alpha}{M(\alpha)} \left[t_p^{1-\beta} \widetilde{I}_R(\tau, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \right. \\
 &\quad \left. - t_{p-1}^{1-\beta} \widetilde{I}_R(\tau, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \right] \\
 &\quad + \frac{\alpha}{M(\alpha)} \int_{t_p}^{t_{p+1}} \widetilde{I}_R(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) \tau^{1-\beta} d\tau \\
 I_T(t_{p+1}) &= I_T(t_p) + \frac{1-\alpha}{M(\alpha)} \left[t_p^{1-\beta} \widetilde{I}_T(\tau, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \right. \\
 &\quad \left. - t_{p-1}^{1-\beta} \widetilde{I}_T(\tau, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \right] \\
 &\quad + \frac{\alpha}{M(\alpha)} \int_{t_p}^{t_{p+1}} \widetilde{I}_T(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) \tau^{1-\beta} d\tau \\
 R(t_{p+1}) &= R(t_p) + \frac{1-\alpha}{M(\alpha)} \left[t_p^{1-\beta} \widetilde{R}(\tau, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \right. \\
 &\quad \left. - t_{p-1}^{1-\beta} \widetilde{R}(\tau, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \right] \\
 &\quad + \frac{\alpha}{M(\alpha)} \int_{t_p}^{t_{p+1}} \widetilde{R}(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) \tau^{1-\beta} d\tau \\
 D(t_{p+1}) &= D(t_p) + \frac{1-\alpha}{M(\alpha)} \left[t_p^{1-\beta} \widetilde{D}(\tau, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \right. \\
 &\quad \left. - t_{p-1}^{1-\beta} \widetilde{D}(\tau, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \right] \\
 &\quad + \frac{\alpha}{M(\alpha)} \int_{t_p}^{t_{p+1}} \widetilde{D}(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) \tau^{1-\beta} d\tau \\
 V(t_{p+1}) &= V(t_p) + \frac{1-\alpha}{M(\alpha)} \left[t_p^{1-\beta} \widetilde{V}(\tau, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \right. \\
 &\quad \left. - t_{p-1}^{1-\beta} \widetilde{V}(\tau, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \right] \\
 &\quad + \frac{\alpha}{M(\alpha)} \int_{t_p}^{t_{p+1}} \widetilde{V}(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) \tau^{1-\beta} d\tau
 \end{aligned}
 \tag{5.26}$$

We can have the following scheme for this model

$$\begin{aligned}
 S^{p+1} &= S^p + \frac{1-\alpha}{M(\alpha)} \left[t_p^{1-\beta} \widetilde{S}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \right. \\
 &\quad \left. - t_{p-1}^{1-\beta} \widetilde{S}(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \right] \\
 &\quad + \frac{\alpha}{M(\alpha)} \left\{ \begin{aligned} &\frac{23}{12} t_p^{1-\beta} \widetilde{S}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \Delta t \\ &-\frac{4}{3} t_{p-1}^{1-\beta} \widetilde{S}(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \Delta t \\ &+\frac{5}{12} t_{p-2}^{1-\beta} \widetilde{S}(t_{p-2}, S^{p-2}, I^{p-2}, I_A^{p-2}, I_D^{p-2}, I_R^{p-2}, I_T^{p-2}, R^{p-2}, D^{p-2}, V^{p-2}) \Delta t \end{aligned} \right\} \\
 I^{p+1} &= I^p + \frac{1-\alpha}{M(\alpha)} \left[t_p^{1-\beta} \widetilde{I}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \right. \\
 &\quad \left. - t_{p-1}^{1-\beta} \widetilde{I}(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \right] \\
 &\quad + \frac{\alpha}{M(\alpha)} \left\{ \begin{aligned} &\frac{23}{12} t_p^{1-\beta} \widetilde{I}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \Delta t \\ &-\frac{4}{3} t_{p-1}^{1-\beta} \widetilde{I}(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \Delta t \\ &+\frac{5}{12} t_{p-2}^{1-\beta} \widetilde{I}(t_{p-2}, S^{p-2}, I^{p-2}, I_A^{p-2}, I_D^{p-2}, I_R^{p-2}, I_T^{p-2}, R^{p-2}, D^{p-2}, V^{p-2}) \Delta t \end{aligned} \right\} \\
 I_A^{p+1} &= I_A^p + \frac{1-\alpha}{M(\alpha)} \left[t_p^{1-\beta} \widetilde{I}_A(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \right. \\
 &\quad \left. - t_{p-1}^{1-\beta} \widetilde{I}_A(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \right] \\
 &\quad + \frac{\alpha}{M(\alpha)} \left\{ \begin{aligned} &\frac{23}{12} t_p^{1-\beta} \widetilde{I}_A(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \Delta t \\ &-\frac{4}{3} t_{p-1}^{1-\beta} \widetilde{I}_A(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \Delta t \\ &+\frac{5}{12} t_{p-2}^{1-\beta} \widetilde{I}_A(t_{p-2}, S^{p-2}, I^{p-2}, I_A^{p-2}, I_D^{p-2}, I_R^{p-2}, I_T^{p-2}, R^{p-2}, D^{p-2}, V^{p-2}) \Delta t \end{aligned} \right\}
 \end{aligned}
 \tag{5.27}$$

$$I_D^{p+1} = I_D^p + \frac{1-\alpha}{M(\alpha)} \left[t_p^{1-\beta} \widetilde{I}_D(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) - t_{p-1}^{1-\beta} \widetilde{I}_D(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \right] + \frac{\alpha}{M(\alpha)} \left\{ \begin{aligned} & \frac{23}{12} t_p^{1-\beta} \widetilde{I}_D(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \Delta t \\ & - \frac{4}{3} t_{p-1}^{1-\beta} \widetilde{I}_D(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \Delta t \\ & + \frac{5}{12} t_{p-2}^{1-\beta} \widetilde{I}_D(t_{p-2}, S^{p-2}, I^{p-2}, I_A^{p-2}, I_D^{p-2}, I_R^{p-2}, I_T^{p-2}, R^{p-2}, D^{p-2}, V^{p-2}) \Delta t \end{aligned} \right\}$$

$$I_R^{p+1} = I_R^p + \frac{1-\alpha}{M(\alpha)} \left[t_p^{1-\beta} \widetilde{I}_R(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) - t_{p-1}^{1-\beta} \widetilde{I}_R(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \right] + \frac{\alpha}{M(\alpha)} \left\{ \begin{aligned} & \frac{23}{12} t_p^{1-\beta} \widetilde{I}_R(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \Delta t \\ & - \frac{4}{3} t_{p-1}^{1-\beta} \widetilde{I}_R(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \Delta t \\ & + \frac{5}{12} t_{p-2}^{1-\beta} \widetilde{I}_R(t_{p-2}, S^{p-2}, I^{p-2}, I_A^{p-2}, I_D^{p-2}, I_R^{p-2}, I_T^{p-2}, R^{p-2}, D^{p-2}, V^{p-2}) \Delta t \end{aligned} \right\}$$

$$I_T^{p+1} = I_T^p + \frac{1-\alpha}{M(\alpha)} \left[t_p^{1-\beta} \widetilde{I}_T(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) - t_{p-1}^{1-\beta} \widetilde{I}_T(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \right] + \frac{\alpha}{M(\alpha)} \left\{ \begin{aligned} & \frac{23}{12} t_p^{1-\beta} \widetilde{I}_T(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \Delta t \\ & - \frac{4}{3} t_{p-1}^{1-\beta} \widetilde{I}_T(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \Delta t \\ & + \frac{5}{12} t_{p-2}^{1-\beta} \widetilde{I}_T(t_{p-2}, S^{p-2}, I^{p-2}, I_A^{p-2}, I_D^{p-2}, I_R^{p-2}, I_T^{p-2}, R^{p-2}, D^{p-2}, V^{p-2}) \Delta t \end{aligned} \right\}$$

$$R^{p+1} = R^p + \frac{1-\alpha}{M(\alpha)} \left[t_p^{1-\beta} \widetilde{R}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) - t_{p-1}^{1-\beta} \widetilde{R}(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \right] + \frac{\alpha}{M(\alpha)} \left\{ \begin{aligned} & \frac{23}{12} t_p^{1-\beta} \widetilde{R}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \Delta t \\ & - \frac{4}{3} t_{p-1}^{1-\beta} \widetilde{R}(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \Delta t \\ & + \frac{5}{12} t_{p-2}^{1-\beta} \widetilde{R}(t_{p-2}, S^{p-2}, I^{p-2}, I_A^{p-2}, I_D^{p-2}, I_R^{p-2}, I_T^{p-2}, R^{p-2}, D^{p-2}, V^{p-2}) \Delta t \end{aligned} \right\}$$

$$D^{p+1} = D^p + \frac{1-\alpha}{M(\alpha)} \left[t_p^{1-\beta} \widetilde{D}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) - t_{p-1}^{1-\beta} \widetilde{D}(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \right] + \frac{\alpha}{M(\alpha)} \left\{ \begin{aligned} & \frac{23}{12} t_p^{1-\beta} \widetilde{D}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \Delta t \\ & - \frac{4}{3} t_{p-1}^{1-\beta} \widetilde{D}(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \Delta t \\ & + \frac{5}{12} t_{p-2}^{1-\beta} \widetilde{D}(t_{p-2}, S^{p-2}, I^{p-2}, I_A^{p-2}, I_D^{p-2}, I_R^{p-2}, I_T^{p-2}, R^{p-2}, D^{p-2}, V^{p-2}) \Delta t \end{aligned} \right\}$$

$$V^{p+1} = V^p + \frac{1-\alpha}{M(\alpha)} \left[t_p^{1-\beta} \widetilde{V}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) - t_{p-1}^{1-\beta} \widetilde{V}(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \right] + \frac{\alpha}{M(\alpha)} \left\{ \begin{aligned} & \frac{23}{12} t_p^{1-\beta} \widetilde{V}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \Delta t \\ & - \frac{4}{3} t_{p-1}^{1-\beta} \widetilde{V}(t_{p-1}, S^{p-1}, I^{p-1}, I_A^{p-1}, I_D^{p-1}, I_R^{p-1}, I_T^{p-1}, R^{p-1}, D^{p-1}, V^{p-1}) \Delta t \\ & + \frac{5}{12} t_{p-2}^{1-\beta} \widetilde{V}(t_{p-2}, S^{p-2}, I^{p-2}, I_A^{p-2}, I_D^{p-2}, I_R^{p-2}, I_T^{p-2}, R^{p-2}, D^{p-2}, V^{p-2}) \Delta t \end{aligned} \right\}$$

For Mittag-Leffler kernel, we can have the following

$$S^{p+1} = S^p + \frac{1-\alpha}{AB(\alpha)} t_p^{1-\beta} \widetilde{S}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) + \frac{\alpha}{AB(\alpha)\Gamma(\alpha)} \sum_{r=2}^p \int_{t_r}^{t_{r+1}} \widetilde{S}(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) (t_{p+1} - \tau)^{\alpha-1} \tau^{1-\beta} d\tau$$

$$\begin{aligned}
 I^{p+1} &= I^p + \frac{1-\alpha}{AB(\alpha)} t_p^{1-\beta} \tilde{I}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\
 &\quad + \frac{\alpha}{AB(\alpha)\Gamma(\alpha)} \sum_{r=2}^p \int_{t_r}^{t_{r+1}} \tilde{I}(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) (t_{p+1} - \tau)^{\alpha-1} \tau^{1-\beta} d\tau \\
 I_A^{p+1} &= I_A^p + \frac{1-\alpha}{AB(\alpha)} t_p^{1-\beta} \tilde{I}_A(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\
 &\quad + \frac{\alpha}{AB(\alpha)\Gamma(\alpha)} \sum_{r=2}^p \int_{t_r}^{t_{r+1}} \tilde{I}_A(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) (t_{p+1} - \tau)^{\alpha-1} d\tau \\
 I_D^{p+1} &= I_D^p + \frac{1-\alpha}{AB(\alpha)} t_p^{1-\beta} \tilde{I}_D(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\
 &\quad + \frac{\alpha}{AB(\alpha)\Gamma(\alpha)} \sum_{r=2}^p \int_{t_r}^{t_{r+1}} \tilde{I}_D(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) (t_{p+1} - \tau)^{\alpha-1} \tau^{1-\beta} d\tau \\
 I_R^{p+1} &= I_R^p + \frac{1-\alpha}{AB(\alpha)} t_p^{1-\beta} \tilde{I}_R(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\
 &\quad + \frac{\alpha}{AB(\alpha)\Gamma(\alpha)} \sum_{r=2}^p \int_{t_r}^{t_{r+1}} \tilde{I}_R(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) (t_{p+1} - \tau)^{\alpha-1} d\tau \\
 I_T^{p+1} &= I_T^p + \frac{1-\alpha}{AB(\alpha)} t_p^{1-\beta} \tilde{I}_T(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\
 &\quad + \frac{\alpha}{AB(\alpha)\Gamma(\alpha)} \sum_{r=2}^p \int_{t_r}^{t_{r+1}} \tilde{I}_T(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) (t_{p+1} - \tau)^{\alpha-1} \tau^{1-\beta} d\tau \\
 R^{p+1} &= R^p + \frac{1-\alpha}{AB(\alpha)} t_p^{1-\beta} \tilde{R}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\
 &\quad + \frac{\alpha}{AB(\alpha)\Gamma(\alpha)} \sum_{r=2}^p \int_{t_r}^{t_{r+1}} \tilde{R}(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) (t_{p+1} - \tau)^{\alpha-1} d\tau \\
 D^{p+1} &= D^p + \frac{1-\alpha}{AB(\alpha)} t_p^{1-\beta} \tilde{D}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\
 &\quad + \frac{\alpha}{AB(\alpha)\Gamma(\alpha)} \sum_{r=2}^p \int_{t_r}^{t_{r+1}} \tilde{D}(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) (t_{p+1} - \tau)^{\alpha-1} \tau^{1-\beta} d\tau \\
 V^{p+1} &= V^p + \frac{1-\alpha}{AB(\alpha)} t_p^{1-\beta} \tilde{V}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\
 &\quad + \frac{\alpha}{AB(\alpha)\Gamma(\alpha)} \sum_{r=2}^p \int_{t_r}^{t_{r+1}} \tilde{V}(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) (t_{p+1} - \tau)^{\alpha-1} \tau^{1-\beta} d\tau
 \end{aligned} \tag{5.28}$$

We can get the following numerical scheme

$$\begin{aligned}
 S^{p+1} &= \frac{1-\alpha}{AB(\alpha)} t_p^{1-\beta} \tilde{S}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\
 &+ \frac{\alpha(\Delta t)^\alpha}{AB(\alpha)\Gamma(\alpha+1)} \sum_{r=2}^p t_{r-2}^{1-\beta} \tilde{S}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \\
 &\times [(p-r+1)^\alpha - (p-r)^\alpha] \\
 &+ \frac{\alpha(\Delta t)^\alpha}{AB(\alpha)\Gamma(\alpha+2)} \sum_{r=2}^p \left[\begin{array}{l} t_{r-1}^{1-\beta} \tilde{S}(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ -t_{r-2}^{1-\beta} \tilde{S}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha (p-r+3+2\alpha) \\ -(p-r)^\alpha (p-r+3+3\alpha) \end{array} \right] \\
 &+ \frac{\alpha(\Delta t)^\alpha}{2AB(\alpha)\Gamma(\alpha+3)} \sum_{r=2}^p \left[\begin{array}{l} t_r^{1-\beta} \tilde{S}(t_r, S^r, I^r, I_A^r, I_D^r, I_R^r, I_T^r, R^r, D^r, V^r) \\ -2t_{r-1}^{1-\beta} \tilde{S}(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ +t_{r-2}^{1-\beta} \tilde{S}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (3\alpha+10)(p-r) \\ +2\alpha^2 + 9\alpha + 12 \end{array} \right] \\ -(p-r)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (5\alpha+10)(p-r) \\ +6\alpha^2 + 18\alpha + 12 \end{array} \right] \end{array} \right]
 \end{aligned}
 \tag{5.29}$$

$$\begin{aligned}
 I^{p+1} &= \frac{1-\alpha}{AB(\alpha)} t_p^{1-\beta} \tilde{I}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\
 &+ \frac{\alpha(\Delta t)^\alpha}{AB(\alpha)\Gamma(\alpha+1)} \sum_{r=2}^p t_{r-2}^{1-\beta} \tilde{I}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \\
 &\times [(p-r+1)^\alpha - (p-r)^\alpha] \\
 &+ \frac{\alpha(\Delta t)^\alpha}{AB(\alpha)\Gamma(\alpha+2)} \sum_{r=2}^p \left[\begin{array}{l} t_{r-1}^{1-\beta} \tilde{I}(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ -t_{r-2}^{1-\beta} \tilde{I}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha (p-r+3+2\alpha) \\ -(p-r)^\alpha (p-r+3+3\alpha) \end{array} \right] \\
 &+ \frac{\alpha(\Delta t)^\alpha}{2AB(\alpha)\Gamma(\alpha+3)} \sum_{r=2}^p \left[\begin{array}{l} t_r^{1-\beta} \tilde{I}(t_r, S^r, I^r, I_A^r, I_D^r, I_R^r, I_T^r, R^r, D^r, V^r) \\ -2t_{r-1}^{1-\beta} \tilde{I}(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ +t_{r-2}^{1-\beta} \tilde{I}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (3\alpha+10)(p-r) \\ +2\alpha^2 + 9\alpha + 12 \end{array} \right] \\ -(p-r)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (5\alpha+10)(p-r) \\ +6\alpha^2 + 18\alpha + 12 \end{array} \right] \end{array} \right]
 \end{aligned}$$

$$\begin{aligned}
 I_A^{p+1} &= \frac{1-\alpha}{AB(\alpha)} t_p^{1-\beta} \widetilde{I}_A(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\
 &+ \frac{\alpha(\Delta t)^\alpha}{AB(\alpha)\Gamma(\alpha+1)} \sum_{r=2}^p t_{r-2}^{1-\beta} \widetilde{I}_A(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \\
 &\times [(p-r+1)^\alpha - (p-r)^\alpha] \\
 &+ \frac{\alpha(\Delta t)^\alpha}{AB(\alpha)\Gamma(\alpha+2)} \sum_{r=2}^p \left[\begin{array}{l} t_{r-1}^{1-\beta} \widetilde{I}_A(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ -t_{r-2}^{1-\beta} \widetilde{I}_A(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha (p-r+3+2\alpha) \\ -(p-r)^\alpha (p-r+3+3\alpha) \end{array} \right] \\
 &+ \frac{\alpha(\Delta t)^\alpha}{2AB(\alpha)\Gamma(\alpha+3)} \sum_{r=2}^p \left[\begin{array}{l} t_r^{1-\beta} \widetilde{I}_A(t_r, S^r, I^r, I_A^r, I_D^r, I_R^r, I_T^r, R^r, D^r, V^r) \\ -2t_{r-1}^{1-\beta} \widetilde{I}_A(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ +t_{r-2}^{1-\beta} \widetilde{I}_A(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (3\alpha+10)(p-r) \\ +2\alpha^2 + 9\alpha + 12 \end{array} \right] \\ -(p-r)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (5\alpha+10)(p-r) \\ +6\alpha^2 + 18\alpha + 12 \end{array} \right] \end{array} \right]
 \end{aligned}$$

$$\begin{aligned}
 I_D^{p+1} &= \frac{1-\alpha}{AB(\alpha)} t_p^{1-\beta} \widetilde{I}_D(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\
 &+ \frac{\alpha(\Delta t)^\alpha}{AB(\alpha)\Gamma(\alpha+1)} \sum_{r=2}^p t_{r-2}^{1-\beta} \widetilde{I}_D(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \\
 &\times [(p-r+1)^\alpha - (p-r)^\alpha] \\
 &+ \frac{\alpha(\Delta t)^\alpha}{AB(\alpha)\Gamma(\alpha+2)} \sum_{r=2}^p \left[\begin{array}{l} t_{r-1}^{1-\beta} \widetilde{I}_D(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ -t_{r-2}^{1-\beta} \widetilde{I}_D(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha (p-r+3+2\alpha) \\ -(p-r)^\alpha (p-r+3+3\alpha) \end{array} \right] \\
 &+ \frac{\alpha(\Delta t)^\alpha}{2AB(\alpha)\Gamma(\alpha+3)} \sum_{r=2}^p \left[\begin{array}{l} t_r^{1-\beta} \widetilde{I}_D(t_r, S^r, I^r, I_A^r, I_D^r, I_R^r, I_T^r, R^r, D^r, V^r) \\ -2t_{r-1}^{1-\beta} \widetilde{I}_D(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ +t_{r-2}^{1-\beta} \widetilde{I}_D(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (3\alpha+10)(p-r) \\ +2\alpha^2 + 9\alpha + 12 \end{array} \right] \\ -(p-r)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (5\alpha+10)(p-r) \\ +6\alpha^2 + 18\alpha + 12 \end{array} \right] \end{array} \right]
 \end{aligned}$$

$$\begin{aligned}
 I_R^{p+1} &= \frac{1-\alpha}{AB(\alpha)} t_p^{1-\beta} \widetilde{I}_R(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\
 &+ \frac{\alpha(\Delta t)^\alpha}{AB(\alpha)\Gamma(\alpha+1)} \sum_{r=2}^p t_{r-2}^{1-\beta} \widetilde{I}_R(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \\
 &\times [(p-r+1)^\alpha - (p-r)^\alpha] \\
 &+ \frac{\alpha(\Delta t)^\alpha}{AB(\alpha)\Gamma(\alpha+2)} \sum_{r=2}^p \left[\begin{array}{l} t_{r-1}^{1-\beta} \widetilde{I}_R(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ -t_{r-2}^{1-\beta} \widetilde{I}_R(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha (p-r+3+2\alpha) \\ -(p-r)^\alpha (p-r+3+3\alpha) \end{array} \right] \\
 &+ \frac{\alpha(\Delta t)^\alpha}{2AB(\alpha)\Gamma(\alpha+3)} \sum_{r=2}^p \left[\begin{array}{l} t_r^{1-\beta} \widetilde{I}_R(t_r, S^r, I^r, I_A^r, I_D^r, I_R^r, I_T^r, R^r, D^r, V^r) \\ -2t_{r-1}^{1-\beta} \widetilde{I}_R(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ +t_{r-2}^{1-\beta} \widetilde{I}_R(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (3\alpha+10)(p-r) \\ +2\alpha^2 + 9\alpha + 12 \end{array} \right] \\ -(p-r)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (5\alpha+10)(p-r) \\ +6\alpha^2 + 18\alpha + 12 \end{array} \right] \end{array} \right]
 \end{aligned}$$

$$\begin{aligned}
 I_T^{p+1} &= \frac{1-\alpha}{AB(\alpha)} t_p^{1-\beta} \widetilde{I}_T(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\
 &+ \frac{\alpha(\Delta t)^\alpha}{AB(\alpha)\Gamma(\alpha+1)} \sum_{r=2}^p t_{r-2}^{1-\beta} \widetilde{I}_T(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \\
 &\times [(p-r+1)^\alpha - (p-r)^\alpha] \\
 &+ \frac{\alpha(\Delta t)^\alpha}{AB(\alpha)\Gamma(\alpha+2)} \sum_{r=2}^p \left[\begin{array}{l} t_{r-1}^{1-\beta} \widetilde{I}_T(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ -t_{r-2}^{1-\beta} \widetilde{I}_T(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha (p-r+3+2\alpha) \\ -(p-r)^\alpha (p-r+3+3\alpha) \end{array} \right] \\
 &+ \frac{\alpha(\Delta t)^\alpha}{2AB(\alpha)\Gamma(\alpha+3)} \sum_{r=2}^p \left[\begin{array}{l} t_r^{1-\beta} \widetilde{I}_T(t_r, S^r, I^r, I_A^r, I_D^r, I_R^r, I_T^r, R^r, D^r, V^r) \\ -2t_{r-1}^{1-\beta} \widetilde{I}_T(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ +t_{r-2}^{1-\beta} \widetilde{I}_T(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (3\alpha+10)(p-r) \\ +2\alpha^2 + 9\alpha + 12 \end{array} \right] \\ -(p-r)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (5\alpha+10)(p-r) \\ +6\alpha^2 + 18\alpha + 12 \end{array} \right] \end{array} \right]
 \end{aligned}$$

$$\begin{aligned}
 R^{p+1} &= \frac{1-\alpha}{AB(\alpha)} t_p^{1-\beta} \tilde{R}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\
 &+ \frac{\alpha(\Delta t)^\alpha}{AB(\alpha)\Gamma(\alpha+1)} \sum_{r=2}^p t_{r-2}^{1-\beta} \tilde{R}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \\
 &\times [(p-r+1)^\alpha - (p-r)^\alpha] \\
 &+ \frac{\alpha(\Delta t)^\alpha}{AB(\alpha)\Gamma(\alpha+2)} \sum_{r=2}^p \left[\begin{array}{l} t_{r-1}^{1-\beta} \tilde{R}(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ -t_{r-2}^{1-\beta} \tilde{R}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha (p-r+3+2\alpha) \\ -(p-r)^\alpha (p-r+3+3\alpha) \end{array} \right] \\
 &+ \frac{\alpha(\Delta t)^\alpha}{2AB(\alpha)\Gamma(\alpha+3)} \sum_{r=2}^p \left[\begin{array}{l} t_r^{1-\beta} \tilde{R}(t_r, S^r, I^r, I_A^r, I_D^r, I_R^r, I_T^r, R^r, D^r, V^r) \\ -2t_{r-1}^{1-\beta} \tilde{R}(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ +t_{r-2}^{1-\beta} \tilde{R}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (3\alpha+10)(p-r) \\ +2\alpha^2 + 9\alpha + 12 \end{array} \right] \\ -(p-r)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (5\alpha+10)(p-r) \\ +6\alpha^2 + 18\alpha + 12 \end{array} \right] \end{array} \right]
 \end{aligned}$$

$$\begin{aligned}
 D^{p+1} &= \frac{1-\alpha}{AB(\alpha)} t_p^{1-\beta} \tilde{D}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\
 &+ \frac{\alpha(\Delta t)^\alpha}{AB(\alpha)\Gamma(\alpha+1)} \sum_{r=2}^p t_{r-2}^{1-\beta} \tilde{D}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \\
 &\times [(p-r+1)^\alpha - (p-r)^\alpha] \\
 &+ \frac{\alpha(\Delta t)^\alpha}{AB(\alpha)\Gamma(\alpha+2)} \sum_{r=2}^p \left[\begin{array}{l} t_{r-1}^{1-\beta} \tilde{D}(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ -t_{r-2}^{1-\beta} \tilde{D}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha (p-r+3+2\alpha) \\ -(p-r)^\alpha (p-r+3+3\alpha) \end{array} \right] \\
 &+ \frac{\alpha(\Delta t)^\alpha}{2AB(\alpha)\Gamma(\alpha+3)} \sum_{r=2}^p \left[\begin{array}{l} t_r^{1-\beta} \tilde{D}(t_r, S^r, I^r, I_A^r, I_D^r, I_R^r, I_T^r, R^r, D^r, V^r) \\ -2t_{r-1}^{1-\beta} \tilde{D}(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ +t_{r-2}^{1-\beta} \tilde{D}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (3\alpha+10)(p-r) \\ +2\alpha^2 + 9\alpha + 12 \end{array} \right] \\ -(p-r)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (5\alpha+10)(p-r) \\ +6\alpha^2 + 18\alpha + 12 \end{array} \right] \end{array} \right]
 \end{aligned}$$

$$\begin{aligned}
 V^{p+1} &= \frac{1-\alpha}{AB(\alpha)} t_p^{1-\beta} \tilde{V}(t_p, S^p, I^p, I_A^p, I_D^p, I_R^p, I_T^p, R^p, D^p, V^p) \\
 &+ \frac{\alpha(\Delta t)^\alpha}{AB(\alpha)\Gamma(\alpha+1)} \sum_{r=2}^p t_{r-2}^{1-\beta} \tilde{V}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \\
 &\times [(p-r+1)^\alpha - (p-r)^\alpha] \\
 &+ \frac{\alpha(\Delta t)^\alpha}{AB(\alpha)\Gamma(\alpha+2)} \sum_{r=2}^p \left[\begin{array}{l} t_{r-1}^{1-\beta} \tilde{V}(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ - t_{r-2}^{1-\beta} \tilde{V}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha (p-r+3+2\alpha) \\ -(p-r)^\alpha (p-r+3+3\alpha) \end{array} \right] \\
 &+ \frac{\alpha(\Delta t)^\alpha}{2AB(\alpha)\Gamma(\alpha+3)} \sum_{r=2}^p \left[\begin{array}{l} t_r^{1-\beta} \tilde{V}(t_r, S^r, I^r, I_A^r, I_D^r, I_R^r, I_T^r, R^r, D^r, V^r) \\ - 2t_{r-1}^{1-\beta} \tilde{V}(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ + t_{r-2}^{1-\beta} \tilde{V}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (3\alpha+10)(p-r) \\ +2\alpha^2 + 9\alpha + 12 \end{array} \right] \\ -(p-r)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (5\alpha+10)(p-r) \\ +6\alpha^2 + 18\alpha + 12 \end{array} \right] \end{array} \right]
 \end{aligned}$$

For power-law kernel, we can have the following

$$\begin{aligned}
 S^{p+1} &= \frac{1}{\Gamma(\alpha)} \sum_{r=2}^p \int_{t_r}^{t_{r+1}} \tilde{S}(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) (t_{p+1} - \tau)^{\alpha-1} \tau^{1-\beta} d\tau \\
 I^{p+1} &= \frac{1}{\Gamma(\alpha)} \sum_{r=2}^p \int_{t_r}^{t_{r+1}} \tilde{I}(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) (t_{p+1} - \tau)^{\alpha-1} \tau^{1-\beta} d\tau \\
 I_A^{p+1} &= \frac{1}{\Gamma(\alpha)} \sum_{r=2}^p \int_{t_r}^{t_{r+1}} \tilde{I}_A(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) (t_{p+1} - \tau)^{\alpha-1} \tau^{1-\beta} d\tau \\
 I_D^{p+1} &= \frac{1}{\Gamma(\alpha)} \sum_{r=2}^p \int_{t_r}^{t_{r+1}} \tilde{I}_D(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) (t_{p+1} - \tau)^{\alpha-1} \tau^{1-\beta} d\tau \\
 I_R^{p+1} &= \frac{1}{\Gamma(\alpha)} \sum_{r=2}^p \int_{t_r}^{t_{r+1}} \tilde{I}_R(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) (t_{p+1} - \tau)^{\alpha-1} \tau^{1-\beta} d\tau \\
 I_T^{p+1} &= \frac{1}{\Gamma(\alpha)} \sum_{r=2}^p \int_{t_r}^{t_{r+1}} \tilde{I}_T(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) (t_{p+1} - \tau)^{\alpha-1} \tau^{1-\beta} d\tau \\
 R^{p+1} &= \frac{1}{\Gamma(\alpha)} \sum_{r=2}^p \int_{t_r}^{t_{r+1}} \tilde{R}(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) (t_{p+1} - \tau)^{\alpha-1} \tau^{1-\beta} d\tau \\
 D^{p+1} &= \frac{1}{\Gamma(\alpha)} \sum_{r=2}^p \int_{t_r}^{t_{r+1}} \tilde{D}(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) (t_{p+1} - \tau)^{\alpha-1} \tau^{1-\beta} d\tau \\
 V^{p+1} &= \frac{1}{\Gamma(\alpha)} \sum_{r=2}^p \int_{t_r}^{t_{r+1}} \tilde{V}(\tau, S, I, I_A, I_D, I_R, I_T, R, D, V) (t_{p+1} - \tau)^{\alpha-1} \tau^{1-\beta} d\tau
 \end{aligned} \tag{5.30}$$

We can get the following numerical scheme

$$\begin{aligned}
 S^{p+1} &= \frac{(\Delta t)^\alpha}{\Gamma(\alpha+1)} \sum_{r=2}^p t_{r-2}^{1-\beta} \tilde{S}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \\
 &\quad \times [(p-r+1)^\alpha - (p-r)^\alpha] \\
 &\quad + \frac{(\Delta t)^\alpha}{\Gamma(\alpha+2)} \sum_{r=2}^p \left[\begin{aligned} &t_{r-1}^{1-\beta} \tilde{S}(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ &- t_{r-2}^{1-\beta} \tilde{S}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{aligned} \right] \\
 &\quad \times \left[\begin{aligned} &(p-r+1)^\alpha (p-r+3+2\alpha) \\ &-(p-r)^\alpha (p-r+3+3\alpha) \end{aligned} \right] \\
 &\quad + \frac{(\Delta t)^\alpha}{2\Gamma(\alpha+3)} \sum_{r=2}^p \left[\begin{aligned} &t_r^{1-\beta} \tilde{S}(t_r, S^r, I^r, I_A^r, I_D^r, I_R^r, I_T^r, R^r, D^r, V^r) \\ &- 2t_{r-1}^{1-\beta} \tilde{S}(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ &+ t_{r-2}^{1-\beta} \tilde{S}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{aligned} \right] \\
 &\quad \times \left[\begin{aligned} &(p-r+1)^\alpha \left[\begin{aligned} &2(p-r)^2 + (3\alpha+10)(p-r) \\ &+ 2\alpha^2 + 9\alpha + 12 \end{aligned} \right] \\ &-(p-r)^\alpha \left[\begin{aligned} &2(p-r)^2 + (5\alpha+10)(p-r) \\ &+ 6\alpha^2 + 18\alpha + 12 \end{aligned} \right] \end{aligned} \right]
 \end{aligned} \tag{5.31}$$

$$\begin{aligned}
 I^{p+1} &= \frac{(\Delta t)^\alpha}{\Gamma(\alpha+1)} \sum_{r=2}^p t_{r-2}^{1-\beta} \tilde{I}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \\
 &\quad \times [(p-r+1)^\alpha - (p-r)^\alpha] \\
 &\quad + \frac{(\Delta t)^\alpha}{\Gamma(\alpha+2)} \sum_{r=2}^p \left[\begin{aligned} &t_{r-1}^{1-\beta} \tilde{I}(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ &- t_{r-2}^{1-\beta} \tilde{I}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{aligned} \right] \\
 &\quad \times \left[\begin{aligned} &(p-r+1)^\alpha (p-r+3+2\alpha) \\ &-(p-r)^\alpha (p-r+3+3\alpha) \end{aligned} \right] \\
 &\quad + \frac{(\Delta t)^\alpha}{2\Gamma(\alpha+3)} \sum_{r=2}^p \left[\begin{aligned} &t_r^{1-\beta} \tilde{I}(t_r, S^r, I^r, I_A^r, I_D^r, I_R^r, I_T^r, R^r, D^r, V^r) \\ &- 2t_{r-1}^{1-\beta} \tilde{I}(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ &+ t_{r-2}^{1-\beta} \tilde{I}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{aligned} \right] \\
 &\quad \times \left[\begin{aligned} &(p-r+1)^\alpha \left[\begin{aligned} &2(p-r)^2 + (3\alpha+10)(p-r) \\ &+ 2\alpha^2 + 9\alpha + 12 \end{aligned} \right] \\ &-(p-r)^\alpha \left[\begin{aligned} &2(p-r)^2 + (5\alpha+10)(p-r) \\ &+ 6\alpha^2 + 18\alpha + 12 \end{aligned} \right] \end{aligned} \right]
 \end{aligned}$$

$$\begin{aligned}
 I_A^{p+1} &= \frac{(\Delta t)^\alpha}{\Gamma(\alpha+1)} \sum_{r=2}^p t_{r-2}^{1-\beta} \tilde{I}_A(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \\
 &\quad \times [(p-r+1)^\alpha - (p-r)^\alpha] \\
 &\quad + \frac{(\Delta t)^\alpha}{\Gamma(\alpha+2)} \sum_{r=2}^p \left[\begin{aligned} &t_{r-1}^{1-\beta} \tilde{I}_A(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ &- t_{r-2}^{1-\beta} \tilde{I}_A(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{aligned} \right] \\
 &\quad \times \left[\begin{aligned} &(p-r+1)^\alpha (p-r+3+2\alpha) \\ &-(p-r)^\alpha (p-r+3+3\alpha) \end{aligned} \right] \\
 &\quad + \frac{(\Delta t)^\alpha}{2\Gamma(\alpha+3)} \sum_{r=2}^p \left[\begin{aligned} &t_r^{1-\beta} \tilde{I}_A(t_r, S^r, I^r, I_A^r, I_D^r, I_R^r, I_T^r, R^r, D^r, V^r) \\ &- 2t_{r-1}^{1-\beta} \tilde{I}_A(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ &+ t_{r-2}^{1-\beta} \tilde{I}_A(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{aligned} \right] \\
 &\quad \times \left[\begin{aligned} &(p-r+1)^\alpha \left[\begin{aligned} &2(p-r)^2 + (3\alpha+10)(p-r) \\ &+ 2\alpha^2 + 9\alpha + 12 \end{aligned} \right] \\ &-(p-r)^\alpha \left[\begin{aligned} &2(p-r)^2 + (5\alpha+10)(p-r) \\ &+ 6\alpha^2 + 18\alpha + 12 \end{aligned} \right] \end{aligned} \right]
 \end{aligned}$$

$$\begin{aligned}
 I_D^{p+1} &= \frac{(\Delta t)^\alpha}{\Gamma(\alpha+1)} \sum_{r=2}^p t_{r-2}^{1-\beta} \widetilde{I}_D(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \\
 &\times [(p-r+1)^\alpha - (p-r)^\alpha] \\
 &+ \frac{(\Delta t)^\alpha}{\Gamma(\alpha+2)} \sum_{r=2}^p \left[\begin{array}{l} t_{r-1}^{1-\beta} \widetilde{I}_D(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ -t_{r-2}^{1-\beta} \widetilde{I}_D(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha (p-r+3+2\alpha) \\ -(p-r)^\alpha (p-r+3+3\alpha) \end{array} \right] \\
 &+ \frac{(\Delta t)^\alpha}{2\Gamma(\alpha+3)} \sum_{r=2}^p \left[\begin{array}{l} t_r^{1-\beta} \widetilde{I}_D(t_r, S^r, I^r, I_A^r, I_D^r, I_R^r, I_T^r, R^r, D^r, V^r) \\ -2t_{r-1}^{1-\beta} \widetilde{I}_D(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ +t_{r-2}^{1-\beta} \widetilde{I}_D(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (3\alpha+10)(p-r) \\ +2\alpha^2 + 9\alpha + 12 \end{array} \right] \\ -(p-r)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (5\alpha+10)(p-r) \\ +6\alpha^2 + 18\alpha + 12 \end{array} \right] \end{array} \right]
 \end{aligned}$$

$$\begin{aligned}
 I_R^{p+1} &= \frac{(\Delta t)^\alpha}{\Gamma(\alpha+1)} \sum_{r=2}^p t_{r-2}^{1-\beta} \widetilde{I}_R(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \\
 &\times [(p-r+1)^\alpha - (p-r)^\alpha] \\
 &+ \frac{(\Delta t)^\alpha}{\Gamma(\alpha+2)} \sum_{r=2}^p \left[\begin{array}{l} t_{r-1}^{1-\beta} \widetilde{I}_R(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ -t_{r-2}^{1-\beta} \widetilde{I}_R(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha (p-r+3+2\alpha) \\ -(p-r)^\alpha (p-r+3+3\alpha) \end{array} \right] \\
 &+ \frac{(\Delta t)^\alpha}{2\Gamma(\alpha+3)} \sum_{r=2}^p \left[\begin{array}{l} t_r^{1-\beta} \widetilde{I}_R(t_r, S^r, I^r, I_A^r, I_D^r, I_R^r, I_T^r, R^r, D^r, V^r) \\ -2t_{r-1}^{1-\beta} \widetilde{I}_R(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ +t_{r-2}^{1-\beta} \widetilde{I}_R(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (3\alpha+10)(p-r) \\ +2\alpha^2 + 9\alpha + 12 \end{array} \right] \\ -(p-r)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (5\alpha+10)(p-r) \\ +6\alpha^2 + 18\alpha + 12 \end{array} \right] \end{array} \right]
 \end{aligned}$$

$$\begin{aligned}
 I_T^{p+1} &= \frac{\alpha(\Delta t)^\alpha}{\Gamma(\alpha+1)} \sum_{r=2}^p t_{r-2}^{1-\beta} \widetilde{I}_T(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \\
 &\times [(p-r+1)^\alpha - (p-r)^\alpha] \\
 &+ \frac{(\Delta t)^\alpha}{\Gamma(\alpha+2)} \sum_{r=2}^p \left[\begin{array}{l} t_{r-1}^{1-\beta} \widetilde{I}_T(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ -t_{r-2}^{1-\beta} \widetilde{I}_T(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha (p-r+3+2\alpha) \\ -(p-r)^\alpha (p-r+3+3\alpha) \end{array} \right] \\
 &+ \frac{(\Delta t)^\alpha}{2\Gamma(\alpha+3)} \sum_{r=2}^p \left[\begin{array}{l} t_r^{1-\beta} \widetilde{I}_T(t_r, S^r, I^r, I_A^r, I_D^r, I_R^r, I_T^r, R^r, D^r, V^r) \\ -2t_{r-1}^{1-\beta} \widetilde{I}_T(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ +t_{r-2}^{1-\beta} \widetilde{I}_T(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (3\alpha+10)(p-r) \\ +2\alpha^2 + 9\alpha + 12 \end{array} \right] \\ -(p-r)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (5\alpha+10)(p-r) \\ +6\alpha^2 + 18\alpha + 12 \end{array} \right] \end{array} \right]
 \end{aligned}$$

$$\begin{aligned}
 R^{p+1} &= \frac{(\Delta t)^\alpha}{\Gamma(\alpha+1)} \sum_{r=2}^p t_{r-2}^{1-\beta} \tilde{R}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \\
 &\times [(p-r+1)^\alpha - (p-r)^\alpha] \\
 &+ \frac{(\Delta t)^\alpha}{\Gamma(\alpha+2)} \sum_{r=2}^p \left[\begin{array}{l} t_{r-1}^{1-\beta} \tilde{R}(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ -t_{r-2}^{1-\beta} \tilde{R}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha (p-r+3+2\alpha) \\ -(p-r)^\alpha (p-r+3+3\alpha) \end{array} \right] \\
 &+ \frac{(\Delta t)^\alpha}{2\Gamma(\alpha+3)} \sum_{r=2}^p \left[\begin{array}{l} t_r^{1-\beta} \tilde{R}(t_r, S^r, I^r, I_A^r, I_D^r, I_R^r, I_T^r, R^r, D^r, V^r) \\ -2t_{r-1}^{1-\beta} \tilde{R}(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ +t_{r-2}^{1-\beta} \tilde{R}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (3\alpha+10)(p-r) \\ +2\alpha^2 + 9\alpha + 12 \end{array} \right] \\ -(p-r)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (5\alpha+10)(p-r) \\ +6\alpha^2 + 18\alpha + 12 \end{array} \right] \end{array} \right]
 \end{aligned}$$

$$\begin{aligned}
 D^{p+1} &= \frac{(\Delta t)^\alpha}{\Gamma(\alpha+1)} \sum_{r=2}^p t_{r-2}^{1-\beta} \tilde{D}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \\
 &\times [(p-r+1)^\alpha - (p-r)^\alpha] \\
 &+ \frac{(\Delta t)^\alpha}{\Gamma(\alpha+2)} \sum_{r=2}^p \left[\begin{array}{l} t_{r-1}^{1-\beta} \tilde{D}(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ -t_{r-2}^{1-\beta} \tilde{D}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha (p-r+3+2\alpha) \\ -(p-r)^\alpha (p-r+3+3\alpha) \end{array} \right] \\
 &+ \frac{(\Delta t)^\alpha}{2\Gamma(\alpha+3)} \sum_{r=2}^p \left[\begin{array}{l} t_r^{1-\beta} \tilde{D}(t_r, S^r, I^r, I_A^r, I_D^r, I_R^r, I_T^r, R^r, D^r, V^r) \\ -2t_{r-1}^{1-\beta} \tilde{D}(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ +t_{r-2}^{1-\beta} \tilde{D}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (3\alpha+10)(p-r) \\ +2\alpha^2 + 9\alpha + 12 \end{array} \right] \\ -(p-r)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (5\alpha+10)(p-r) \\ +6\alpha^2 + 18\alpha + 12 \end{array} \right] \end{array} \right]
 \end{aligned}$$

$$\begin{aligned}
 V^{p+1} &= \frac{(\Delta t)^\alpha}{\Gamma(\alpha+1)} \sum_{r=2}^p t_{r-2}^{1-\beta} \tilde{V}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \\
 &\times [(p-r+1)^\alpha - (p-r)^\alpha] \\
 &+ \frac{(\Delta t)^\alpha}{\Gamma(\alpha+2)} \sum_{r=2}^p \left[\begin{array}{l} t_{r-1}^{1-\beta} \tilde{V}(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ -t_{r-2}^{1-\beta} \tilde{V}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha (p-r+3+2\alpha) \\ -(p-r)^\alpha (p-r+3+3\alpha) \end{array} \right] \\
 &+ \frac{(\Delta t)^\alpha}{2\Gamma(\alpha+3)} \sum_{r=2}^p \left[\begin{array}{l} t_r^{1-\beta} \tilde{V}(t_r, S^r, I^r, I_A^r, I_D^r, I_R^r, I_T^r, R^r, D^r, V^r) \\ -2t_{r-1}^{1-\beta} \tilde{V}(t_{r-1}, S^{r-1}, I^{r-1}, I_A^{r-1}, I_D^{r-1}, I_R^{r-1}, I_T^{r-1}, R^{r-1}, D^{r-1}, V^{r-1}) \\ +t_{r-2}^{1-\beta} \tilde{V}(t_{r-2}, S^{r-2}, I^{r-2}, I_A^{r-2}, I_D^{r-2}, I_R^{r-2}, I_T^{r-2}, R^{r-2}, D^{r-2}, V^{r-2}) \end{array} \right] \\
 &\times \left[\begin{array}{l} (p-r+1)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (3\alpha+10)(p-r) \\ +2\alpha^2 + 9\alpha + 12 \end{array} \right] \\ -(p-r)^\alpha \left[\begin{array}{l} 2(p-r)^2 + (5\alpha+10)(p-r) \\ +6\alpha^2 + 18\alpha + 12 \end{array} \right] \end{array} \right]
 \end{aligned}$$

Now, we handle the following model

$$\begin{aligned}
 {}_0^{FFM}D_t^{\alpha,\beta}S &= \Lambda - (\alpha(x) + \gamma_1 + \mu_1)S \\
 {}_0^{FFM}D_t^{\alpha,\beta}I &= \alpha(x)S - (\varepsilon + \xi + \lambda + \mu_1)I \\
 {}_0^{FFM}D_t^{\alpha,\beta}I_A &= \xi I - (\theta + \mu + \chi + \mu_1)I_A \\
 {}_0^{FFM}D_t^{\alpha,\beta}I_D &= \varepsilon I - (\eta + \varphi + \mu_1)I_D \\
 {}_0^{FFM}D_t^{\alpha,\beta}I_R &= \eta I_D + \theta I_A - (v + \xi + \mu_1)I_R \\
 {}_0^{FFM}D_t^{\alpha,\beta}I_T &= \mu I_A + v I_R - (\sigma + \tau + \mu_1)I_T \\
 {}_0^{FFM}D_t^{\alpha,\beta}R &= \lambda I + \varphi I_D + \chi I_A + \xi I_R + \sigma I_T - (\Phi + \mu_1)R \\
 {}_0^{FFM}D_t^{\alpha,\beta}D &= \tau I_T - \mu_1 D \\
 {}_0^{FFM}D_t^{\alpha,\beta}V &= \gamma_1 S + \Phi R - \mu_1 V - \tau I_T
 \end{aligned} \tag{5.32}$$

where initial conditions are

$$\begin{aligned}
 S(0) &= 57780000, I(0) = 1, I_A(0) = 1, I_D(0) = 1, I_R(0) = 1, \\
 I_T(0) &= 1, R(0) = 0.1, D(0) = 0.2, V(0) = 0.1.
 \end{aligned} \tag{5.33}$$

Also the parameters are chosen as

$$\begin{aligned}
 \Lambda &= 57000000, k = 3, p = 0.5, \eta = 0.12, \chi = 0.015, v = 0.027, x = 0.4, \theta = 0.301, \\
 \gamma &= 0.09, \beta = 0.013, \gamma_1 = 0.4, \mu_1 = 0.3, \varepsilon = 0.161, \xi = 0.015, \sigma = 0.015, \\
 \tau &= 0.0199, \Phi = 0.2, \lambda = 0.0345, \varphi = 0.0345, \delta_1 = 0.01.
 \end{aligned} \tag{5.34}$$

We present numerical simulation for COVID-19 model in Figure 41 and 42.




Figure 41. Numerical visualization for COVID-19 model in South Africa for $\alpha = 0.9, \beta = 0.75$.




Figure 42. Numerical visualization for COVID-19 model in South Africa for $\alpha = 0.7, \beta = 0.75$.

For same model, initial conditions are chosen

$$\begin{aligned} S(0) &= 81000000, I(0) = 1, I_A(0) = 1, I_D(0) = 1, I_R(0) = 1, \\ I_T(0) &= 1, R(0) = 0.1, D(0) = 0.2, V(0) = 0.1. \end{aligned} \quad (5.35)$$

Also the parameters are

$$\begin{aligned} \Lambda &= 57000000, k = 2, p = 0.5, \eta = 0.12, \chi = 0.015, v = 0.027, x = 0.4, \theta = 0.301, \\ \gamma &= 0.09, \beta = 0.013, \gamma_1 = 0.4, \mu_1 = 0.3, \varepsilon = 0.161, \xi = 0.015, \sigma = 0.015, \\ \tau &= 0.0199, \Phi = 0.2, \lambda = 0.0345, \varphi = 0.0345, \delta_1 = 0.01. \end{aligned} \quad (5.36)$$

We present numerical simulation for COVID-19 model in Figure 43 and 44.




Figure 43. Numerical visualization for COVID-19 model in Turkey for $\alpha = 0.9, \beta = 0.75$.




Figure 44. Numerical visualization for COVID-19 model in Turkey for $\alpha = 0.7, \beta = 0.75$.

6 Optimal control for COVID-19 model

Optimal control theory provides us important contributions in controlling COVID-19 outbreak. In this section, we will use 7 control variables as 7 possible control strategies to perform our aim. The control variable u_1 is the partial lockdown of schools, travels, universities, some businesses in Turkey. Also government apply partial lockdown by age of people and sometimes states where spread of virus is high. The control variable u_2 is the vaccination which is applied to susceptible individuals. The control variable u_3 is the information campaign to people that have symptoms but not have been tested. The control variable u_4 is the treatment for the infected individuals. The control variable u_5 is the personal protection which is performed with masks, sanitizer and other stuffs. The control variable u_6 is the self-quarantine exposed people. The control variable u_7 is the isolation of infected people.

We modify our model by adding these control variables such as;

$$\begin{aligned}
 \dot{S} &= \Lambda - \left(\frac{k_1 p e^{-x}}{N} (1 - u_1) (I + w (\beta I_D + \gamma I_A + \delta_1 I_R)) + \gamma_1 + \mu_1 \right) S - u_5 S - u_2 S + u_7 I_R - u_6 S \\
 \dot{I} &= \frac{k_1 p e^{-x}}{N} (1 - u_1) (I + w (\beta I_D + \gamma I_A + \delta_1 I_R)) S - (\varepsilon + \xi + \lambda + \mu_1) I \\
 \dot{I}_A &= \xi I - (\theta + \mu + \chi + \mu_1) I_A - u_3 I_A \\
 \dot{I}_D &= \varepsilon I - (\eta + \varphi + \mu_1) I_D \\
 \dot{I}_R &= \eta I_D + \theta I_A - (v + \xi + \mu_1) I_R - u_7 I_R - u_4 I_R + u_6 S \\
 \dot{I}_T &= \mu I_A + v I_R - (\sigma + \tau + \mu_1) I_T \\
 \dot{R} &= \lambda I + \varphi I_D + \chi I_A + \xi I_R + \sigma I_T - (\Phi + \mu_1) R + u_4 I_R + u_5 S + u_3 I_A \\
 \dot{D} &= \tau I_T - \mu_1 D \\
 \dot{V} &= \gamma_1 S + \Phi R - \mu_1 V - \tau I_T + u_2 S.
 \end{aligned} \tag{6.1}$$

In this paper, we aim to minimize susceptible, infected, critically infected, asymptomatic people and to maximize recovered people while minimizing the costs caused by the partial lockdown, vaccination, information campaign, treatment, personal protection, self-quarantine and isolation. Thus, we construct the cost functional as follows;

$$\min_{(u_1, u_2, u_3, u_4, u_5, u_6, u_7) \in U} J(u_1, u_2, u_3, u_4, u_5, u_6, u_7) = \int_0^T \left(\begin{aligned} &\rho_1 S + \rho_2 I + \rho_3 I_R + \rho_4 I_A - \rho_5 R + \pi_1 u_1^2 + \pi_2 u_2^2 \\ &+ \pi_3 u_3^2 + \pi_4 u_4^2 + \pi_5 u_5^2 + \pi_6 u_6^2 + \pi_7 u_7^2 \end{aligned} \right) dt \tag{6.2}$$

on the set of admissible controls

$$U = \left\{ \begin{aligned} &(u_1, u_2, u_3, u_4, u_5, u_6, u_7) \in L^\infty(0, T) \times L^\infty(0, T) \times L^\infty(0, T) \\ &\quad \times L^\infty(0, T) \times L^\infty(0, T) \times L^\infty(0, T) \times L^\infty(0, T) : \\ &0 \leq u_1(t) \leq \tilde{u}_1, 0 \leq u_2(t) \leq \tilde{u}_2, 0 \leq u_3(t) \leq \tilde{u}_3, \\ &0 \leq u_4(t) \leq \tilde{u}_4, 0 \leq u_5(t) \leq \tilde{u}_5, 0 \leq u_6(t) \leq \tilde{u}_6, 0 \leq u_7(t) \leq \tilde{u}_7 \end{aligned} \right\}. \tag{6.3}$$

The parameters $\rho_1, \rho_2, \rho_3, \rho_4, \rho_5, \pi_1, \pi_2, \pi_3, \pi_4, \pi_5, \pi_6, \pi_7$ represent the weighted parameters.

To show the existence of the optimal control for the problem under consideration, we notice that the set of admissible controls U is, by definition, closed and bounded. It is obvious that there is an admissible pair $(u_1, u_2, u_3, u_4, u_5, u_6, u_7)$ for the problem. Hence, the existence of the optimal control comes as a direct result from the Filippove-Cesari theorem [11,12]. We therefore, have the following result:

We prove that the existence of an optimal control is guaranteed by providing the following conditions.

- The set of admissible controls is convex, bounded and closed.
- The set of controls and corresponding state variables is nonempty.
- The right-hand side of the state ODE system is bounded by a linear function in the state and control variables.

- The convexity of the integrand of cost functional with respect to u on the set U . The Hessian matrix of this functional is given by;

$$H = \begin{bmatrix} 2\pi_1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 2\pi_2 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 2\pi_3 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 2\pi_4 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 2\pi_5 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 2\pi_6 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 2\pi_7 \end{bmatrix}. \quad (6.4)$$

Since the Hessian of of this functional is everywhere positive definite, then the functional $J(u_1, u_2, u_3, u_4, u_5, u_6, u_7)$ is strictly convex.

There exist constants $\pi = \min\{\pi_1, \pi_2, \pi_3, \pi_4, \pi_5, \pi_6, \pi_7\} > 0$ such that the integrand of the cost functional holds

$$\begin{aligned} \tilde{J}(U) &= \pi_1 u_1^2 + \pi_2 u_2^2 + \pi_3 u_3^2 + \pi_4 u_4^2 + \pi_5 u_5^2 + \pi_6 u_6^2 + \pi_7 u_7^2 \\ &\quad + \rho_1 S + \rho_2 I + \rho_3 I_R + \rho_4 I_A - \rho_5 R \\ &\geq \pi_1 u_1^2 + \pi_2 u_2^2 + \pi_3 u_3^2 + \pi_4 u_4^2 + \pi_5 u_5^2 + \pi_6 u_6^2 + \pi_7 u_7^2 \\ &\geq \pi (u_1^2 + u_2^2 + u_3^2 + u_4^2 + u_5^2 + u_6^2 + u_7^2) \end{aligned} \quad (6.5)$$

under the condition $\rho_1 S + \rho_2 I + \rho_3 I_R + \rho_4 I_A > \rho_5 R$. Applying the Pontryagin's maximum principle, we present the first order necessary conditions for optimal solution for the considered optimal control problem. To achieve this, we construct the Hamiltonian H which is given as

$$\begin{aligned} H &= \pi_1 u_1^2 + \pi_2 u_2^2 + \pi_3 u_3^2 + \pi_4 u_4^2 + \pi_5 u_5^2 + \pi_6 u_6^2 + \pi_7 u_7^2 \\ &\quad + \rho_1 S + \rho_2 I + \rho_3 I_R + \rho_4 I_A - \rho_5 R \\ &\quad + \lambda_1 \left(\begin{array}{c} \Lambda - (\delta(x)(1-u_1)(I + w(\beta I_D + \gamma I_A + \delta_1 I_R)) + \gamma_1 + \mu_1) S \\ -u_5 S - u_2 S + u_7 I_R - u_6 S \end{array} \right) \\ &\quad + \lambda_2 (\delta(x)(1-u_1)(I + w(\beta I_D + \gamma I_A + \delta_1 I_R)) S - (\varepsilon + \xi + \lambda + \mu_1) I) \\ &\quad + \lambda_3 (\xi I - (\theta + \mu + \chi + \mu_1) I_A - u_3 I_A) \\ &\quad + \lambda_4 (\varepsilon I - (\eta + \varphi + \mu_1) I_D) \\ &\quad + \lambda_5 (\eta I_D + \theta I_A - (v + \xi + \mu_1) I_R - u_7 I_R - u_4 I_R + u_6 S) \\ &\quad + \lambda_6 (\mu I_A + v I_R - (\sigma + \tau + \mu_1) I_T) \\ &\quad + \lambda_7 (\lambda I + \varphi I_D + \chi I_A + \xi I_R + \sigma I_T - (\Phi + \mu_1) R + u_4 I_R + u_5 S + u_3 I_A) \\ &\quad + \lambda_8 (\tau I_T - \mu_1 D) \\ &\quad + \lambda_9 (\gamma_1 S + \Phi R - \mu_1 V - \tau I_T + u_2 S). \end{aligned} \quad (6.6)$$

Then there exists $\lambda \in \mathbb{R}^9$ such that the first order necessary conditions for the existence of optimal control are given by the equations

$$\begin{aligned}
 \frac{d\lambda_1}{dt} &= -\frac{\partial H}{\partial S} = - \left\{ \begin{aligned} &\rho_1 - \lambda_1 \left(\begin{aligned} &(\delta(x)(1-u_1)(I+w(\beta I_D + \gamma I_A + \delta_1 I_R)) + \gamma_1 + \mu_1) \\ &-u_5 - u_2 - u_6 \end{aligned} \right) \\ &+ \lambda_2 (\delta(x)(1-u_1)(I+w(\beta I_D + \gamma I_A + \delta_1 I_R))) \\ &+ \lambda_5 u_6 + \lambda_7 u_5 + \lambda_9 (u_2 + \gamma_1) \end{aligned} \right\} \\
 \frac{d\lambda_2}{dt} &= -\frac{\partial H}{\partial I} = - \left\{ \begin{aligned} &\rho_2 + \lambda_1 (\delta(x)(1-u_1)S) + \lambda_2 (\delta(x)(1-u_1)S - (\varepsilon + \xi + \lambda + \mu_1)) \\ &+ \lambda_3 \xi + \lambda_4 \varepsilon + \lambda_7 \lambda I \end{aligned} \right\} \\
 \frac{d\lambda_3}{dt} &= -\frac{\partial H}{\partial I_A} = - \left\{ \begin{aligned} &\rho_4 + (\lambda_1 + \lambda_2) \delta(x)(1-u_1)w\gamma S + \lambda_3 (-(\theta + \mu + \chi + \mu_1) - u_3) \\ &+ \lambda_5 \theta + \lambda_6 \mu + \lambda_7 (\chi + u_3) \end{aligned} \right\} \\
 \frac{d\lambda_3}{dt} &= -\frac{\partial H}{\partial I_D} = - \{(\lambda_1 + \lambda_2) \delta(x)(1-u_1)w\beta S - \lambda_4 (\eta + \varphi + \mu_1) + \lambda_5 \eta + \lambda_7 \varphi\} \\
 \frac{d\lambda_3}{dt} &= -\frac{\partial H}{\partial I_R} = - \left\{ \begin{aligned} &\rho_3 + (\lambda_1 + \lambda_2) \delta(x)(1-u_1)w\delta_1 S + \lambda_1 u_7 \\ &- \lambda_5 ((v + \xi + \mu_1) + u_7 + u_4) + \lambda_6 v + \lambda_7 (\xi + u_4) \end{aligned} \right\} \\
 \frac{d\lambda_3}{dt} &= -\frac{\partial H}{\partial I_T} = - \{-\lambda_6 (\sigma + \tau + \mu_1) + \lambda_7 \sigma - \lambda_9 \tau\} \\
 \frac{d\lambda_3}{dt} &= -\frac{\partial H}{\partial R} = - \{-\rho_5 - \lambda_7 (\Phi + \mu_1) + \lambda_9 \Phi\} \\
 \frac{d\lambda_3}{dt} &= -\frac{\partial H}{\partial D} = - \{-\lambda_8 \mu_1\} \\
 \frac{d\lambda_4}{dt} &= -\frac{\partial H}{\partial V} = - \{-\lambda_9 \mu_1\}
 \end{aligned} \tag{6.7}$$

Hence the optimal controls are given as

$$\begin{aligned}
 u_1 &= \frac{I(t)S(t)(\lambda_1 + \lambda_2)}{2\pi_1} \\
 u_2 &= \frac{S(t)(\lambda_1 - \lambda_9)}{2\pi_2} \\
 u_3 &= \frac{I_A(t)(\lambda_3 - \lambda_7)}{2\pi_3} \\
 u_4 &= \frac{I_R(t)(\lambda_5 - \lambda_7)}{2\pi_4} \\
 u_5 &= \frac{S(t)(\lambda_1 - \lambda_7)}{2\pi_5} \\
 u_6 &= \frac{S(t)(\lambda_1 - \lambda_5)}{2\pi_6} \\
 u_7 &= \frac{I_R(t)(\lambda_5 - \lambda_1)}{2\pi_7}
 \end{aligned} \tag{6.8}$$

and optimality conditions are given by

$$\begin{aligned}
 u_1^* &= \min \left\{ \tilde{u}_1, \max \left\{ 0, \frac{I(t) S(t) (\lambda_1 + \lambda_2)}{2\pi_1} \right\} \right\} \\
 u_2^* &= \min \left\{ \tilde{u}_2, \max \left\{ 0, \frac{S(t) (\lambda_1 - \lambda_9)}{2\pi_2} \right\} \right\} \\
 u_3^* &= \min \left\{ \tilde{u}_3, \max \left\{ 0, \frac{I_A(t) (\lambda_3 - \lambda_7)}{2\pi_3} \right\} \right\} \\
 u_4^* &= \min \left\{ \tilde{u}_4, \max \left\{ 0, \frac{I_R(t) (\lambda_5 - \lambda_7)}{2\pi_4} \right\} \right\} \\
 u_5^* &= \min \left\{ \tilde{u}_5, \max \left\{ 0, \frac{S(t) (\lambda_1 - \lambda_7)}{2\pi_5} \right\} \right\} \\
 u_6^* &= \min \left\{ \tilde{u}_6, \max \left\{ 0, \frac{S(t) (\lambda_1 - \lambda_5)}{2\pi_6} \right\} \right\} \\
 u_7^* &= \min \left\{ \tilde{u}_7, \max \left\{ 0, \frac{I_R(t) (\lambda_5 - \lambda_1)}{2\pi_7} \right\} \right\}.
 \end{aligned} \tag{6.9}$$

7 Discussion, recommendations and conclusion

The COVID-19 fatality on mankind prompted them to undertake serious investigations covering various aspects within several fields of science, technology and engineering in the last 4 months. While researchers have obtained some successful results, they are still struggling to get an effective vaccine that could prevent the spread of the deadly COVID-19 among human beings. From December 2019 to 30 April 2020, there are 3441767 confirmed infected cases, 1097858 recovered and 243922 deaths worldwide. Among which, 6336 confirmed cases, 2549 recovered and 123 deaths are recorded in South Africa; and 124 375 confirmed cases, 58259 recovered and 3336 deaths in Turkey. South Africa registered its first confirmed case of COVID-19 before Turkey on 5 March 2020; while Turkey witnessed its first case on 11 March. The unfolding of the spread of COVID-19 in both countries has defeated the general expectations that South Africa would record more infections and deaths comparing to Turkey. As a result, endless scientific questions were asked within different fields of science; which impelled the compilation of this paper to present critical and comprehensive studies with cases studied in South Africa and Turkey in particular. Although both countries have put in place severe measures to protect their citizens; the statistical predictions from the suggested mathematical models and predictions from statistical analysis show two different patterns for both countries. For instance, in Turkey, a high and exponential growth in new infected numbers from 11 March to 11 April 2020 was observed due to late implementation of the lockdown regulations; however, from 12 April to 02 May 2020 this country has observed an exponential decay in the daily numbers of new infected cases. Thus, Turkey curve seems to follow a lognormal distribution, which of course could mean that they are winning the COVID-19 war; or they took control of the situation. As a result, it is possible that Turkey in the next few months could end the spread of COVID-19, if they maintain the energy and adhere to the measures in place to combat this virus. However, if they relax, the prediction from reliability level method indicated that Turkey could see a very rapid exponential growth in numbers of daily deaths and new infections. Furthermore, it is observed that the exponential decay in the daily number of new infected and death cases corresponds to the period of lockdown implementation and the stringent rules put in place by the Turkish government; by which the contravening of the rule is punishable with a monetary fine. On the other hand in South Africa, although the numbers are not as high as those of Turkey, three phases are observed from statistical results. The first phase goes from 5 March 2020 to 27 March 2020, where the country witnessed an exponential growth in numbers of new infected and deaths daily; and it corresponds to a pre-lockdown period. The second phase began on 28 March until 18 April 2020, in which the country observed a slowly increase of new infected and death daily; a period corresponding to lockdown period enforced with the presence of South Africa Defence Force. While the last phase ranges from 19 April to 02 May 2020, in which the country observed an exponential growth in numbers of new infected and deaths per day. This exponential growth is attributed to the relaxation and disobedience of lockdown regulations; probably due to economic breakdown, increasing poverty effects among the larger population and also due to migration from level five to level four on the 1 st of May 2020. Therefore, as the provision of a suitable vaccine to save and protect human beings the wrath and the fatality of

the deadly COVID-19, which breakout in Wuhan, China, December 2019 is delayed; it is clearly evident that the exponential growth in numbers of new infected can only be stopped or halted by enforcing the implementation of social distancing and ensuring that people do frequently wash their hands upon touching any object or even animals; whether infected with COVID-19 or not. Additionally, the wearing of masks should be adopted in public places to avoid the spreading of the virus, in case the social distancing rules is not being kept. It is further paramount that the medical workers in charge of COVID-19 patients are well protected to minimize the contraction of the virus from the patients and passing it on to the general public. In addition to the prohibition of alcohol sales and usage in public places; public smoking should also be prohibited in the effort to combat the spread of COVID-19. Moreover, the statistical analysis results specifically the reliability level prediction and the results obtained from suggested mathematical models; indicated that without social distancing restrictions or clear implementation of lockdown regulations, it will be impossible for countries to control the spread of COVID-19. This implies that the number of new infected and deaths per day would be difficult to contain, resulting in the fight against the virus to get out of hand. These outcomes from reliability level are therefore indicated in blue lines in Figures 18, 19, 20, 29 and 30 those from reliability level. The suggested mathematical models with different differential operators including classical and nonlocal operators in the last 12 Figures as case with lockdown and no-lockdown presented for different fractional orders also confirmed the results obtained from the reliability level. In consideration of all prediction results, it is concluded that South Africa has not yet won the war against COVID-19 and serious outbreak are expected in the near future as the climate season changes to winter. Cold seasons are scientifically proven to be thriving climate for the survival of corona virus. Therefore to avoid this foreseen crisis, social distancing must be a responsibility of each person living within the Republic of South Africa, and the transition from level 5 to level 1 should be implemented very wisely.

References

- [1] Giordano, G., Blanchini, F., Bruno, R. et al. Modelling the COVID-19 epidemic and implementation of population-wide interventions in Italy. *Nat Med* (2020).
- [2] MA Khan, A Atangana, Modeling the dynamics of novel coronavirus (2019-nCov) with fractional derivative, *Alexandria Engineering Journal*, 2020.
- [3] KM Owolabi, A Atangana, Mathematical analysis and computational experiments for an epidemic system with nonlocal and nonsingular derivative, *Chaos, Solitons & Fractals* 126, 2019, 41-49.
- [4] Tuite AR., Fisman DN., Greer LA., Mathematical modelling of COVID-19 transmission and mitigation strategies in the population of Ontario, Canada, *CMAJ*, 2020.
- [5] Anastassopoulou C., Russo L., Tsakris A., Siettos C., Data-Based Analysis, Modelling and Forecasting of the COVID-19 outbreak, 2020, medRxiv.
- [6] Kucharski AJ., Russell TW., Diamond C., Liu Y., Edmunds J., Funk S., Eggo RM., Early dynamics of transmission and control of COVID-19: a mathematical modelling study, *Lancet Infect Dis.* 2020.
- [7] COVID-19 pandemic in South Africa, https://en.wikipedia.org/wiki/COVID-19_pandemic_in_South_Africa, 2020.
- [8] LaSalle JP., *The stability of dynamical systems*, SIAM Press, 1976.
- [9] Driessche P., Watmough J., (2002), Reproduction numbers and sub-threshold endemic equilibria for compartmental models of disease transmission, *Mathematical Biosciences*, 180 (1), 29-48.
- [10] Atangana A, Araz Sİ., New numerical method for ordinary differential equations: Newton polynomial, *Journal of Computational and Applied Mathematics* 372, 2020, 112622.
- [11] S. Nababan ; A flippov-type lemma for functions involving delays and its application to time delayed optimal control problems, *Optim. Theory appl.* 27 3 (1979), 357-376.
- [12] Alvarez FE., Argente D., Lippi F., A Simple Planning Problem for COVID-19 Lockdown, National Bureau of Economic Research, 2020.