

1 **Post-Stroke Physical Activity Improves Middle Cerebral Artery Dynamic Response**

2 Sandra A. Billinger, PT, PhD^{1,2,3,4}; Alicen A. Whitaker, DPT¹; Allegra Morton, BS¹; Carolyn S.
3 Kaufman, BS^{1,4}; Sophy J Perdomo, PhD¹; Jaimie L. Ward, MS¹; Sarah M. Eickmeyer, MD²;
4 Stephen X. Bai, MD²; Luke Ledbetter, MD⁵; Michael G. Abraham, MD^{3,6}

5
6 ¹Department of Physical Therapy and Rehabilitation Science, University of Kansas Medical
7 Center, Kansas City, KS

8 ²Department of Physical Medicine and Rehabilitation, University of Kansas Medical Center,
9 Kansas City, KS

10 ³Department of Neurology, University of Kansas Medical Center, Kansas City, KS

11 ⁴Department of Integrative and Molecular Physiology, University of Kansas Medical Center,
12 Kansas City, KS

13 ⁵Department of Diagnostic Radiology, University of Kansas Medical Center, Kansas City, KS

14 ⁶Department of Interventional Radiology, University of Kansas Medical Center, Kansas City, KS

15

16 **Corresponding Author:**

17 Sandra A. Billinger, PT, PhD

18 Address: 3901 Rainbow Blvd, Mail Stop 2002, Kansas City, KS 66160

19 Phone number: 913-945-6685

20 Fax: 913-588-9428

21 Email: sbillinger@kumc.edu

22 **Abstract**

23 **Background and Purpose:** The primary aim was to characterize the middle cerebral artery
24 blood velocity (MCAv) dynamic response to an acute bout of exercise at 3- and 6-months post
25 stroke. As a secondary objective, we grouped individuals according to the MCAv dynamic
26 response to the exercise bout as responder or non-responder. We tested whether physical activity,
27 aerobic fitness and exercise mean arterial blood pressure (MAP) differed between groups.

28 **Methods:** Transcranial Doppler ultrasound measured MCAv during a 90-second baseline (BL)
29 followed by a 6-minute moderate intensity exercise bout. Heart rate (HR), MAP and end tidal
30 CO₂ (P_{ET}CO₂) were additional variables of interest. The MCAv dynamic response variables
31 included: BL, time delay (TD), amplitude and time constant (τ).

32 **Results:** Individuals enrolled in the study at 3 months post-stroke and the follow up visit
33 commenced at 6 months post-stroke. Linear mixed model revealed no significant differences in
34 our selected outcomes across between 3- and 6-months post-stroke. Individuals characterized as
35 responders demonstrated a faster TD, higher amplitude, reported higher levels of physical
36 activity and aerobic fitness when compared to the non-responders. No between group
37 differences were identified for BL, τ or exercise MAP. In the non-responders, we observed an
38 immediate rise in MCAv following exercise onset followed by an immediate decline to near BL
39 values while the responders showed an exponential rise until steady state was reached.

40 **Conclusions:** The MCAv dynamic response profile has the potential to provide valuable
41 information during an acute exercise bout following stroke. Individuals with a greater MCAv
42 response to the exercise stimulus reported regular participation in exercise than those who
43 reported being sedentary.

44 **Introduction**

45 Impaired cerebrovascular hemodynamic response during exercise has been reported in
46 individuals within one year post-stroke.¹ We previously demonstrated individuals with middle
47 cerebral artery (MCA) stroke showed no significant differences in resting MCA blood velocity
48 (MCAv) when compared to age- and sex-matched peers.² However, the MCAv dynamic
49 response during exercise was significantly lower for those post-stroke. Despite the impaired
50 cerebrovascular response to exercise, a six-month aerobic exercise treadmill training program in
51 chronic stroke survivors conferred benefit on cerebrovascular reserve for those randomized to the
52 intervention when compared to controls.³ This evidence, albeit limited, suggests potential
53 impairment in cerebrovascular control mechanisms following stroke.

54

55 The primary objective of this study was to characterize the MCAv dynamic response to a
56 moderate intensity acute exercise bout at 3- and 6-months post stroke. We hypothesized that the
57 dynamic response profile would improve from 3- to 6-months post-stroke. In our previous work
58 comparing young and older adults, we excluded one older female participant from analysis as her
59 data did not fit the exponential model.⁴ Rather than exclude participants with stroke whose data
60 may not fit the exponential model, we decided further scientific inquiry was warranted. As a
61 secondary objective, we divided participants into responders and non-responders based on the
62 model fit. We compared physical activity levels and hemodynamic exercise response. We
63 hypothesized that individuals in the responder group would report greater physical activity,
64 higher estimated $\dot{V}\dot{O}_2$ maximum and higher mean arterial blood pressure (MAP) during
65 exercise when compared to non-responders.

66

67 **Methods**

68 *Participants*

69 We identified individuals post-stroke during the acute or inpatient rehabilitation stay at the
70 University of Kansas Health System. Inclusion criteria were: 1) unilateral ischemic stroke, 2)
71 stenosis in the common and internal carotid artery was <70%.^{3,5} A radiologist (LL) made this
72 interpretation using ultrasound imaging or angiogram in the electronic medical record, 3) 35-95
73 years of age, 4) physician approval for exercise participation, 5) able to walk >10 meters without
74 physical assistance and 6) able to travel to the university for the experimental protocol.
75 Exclusion criteria were: 1) unable to consent, 2) inability to perform the exercise movements on
76 the seated recumbent stepper (T5XR NuStep, Inc. Ann Arbor, MI), 3) diagnosis of other
77 neurologic disease, and 4) use of supplemental oxygen.

78

79 *Experimental Procedure*

80 Study visits occurred in a laboratory setting at 3- and 6-months post-stroke. Participants were
81 asked to abstain from caffeine for a minimum of six hours, a meal for two hours, and vigorous
82 exercise at least twelve hours prior to the study visit.^{4,6} The University of Kansas Medical Center
83 Human Subjects Committee approved all experimental procedures. Institutionally approved
84 written informed consent was obtained from each individual prior to study participation.

85

86 *Familiarization*

87 The laboratory room in which the experimental protocol took place was dimly lit, quiet and
88 maintained a constant temperature between 22-24°C.⁶ All external stimuli were kept to a minimum.

89 Participants completed the familiarization session as previously described.^{2,4,6} Following
90 familiarization, we obtained height, weight and completed remaining study documents.

91

92 *Protocol Set Up*

93 Individuals were instrumented with the following equipment: 1) transcranial Doppler ultrasound
94 (TCD) (Multigon Industries Inc. Yonkers, NY) and 2-MHz TCD probes for acquisition of
95 MCAv ($\text{cm}\cdot\text{sec}^{-1}$). The TCD sonographer was blinded to the side of stroke; 2) A 5- lead
96 electrocardiogram (ECG) (Cardiocard, Nasiff Associates, Central Square, NY) recorded HR; 3)
97 beat to beat MAP was obtained from the left middle finger (Finometer, Finapres Medical
98 Systems, Amsterdam, The Netherlands); and 4) nasal cannula and capnograph (BCI Capnocheck
99 Sleep 9004 Smiths Medical, Dublin, OH) assessed end-tidal carbon dioxide (P_{ETCO_2}). We
100 monitored P_{ETCO_2} to ensure participants did not hyperventilate. Raw data acquisition occurred
101 through an analog-to-digital unit (NI-USB-6212, National Instruments) and custom written
102 software operating in MATLAB (v2014a, The Mathworks Inc. Natick, MA).

103

104 *Exercise Protocol*

105 We used moderate intensity exercise (45% to 55% of the participant's heart rate reserve) for the
106 acute exercise bout. We determined the HR range from one of two equations: 1) age-predicted
107 (220- age) HR maximum (HR_{max}) or 2) $164 - 0.72 \times \text{age}$ for participants using beta-blocker
108 medication.⁷

109

110 The baseline (BL) recording lasted 90 seconds followed by 6 minutes of moderate intensity
111 exercise. Following the 6-minute exercise bout, the participant engaged in an active cool down

112 for 2 minutes followed by rest until HR, MCA_v and MAP returned to BL values. Participants
113 performed a second exercise bout and data points were averaged to optimize signal to noise
114 ratio.⁶ All variables were sampled at 500 Hz and then interpolated to 2.0 Hz. Three-second
115 averages were calculated and then smoothed using a 9-second sliding window average. We used
116 R version 3.2.4 (R Core team, Vienna, Austria)⁸ with the ‘nls’ function package to model the
117 response. To ensure data quality, data with R-to-R intervals greater than 5 Hz or changes in peak
118 blood velocity greater than 10 cm/s in a single cardiac cycle were considered artifact and
119 censored. If acquisitions had more than 15% of data points censored, they were discarded.

120

121 *MCA_v and Dynamic Response Profile*

122 Kinetics were modeled for MCA_v over the entire exercise bout with a mono-exponential model:

$$123 \quad \text{MCA}_v(t) = \text{BL} * (t \leq \text{TD}) + (\text{BL} + \text{Amp}(1 - e^{-(t - \text{TD})/\tau})) * (t > \text{TD})$$

124 where MCA_v(*t*) is the MCA_v at any point in time, BL is baseline before exercise onset, Amp is
125 the peak amplitude of the response above resting BL, TD is the time delay preceding the
126 exponential increase in MCA_v, and τ is the time constant or time-to-63% of the peak amplitude.

127

128 *Six-Minute Walk Test (6MWT)*

129 Participants were given a 20-minute rest following the exercise session. HR and blood pressure
130 were taken to ensure the participant sufficiently recovered from the exercise bout. The 6MWT
131 was performed in a hallway with no distractions or foot traffic according to guidelines outlined
132 by the American Thoracic Society.⁹

133

134 *Statistical Analysis*

135 Data are presented as mean \pm standard deviation. To account for the missing data across study
136 visits, we used a linear mixed model to test for differences between 3- and 6-months post stroke.
137 To test for between group differences (responders vs non-responders), we performed one-way
138 ANOVA or Mann Whitney U as appropriate following visual inspection of probability plots and
139 Shapiro-Wilk tests. Fisher exact tests were used for categorical data. Differences were
140 considered significant when $p < 0.05$.

141

142 **Results**

143 *Participant Characteristics*

144 We consented 27 participants into the study. One person was consented twice as this individual
145 experienced 2 separate stroke events and has been published as a case report.¹⁰ Data from the
146 second stroke was not included. Participant demographics (n=26) at enrollment are presented in
147 Table 1. No serious adverse events occurred during any study visit.

148

149 Eighteen participants (72.2% male) returned for the 6-month visit. Participants were 65.7 ± 14.1
150 years of age and 188.4 ± 14.1 days post-stroke. 83.3% identified as Caucasian, 0% Hispanic,
151 22.2% African American and 5.6% Native American. Multiple race categories were selected.
152 Reasons for loss to follow up included: unable to contact (n = 2); participant's son had a stroke
153 (n = 1); hospitalized (n = 1); visit scheduled but did not attend study visit (n = 2); no
154 transportation (n = 1) and participant died (n = 1).

155

156 *MCAv Dynamic Response Across Time*

157 3-Month Visit. Of the 26 participants, four individuals were excluded from analysis due to: a
158 valid MCAv signal was not acquired, n = 3 and poor MCAv signal acquisition during exercise,
159 n=1. Data are presented for 22 participants.

160
161 6-Month Visit. Of the 18 individuals who returned, we excluded 3 datasets from analysis.
162 Reasons for exclusion included an invalid MCAv signal, n = 2 and poor MCAv signal
163 acquisition during exercise, n=1. Statistical analyses were conducted on 15 datasets.

164
165 The MCAv BL and kinetic measures during moderate intensity exercise showed no differences
166 over time for the stroke affected and non-stroke affected MCAv (Table 2). We found no
167 differences between race (p = 0.09) or sex (p = 0.21) for all MCAv data and hemodynamic data
168 (blood pressure, heart rate and P_{ET}CO₂) at rest and during exercise at either 3- or 6-month visits.

169

170 *Responders and Non-Responders (3 Months Post-stroke)*

171 MCAv Dynamic Response

172 Table 3 presents demographics for each group. We observed 12 participants whose MCAv
173 increased in a close-to-exponential pattern being well-fit by a delay + exponential function
174 (responders). The remaining 10 individuals (non-responders) demonstrated poor model fit with
175 visual inspection (n = 5), non-exponential response (n = 4) or MCAv was below BL (n = 1) after
176 exercise onset on the stroke affected side. On the non-stroke affected side, MCAv decreased
177 below BL after exercise onset for the same individual (n = 1) and we observed a non-exponential
178 response in the other participants (n = 9). No between group differences were reported for age,
179 body mass index, 6MWT or the MoCA. The responders were more likely to report taking a statin

180 medication, had greater physical activity levels and had higher estimated $\dot{V}\dot{O}_2$ maximum values
181 ($p < 0.05$).

182
183 Figure 1 shows the BL and MCAv response after the onset of moderate intensity exercise for
184 responders and non-responders. Data from the MCAv kinetics analysis are presented in Table 4.
185 During seated rest prior to exercise onset, BL data for the stroke affected and non-stroke affected
186 MCAv were not significantly different between groups. After exercise onset, MCAv amplitude
187 on the stroke affected side was significantly reduced in the non-responders. This finding is
188 expected based on group assignment (responder/non-responder). For the 5 participants with poor
189 model fit, we observed a significantly slower TD response for the non-responders on the stroke
190 affected side while the τ was not different. Only one person in the non-responder group had data
191 for analysis that fit the model, although a negative amplitude (below BL). The remaining
192 participants ($n = 9$) showed a non-exponential response.

193
194 Baseline and Exercise Response in MAP, $P_{ET}CO_2$ and HR

195 The baseline and exercise-induced responses (Δ) in MAP, $P_{ET}CO_2$ and HR are shown in Table 4.
196 The BL and steady state exercise for MAP, $P_{ET}CO_2$ and HR were not significantly different. We
197 found between group differences for ΔMAP and ΔHR ($p < 0.025$) but not $\Delta P_{ET}CO_2$ ($p = 0.22$).

198
199 *Responders and Non-Responders (6 Months Post-stroke)*

200 All individuals characterized as a responder at the 3-month visit remained at the 6-month visit.
201 Three individuals initially characterized as a non-responder met the criteria for responder. Of
202 these three individuals, two increased their physical activity score from a 2 (low intensity

203 walking for 10 minutes or less) to a 5 (vigorous activity). The other individual remained in the
204 category of 2 for both visits. Three individuals changed medications related to blood pressure,
205 anticoagulation or antiplatelet therapy that may have influenced MCAv response or other
206 hemodynamic measures at rest or during exercise. In the responder group, one person
207 discontinued clopidogrel. In the non-responder group, lisinopril was added to the medication list
208 while another person reported discontinued use of lisinopril, furosemide and clopidogrel.

209
210 There were no group differences between responders (n = 10) and non-responders (n = 5) for age
211 (p = 0.06), time post-stroke (6-month visit, p = 0.47), BMI (p = 0.58), MoCA score (p = 0.54),
212 mRS (p = 0.40) or self-reported physical activity score (p = 0.10). We report significant
213 differences for the non-exercise estimate for $V\dot{V}O_2$ maximum (p < 0.01) and six-minute walk test
214 (p = 0.02).

215

216 MCAv Dynamic Response

217 Responders worked at a higher workload (p = 0.04) with no significant differences in the rate of
218 perceived exertion (p = 0.96). BL MCAv was not different between groups for the stroke
219 affected (p = 0.58) and non-stroke affected MCA (p = 0.87). For the stroke affected MCA, the
220 non-responders showed a significantly slower TD (p = 0.04) and lower amplitude (p = 0.04)
221 while τ was not different (p = 0.93). For the non-stroke affected MCA, no between group
222 significant differences were found for any MCAv dynamic response variables.

223

224 Baseline and Exercise Response in MAP, $P_{ET}CO_2$ and HR

225 No between group differences were found for resting and exercise hemodynamic responses,
226 Δ MAP, Δ HR or Δ P_{ET}CO₂.

227

228 **Discussion**

229 Aerobic exercise, identical to the intensity used in this study, is recommended for overall health
230 post-stroke.¹¹ This study provides novel and relevant information regarding cerebrovascular
231 response to moderate intensity exercise for individuals at 3- and 6-months post-stroke. The main
232 primary findings of the current investigation are: 1) MCAv kinetic response was stable from 3 to
233 6 months post-stroke for bilateral MCA and 2) MCAv kinetic response (amplitude and TD)
234 significantly differed for those with higher estimated $\dot{V}\dot{O}_2$ maximum. This finding was
235 consistent for the stroke affected MCA at 3- and 6-months.

236

237 *MCAv Dynamic Response Across Time*

238 The primary objective of this study was to characterize the MCAv response at 3- and 6- months
239 post-stroke. To our knowledge, this longitudinal cerebrovascular assessment at rest and in
240 response to exercise had not been previously conducted. We hypothesized that the MCAv
241 dynamic response profile would improve from 3 to 6 months post-stroke. This rationale was
242 developed from published work highlighting improved longitudinal changes in aerobic fitness at
243 1, 3 and 6 months post-stroke¹² and aerobic fitness positively influences peripheral vascular
244 health measures in individuals post-stroke.¹³ Our hypothesis that MCAv dynamic response
245 would improve with time was not supported by our findings. We have reported high
246 reproducibility within older adults for resting and exercise MCAv¹⁴ and bilateral reproducibility

247 for the MCAv kinetic measures.⁶ Therefore, measurement error from our sonographers (SP, CK)
248 was unlikely.

249

250 *MCAv Dynamic and Hemodynamic Responses 3 Months Post-Stroke*

251 During exercise onset, neural activation increases the metabolic requirements of the brain above
252 resting values in order to maintain normal function¹⁵⁻¹⁸ and this demand is met with increased
253 cerebral blood flow. Our laboratory^{6,19} and others^{17,18,20} have shown that moderate intensity
254 continuous exercise produces the greatest increase in MCAv compared to low and high intensity
255 (above anaerobic threshold) continuous exercise.

256

257 Ten of the twenty-one individuals enrolled demonstrated a non-exponential MCAv response
258 (non-responders) following exercise onset. This could be indicative of slowed vasomotor control
259 within the cerebrovasculature as both sides showed slower MCAv kinetics or non-exponential
260 MCAv increase.⁴ When considering potential factors, the data in Table 3 show no between group
261 differences for age, sex, race or ethnicity, time of study visit, or the calculated lesion size (SXB)
262 from the electronic medical record.^{21,22} We found responders had greater statin use, self-reported
263 physical activity levels and estimated $V\dot{V}O_2$ maximum values. One previous study grouped
264 healthy young adults based on self-reported physical activity levels into low and high fitness.²³
265 Those in the high fitness group demonstrated a greater change in MCAv in response to
266 rebreathing-induced hypercapnia while no resting MCAv differences were found. This finding,
267 combined with the present study, underscores the value in considering dynamic challenges such
268 as exercise. Further, others have reported that physically active individuals with higher $V\dot{V}O_2$
269 maximum values exhibit better cerebral hemodynamics than sedentary individuals.^{24,25} Although

270 the evidence reporting the dynamic cerebrovascular response to exercise in stroke is
271 limited,^{1,2,6,10} the present study suggests that physical activity and exercise post-stroke may
272 confer some benefit to overall cerebrovascular health. We acknowledge individuals in the
273 responder group also reported a significantly higher use of statin medications. One study showed
274 3 months of statin treatment after stroke improved cerebrovascular reactivity compared to the
275 control group while no differences in resting cerebral blood flow were reported.²⁶ Our planned
276 future work aims to test whether aerobic exercise interventions can increase the MCAv kinetic
277 response profile and other cerebrovascular outcomes after stroke.

278
279 The BL values and steady state hemodynamic responses during the exercise bout were not
280 significantly different between groups. We report between group differences for Δ MAP and
281 Δ HR. We acknowledge that these differences may be largely driven by the significantly higher
282 workload in the responder group, which likely influenced the MCAv amplitude response. One
283 individual (responder) who exercised at 110 watts largely influenced the difference in work rate.
284 He was an avid cyclist prior to his stroke and interestingly, had the largest stroke lesion at
285 admission to the hospital. We report no between group differences for BL, steady state or
286 Δ P_{ET}CO₂. We are confident that the lower MCAv amplitude in the non-responders is not the
287 result of hyperventilation. We closely monitor P_{ET}CO₂ during the exercise bout to ensure our
288 participants stay within the prescribed HR range without hyperventilation.⁶

289
290 The literature is well-established that MAP increases with age²⁷ and after stroke,¹¹ which
291 provides unique opportunities to study exercise-induced changes in MAP and HR. The
292 cerebrovasculature and the associated neuroprotective mechanisms have the ability to

293 autoregulate and buffer the sudden acute increases in MAP associated across various exercise
294 intensities.²⁸ As observed in Figure 1, the non-responders show a rapid rise in the MCAv
295 response to exercise followed by an immediate decline with steady state values near BL. One
296 proposed hypothesis for this unexpected response may be an impairment in the cerebrovascular
297 control mechanisms following stroke.⁶ Despite the increased neural activation with exercise and
298 demand for energy, MCAv did not increase. Another plausible explanation may be an
299 impairment in the feed-forward signals coming from the motor cortex to increase MCAv
300 concomitantly with neuromuscular innervation.^{4,29} Our data extend current knowledge regarding
301 cerebrovascular response to exercise and is timely given the increased interest and concern for
302 the role of high-intensity interval training on cerebrovascular response, especially for those with
303 neurological conditions such as stroke.^{28,30,31} The authors of these review papers^{28,31} and a recent
304 study describing the cerebrovascular response to a 30-second bout of high intensity exercise³⁰
305 suggest that caution should be taken when implementing this type of protocol, as the evidence
306 for safety with respect to the brain and its delicate cerebrovasculature has not been tested. A
307 recently published clinical practice guideline for physical therapy suggests higher locomotor
308 intensities benefit walking speed and endurance in chronic stroke.³² The current knowledge of
309 cerebrovascular control during an acute exercise bout, especially following stroke, is still in its
310 infancy and warrants further investigation.

311

312 *MCAv Dynamic and Hemodynamic Responses 6 Months Post-Stroke*

313 In the chronic phase of stroke recovery, the responders continued to exercise at higher
314 workloads, which may have influenced cerebrovascular response during the exercise bout. At
315 this study visit, only differences in TD and amplitude on the stroke affected vessel remained. All

316 other MCAv kinetic variables and hemodynamic responses were not significant. The reduced
317 sample size warrants caution when interpreting the findings. A noteworthy observation is two of
318 the three newly characterized responders had increased their self-reported physical activity.
319 Whether the increase in physical activity was the primary driver of the higher MCAv amplitude
320 responses remains an important query for future work.

321
322 Several limitations in the study design should be considered. First, as with all studies using
323 transcranial Doppler ultrasound, the assumption of constant MCA diameter is important in order
324 for the MCA_v to be used as a direct proxy for cerebral blood flow.⁶ Second, maximal exercise
325 testing was not performed to determine maximal heart rate to more accurately prescribe the
326 moderate intensity exercise session. However, as evidenced by the increase in P_{ET}CO₂ during
327 exercise, it is unlikely that individuals exercised above anaerobic levels. Third, we observed an
328 increase in BL MCAv prior to exercise onset. This could be an anticipatory response or
329 nervousness to our verbal instructions 15 seconds prior to exercise onset.

330

331 **Conclusion**

332 The cerebrovascular response to acute exercise remained stable between 3- and 6-months post-
333 stroke. Physical activity participation may positively influence the MCAv dynamic response
334 during exercise. Further exploration into the cerebrovascular control mechanisms or other stroke-
335 specific characteristics that may result in a diminished response (non-responders) to exercise and
336 various intensities is needed. The MCAv kinetic response may better inform future exercise trials
337 designed to benefit overall brain health.

338

339 **Acknowledgments**

340 We would like to thank Yumei Liu, MD, PhD and Emily Nardo for their assistance with data
341 collection and Madison Henry, SPT for her assistance with data organization.

342

343 **Sources of funding**

344 Dr. Billinger was supported by the Eunice Kennedy Shriver National Institute of Child Health
345 and Human Development (K01HD067318) and the Wohlgemuth Faculty Scholar Award. Dr.
346 Whitaker and Ms. Morton were supported in part by T32HD057850 from the Eunice Kennedy
347 Shriver National Institute of Child Health and Human Development. Ms. Morton received
348 support from a Student Scholarship in Cerebrovascular Disease and Stroke (American Heart
349 Association). Dr. Perdomo received partial support from the University of Kansas Alzheimer's
350 Disease Center (P30AG035982), the KU Medical Center Biomedical Research Training Program
351 and from the National Institute on Aging Diversity Supplement to R01 AG058162. REDCap at
352 University of Kansas Medical Center is supported by CTSA Award (UL1TR000001) from
353 NCCR and NCATS. This work was supported from NCATS awarded to the University of
354 Kansas for Frontiers: Clinical and Translational Science Institute (UL1TR002366) The contents
355 are solely the responsibility of the authors and do not necessarily represent the official views of
356 the NIH or NCATS.

357

358 **Disclosures**

359 SAB reports a patent pending (18KU028M-02). MA reports consulting for Stryker
360 Neurovascular and Penumbra Inc. outside the submitted work. AW, AM, CK, SP, JW, SE, SB,
361 and LL report no conflicts of interest.

362 **References**

- 363 1. Robertson AD, Atwi S, Kostoglou K, et al. Cerebrovascular Pulsatility During Rest and
364 Exercise Reflects Hemodynamic Impairment in Stroke and Cerebral Small Vessel
365 Disease. *Ultrasound in medicine & biology*. 2019;45(12):3116-3127.
- 366 2. Kempf KS, Whitaker AA, Lui Y, et al. The Effect of Stroke on Middle Cerebral Artery
367 Blood Flow Velocity Dynamics During Exercise. *J Neurol Phys Ther*. 2019;43(4):212-
368 219.
- 369 3. Ivey FM, Ryan AS, Hafer-Macko CE, Macko RF. Improved cerebral vasomotor
370 reactivity after exercise training in hemiparetic stroke survivors. *Stroke*.
371 2011;42(7):1994-2000.
- 372 4. Ward JL, Craig JC, Liu Y, et al. Effect of Healthy Aging and Sex on Middle Cerebral
373 Artery Blood Velocity Dynamics During Moderate Intensity Exercise. *American journal*
374 *of physiology Heart and circulatory physiology*. 2018.
- 375 5. Guo ZN, Liu J, Xing Y, et al. Dynamic cerebral autoregulation is heterogeneous in
376 different subtypes of acute ischemic stroke. *PLoS One*. 2014;9(3):e93213.
- 377 6. Billinger SA, Craig JC, Kwapiszeski SJ, Sisante J-FV, Vidoni ED, Maletsky R.
378 Dynamics of middle cerebral artery blood flow velocity during moderate intensity
379 exercise. In. *Journal of Applied Physiology*2017.
- 380 7. Brawner CA, Ehrman JK, Schairer JR, Cao JJ, Keteyian SJ. Predicting maximum heart
381 rate among patients with coronary heart disease receiving beta-adrenergic blockade
382 therapy. *Am Heart J*. 2004;148(5):910-914.
- 383 8. *A language and environment for statistical computing*. [computer program]. Vienna,
384 Austria: R Foundation for Statistical Computing; 2013.

- 385 9. Crapo RO, Casaburi R, Coates AL, et al. ATS Statement: Guidelines for Six-Minute
386 Walk Test. *American Journal of Respiratory and Critical Care Medicine*. 2002;166:111 -
387 117.
- 388 10. Kaufman CS, Bai SX, Ward JL, Eickmeyer SM, Billinger SA. Middle cerebral artery
389 velocity dynamic response profile during exercise is attenuated following multiple
390 ischemic strokes: a case report. *Physiol Rep*. 2019;7(21):e14268.
- 391 11. Billinger SA, Arena R, Bernhardt J, et al. Physical Activity and Exercise
392 Recommendations for Stroke Survivors: A Statement for Healthcare Professionals From
393 the American Heart Association/American Stroke Association. *Stroke*. 2014;45(8):2532-
394 2553.
- 395 12. Mackay-Lyons MJ, Makrides L. Longitudinal changes in exercise capacity after stroke.
396 *Arch Phys Med Rehabil*. 2004;85(10):1608-1612.
- 397 13. Billinger SA, Mattlage AE, Ashenden AL, Lentz AA, Harter G, Rippee MA. Aerobic
398 exercise in subacute stroke improves cardiovascular health and physical performance. *J*
399 *Neurol Phys Ther*. 2012;36(4):159-165.
- 400 14. Perdomo SJ, Ward J, Liu Y, et al. Cardiovascular Disease Risk Is Associated With
401 Middle Cerebral Artery Blood Flow Velocity in Older Adults. *Cardiopulmonary Physical*
402 *Therapy Journal*. 2020;31(2):38-46.
- 403 15. Gonzalez-Alonso J, Dalsgaard MK, Osada T, et al. Brain and central haemodynamics and
404 oxygenation during maximal exercise in humans. *J Physiol*. 2004;557(Pt 1):331-342.
- 405 16. Jorgensen LG, Perko M, Hanel B, Schroeder TV, Secher NH. Middle cerebral artery flow
406 velocity and blood flow during exercise and muscle ischemia in humans. *Journal of*
407 *applied physiology*. 1992;72(3):1123-1132.

- 408 17. Linkis P, Jorgensen LG, Olesen HL, Madsen PL, Lassen NA, Secher NH. Dynamic
409 exercise enhances regional cerebral artery mean flow velocity. *Journal of applied*
410 *physiology*. 1995;78(1):12-16.
- 411 18. Moraine JJ, Lamotte M, Berre J, Niset G, Leduc A, Naeije R. Relationship of middle
412 cerebral artery blood flow velocity to intensity during dynamic exercise in normal
413 subjects. *European journal of applied physiology and occupational physiology*.
414 1993;67(1):35-38.
- 415 19. Witte E, Liu Y, Ward JL, et al. Exercise intensity and middle cerebral artery dynamics in
416 humans. *Respir Physiol Neurobiol*. 2019;262:32-39.
- 417 20. Ide K, Horn A, Secher NH. Cerebral metabolic response to submaximal exercise. *Journal*
418 *of applied physiology*. 1999;87(5):1604-1608.
- 419 21. Newman GC. Clarification of abc/2 rule for ICH volume. *Stroke*. 2007;38(3):862.
- 420 22. Majersik JJ, Cole JW, Golledge J, et al. Recommendations from the international stroke
421 genetics consortium, part 1: standardized phenotypic data collection. *Stroke*.
422 2015;46(1):279-284.
- 423 23. Hwang J, Kim K, Brothers RM, Castelli DM, Gonzalez-Lima F. Association between
424 aerobic fitness and cerebrovascular function with neurocognitive functions in healthy,
425 young adults. *Exp Brain Res*. 2018;236(5):1421-1430.
- 426 24. Ainslie PN, Cotter JD, George KP, et al. Elevation in cerebral blood flow velocity with
427 aerobic fitness throughout healthy human ageing. *J Physiol*. 2008;586(16):4005-4010.
- 428 25. Bailey DM, Marley CJ, Brugniaux JV, et al. Elevated aerobic fitness sustained
429 throughout the adult lifespan is associated with improved cerebral hemodynamics. *Stroke*.
430 2013;44(11):3235-3238.

- 431 26. Pretnar-Oblak J, Sabovic M, Sebestjen M, Pogacnik T, Zaletel M. Influence of
432 atorvastatin treatment on L-arginine cerebrovascular reactivity and flow-mediated
433 dilatation in patients with lacunar infarctions. *Stroke*. 2006;37(10):2540-2545.
- 434 27. Hart EC, Joyner MJ, Wallin BG, Charkoudian N. Sex, ageing and resting blood pressure:
435 gaining insights from the integrated balance of neural and haemodynamic factors. *J*
436 *Physiol*. 2012;590(9):2069-2079.
- 437 28. Lucas SJ, Cotter JD, Brassard P, Bailey DM. High-intensity interval exercise and
438 cerebrovascular health: curiosity, cause, and consequence. *J Cereb Blood Flow Metab*.
439 2015;35(6):902-911.
- 440 29. Sato K, Sadamoto T, Ueda-Sasahara C, et al. Central command and the increase in
441 middle cerebral artery blood flow velocity during static arm exercise in women.
442 *Experimental physiology*. 2009;94(11):1132-1138.
- 443 30. Labrecque L, Drapeau A, Rahimaly K, Imhoff S, Billaut F, Brassard P. Comparable
444 blood velocity changes in middle and posterior cerebral arteries during and following
445 acute high-intensity exercise in young fit women. *Physiol Rep*. 2020;8(9):e14430.
- 446 31. Calverley TA, Ogoh S, Marley CJ, et al. HIITing the brain with exercise; mechanisms,
447 consequences and practical recommendations. *J Physiol*. 2020.
- 448 32. Hornby TG, Reisman DS, Ward IG, et al. Clinical Practice Guideline to Improve
449 Locomotor Function Following Chronic Stroke, Incomplete Spinal Cord Injury, and
450 Brain Injury. *J Neurol Phys Ther*. 2020;44(1):49-100.

451

Table 1. Participant Demographics (n = 26)

Age (years)	63.0 ± 14.0
Sex % (n)	
Female	31% (8)
Male	69% (18)
Race/Ethnicity* (n)	
White	18
Black	8
Native American	1
Hispanic/Latino	0
Asian	1
Pacific Islander	0
tPA (n)	14
Thrombectomy (n)	9
tPA and Thrombectomy (n)	3
Resting SBP (mm Hg)	138.5 ± 18.3
Resting DBP (mm Hg)	80.4 ± 11.2
BMI (kg*(m ²) ⁻¹)	29.8 ± 5.5
Time Post Stroke to 3-month visit (days)	96.3 ± 12.6
Lesion Size (cm ³)	40.2 ± 76.5
Estimated $V\dot{V}O_2$ maximum (ml*kg ⁻¹ *min ⁻¹)	23.2 ± 8.9
Medications (n)	
Beta Blocker	17
Calcium Channel Blocker	7
ACE Inhibitor	5
Angiotensin II Receptor Blocker	5
Statins	19
Diuretics	3
Blood Thinners	12
Aspirin	12
Metformin	5
Insulin	6
Antidepressants	9

Data are presented as Mean ± SD unless stated otherwise. *Multiple race categories selected. tPA = Tissue Plasminogen Activator; SBP = Resting Systolic Blood Pressure; DBP = Resting Diastolic Blood Pressure; BMI = Body Mass Index; $V\dot{V}O_2$ = oxygen uptake; ACE= Angiotensin Converting Enzyme.

Table 2. Kinetic Response Across Time

	3 months (n=22)	3 months 95 CI	6 months (n=15)	6 months 95 CI	F-test	P-value
Stroke Affected MCAv						
BL (cm*s ⁻¹)	51.1 ± 16.3	43.9-58.4	50.2 ± 12.5	43.3-57.1	1.9	0.19
TD (s)	55.6 ± 57.4	26.1-85.1	59.0 ± 52.6	25.5-92.4	0.2	0.70
Amp (cm*s ⁻¹)	5.5 ± 5.8	2.6-8.3	5.8 ± 5.2	2.5-9.0	0.3	0.63
τ (s)	34.6 ± 34.6	16.8-52.4	53.9 ± 53.9	19.6-88.1	1.6	0.24
Non-Stroke Affected MCAv						
BL (cm*s ⁻¹)	62.1 ± 23.7	50.7-73.6	56.2 ± 12.8	48.8-63.6	1.3	0.27
TD (s)	42.2 ± 38.6	17.7-66.7	62.7 ± 38.8	38.0-87.4	3.0	0.11
Amp (cm*s ⁻¹)	7.2 ± 5.8	3.5-10.90	7.3 ± 4.7	4.3-10.30	0.4	0.56
τ (s)	25.4 ± 19.1	13.3-37.5	21.0 ± 27.3	3.6-38.3	0.0	0.91
Data are presented as Mean ± SD. CI= Confidence Interval; BL= Baseline MCAv; TD= Time Delay; Amp= Amplitude; τ = Time Constant						

Table 3. Characteristics of Responders and Non-Responders

	Responders (n=12)	Responders 95 CI	Non- Responders (n=10)	Non- Responders 95 CI	F-test	P-value
Age (years)	59.4 ± 12.5	51.5-67.4	67.3 ± 16.1	55.8-78.8	1.7	0.21
Sex % (n)						
Female	63% (5)		37% (3)			
Male	50% (7)		50% (7)			0.68
Race/Ethnicity* (n)						
White	10		5			0.17
Black	2		5			0.17
Native American	0		1			0.46
Hispanic/Latino	0		0			
Asian	0		1			0.46
Pacific Islander	0		0			
Time Post Stroke to 3-month visit (days)	94.8 ± 8.3	89.5-100.0	99.5 ± 18.0	86.6-112.4	0.7	0.42
BMI (kg*(m ²) ⁻¹)	31.2 ± 6.2	27.3-35.2	28.2 ± 4.9	24.7-31.8	1.5	0.24
Lesion Size (cm ³)	70.6 ± 97.4	-4.2-145.5	20.7 ± 52.5	-23.2-64.5	1.7	0.22
Estimated $\dot{V}\dot{O}_2$ maximum (ml*kg ⁻¹ *min ⁻¹)	26.6 ± 8.9	20.9-32.3	18.3 ± 8.0	12.6-24.0	5.2	0.03
Physical Activity Score	2.8 ± 1.3	1.9-3.6	1.6 ± 0.7	1.1-2.1	6.4	0.02
MoCA	22.0 ± 5.8	18.3-25.7	23.8 ± 3.3	21.4-26.2	0.8	0.39
mRS	2.1 ± 0.9	1.5-2.5	2.2 ± 1.0	1.5-2.9	0.3	0.62
6-minute Walk Test (m)	338.4 ± 128.0	257.1-419.8	264.6 ± 128.9	165.5-363.6	1.7	0.21
Thrombectomy (n)	4		1			0.32
tPA and Thrombectomy (n)	2		1			1.00
tPA (n)	3		6			0.19
Medications (n)						
Beta Blocker	8		7			1.00
Calcium Channel Blocker	5		2			0.38
ACE Inhibitor	4		0			0.10

Angiotensin II Receptor Blocker	3	2	1.00
Statins	11	6	0.03
Diuretics	1	1	1.00
Blood Thinners	7	4	0.67
Aspirin	4	5	0.67
Insulin	1	4	0.14
Metformin	1	2	0.57
Antidepressants	4	4	1.00


Data are presented as Mean \pm SD unless stated otherwise. *Multiple race categories selected. BMI = Body Mass Index; $V\dot{O}_2$ = oxygen uptake; MoCA= Montreal Cognitive Assessment; mRS= Modified Rankin Scale; tPA = Tissue Plasminogen Activator; ACE= Angiotensin Converting Enzyme.

Table 4. MCAv Kinetics and Exercise Parameters For Responders and Non-Responders

	Responders (n=12)	Responders 95 CI	Non-Responders (n=10)	Non-Responders 95 CI	F-test	P-value
Stoke Affected MCAv						
BL (cm*sec ⁻¹)	54.2 ± 15.0	44.6 – 63.7	47.5 ± 17.7	34.8 - 60.2	0.9	0.35
TD (s)	32.9 ± 40.5	7.1 – 58.6	110.1 ± 58.4±	37.6 - 182.6	10.0	0.006
Amp (cm*sec ⁻¹)	8.2 ± 4.8	5.1 - 11.2	0.10 ± 3.2±	-3.2 - 3.4	13.7	0.002
τ (s)	30.2 ± 24.7	14.5 – 45.8	45.2 ± 53.8±	-21.7 - 112.0	0.7	0.43
Non-Stroke Affected MCAv						
BL (cm*sec ⁻¹)	59.9 ± 15.0	49.8 – 69.9	54.4 ± 13.5	41.9 - 66.9	0.23	0.64
TD (s)	36.2 ± 34.1	13.3-59.1	107.8*	-	-	-
Amp (cm*sec ⁻¹)	8.3 ± 4.4	5.3 - 11.3	-5.2*	-	-	-
τ (s)	22.7 ± 17.5	11.0 – 34.6	54.6*	-	-	-
BL MAP (mmHg)	77.6 ± 19.1	65.3 - 89.7	80.6 ± 19.0	67.0 - 94.2	0.14	0.71
SS MAP (mmHg)	98.4 ± 16.8	87.7 - 109.1	91.2 ± 25.6	72.9 - 109.6	0.6	0.44
Δ MAP (mmHg)	20.8 ± 9.6	14.7 – 26.9	10.6 ± 12.0	2.0 - 19.2	7.3	0.04
BL P _{ET} CO ₂ (mmHg)	34.6 ± 3.6	32.3 - 37.0	35.1 ± 5.1	31.2 - 39.1	0.1	0.81
SS P _{ET} CO ₂ (mmHg)	39.3 ± 4.4	36.5 - 41.6	38.2 ± 4.4	34.9 - 41.6	0.3	0.61
Δ P _{ET} CO ₂ (mmHg)	4.6 ± 2.3	3.2 – 6.1	3.1 ± 1.8	1.7 - 4.5	1.6	0.13
BL HR (bpm)	65.4 ± 8.9	59.8 - 71.1	69.9 ± 11.3	61.8 - 77.9	1.1	0.31
SS HR (bpm)	94.5 ± 13.1	86.2 - 102.8	87.6 ± 18.1	74.6 - 100.6	1.1	0.31
Δ HR (bpm)	29.1 ± 9.5	23.1 - 35.1	17.7 ± 11.9	9.2 - 26.3	6.2	0.02
Workload (watts)	52.9 ± 25.4	36.8-69.1	36.0 ± 12.0	27.4-44.6	3.7	0.07
RPE	12.8 ± 2.2	11.3-14.2	13.4 ± 3.1	11.2-15.6	0.3	0.58

Data are presented as Mean \pm SD. CI= Confidence Interval; BL= Baseline MCAv; TD= Time Delay; Amp= Amplitude; τ = Time Constant; BL= Baseline; SS= Steady-State Exercise; MAP= Mean Arterial Pressure; P_{ET}CO₂= End Tidal Carbon Dioxide; HR= Heart Rate; RPE= Rate of Perceived Exertion. \pm Data represent n = 5. *Data represents 1 dataset that modeled an exponential response.

452 **Figure 1.** Responders versus Non-Responders, Stroke Affected and Non-Stroke Affected MCAv
453 response to moderate intensity exercise
454



455