

Prevalence and Risk Factors of Diabetes Mellitus in People over 40 Years Old in Rural Areas of Central Hunan Province

Lan Li^{1,2#}, Yuan Liao^{1,2,3#}, Hao Xiao^{1,2}, Tiantian Wang^{1,2}, Jing Liu^{1,2},
Guijuan Zhou^{1,2}, Guanghua Sun^{1,2}, Kang Li^{1,2}, Fujin Huang^{1,2}, Weibin
Feng^{1,2}, Jun Zhou^{1,2*}

Affiliations

1. Department of Rehabilitation, The First Affiliated Hospital of University of South China, Hengyang, Hunan, People's Republic of China, 421001

2. Rehabilitation Laboratory, The First Affiliated Hospital of University of South China, Hengyang, Hunan, People's Republic of China, 421001

3. Clinical Medical College of Acupuncture Moxibustion and Rehabilitation, Guangzhou University of Traditional Chinese Medicine, Guangzhou, Guangdong, People's Republic of China, 510006

*Correspondence

Jun Zhou

Tel: +86-13575239169; E-mail address: zhoujun8005@163.com

#These authors contributed equally to this work and should be considered co-first authors.

Acknowledgements

We would like to thank Shiju Chen MD and Chengxiao Fu MD for their assistance with language editing of this manuscript. And this study was supported by the Natural Science Foundation of Hunan Province (2018JJ2358).

Abstract

Objective To explore the prevalence of diabetes mellitus in rural areas of central Hunan province, and to analyze the risk factors related to diabetes mellitus (DM). **Methods** A cross-sectional study was conducted by using questionnaires (gender, age, occupation, degree of education, exercise habits, smoke status, alcohol consumption, hypertension, cardiovascular disease, family history of diabetes), anthropometric measurements (height, weight, waist circumference, blood pressure) and biochemical indexes test (fasting blood-glucose, glycosylated hemoglobin, blood lipids). Villagers over 40 years old from a community in rural areas of central Hunan Province were investigated.

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

Results A total of 410 clinical data were collected. The prevalence of diabetes mellitus in 410 (177 in male and 233 in female) villagers was 13.41%, including 13 males with a prevalence rate of 7.34% and 42 females with a prevalence rate of 15.16%. The prevalence of impaired fasting glucose (IFG) was 11.95%, 21 males (11.86% of males) and 28 females (10.11% of females). And the results of glycosylated hemoglobin test showed that 64% villagers with diabetes mellitus had hemoglobin A1c(HbA1c) above 6.5%. Univariate analysis suggested that gender, smoke status, alcohol consumption, family history of diabetes mellitus, hypercholesterolemia and hypertension were involved in diabetes mellitus ($P < 0.05$). Multivariate logistic regression analysis showed that family history of diabetes (OR: 1.759; 95% CI: 1.010-3.065), hypercholesterolemia (OR: 3.819; 95% CI: 1.27-11.486) and hypertension (OR: 2.074; 95% CI: 1.130-3.809) were independent risk factors for diabetes mellitus, and the differences were statistically significant ($P < 0.05$). **Conclusion** The prevalence of diabetes mellitus in rural areas of central Hunan Province is higher. Family history of diabetes, hypercholesterolemia and hypertension are major risk factors for diabetes mellitus. The knowledge of diabetes should be strengthened. Related interventions should be given based on the diabetes epidemic status of local.

Key Words: Diabetes Mellitus; Prevalence; Risk factors; Rural areas

Introduction

Diabetes mellitus, as well as hyperglycemia, is considered one of the most well-known disease affecting various body systems. The prevalence of Diabetes mellitus increases greatly with the aged tendency of population and change of life style, which severely threatens people's health and affects the quality of life of adults even children^[1]. Diabetes mellitus has been one of the biggest public health problems in the modern world, but people's awareness of diabetes is not high. At present, domestic data on this aspect are mostly based on first-tier cities^[2-4], while there is rarely systematic and in-depth investigation on low-income areas such as rural areas in central Hunan province. The purposes of this study were to investigate the prevalence of diabetes and explore its risk factors among people over 40 years old in rural areas of central Hunan

province, in order to enrich the prevalence of diabetes among residents in different regions in China and provide scientific evidence for the formulation of national health policies.

Methods

Study participants

In January 2019, a representative sample of the general participants aged ≥ 40 years was selected in order to investigate the prevalence and related risk factors in rural areas of central Hunan province. The study adopted the scheme of random sampling, and selected Jiao Shan Community in Hengyang city, Hunan province. All eligible permanent residents aged ≥ 40 years in each village were invited to participate in the study. The flow chart of this study was shown in **Fig. 1**. This study was subject to approval by the Ethics Committee of the First Affiliated Hospital of University of South China, Hengyang, Hunan. And clinical trial registration in Chinese Clinical Trial Registry Center, ChiCTR1800020006. All procedures were carried out in accordance with the ethical standards. Additionally, all participants, after being properly informed about the study procedures and potential risks, signed a free informed consent document.


Data collection and measurements

The methods of investigation include questionnaires (such as gender, age, occupation, degree of education, exercise habits, smoke status, alcohol consumption, hypertension, cardiovascular disease, family history of diabetes), anthropometric measurements (include height, weight, waist circumference, blood pressure) and biochemical indexes test (fasting blood-glucose, glycosylated hemoglobin, blood lipids and so on). The participants were instructed to take off their shoes and wear lightweight clothes When measuring height and weight. Height and waist circumference were accurate to 0.1cm and weight was accurate to 0.1kg. Waist circumference (WC) was measured at the umbilical level (to the nearest 0.1cm). The body mass index (BMI) was calculated as $\text{weight}(\text{kg})/\text{height}(\text{m}^2)$.

Blood pressure was measured two times at one-minute intervals after at least five minutes of rest using a standardized electronic automatic sphygmomanometer (TSB-

615B-T; Chaosi, Beijing, China). the average of the two blood pressure measurements was calculated at each examination.

Fig.1 Process of the study



Fasting venous blood was drawn and placed into a vacuum tube containing EDTA (Becton Dickinson, Franklin Lakes, NJ, USA) after fasting for at least 8 hours. Fasting plasma glucose (FPG) were measured using a blood glucose test paper (Sinocare Inc.). The correlation analyses of biochemical indices (total cholesterol, triglycerides, low-density lipoprotein cholesterol and high-density lipoprotein cholesterol) were identified by the Cadillac detector (SN:3222597, Indianapolis, USA). Centrifugation was done within 2 hours after collection and hemoglobin A1c was determined within 24 hours using a specific protein analyzer HP-AFS/3, Hebei, China). All laboratory equipment was calibrated and blinded duplicate samples were used.

Definitions

According to the 2017 American Diabetes Association (ADA) guidelines diabetes

diagnosis^[5] was that FPG \geq 7.0 mmol/L or using insulin or oral hypoglycaemic drugs or history of diabetes diagnosis. And IFG was defined FPG $>$ 6.1mmol/L and $<$ 7.0mmol/L. The diagnostic criteria for Dyslipidemia were based on the 2007 Chinese Guideline for Prevention and Treatment of Dyslipidemia in Adult Patients: total cholesterol (TC) \geq 6.2mmol/L; or triglycerides (TG) \geq 2.3mmol/L; or low-density lipoprotein cholesterol (LDL-C) \geq 4.1mmol/L; or high-density lipoprotein cholesterol (HDL-C) $<$ 1.0mmol/L was defined Dyslipidemia. We categorized participants into four groups according to their BMI: obese cases were BMI \geq 28.0 kg/m², overweight cases were BMI $<$ 28.0Kg/m² and BMI \geq 24.0Kg/m²; normal weight cases were BMI \leq 18.5Kg/m² and BMI $>$ 24.0Kg/m² and underweight cases were BMI $<$ 18.5Kg/m²^[6]. Abdominal obesity^[7] was defined as a waist circumference in men \geq 90cm, women \geq 80cm. In this study, smoke status was defined as ever smokers (former or current smoking). Alcohol consumption (liquor) \geq 3 times per week and each \geq 400g were defined heavy drinking; otherwise it's a low drinking. Regular exercise was defined as exercising 3 to 5 times or more per week. Family history of diabetes was defined as having at least one parent or sibling with diabetes. What's more, Hypertension was defined as systolic blood pressure \geq 140 mmHg and/or diastolic blood pressure \geq 90 mmHg, or taking antihypertensive drugs^[8]. Coronary heart disease (CHD) was defined as previous history of coronary artery disease.

Statistical analysis

All data were analyzed using the IBM SPSS Statistics for Windows version 22. A univariate analysis of risk factors was conducted with gender, age, occupation, degree of education, exercise habits, smoke status, alcohol consumption, hypertension, cardiovascular disease, family history of diabetes, BMI, waist circumference and blood lipid level as the independent variables. T-test was used for measurement data, chi-square(χ^2) test was used for counting data, and Risk factors with statistical significance in univariate analysis were subjected to multivariate logistic regression analysis. $P < 0.05$ was considered statistically significant.

Results

Univariate analysis

A total of 410 participants [males:177(43.2%), females:233(56.8%)] were investigated in the study. The biochemical characteristics and risk factors of the studied group are shown in **Table 1**. The prevalence of diabetes mellitus in 410 villagers was 13.41%, including 13 males with a prevalence rate of 7.34% and 42 females with a prevalence rate of 15.16%. And the prevalence of IFG was 11.95%, 21 males (11.86% of males) and 28 females (10.11% of females) (**Table 2**).

Table 1 The biochemical characteristics and risk factors of the population in rural areas of central Hunan Province

Factor	Total	Men	Women	χ^2	P-value
Age group(years)	410			6.69	0.154
40-49	33	14	19		
50-59	134	54	80		
60-69	185	83	102		
70-79	42	19	23		
≥80	16	7	9		
BMI (kg/m²) *	341			2.006	0.571
≥28	17	10	7		
24-27.9	114	55	59		
18.5-23.9	193	77	116		
<18.5	17	3	14		
Education level *	397			5.630	0.131
pre-school or Primary school	269	95	174		
Secondary school	107	60	47		
Tertiary education	19	14	5		
Occupation*	399			0.207	0.649
Crop and Pasture Production	330	131	199		
other	69	43	26		
Smoke status *	408			6.958	0.008
yes	121	116			
no	287	59			
Alcohol consumption *	407			7.615	0.022
no	283	87	196		

light	91	56	35		
heavy	33	31	2		
Exercise *	<i>403</i>			0.045	0.832
often	305	133	172		
rarely	98	41	57		
Waist circumference (cm) *	<i>400</i>			0.246	0.620
≥90 in men, ≥80 in women	129	38	91		
<90 in men, <80 in women	271	134	137		
Hypertriglyceridemia*	<i>395</i>			1.603	0.206
yes	99	46	53		
no	296	124	172		
Hypercholesterolemia*	<i>395</i>			8.235	0.004
yes	16	4	12		
no	379	166	213		
High LDL cholesterolemia*	<i>365</i>			1.003	0.317
yes	13	7	6		
no	352	149	203		
Low HDL cholesterolemia*	<i>395</i>			0.170	0.680
yes	83	47	36		
no	312	123	189		
Hypertension	<i>410</i>			6.431	0.11
yes	160	72	88		
no	250	105	145		
Coronary heart disease *	<i>348</i>			0.073	0.787
yes	32	14	18		
no	316	136	180		
Family history of diabetes	<i>410</i>			16.5	0.000
yes	33	11	22		
no	363	160	203		

* missing values


It's known that hemoglobin A1c above 6.5 indicates poor glycemic control in the last 2-3 months ^[9]. In our study, the results of glycosylated hemoglobin test showed that 35 patients with diabetes mellitus had no blood glucose control in the past 2-3 months, and the rate of non-compliance reached 64% (**Table 2**).

Table 2 The prevalence of diabetes mellitus and IFG and their HbA1c level among people over 40 years old in rural areas of central Hunan province

Variable	DM		IFG	
	Number of DM participants	Prevalence (%)	Number of IFG participants	Prevalence (%)
Men(n=177)	13	7.34	21	11.86
Women(n=233)	42	15.16	28	10.11
Total(n=410)	55	13.41	49	11.95
HbA1c>6.5%	35	64	3	6.1

Hypercholesterolemia in dyslipidemia is major risk factors for diabetes mellitus in both men and women (**Table 1**). Furthermore, univariate analysis suggested that gender, smoke status, alcohol consumption, family history of diabetes mellitus, hypercholesterolemia and hypertension were involved in diabetes mellitus ($P<0.05$) (**Table 1**). The prevalence of diabetes generally increased with age, which was more noticeable in females than in males (**Table 2 and Fig. 2**). The prevalence of diabetes in 40-49 years old was less than that of IFG, while the prevalence of IFG was lower than that of diabetes in over 50 years old (**Fig. 2**).

Fig.2 The prevalence of diabetes mellitus and IFG among villagers of different ages



Multivariate analysis

Subsequently, that gender, smoke status, alcohol consumption, family history of diabetes mellitus, hypercholesterolemia and hypertension had statistical significance in univariate analysis was incorporated into the multivariate analysis. The result of multivariate logistic regression analysis is showed that family history of diabetes (OR: 1.759; 95% CI: 1.010-3.065), hypercholesterolemia (OR: 3.819; 95% CI: 1.270-11.486) and hypertension (OR: 2.074; 95% CI: 1.130-3.809) were independent risk factors for diabetes mellitus, and the differences were statistically significant ($P < 0.05$) (Table 3).

Table 3 Multivariate analysis for diabetes mellitus risk factors

Variable	Odd ratio	95%CI	P-value
Gender	0.639	0.261-1.566	0.328
Smoke status	0.458	0.147-1.424	0.177
Alcohol consumption	0.646	0.333-1.256	0.198
Hypercholesterolemia	3.819	1.27-11.486	0.017
Hypertension	2.074	1.130-3.809	0.019
Family history of diabetes	1.759	1.010-3.056	0.046

Discussion

Diabetes mellitus poses serious public health problems, closely related to a variety of diseases, such as cerebral infarction and coronary heart disease, affecting individuals and society in terms of life quality and economic burden^[10-12]. It is reported that women may exercise less than men, and a series of physiological changes associated with the relatively lower level of sex hormones after menopause in women affect blood glucose levels in the body^[13].

In this study, the prevalence of diabetes generally increased with age, which was more noticeable in females than in males. This study was a community-based, cross-sectional survey carried out in rural areas of central Hunan Province, China. And the prevalence of the areas is 13.41%, which is higher than Liaoning and Sichuan province^[4, 14, 15]. Recently, more and more people began to occupied in sedentary lifestyle instead

of manual labor, and the risk of diabetes increases with lack of physical activity^[16]. Adequate physical activity can increase insulin sensitivity, and is important to preventing the development of many chronic diseases such as diabetes, hypertension and obesity^[17, 18]. The presented study shows that their occupation is not a risk factor for diabetes, which is related to the fact that most villagers have been engaged in crop and pasture production (moderate to severe physical labor) for a long time, indicating that physical labor can reduce the occurrence of diabetes in rural China.

It is known that IFG is pre-diabetes^[19], and the prevalence of IFG in this study was 11.95%, which is slightly lower than diabetes mellitus. If they are not taken seriously, it is easy for them to further develop into diabetes, which in turn increases the prevalence of diabetes. Actually, most villagers have few symptoms that may not be recognized, are not aware of their pre-diabetes, and do not control diet and strengthen exercise, which may be one of the reasons that the prevalence of diabetes in this region is higher than in other regions. In addition, diabetes control in these diabetic villagers is not adequate as showed by high rates of poor control of HbA1c, because of the low education level of the villagers, the weak awareness of diabetes prevention and control, and the poor treatment compliance.

Some, but not all, studies^[20, 21] have shown that smoking is associated with an increase in systemic insulin resistance. Maddatu et al.^[22] also showed that compared with non-smokers, long-term smokers can inhibit insulin signaling pathways, strengthen fat infiltration in muscles, reduce adiponectin levels, increase retinol binding protein 4 and increase insulin resistance in skeletal muscles. In this research, alcohol consumption increase the risk of diabetes mellitus, which is different from one study that reported that light alcohol consumption can reduces the risk of diabetes^[23]. Multi-center studies with large sample size are needed for further analysis and verification.

Both univariate and multivariate analyses suggested that hypercholesterolemia increased the risk of diabetes. Sabag et al.^[24] reported that exercise can effectively improve obesity and abnormal lipid metabolism in Type 2 diabetes mellitus(T2DM) patients. Moreover, residents with positive family history of diabetes had a higher prevalence than those without family history of diabetes, which is related to family

dietary habits, lifestyle similarity and genetic factors.

Increased adipokine secretion, lipid deposition, inflammatory response, oxidative stress, and metabolic abnormalities associated with insulin resistance are the common pathophysiological basis of obesity, diabetes, and coronary heart disease^[25]. Increased adipose tissue accelerates oxidative stress response in obese patients, and the accumulation of intracellular reactive oxygen leads to decreased insulin sensitivity through protein kinase C, mitokinase-activated protein kinase p38 and c-jun amino-terminal kinase^[26, 27]. A series of metabolic disorders such as hyperglycemia and increased free fatty acids which caused by decreased insulin sensitivity has a direct toxic effect on cells^[28], promoting the occurrence of diabetes.

There are common risk factors for diabetes and hypertension, which may act on both diseases^[29]. Diabetes is a risk factor for hypertension, and it highlighted separately in guidelines on hypertension in China. As a risk factor for diabetes, hypertension has a great influence on the occurrence and development of diabetes, closely to abnormal blood pressure regulation, endothelial dysfunction and lipid metabolism disorder^[30].

The study has great significance to the formation of diabetes prevention and control strategies in central Hunan province and even the country, but there are also some deficiencies: First, some villagers over 40 years old are not included in the survey because they are out of town or far away from the survey site, and the study subjects may have gender deviation. Second, although type 2 diabetes is the dominant type in adults over the age of 40, but type 1 and type 2 diabetes were not classified in this study. Third, the risk factors in this study did not include recognized risk factors such as obesity and exercise, which may be related to insufficient sample and the erroneous definition of exercise. Additionally, dietary habits were not included in this study because of the unreasonable questionnaire design. In the future, more large-sample size is needed to obtain more scientific evidences.

Conclusion

The prevalence of diabetes mellitus in rural areas of central Hunan Province is higher, and most diabetic villagers have failed to achieve glycemic control in the past

2-3 months. Family history of diabetes, hypercholesterolemia and hypertension are major risk factors for diabetes mellitus. The knowledge of diabetes should be strengthened. Related interventions should be given based on the diabetes epidemic status of local.

Reference

- [1] W. Xu, M. Tian and Y. Zhou. The relationship between insulin resistance, adiponectin and C-reactive protein and vascular endothelial injury in diabetic patients with coronary heart disease [J]. *Experimental and therapeutic medicine*, 2018, 16(3): 2022-6.
- [2] J. H. Wang, H. L. Zheng, M. Liu, Y. He, L. Wu, Y. Y. Wang, D. Zhang, J. Zeng and Y. Yao. [Prevalence and control status of diabetes in elderly community residents and risk factors] [J]. *Zhonghua liu xing bing xue za zhi = Zhonghua liuxingbingxue zazhi*, 2016, 37(9): 1220-2.
- [3] Y. Ruan, Q. H. Yan, J. Y. Xu, Q. D. Yang, H. H. Yao, R. Li and Y. Shi. Epidemiology of Diabetes in Adults Aged 35 and Older from Shanghai, China [J]. *Biomedical and environmental sciences : BES*, 2016, 29(6): 408-16.
- [4] G. Cheng, T. Dai, Y. Liu, X. Liao, J. Liu, C. Ren and X. Zhang. [Prevalence and related risk factors among residents with diabetes in urban and rural areas in Chengdu] [J]. *Zhonghua liu xing bing xue za zhi = Zhonghua liuxingbingxue zazhi*, 2014, 35(4): 386-9.
- [5] Standards of Medical Care in Diabetes-2017 Abridged for Primary Care Providers [J]. *Clinical diabetes : a publication of the American Diabetes Association*, 2017, 35(1): 5-26.
- [6] B. F. Zhou. Predictive values of body mass index and waist circumference for risk factors of certain related diseases in Chinese adults--study on optimal cut-off points of body mass index and waist circumference in Chinese adults [J]. *Biomedical and environmental sciences : BES*, 2002, 15(1): 83-96.
- [7] K. G. Alberti, P. Zimmet and J. Shaw. The metabolic syndrome--a new worldwide definition [J]. *Lancet (London, England)*, 2005, 366(9491): 1059-62.
- [8] A. V. Chobanian, G. L. Bakris, H. R. Black, W. C. Cushman, L. A. Green, J. L. Izzo, Jr., D. W. Jones, B. J. Materson, S. Oparil, J. T. Wright, Jr. and E. J. Roccella. The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure: the JNC 7 report [J]. *Jama*, 2003, 289(19): 2560-72.
- [9] Y. Kim and C. W. Park. New therapeutic agents in diabetic nephropathy [J]. *The Korean journal of internal medicine*, 2017, 32(1): 11-25.
- [10] P. Zimmet, K. G. Alberti, D. J. Magliano and P. H. Bennett. Diabetes mellitus statistics on prevalence and mortality: facts and fallacies [J]. *Nature reviews. Endocrinology*, 2016, 12(10): 616-22.
- [11] N. Laiteerapong and A. S. Cifu. Screening for Prediabetes and Type 2 Diabetes Mellitus [J]. *Jama*, 2016, 315(7): 697-8.
- [12] J. da Rocha Fernandes, K. Ogurtsova, U. Linnenkamp, L. Guariguata, T. Seuring, P. Zhang, D. Cavan and L. E. Makaroff. IDF Diabetes Atlas estimates of 2014 global health expenditures on diabetes [J]. *Diabetes research and clinical practice*, 2016, 117(48-54).
- [13] Z. B. Zhang, Z. X. Xue, H. Wang, T. M. Wang, Y. H. Li, X. L. Song, X. F. Chao, G. Wang, X. J. Wu, N. Nazibam, B. Ayxamgul, E. Gulbahar, Z. Y. Zhou, B. S. Sun, Y. Z. Wang and M. Wang. [Prevalence

of diabetes mellitus and associated risk factors in Uygur population in Kashgar area of Xinjiang Uygur Autonomous Region] [J]. *Zhonghua liu xing bing xue za zhi = Zhonghua liuxingbingxue zazhi*, 2017, 38(6): 696-701.

[14] X. Zhou, H. Guan, L. Zheng, Z. Li, X. Guo, H. Yang, S. Yu, G. Sun, W. Li, W. Hu, L. Guo, G. Pan, L. Xing, Y. Zhang and Y. Sun. Prevalence and awareness of diabetes mellitus among a rural population in China: results from Liaoning Province [J]. *Diabetic medicine : a journal of the British Diabetic Association*, 2015, 32(3): 332-42.

[15] Q. Sun, X. F. Chen, X. P. Wu, G. J. Luo, J. G. Li, H. Wang, X. F. Chen, Z. Bian and Y. Guo. [Prevalence of diabetes and related factors in Pengzhou city, Sichuan province] [J]. *Zhonghua liu xing bing xue za zhi = Zhonghua liuxingbingxue zazhi*, 2016, 37(8): 1091-4.

[16] R. Siqueira Reis, A. A. Hino, C. Ricardo Rech, J. Kerr and P. Curi Hallal. Walkability and physical activity: findings from Curitiba, Brazil [J]. *American journal of preventive medicine*, 2013, 45(3): 269-75.

[17] C. Handschin and B. M. Spiegelman. The role of exercise and PGC1alpha in inflammation and chronic disease [J]. *Nature*, 2008, 454(7203): 463-9.

[18] A. Westheim, I. Os and I. Hjermann. [Physical activity and the metabolic syndrome] [J]. *Tidsskrift for den Norske laegeforening : tidsskrift for praktisk medicin, ny raekke*, 1993, 113(6): 723-4.

[19] Z. Sheng, J. Y. Cao, Y. C. Pang, H. C. Xu, J. W. Chen, J. H. Yuan, R. Wang, C. S. Zhang, L. X. Wang and J. Dong. Effects of Lifestyle Modification and Anti-diabetic Medicine on Prediabetes Progress: A Systematic Review and Meta-Analysis [J]. *Frontiers in endocrinology*, 2019, 10(455).

[20] Y. Wu, P. Song, W. Zhang, J. Liu, X. Dai, Z. Liu, Q. Lu, C. Ouyang, Z. Xie, Z. Zhao, X. Zhuo, B. Viollet, M. Foretz, J. Wu, Z. Yuan and M. H. Zou. Activation of AMPKalpha2 in adipocytes is essential for nicotine-induced insulin resistance in vivo [J]. *Nature medicine*, 2015, 21(4): 373-82.

[21] A. A. Saeed. Association of tobacco products use and diabetes mellitus-results of a national survey among adults in saudi arabia [J]. *Balkan medical journal*, 2012, 29(3): 247-51.

[22] J. Maddatu, E. Anderson-Baucum and C. Evans-Molina. Smoking and the risk of type 2 diabetes [J]. *Translational research : the journal of laboratory and clinical medicine*, 2017, 184(101-7).

[23] Y. J. Lai, H. Y. Hu, Y. L. Lee, M. C. Ko, P. W. Ku, Y. F. Yen and D. Chu. Frequency of alcohol consumption and risk of type 2 diabetes mellitus: A nationwide cohort study [J]. *Clinical nutrition (Edinburgh, Scotland)*, 2019, 38(3): 1368-72.

[24] A. Sabag, K. L. Way, S. E. Keating, R. N. Sultana, H. T. O'Connor, M. K. Baker, V. H. Chuter, J. George and N. A. Johnson. Exercise and ectopic fat in type 2 diabetes: A systematic review and meta-analysis [J]. *Diabetes & metabolism*, 2017, 43(3): 195-210.

[25] A. R. Saltiel and J. M. Olefsky. Inflammatory mechanisms linking obesity and metabolic disease [J]. *The Journal of clinical investigation*, 2017, 127(1): 1-4.

[26] S. K, G. P. Senthilkumar, P. Sankar and Z. Bobby. Attenuation of oxidative stress, inflammation and insulin resistance by allium sativum in fructose-fed male rats [J]. *Journal of clinical and diagnostic research : JCDR*, 2013, 7(9): 1860-2.

[27] X. Lin, Z. Zhang, J. M. Chen, Y. Y. Xu, H. R. Ye, J. Cui, Y. Fang, Y. Jin, D. R. Zhu and L. Yuan. Role of APN and TNF-alpha in type 2 diabetes mellitus complicated by nonalcoholic fatty liver disease [J]. *Genetics and molecular research : GMR*, 2015, 14(2): 2940-6.

[28] S. Pereira, E. Park, Y. Mori, C. A. Haber, P. Han, T. Uchida, L. Stavar, A. I. Oprescu, K. Koulajian,

A. Ivovic, Z. Yu, D. Li, T. A. Bowman, J. Dewald, J. El-Benna, D. N. Brindley, R. Gutierrez-Juarez, T. K. Lam, S. M. Najjar, R. A. McKay, S. Bhanot, I. G. Fantus and A. Giacca. FFA-induced hepatic insulin resistance in vivo is mediated by PKCdelta, NADPH oxidase, and oxidative stress [J]. American journal of physiology. Endocrinology and metabolism, 2014, 307(1): E34-46.

[29] Y. Shen, C. Chang, J. Zhang, Y. Jiang, B. Ni and Y. Wang. Prevalence and risk factors associated with hypertension and prehypertension in a working population at high altitude in China: a cross-sectional study [J]. Environmental health and preventive medicine, 2017, 22(1): 19.

[30] R. Chilton, I. Tikkanen, U. Hehnke, H. J. Woerle and O. E. Johansen. Impact of empagliflozin on blood pressure in dipper and non-dipper patients with type 2 diabetes mellitus and hypertension [J]. Diabetes, obesity & metabolism, 2017, 19(11): 1620-4.