1

ACE2 polymorphisms as potential players in COVID-19 outcome

2

Short title: ACE2 polymorphisms in COVID-19 outcome

André Salim Khayat^{1,2,*}, Paulo Pimentel de Assumpção², Bruna Claudia Meireles Khayat¹, Taíssa
Maíra Thomaz Araújo², Jéssica Almeida Batista-Gomes¹, Luciana Carvalho Imbiriba², Geraldo
Ishak², Paula Baraúna de Assumpção², Fabiano Cordeiro Moreira², Rommel Rodriguez
Burbano^{2,3}, André Ribeiro-dos-Santos¹, Ândrea Kelly Ribeiro-dos-Santos^{1,2}, Ney Pereira
Carneiro dos Santos^{1,2}, Sidney Emmanuel Batista dos Santos^{1,2}.

8 ¹Instituto de Ciências Biológicas, Universidade Federal do Pará, Belém, Pará, 66075-110, Brazil.

9 ²Núcleo de Pesquisas em Oncologia, Universidade Federal do Pará, Belém, Pará, 66073-000, Brazil.

10 ³Laboratório de Biologia Molecular, Hospital Ophir Loyola, Belém, Pará, 66060-281, Brazil.

11 *Corresponding author (E-mail: <u>khayatas@gmail.com</u>)

12

13 Abstract

The clinical condition COVID-19, caused by SARS-CoV-2, was declared a pandemic by the 14 WHO in March 2020. Currently, there are more than 5 million cases worldwide, and the 15 pandemic has increased exponentially in many countries, with different incidences and death 16 rates among regions/ethnicities and, intriguingly, between sexes. In addition to the many factors 17 that can influence these discrepancies, we suggest a biological aspect, the genetic variation at the 18 19 viral S protein receptor in human cells, ACE2 (angiotensin I-converting enzyme 2), which may contribute to the worse clinical outcome in males and in some regions worldwide. We performed 20 21 exomics analysis in native and admixed South American populations, and we also conducted in 22 silico genomics databank investigations in populations from other continents. Interestingly, at least ten polymorphisms in coding, noncoding and regulatory sites were found that can shed light 23 24 on this issue and offer a plausible biological explanation for these epidemiological differences. In 25 conclusion, ACE2 polymorphisms should influence epidemiological discrepancies observed 26 among ancestry and, moreover, between sexes.

27 Introduction

At the end of 2019, a new outbreak caused by SARS-CoV-2 (a coronavirus) started in Hubei Province, China. The clinical condition, COVID-19, probably arose from natural selection in bat reservoirs [1]. There are currently more than five million cases worldwide, and the pandemic has been increasing exponentially in many countries since the disease was deemed a pandemic by WHO in March 2020 [2].

The lethality rate is influenced by the speed of contagion, idiosyncrasies of the affected populations according to the containment policies adopted, socioeconomic conditions and the absorption limit of the health system [3]. Due to the high rate of transmission of the virus by air and the novelty of the infection to humans, the disease has become a global emergency problem, forcing periods of social confinement to contain the pandemic, in addition to hygiene habits [4].

Epidemiological data show a tendency towards slightly greater contamination in men; however, male mortality is significantly higher [5], representing an increase from 20% to 70% in European countries, approximately 65% in some Asian countries, and, even more peculiarly, Dominican Republic citizens showed three times more deaths in men than women [6,2,7], even considering factors related to behavioral issues. Certainly, there are biological aspects that contribute to this more adverse clinical condition.

44 Studies indicate that cell–virus interaction is mediated by the connection of the 45 transmembrane glycoprotein spike (S), present in the form of homotrimers on the viral surface, to 46 angiotensin I-converting enzyme 2 (ACE2, also called hACE2), which is responsible primarily 47 for inducing vasodilation [8].

The level and expression pattern of ACE2 in different tissues and cells can be critical to 48 the susceptibility and symptoms resulting from SARS-CoV-2 infection [9]. Zhou and 49 collaborators [10], using scRNA-seq datasets, classified organs vulnerable to infection as high 50 and low risk based on their expression levels of ACE2. At the clinical level, the symptoms of 51 52 COVID-19 may be related to the entry and affinity of the virus in these organs, as observed in heart failure disease and increased ACE2 expression, in which viral infection is related to a 53 higher risk of heart attack and worse ill condition [11]. For Li and collaborators [12], ACE2 54 genetic variations could be crucial to the susceptibility in different cohorts and to clinical 55 outcomes of COVID-19. 56

57 Currently, investigations of potential genetic variations that may favor or hinder 58 interactions between the virus and the host have been conducted [13,14,15], showing a high 59 number of codons that can, if altered, interfere with the complexity of the virus–cell interaction. 60 It is noteworthy that the ACE2 is located on the X chromosome, causing the impossibility of 61 heterozygosity in men. Therefore, polymorphisms in their single copy could be related to the 62 worst outcomes observed in males [16].

Considering the above, we sought explanations for an intrinsic factor that differed between sexes and populations that may justify the differences observed in the incidence and lethality of SARS-CoV-2 infection among the different regions of the world, as well as between sexes. We analyzed global data in the 1000 Genomes Database and, in addition, we conducted studies of exomes in two population groups in the Brazilian Amazon (Indians and miscegenated), without description in public genomic banks, and we compared this information with a public databank from a population in southeastern Brazil. These comparisons are important because

Brazil has a continental size and an admixed population in the North (more Amerindians among
all regions), Northeast (more Africans), and South and Southeast (more Europeans) [17].

72 Materials and Methods

73 Analyses in the 1000 Genomes Project

The analysis was performed on data from the 1000 Genomes Phase 3 database (1000G), which comprises 84,4 million variants in 2,504 individuals from 26 different populations [18]. These populations were concentrated in five large groups: African (AFR), Ad Mixed American (AMR), East Asian (EAS), European (EUR), and South Asian (SAS).

Through the complete sequence of the X chromosome, a region between nucleotides 78 15620281 and 15512643 was selected in revision GRCh37.p13 (15602158 and 15494520 in 79 GRCh38.p13) since the angiotensin I-converting enzyme 2 gene (ACE2, Gene ID: 59272, 80 updated on 22-Mar-2020) is located on the complementary strand, covering 107639 bp. 81 82 Additionally, a region of ten thousand base pairs upstream to the gene was included in the analysis so that the initial search site became 15630281 (GRCh37.p13) [18,19], aiming to search 83 for modifications in noncanonical sites to locate minor allele frequency (MAF) and allelic 84 85 differences that could be relevant. The index considered was the difference in allele frequency among all polymorphisms (SNP, INDEL and SV) contained in the study region, regardless of 86 their global or populational frequency. 87

88 Analyses in Amazon Natives and Admixed Population

For the allelic comparison between the populations cataloged in the 1000 Genomesdatabase and the population not described in the respective project, we investigated a population

91 composed of 64 Amerindians and 82 admixed individuals from the Amazon region of northern 92 Brazil. This study was approved by the National Committee for Ethics in Research (CONEP) and the Research Ethics Committee of the UFPA Tropical Medicine Center under CAAE number 93 20654313.6.0000.5172. The Amerindians represent 10 different Amazonian ethnic groups that 94 were grouped together as the Native American (NAM) group. Tribe names and geographic 95 96 coordinates of the Brazilian Amazon Indian populations are presented in Table S4. The 82 97 admixed individuals (Brazilian Admixed Population - BAP) live in Belém city, located in northern Brazil, where, due to the colonization process, are characterized by three ancestral 98 genetic components: European, Native American and African. This sample group is also enrolled 99 100 in a broad project. Furthermore, we also compared our findings to a database of variants analyzed in a Southeast Brazilian population, the Online Archive of Brazilian Mutations (ABraOM, we 101 102 represent here as ABM) [20].

103 DNA Extraction and Exome Library

The DNA was extracted as described by Sambrook and collaborators [21]. The genetic material was quantified using a Nanodrop spectrophotometer (Thermo Fisher Scientific Inc., USA). The libraries were prepared using the Nextera Rapid Capture Exome (Illumina) and SureSelect Human All Exon V6 (Agilent) Kits. Sequencing reactions were run using the NextSeq 500 High-output v2 300 Cycle Kit (Illumina®, USA) on the NextSeq 500® platform (Illumina®, USA).

110 Exomic Bioinformatics

111 The quality of the FASTQ reads was analyzed (FastQC v.0.11112 http://www.bioinformatics.babraham.ac.uk/projects/fastqc/), and the samples were filtered to

eliminate low-quality readings (fastx tools v.0.13 - http://hannonlab.cshl.edu/fastx toolkit/). The 113 114 sequences were aligned with the reference genome (GRCh37) using the BWA v.0.7 tool 115 (http://bio-bwa.sourceforge.net/). The file was indexed and sorted (SAMtools v.1.2 http://sourceforge.net/projects/samtools/). Subsequently, the alignment was processed (duplicate 116 117 PCR removal) (Picard Tools v.1.129 - http://broadinstitute.github.io/picard/), and mapping quality recalibration and local realignment (GATK v.3.2 - https://www.broadinstitute.org/gatk/) 118 were performed. The results were processed to determine the variants (GATK v.3.2) from the 119 reference genome. SnpEff v.4.3t, Ensembl Variant Effect Predictor (Ensembl release 99) and 120 121 ClinVar (v.2018-10) were used for variant annotations.

122 Databank Analysis

123 Information from the databank and exomes were analyzed using descriptive statistics, 124 considering the values of allele frequencies of populations and subpopulations as the data 125 explored comparatively. Genotypic differences between sexes in the homo/hemizygous state 126 were calculated based on the premise that populations are in Hardy-Weinberg equilibrium.

127 **Results**

Analyzing the polymorphisms contained in the ACE2 locus, in addition to ten thousand base pairs upstream, we found 2266 polymorphisms, of which 199 were contained in the region 5' upstream of the gene, 85 were located in exonic regions, and the others were located in the introns.

132 Polymorphisms in exonic regions that may influence disease outcome

In the exonic region, 15 SNPs of the 85 polymorphisms found in 1000G present differences greater than 1% between some of the populations (all between exons 17 to 21). Another three polymorphisms (rs889263894, rs1027571965, rs147464721) appear mainly in the Brazilian population. Nine of these exonic polymorphisms are present in the most common isomorphs (v1 and v2) and show important differences in populational frequency (Fig 1, additional data on S1 Table).



139



144	As noted, some of these polymorphisms have very significant differences between MAFs
145	from different populations. rs35803318 is absent in Asians (virtually absent in AFR) and has an
146	average MAF of 0.05 in EUR and ABM, increased to 0.074 in BAP, and has the highest allele
147	frequency in the indigenous population among all populations (MAF=0.121).
148	On the other hand, rs4646179 is absent in indigenous, miscegenated people from the
149	Amazon, Asians and Europeans and is found in the population of southeastern Brazil and
150	Americans, with a MAF=0.023 and an even greater frequency in Africans (MAF=0.074).
151	Interestingly, rs1027571965 and rs889263894 presented allele frequencies exclusively in
152	indigenous people and are not being found in any other world population of 1000G, neither in
153	BAP nor in the Brazilian ABM database, with MAF=0.095 and 0.034, respectively.
154	rs147464721 has MAF=0.014 in the miscegenated population of the Amazon (BAP);
155	however, they did not present any allelic frequency among the indigenous population, as well as
156	in Asians and Europeans, presenting a slightly lower MAF difference when compared to AFR
157	and AMR.

158 **Polymorphisms in upstream regions that may influence disease outcome**

Differences greater than 1% of MAF in the region 10,000 base pairs 5' upstream to thegene were observed in 57 polymorphisms (Fig 2).





Among the highlights, rs5934250, with a change from G to T at approximately 5700 bp upstream to the gene, presented a difference of up to 0.47 in the AFR (0.10), AMR (0.29), EAS (0.01), EUR (0.47) and SAS (0.22) populations for the T allele. This means that the T allele is almost zero in the East Asian population, while it has a MAF in almost half of Europeans.

It is also worth mentioning that the rs2097723 SNP presents a very heterogeneous distribution, oscillating between 7% in Africans, 32% in Americans, 42% in East Asians, 28% in Europeans and 22% in South Asians.

171 Polymorphisms in intronic regions that may influence disease outcome

In intronic polymorphisms, many of them present a very relevant MAF interpopulational
difference (up to 0.46). Two of them deserve to be highlighted, rs2285666 and rs4646140,
because they are near exons.

175	It is important to mention that rs2285666 has the highest frequency of the rarest allele
176	(MAF=0.71) in the indigenous population, with very important MAF differences of 0.17 (EAS),
177	0.23 (SAS), 0.36 (BAP), and 0.37 (AMR) and an average difference of 0.48 for the others (EUR,
178	AFR and ABM). In contrast, rs4646140 has a MAF ranging from zero in Indians to 0.13 in
179	Africans through EUR, AMR, BAP, EAS, ABM and SAS (Fig 3). Furthermore, considering the
180	possibility of influence in determining isoforms v2, it is worth noting rs190614788 on intron 1
181	(with a difference of more than 0.11 between EUR and EAS).

The main findings of our study are concatenated in Table 1 (additional information in S3Table), and they are discussed below.



185 Fig 3. Minor allele frequency of the SNP intronic rs2285666 and rs4646140.

186 Table 1. ACE2 polymorphisms with the potential to influence COVID-19 and their

Population*	RS	Site	MAF*	Others (MAF)	Modification (score)	Men risk	Women risk	M/W
BAP	rs147464721	codon 351	0.014	absent	Synonymous	1.4%	0.0%	71.4
ABM, AMR, AFR	rs4646179	codon 690	0.074	absent	Synonymous	7.4%	0.5%	13.5
NAM	rs35803318	codon 749	0.121	EAS, SAS, AFR (null)	Synonymous	12.1%	1.5%	8.3
EUR	rs41303171	codon 720	0.018	absent	Asn > Asp(0.02)	1.8%	0.0%	55.6
AFR	rs147311723	codon 731	0.017	EAS, SAS, AMR, ABM (null)	Leu > Phe (0.941)	1.7%	0.0%	58.8
NAM	rs1027571965	codon 673	0.095	absent	Ala > Gly (0.045)	9.5%	0.9%	10.5
NAM	rs889263894	codon 541	0.034	absent	Lys > Ile (0.958)	3.4%	0.1%	29.4
EAS	rs182366225	3´UTR	0.018	absent	Upregulation	1.8%	0.0%	55.6
AFR	rs142017934	3´UTR	0.013	absent	Upregulation	1.3%	0.0%	76.9
NAM	rs2285666	intron	0.71	EAS (0.54) to AFR (0.17) SAS	Upregulation (Brain, Nerve)	71.0%	50.4%	1.4
AFR	rs4646140	intron	0.13	(0.085) to NAM (0)	No evidence	13.0%	1.7%	7.7
EAS	rs2097723	upstream	0.42	AMR (0.32) to AFR (0.07)	Upregulation (Brain/Nerve)	42.0%	17.6%	2.4
EUR	rs5934250	upstream	0.47	AMR (0.29) to EAS (0.01)	Downregulation (Brain, Nerve, Artery, Pituitary, Prostate)	47.0%	22.1%	2.1

187 interpopulation and inter-sex differences.

Population*: Population with greater MAF; RS: Reference SNP; Site: genic region; MAF*: minor allele
frequency of population*; Others (MAF): MAF of the other populations (or populations without MAF =
null); Modification (score): Type of molecular consequence and PolyPhen Score [22]; Men risk: based on
hemizygous genotypic frequency (q); Women risk: based at homozygous genotypic frequency (q²); M/W:
risk ratio between sexes of being a mutant genotypically.

193 **Discussion**

194 Polymorphisms of the ACE2 gene are important in the binding region of the viral particle

195 Considering the regions where the virus commonly binds to ACE2 [14,15], our results 196 point to the absence of relevant polymorphisms at these sites because many of those located in 197 coding regions have MAFs close to or less than 0.001, as well as a very low possibility of conferring any global (or population) impact on the destination of the disease. Therefore, there 198 199 does not seem to be any direct mechanism (at the sites of interaction) that could confer some 200 form of resistance or greater propensity to contagion in humans. In view of this finding, other molecular modifications with potential functional repercussions were investigated, including 1) 201 202 changes in sites distant to the viral binding locus, but which may bring about some structural 203 protein change with the potential to influence the cell-virus interaction; 2) changes in translation 204 regulation regions at 3' UTR sites, at points of interaction with miRNA; and 3) modifications in 205 transcription regulation zones in 5' upstream regions and intragenic promoters.

Polymorphisms with the potential to cause changes in the protein structure of ACE2 that may impact virus-cell interactions

The ACE2 gene is mainly composed of two isoforms with 18 or 19 exons (v1 and v2) that encode the same protein (805 amino acids) and three other smaller variants: x1, x2 and x3, which have rarely been studied [19,23]. Thus, we searched for genotypic information in these exons that could allow us to infer a disruption with the potential to culminate in impacts on the disease process.

Fifteen SNPs showed MAF differences greater than 1% among the studied populations, mainly belonging to exons 17, 18, 19, 20 and 21; the last two are terminal exons belonging only to rare isoforms [19].

216 Among these, rs41303171 is a missense SNP, causing the replacement of an asparagine (neutral amino acid) with aspartic acid (electronegative amino acid) at codon 720, which can 217 218 culminate in a conformational disorder of this protein that, directly or indirectly, can change viral interactions. This polymorphic variant C is practically exclusive to Europeans (MAF=0.018), a 219 fact corroborated by Cao and collaborators [9], mainly in British individuals (MAF=0.03). 220 221 Across Asia, this allele is not found, and is only present in a single Asian resident of the UK [18]. 222 In Brazil, ABraOM data point to MAF<0.01. Thus, the possible biological implications of this 223 change may have some clinical-epidemiological consequences in a small niche of European patients with COVID-19 when compared to other regions of the world. 224

The change from leucine to phenylalanine in codon 731 (rs147311723) results in the 225 226 exchange of two nonpolar amino acids that are structurally different (with the presence of an 227 aromatic ring in this last), which may culminate in functional modifications in the ACE2 protein, with the prediction that this will occur equally to 0.941, according to the PolyPhen algorithm 228 229 [22]. This polymorphism is absent in Asians and Europeans, with low frequency in Americans 230 and Southeast Brazilians, and it has MAF=0.017 in Africans, mainly in Nigerians, with 231 MAF=0.043 of allele A. In a study by Cao and colleagues [9], this polymorphism was described as a low frequency SNP in the 1000G database, but without any mention in the China Metabolic 232 233 Analysis Project (ChinaMAP) database, reinforcing its absence in this population group.

234	In a context focused on the Brazilian Amazon, unprecedented data showed the presence of
235	two SNPs absent in all populations of the 1000G. One of them is rs1027571965, which is
236	characterized by an exchange of G>C in exon 16, leading to a substitution of alanine for glycine
237	at codon 673 (MAF=0.095), and the other, rs889263894, an exchange of T>A in exon 13
238	(MAF=0.034), causing an alteration of lysine to isoleucine in codon 541; thus, this last SNP
239	should cause structural differences by the exchange of a polar and electropositive amino acid
240	with a hydrophobic amino acid, possibly resulting in functional changes in ACE2, with a
241	probability of 0.958 that this event will occur [22]. Both SNPs had uncertain significance until
242	now.

It is also noteworthy that in codons 351 (rs147464721), 690 (rs4646179) and 749 (rs35803318), there are just synonymous alterations, without any study of modifications of this enzyme in a genotype-dependent manner.

Polymorphisms with the potential to accentuate ACE2 gene expression or translation that may impact virus-cell interactions

Our data point to a series of polymorphisms in the region upstream of the ACE2 gene, which oscillate markedly among populations and, therefore, in individuals, although some of them may not have relevant frequency at a global level. Among them, there are rs2097723 and rs5934250, with population allele differences of up to 0.35 and 0.47, respectively.

The lower frequency allele (C) of rs2097723 has a normalized effect size (NES) of up to 0.36 [24] for increased expression of the ACE gene in brain tissue. The T allele of rs5934250 has an NES of 0.64 for lower ACE2 expression in this tissue, among others. Thus, when looking at population data, it can be inferred that, according to these two variants, populations in East Asia

would have the worst scenario with regard to increased gene expression and, Europeans orAfricans have the most favorable allelic combination for these two SNPs in the pre-genic region.

258 Another interesting observation involves the exon 19 polymorphisms, which are 259 contained in the 3' UTR region of the two most important isoforms, the canonical miRNA binding site for translation control dependent on this epigenetic mechanism [19,25]. In this 260 261 regard, observing the regions of interaction between the miRNA and nucleotide exchange sites, it 262 can be noted that rs182366225 is included in the site of attachment to the seed region of miR-140-3p.1 and miR-483-3p.2, both with a match of 7mer-1A and Context ++ score percentile of 90 263 and 74, respectively. Thus, the proportion of translation regulation that depends on these 264 miRNAs will be upregulated in the East Asian population, especially in Vietnamese and Chinese 265 266 individuals, who have an average MAF of 0.032 of the C allele, which is not observed in any 267 other population in the world [26]. A reporter assay for miR-483-3p predicted targets by Kemp 268 and collaborators [27] showed that the regulation of ACE2 is dependent on this microRNA.

269 The rs142017934 site includes the connection site for miR-610 and miR-3646, both with 270 8mer of match in ACE2 (context ++ score percentile of 98 and 77, respectively), in addition to 271 the seed regions of miR-3609 and miR-548ah-5p, both with a match of 7mer-m8 and scores of 84 272 and 74, respectively [25]. This polymorphism occurs exclusively in the population of African origin (MAF=0.013), mainly in Nigerians and individuals from Barbados with African ethnicity, 273 274 with an average of MAF=0.026. This higher frequency among people of African ethnicity 275 residing on different continents is probably due to the strong population ancestry of Barbados 276 being from West Africa, a region containing Nigeria [28].

277	Among the intronic polymorphisms, rs2285666 draws much attention because it presents
278	the highest frequency of the rarest allele (MAF=0.71) in the indigenous population, with
279	differences ranging from 0.48 to 0.17 for the other populations, with the Asians being the most
280	similar to the Indians, mainly the Chinese. The high MAF observed in the Chinese population in
281	the present study corroborates the data of Cao and collaborators ⁹ using the ChinaMAP database.
282	The substitution of C for T in intron 4, to only four nucleotides of exon 3 (located in the splicing
283	region), influences the gene expression in brain tissues and tibial nerve in some way so that the T
284	allele is related to a significant increase in the expression of ACE2 [22], thus being a determinant
285	in clinical differences that will be demonstrated in this naturally more vulnerable population.

Another intronic polymorphism, rs4646140, has no MAF in the indigenous population, reaching 0.13 in Africans, mainly in Nigerians (0.17). Some studies show the influence of these two intronic SNPs with hypertension [29,30], since there is a possibility that they will interfere in the ACE2 protein product.

In conclusion, considering this genetic aspect involving ACE2 in the complex relationship
between SARS-CoV-2 and humans, we emphasize the following:

There are genetic markers that could influence the unequal rates of aggravation and death observed between men and women. Considering the polymorphisms as harmful when in homozygosity (genotypic frequency equal to allelic frequency squared) or hemizygosity (genotypic frequency equal to allelic frequency), men would have this conditions from 1.4 to 77 times more (uniallelic) than women (Table 1). Thus, a relevant contribution to the understanding of higher mortality in males is presented, as reported in the various populations affected by COVID-19.

The rates of contagion and death fluctuate greatly; in this sense, ACE2 polymorphisms could contribute to these differences. The rs182366225 and rs2097723 polymorphisms are more frequent in the East Asian population and are potentially unfavorable to the individual, as they increase the expression of the enzyme. These allele frequencies are even higher in Chinese and Vietnamese populations. Such markers are on the order of 30% to 180% more frequently in East Asians than in other populations.

Indigenous populations from Amazon have exclusive genetic polymorphisms (rs1027571965 and rs889263894) or with higher frequencies (rs2285666 and rs35803318) than other populations. These polymorphisms are related to increased expression of the ACE2 gene in brain tissues, among others. This is an extremely relevant finding because they may influence the outcomes in these populations, whose involvement by COVID-19 was recently reported. This population group, due to its genetic peculiarity and less previous exposure to viral infections, represents a major challenge in understanding and handling this pandemic.

Africans have higher rates of three relevant polymorphisms (rs147311723, rs142017934 and rs4646140). Polymorphism in rs142017934 is exclusive to this population and can influence the translation regulation of the ACE2 gene, thus enhancing the expression of this gene, which may present an increased risk for individuals who carry this variant. However, Europeans and some Africans have a higher frequency of an allele (rs5934250) that seems to reduce the expression of ACE2 in some tissues. Individuals with this genotype would have a more protective factor against this infection.

Therefore, the study highlights the importance of this genetic factor in facilitating or restricting infection and, especially, in potential clinical manifestations and outcomes. The

17

investigation of these polymorphisms in patients affected by COVID-19, with different clinical
 conditions and outcomes, has great potential to favor understanding the behavior of this
 pandemic disease.

324 Acknowledgments

325 The authors are grateful to the Federal University of Pará (Propesp), CNPq (National

326 Council for Scientific and Technological Development) and CAPES for supporting the current

327 research; to the 1000 Genomes Database and ABraOM for making their data available to

- 328 researchers; and to all the people who, with their genetics or dedication, collaborated here to
- 329 understand an important aspect of this pandemic.

330 **References**

- 331 01. Andersen K, Rambaut A, Lipkin W, Holmes E, Garry R. The proximal origin of SARS-CoV2. Nature Medicine 2020; 26: 450-452.
- WHO. World Health Organization, member, states and regions. (acessed April 2020).
 Avalaible from: https://who.maps.arcgis.com
- 335 03. Boccia S, Ricciardi, W, Ioannidis, J. What Other Countries Can Learn From Italy During the
 336 COVID-19 Pandemic. JAMA Inter Med 2020. (accessed Abril 2020). Available from:
 337 https://jamanetwork.com/journals/jamainternalmedicine/fullarticle/2764369
 338 doi:10.1001/jamainternmed.2020.1447
- 339 04. Wang Y, Wang Y, Chen Y, Qin Q. Unique epidemiological and clinical features of the 340 emerging2019 novel coronavirus pneumonia (COVID-19) implicate special control 2020. 341 measures. J Med Virol (accessed March 2020). Available from: https://onlinelibrary.wiley.com/doi/full/10.1002/jmv.25748 doi: 10.1002/jmv.25748. 342
- 343 05. Wenham C, Smith J, Morgan R. On behalf of the Gender and COVID-19 Working Group
 344 COVID-19: the gendered impacts of the outbreak. The Lancet 2020; 395 (10227): 846-848.
- 06. Chinese Center for Disease Control and Prevention. The epidemiological characteristics of an
 outbreak of 2019 novel coronavirus diseases (COVID-19) in China. Zhonghua Liu Xing
 Bing Xue Za Zhi 2020; 17;41(2):145-151.
- 348 07. GlobalHealth50/50. Global Health 50/50. (accessed April 2020). Available from:
 349 http://globalhealth5050.org/covid19/

- 08. Kohlstedt K, Shoghi F, Muller-Esterl W, Busse R, Fleming I. CK2 phosphorylates the angiotensin-converting enzyme and regulates its retention in the endothelial cell plasma membrane. Circ. Res 2002; 91: 749–756.
- O9. Cao Y, Li L, Feng Z, et al. Comparative genetic analysis of the novel coronavirus (2019 nCoV/SARS-CoV-2) receptor ACE2 in different populations. Cell Discov 2020; 24: 6:11.
- 355 10. Zhou P, Yang X, Wang X, et al. A pneumonia outbreak associated with a new coronavirus of
 356 probable bat origin. Nature 2020; 579: 270–273.
- 11. Chen L, Li X, Chen M, Feng Y, Xiong C. The ACE2 expression in human heart indicates
 new potential mechanism of heart injury among patients infected with SARS-CoV-2.
 Cardiovasc Res 2020; Mar 30. pii: cvaa078.
- 12. Li Y, Zhou W, Yang L, You R. Physiological and pathological regulation of ACE2, the
 SARS-CoV-2 receptor. Pharmacol Res 2020; Apr 14:104833.
- 362 13. Othman H, Bouslama Z, Brandenburg J, et al. Interaction of the spike protein RBD from SARS-CoV-2 with ACE2: similarity with SARS-CoV, hot-spot analysis and effect of the 363 2020). polymorphism. (accessed March Available from: 364 receptor https://www.biorxiv.org/content/10.1101/2020.03.04.976027v3 365 .doi.org/10.1101/2020.03.04.976027 366
- 367 14. Ortega J, Serrano M, Pujol F, Rangel H.. Role of changes in SARS-coV-2 spike protein in the
 368 interaction with the human ACE2 receptor: an in silico analysis. EXCLI Journal 2020; 19:
 369 410-417.
- 15. Yan R, Zhang Y, Li Y, Xia L, Guo Y, Zhou Q. Structural basis for the recognition of SARSCoV-2 by full-length human ACE2. Science 2020; 367 (6485): 1444-1448.
- 372 16. Walls A, Park Y, Tortorici M, Wall A, McGuire A, Veesler D. Structure, Function, and
 373 Antigenicity of the SARS-CoV-2 Spike Glycoprotein. Cell 2020; pii, S0092-8674(20)30262374 2.
- 375 17. Santos N, Ribeiro-Rodrigues E, Ribeiro-Dos-Santos A, Pereira R, Gusmão L, Amorim
 376 Guerreiro J, Zago MA, Matte C, Hutz M, Santos S. Assessing individual interethnic
 377 admixture and population substructure using a 48-insertion-deletion (INSEL) ancestry378 informative marker (AIM) painel. Hum Mutat 2010; 31(2): 184-90.
- 18. IGSR. The International Genome Sample Resource. IGRS. (accessed April 2020). Available
 from: https://www.internationalgenome.org/
- 19. NCBI. ACE2 angiotensin I converting enzyme 2 [Homo sapiens (human)]. PUBMED.
 (accessed April 2020). Available from: https://www.ncbi.nlm.nih.gov/gene/59272
- 383 20. ABraOM Online Archive of Brazilian Mutations. (accessed April 2020). Available from:
 384 http://abraom.ib.usp.br/
- Sambrook J, Fritsch E, Maniatis T. Molecular cloning: a laboratory manual. New York: Cold
 Spring Harbor Laboratory Press 1989.
- 22. Adzhubei I, Jordan D, Sunyaev S. Predicting functional effect of human missense mutations
 using PolyPhen-2. Curr Protoc Hum Genet 2013; 76: 7.20.1-7.20.41.

- 389 23. Chen J, Jiang Q, Xia X, Liu K, Yu Z, Tao W, Gong Han J. Individual Variation of the SARS-
- CoV2 Receptor ACE2 Gene Expression and Regulation. Preprints 2020. (accessed March 2020). Available from: https://www.preprints.org/manuscript/202003.0191/v1
- 392 24. GTEx Portal. The Genotype-Tissue Expression (GTEx). GTEx Portal. (accessed March 2020). Available from: https://www.gtexportal.org/
- Agarwal V, Bell G, Nam J, Bartel D. Predicting effective microRNA target sites in mammalian mRNAs. Elife 2015; 4: e05005.
- 26. Cunningham F, Achuthan P, Akanni W, et al. Ensembl 2019. Nucleic Acids Res 2019; 47D,
 D745–D751.
- Kemp J, Unal H, Desnoyer R, Yue H, Bhatnagar A, Karnik S. Angiotensin II-regulated
 microRNA 483-3p directly targets multiple components of the renin-angiotensin system. J
 Mol Cell Cardiol 2014; 75, 25-39.
- 401 28. Gouveia M, Borda V, Leal T, et al. Origins, admixture dynamics and homogenization of the
 402 African gene pool in the Americas. Mol Biol Evol 2020; pii, msaa033.
- 29. Zhao Q, Gu D, Kelly T, et al. Association of Genetic Variants in the Apelin–APJ System and
 ACE2 With Blood Pressure Responses to Potassium Supplementation: The GenSalt Study.
 Am J Hypertens 2010; 23(6), 606–613.
- 406 30. Luo Y, Liu C, Guan T, et al. Association of ACE2 genetic polymorphisms with hypertension 407 related target organ damages in south Xinjiang. Hypertens Res 2019; 42(5), 681-689.

408 **Supporting information**

409 S1 Table. The minor allele frequency (MAF) of SNPs in exonic regions of ACE2 that

410 showed differences between the investigated populations. RS: Reference SNP; Ref: reference

- 411 allele; Alt: alternative allele; EUR: European; EAS: East Asian; SAS: South Asian; AMR: Ad Mixed
- 412 American; AFR: African; NAM: Native American; BAP: Brazilian Admixed Population; ABM: Online
- 413 Archive of Brazilian Mutations; GLO: Global (1000 Genomes). Bold exons are contained in common
- 414 isoforms.
- 415 S2 Table. Chromosome position, reference (RS) and minor allele of SNPs (MAF>0.01) in
- 416 ten thousand base pairs 5' upstream of ACE2. *Insertion of an Alu mobile element relative to the
- 417 reference

418 S3 Table. ACE2 polymorphisms with the potential to influence COVID-19 and their

419 **population and sex differences**. Population*: Population with greater MAF; RS: Reference SNP; Site:

420 genic region; MAF*: minor allele frequency of population*; Subpopulation alert (MAF): Subpopulation

421 with greater MAF; Others (MAF): MAF of the other populations (or populations without MAF=null);

422 Modification (score): Type of molecular consequence and PolyPhen Score²³; Hypothetical influence:

423 based on hypothetical biological influence in disease; Men risk: based on hemizygous genotypic

424 frequency (q); Women risk: based on homozygous genotypic frequency (q^2) ; M/W: risk ratio between

425 sexes to be a carrier of the minor allele only.

426 S4 Table. Tribe names and geographic coordinates of the Brazilian Amazon indigenous

427 populations enrolled in this study.