

1 **INTRAREGIONAL PROPAGATION OF COVID-19 CASES IN PARÁ, BRAZIL.**
2 **ASSESSMENT OF ISOLATION REGIME TO LOCKDOWN.**

3 Félix Lélis da Silva ^a, Javier Dias Pita ^b, Maryjane Diniz A. Gomes ^c, Andréa P. Lélis da
4 Silva ^d, Gabriel Lélis P. da Silva ^e

5
6 ^{a, b, c} Federal Institute of Science and Technology of Pará – IFPA Campus Castanhal

7 ^{a, b, b} Research Group for Biosystems Management, Modeling and Experimentation -
8 GEMAbio

9 ^d Metropolitan Regional Hospital, Specialist in Intensive Care Unit

10 ^e Estácio de Sá College of Castanhal, Law undergraduate student.

11 corresponding author (s): Félix Lélis da Silva (felix.jelis@ifpa.edu.br), Javier Dias Pita
12 (javier.pita@ifpa.edu.br), Maryjane Dias A. Gomes (maryjane.gomes@ifpa.edu.br),
13 Andréa P. Lélis da Silva (andlelis@yahoo.com.br), Gabriel Lélis P. da Silva
14 (biellellis@yahoo.com.br).

15

16 **Abstract**

17

18 Due to the high incidence of COVID-19 case numbers internationally, the World Health
19 Organization (WHO) declared a Public Health Emergency of global relevance, advising
20 countries to follow protocols to combat pandemic advance through actions that can
21 reduce spread and consequently avoid a collapse in the local health system. On March
22 18, 2020, Pará notified the first case of COVID-19. After seven weeks, the number of
23 confirmed cases reached 4,756 with 375 deaths. Knowing that infected people may be

24 asymptomatic, the disease symptomatology absence and the population's neglect of
25 isolation influence the spread, and factors such as chronic pneumonia, high age, obesity,
26 chronic kidney diseases and other comorbidities favor the mortality rate. On the other
27 hand, social isolation, quarantine and lockdown seek to contain the intraregional
28 contagion advance. This study analyzes the dynamics of COVID-19 new cases advance
29 among municipalities in the state of Pará, Brazil. The results show it took 49 days for
30 81% of the state's municipalities to register COVID-19 cases. The association between
31 social isolation, quarantine and lockdown as an action to contain the infection was
32 effective in reducing the region's new cases registration of COVID-19 in the short-term.

33

34 Keywords: Coronavirus, syndrome, contagion, advance, control.

35

36 **1. Introduction**

37

38 The COVID-19 pandemic stands out as the main global health crisis (Wu et al.,
39 2020). It started in Wuhan, China in December 2019 (Şahin, 2020; Ahmadi et al., 2020;
40 See et al., 2020). It is a respiratory infection caused by the coronaviruses family (2019-
41 nCoV) (Prata et al., 2020). Its etiological agent is Sars-CoV-2 (Saez et al., 2020; Yang
42 et al., 2020), which causes severe acute respiratory syndrome (SARS) (Solé at al.,
43 2020). The viral infection presents severe clinical symptoms such as fever, dry cough,
44 dyspnea, and pneumonia (Coccia, 2020; Wu et al., 2020) and can cause the death of the
45 infected (Waldecy; David; Wainesten; 2020). Flu is one of the main causes of illnesses
46 and death in the world (Mertz et al., 2013).

47 Epidemiological data associated with COVID-19 infection has shown different
48 dynamics in several countries (Marson; Ortega, 2020). Several studies have shown the

49 reverse effect of rising temperatures and the confirmed number of new cases (Zhu and
50 Xie, 2020; Wu et al. 2020). Brazil, a country with a tropical climate, with an average
51 annual temperature ranging from 16 to 27.4 °C, modeled results show a negative effect
52 on the linear relationship of temperature with the new confirmed cases number (Prata et
53 al., 2020).

54 Several factors have contributed to pandemic advance in Brazil, as the country,
55 in addition to having a deficiency in the number of doctors per inhabitant, the number of
56 ICU beds and ventilators available for urgent and emergency cases, presents several risk
57 groups, such as: elderly over 60 years old, people with prognostic comorbidity,
58 indigenous people and population's great genetic variation (Marson; Ortega, 2020).

59 Government actions to mitigate the COVID-19 epidemic curve have been
60 adopted in several countries. In Spain, according to Saez (2020), measures of social
61 distancing led to a cases curve flattening, after the first few days the cumulative change
62 rate in new cases decreased by an average of 3.059 percentage points daily. In Pará
63 state, high mortality and lethality rates were reported even with social isolation, which
64 raised the curve of new cases and directly affected hospital care due to overcrowding by
65 infected people. On May 6, 2020, a more extreme control policy was decreed by the
66 State of Pará government aimed at containing the pandemic advance, setting the
67 lockdown model in the metropolitan region and neighboring municipalities.

68 People's traffic control measures are fundamental in controlling the pandemic
69 progress. According to Kucharski et al. (2020), the cases' reduction of COVID-19 in
70 Wuhan observed in February 2020, coincided with traveling control measures adopted
71 in the region.

72 Therefore, it has been suggested that environmental factors and social control
73 through social distancing tend to control the new cases and deaths numbers due to


74 COVID-19. As well, the deficiency in the health system associated with the reduced
75 number of intensive-care physicians, low numbers of mechanical ventilators, the
76 minimum number of available beds, and the specific medication absence tend to
77 compromise the diagnosed population care. It is questioned: What are the dynamics of
78 new community cases and mortality records evolution due to COVID-19 in subtropical
79 regions with temperatures between 20 and 35 °C, with social distancing by quarantine
80 and lockdown adoption?

81 **2. Materials and methods**

82 *2.1 Study area*

83 The study was carried out in 114 affected municipalities among the 144 existing
84 in the state of Pará, North region, Brazil (Fig.1). The state has a territorial extension of
85 247,689,515 km² and an estimated population of 8,602,285 people, with 8,191,559
86 residents in urban areas and 2,389,492 in rural areas, with a population density of 6.07
87 inhabitants/km² and an average Human Development Index of (HDI = 0.698) (BIGS,
88 2020).

89



90
91 **Fig. 1.** State of Pará, Northern Brazil and its 144 municipalities.

92 *2.2. Data collection*

93 The cumulative daily number of deaths by age group and sex of COVID-19
94 cases in 114 municipalities in the state of Pará, with a record of infection, were obtained
95 through the daily monitoring of technical bulletins provided by the State of Pará Public
96 Health Secretary (SHSP, 2020). General data for Brazil and its northern region were
97 obtained from the Secretary of Health Surveillance of the Ministry of Health (BRAZIL,
98 2020).

99
100 *2.3. Statistical analysis*

101 Since understanding the dynamics of infection spread can favor more effective
102 public control policies and allow control of transmission in new regions, an exploratory
103 data analysis (EDA) was carried out, with numerical variables described using means,

104 standard deviations, coefficient of variation, distributions and Pareto. Weekly maps
105 were built for the new accumulated cases, in the period of 49 days, to assess the spread
106 of the COVID-19 pandemic in the municipalities of the state of Pará. It was used a
107 classification method using the Jenks algorithm based on the Absolute Deviations over
108 the Median of the Classes.

109

110 **3. Results and Discussion**

111 *3.1 Descriptive analysis*

112


113 In Brazil, on May 5, 2020, 114,715 cases of COVID-19 were registered and
114 7,921 deaths, with a lethality rate of 6.8%, the Southeast and Northeast regions are the
115 most affected counting around 75% of the registered cases (Fig 2B). The most affected
116 regions in the country are the Southeast (64,756; 44.6%), followed by the Northeast
117 (45,724; 31.5%) and North (23,207; 16.0%) (BRASIL, 2020). In the North, the state of
118 Amazonas had the largest number of confirmed infected, 10,727 cases and disease
119 lethality around 874 registered deaths The country on May 6, 2020, registered 114,715
120 cases and 7,921 deaths.

121 In Northern Brazil, the federative units most affected on May 5, 2020, in terms
122 of the spatial distribution of registered cases of COVID-19 with an incidence and
123 mortality rate per 1,000,000 inhabitants, were Amazonas (2327.5 and 251.7) and Pará
124 (627.4 and 49.5). The capital of Pará has the highest incidence (1,816.4/1,000,000
125 inhabitants) Fig. 3A) and mortality (240/1,000,000 inhabitants) with a mortality rate of
126 9.9% (Fig. 3B).

127 Predictive studies in Brazil report smaller records of new cases of COVID-19 at
128 temperatures around 25.8°C, reducing the behavior of curve growth (Prata et al., 2020).

129 Since SARS-CoV-2 may be vulnerable to fluctuations in environmental conditions
130 similar to other coronaviruses (Le et al., 2020). It is important to note that
131 climatological factors variability can interfere with the curve behavior even at higher
132 temperatures.

133 The northern region of Brazil has an equatorial climate ranging from humid to
134 semi-arid, with temperatures ranging from 20 to 35 °C. In the state of Pará, the
135 temperature presents spatial and seasonal homogeneity, with an average variation of 25
136 °C to 35 °C. In the region, there are two distinct periods of temperature ranges,
137 classified as rainier between December to May and less rainy between June to
138 November, according to rainfall variation that occurs in the Amazon. The metropolitan
139 region of Belém city in the period between January 1, 2020, to May 5, 2020, presented
140 temperature variation between 24 °C to 31 °C and relative humidity between 70 to 80%.
141 In this period, the number of new cases and mortality due to COVID-19 has risen since
142 the first case notification on March 18, 2020, with the first death recorded on April 1,
143 2020 (Fig.2).



144

145 **Fig. 2.** Temperature behavior (°C), rainfall behavior, 1st confirmed case and 1st death
146 registration by COVID-19 in the state of Pará.

147

148 The North region was in 3rd place in cases number with approximately 16% of
149 the confirmed cases on May 5, 2020. In the state of Pará, the outbreak of COVID-19
150 registered a record of 4,756 cases with an incidence rate of 788 cases in 49 days after
151 the first infection confirmation. The first confirmed case of the disease was notified on
152 March 18, 2020, in Belém city, capital of the state. These records occurred when the
153 average temperature of the region, for the period considered to be the rainiest in the
154 region, was 27.9 °C and the air relative humidity ranging from 70 to 80%. The first
155 notification of death occurred on April 1, 2020, and since then the curve of new cases
156 and deaths incidence have been frequent and registered with high rates in the region.

157 *A priori* it is suggested that there is no inverse relationship between ambient
158 temperature and new cases of COVID-19 in Pará in the period considered as the rainiest
159 in the Amazon region. Therefore, there is no evidence to suppose that in regions with
160 warm climates, low relative humidity and high rainfall variation, the cases number is
161 lower compared to regions with moderate and/or cold climates. Similar results were
162 obtained by Jahangiri, Jarangiri, Njafgholipour (2020) in the transmission rate
163 evaluation of the new coronavirus in different provinces of Iran.

164 Fig. 3 summarizes the overall cases number compared to diagnosed population
165 sex, the cumulative daily descriptive statistics of newly notified cases of COVID-19,
166 and the age group distribution behavior of diagnosed population with the infection since
167 the first disease case notified in the State of Pará.

168 The analysis was carried out after seven weeks of the first COVID-19 case
169 recorded in Pará, being 4,756 cases reported between March 18 to May 5, 2020. Of the

170 total, 2,420 (50.9%) diagnosed cases were men and had an average age of 45.6 years.
171 The women were 2,320 (48.8%) infected cases with an average age of 47.4 years. It is
172 reported that 15 registered cases did not present notification regarding gender and were
173 considered as unidentified (Fig. 3A). Similar results of higher COVID-19 cases
174 prevalence in men were obtained by Chen et al. (2020) and Nikpouraghdam et al.,
175 (2020).

176 In Pará, 75% were under 58 years old. Only 25% of infected cases were elderly
177 and 1.6% of those diagnosed infected were children from 0 to 10 years old. Only 26
178 (0.5%) registered cases had no age identified and were considered in terms of
179 registration as “unidentified” (Fig. 3B).

180 In terms of the sample of the infected population age range, negative asymmetric
181 behavior was identified for distribution, with ages concentration around the median of
182 34 years old, with 80% of cases diagnosed aged between 20 and 60 years old (Fig. 3C).
183 In Iran, the cases concentration was between 30 and 70 years old for 79.1% of COVID-
184 19 records and approximately 39% for the elderly (Nikpouraghdam et al., 2020).


185 Concerned with the new cases and mortality rates, the government of Pará
186 established a decree for lockdown in the capital and additional nine municipalities to
187 increase the social isolation index, reducing the disease spread and reducing the new
188 cases registration number of COVID-19.

189 The restrictive measure was initially focused on forcing social isolation in the
190 municipalities of Belém, Ananindeua, Marituba, Benevides, Santa Bárbara do Pará,
191 Santa Izabel do Pará, Castanhal, which form the Belém Metropolitan Region (BMR)
192 and inland towns, such as Santo Antonio do Tauá, Vigia de Nazaré and Breves. The
193 lockdown arises when the state of Pará registered a COVID-19 rate in the order of
194 51/100,000 inhabitants, higher than the national rate. The municipalities affected by the


195 restriction presented rates of 75/100,000 inhabitants, higher than that registered by the

196 state.

197



199



200

201 **Fig. 3.** Exploratory statistical analysis of the confirmed cases accumulated of COVID-
 202 19 by sex (A), age group (B) and Pareto distribution of ages (C) of those infected, Pará-
 203 Brazil.


204 Despite the restrictions having focused on only 10 municipal regions, it is worth
 205 mentioning that when the lockdown was set, more than 70% of the municipalities
 206 forming the state of Pará already had notifications of COVID-19 cases.

207 The regional disease expansion and the high registration rates in the BMR have
 208 compromised the structure of the state's public and private healthcare systems due to the
 209 high demand for basic care services and the high need for more complex services
 210 involving hospitalizations and intubation of patients with aggravated cases.

211 In the Pará state, severe cases of COVID-19 and mortality records are related to
 212 several comorbidities, the most commons are associated with heart disease, diabetes,
 213 kidney disease, pneumonia, immunodeficiency, asthma, obesity, neurological disease,
 214 hematological disease and illness hepatic; the most frequent ones being associated with
 215 heart disease and diabetes (Fig. 4).

216 According to Chen et al. (2020), patients infected with COVID-19 tend to have
 217 associated comorbidities with one or more chronic diseases. In Iran, patients diagnosed
 218 with COVID-19 had a strong association with chronic diseases such as diabetes, chronic

219 respiratory diseases, hypertension, cardiovascular diseases, kidney diseases and cancer,
220 these comorbidities were also observed in cases of deaths (Nikpouraghdam et al. 2020).
221 Similar results were verified in infected people in Wuhan, China (Wu et al., 2020),
222 corroborating with the results occurred in this study region.
223



224

225 **Fig. 4.** Common comorbidities in diagnosed cases of COVID-19 in the state of Pará,
226 from March 18 to May 5, 2020.

227

228 These factors associated with the precarious service capacity of hospitals and
229 emergency care units (ECU), the insufficient health professionals staff, lack of rapid
230 tests to identify infected patients, contributed to COVID-19 advance in the region.
231 Associated with these problems, there is still an insufficient hospital beds number
232 available according to the State Department of Public Health of Pará. There are only
233 249 ICU beds available for adults care in the pandemic for a population of 7,581,051
234 inhabitants, which represents a ratio of 1/30,445.98 bed/inhabitants, such fact leads the

235 system to collapse in a short period during a pandemic, contributing to the disease and
236 the number of deaths progress. Collapsing is already experienced in pediatric clinic beds
237 (Table 1).

238 Due to the risk of COVID-19 infected people promoting the disease advance
239 through community transmission to more remote regions of the state, it is essential to
240 understand the dynamics of disease new cases to enable decision-making regarding
241 control, prevention, treatment of infected, identify groups of risks and enable decision-
242 making on resources allocation and allow better planning on the health system in
243 difficult times.

244

245 **Table 1:** Availability and occupation of exclusive hospital beds for COVID-19 in the
246 public healthcare system in the state of Pará.

Hospital bed type	Total	Available	% Occupation
Pediatric	1109	497	55.2
Pediatric clinical	8	8	0.0
adult ICU	249	37	85.2
Pediatric ICU	7	4	42.9
Intermediate unit	26	25	3.9
Neonatal ICU	4	1	75.0

Source: Secretary of Public Health of Pará.

<https://portalarquivos.saude.gov.br/images/pdf/2020/May/09/2020-05-06-BEE15-Boletim-do-COE.pdf>


247

248 Results show that a week after the first notified case, only 0.82% of the
249 municipalities had infected people, in the second week of contagion they already had
250 5.56%, in the third 15.3%, in the fourth 24.3%, in the fifth 46.5%, in the sixth 69.4%
251 and the seventh week after the first notified case, there were already 81.8% of the
252 municipalities in the state of Pará with notified COVID-19 cases (Fig.7). The highest
253 concentration of registered cases was in the Metropolitan Region of Belém, constituted

254 by the capital Belém, Ananindeua, Marituba, Benevides, Santa Bárbara do Pará, Santa
 255 Izabel do Pará, Castanhal.

256 The measure of containment by social isolation in the region did not have an
 257 expected effect on the curve of new cases for the period under study, mainly due to the
 258 low rate of population adherence. However, the lockdown instituted on May 5, 2020,
 259 gave positive percentage results in the short-term on new cases registration in the
 260 following week of decreed intervention, with a 10.07% reduction compared to the
 261 accumulated record in the week previous epidemiologic (Fig. 5). In the long-term, the
 262 reduction of the cases may be more significant, given the greater respect and adherence
 263 of the population to isolation policies.

264



265

266 **Fig. 5.** Propagation of notified COVID-19 cases in the municipalities of the state of
 267 Pará, from March 18 to May 5, 2020, and the 1st post-lockdown period.


268

269 Efforts to prevent infected people from reaching more remote regions are
 270 important to stop disease transmission and prevent new cases (Coccia, 2020). To

271 understand the dynamics of COVID-19 cases advance to remote regions from the
272 capital, 7 maps were adjusted and show the weekly evolution of the accumulated cases
273 registered in the municipalities of the state (Figures 6 to 12).

274 The municipalities that registered cases of infection in the first week were Belém
275 (code 19) with 2 cases and Marabá (code 67) with 1 notified case (Fig. 6). The
276 municipalities that did not present COVID-19 case records in the seven weeks after the
277 first case were: Abel Figueiredo (code 2), Aveiro (code 14), Belterra (code 20), Brasil
278 Novo (code 25), Brejo Grande do Araguaia (code 26), Cachoeira do Arari (code 30),
279 Cumaru do Norte (code 41), Curionópolis (code 42), Eldorado dos Carajas (code 47),
280 Faro (code 48), Araguaia Forest (code 49), Gurupá (code 52), Jacareacanga (code 60),
281 Mojuí dos Campos (code 74), Piçarra (code 93), Plates (code 94), Rio Maria (code 102),
282 Rurópolis (code 104), Santa Luzia do Pará (code 112), Santa Maria das Barreiras (code
283 111), Santana do Araguaia (code 113) Francisco do Pará (code 121), Sapucaia (code
284 128), Soure (code 130), Trairão (code 136) and Vitória do Xingu (code 120) (Fig.12).


285




286

287 **Fig. 6.** Propagation of new cases of COVID-19 accumulated in the municipalities of the
288 state of Pará in the **1st week** after registration of the first case in the region.


289



291 **Fig. 7.** Propagation of new cases of COVID-19 accumulated in the municipalities of the
 292 state of Pará in the **2nd week** after registration of the first case in the region.




294 **Fig. 8.** Propagation of new cases of COVID-19 accumulated in the municipalities of the
 295 state of Pará in the **3rd week** after registration of the first case in the region.




296

297 **Fig. 9.** Propagation of new cases of COVID-19 accumulated in the municipalities of the
298 state of Pará in the 4th week after registration of the first case in the region.




299

300 **Fig. 10.** Propagation of new cases of COVID-19 accumulated in the municipalities of
301 the state of Pará in the 5th week after registration of the first case in the region.



302

303 **Fig. 11.** Propagation of new cases of COVID-19 accumulated in the municipalities of
 304 the state of Pará in the 6th week after registration of the first case in the region.



305

306 **Fig. 12.** Propagation of new cases of COVID-19 accumulated in the municipalities of
 307 the state of Pará in the 7th week after registration of the first case in the region.

308 **4. Conclusions**

309

310 Several factors can influence the spread of the virus and consequently influence
311 the rate of incidence and mortality. In this research, the incidence rate and mortalities in
312 Brazil and the North region were analyzed, as well as the monitoring of dynamics of the
313 COVID-19 progress for the municipalities of the state of Pará. The results indicated that
314 49 days were necessary to 81% out of 144 municipalities over an area of approximately
315 1,248,000 km² were affected by COVID-19 cases. Social isolation was not efficient to
316 contain new cases advance, due to low population adherence. The social isolation and
317 quarantine associated with the adoption of a strict measure of population circulation, the
318 “Lockdown”, and the mandatory use of masks in public environments were effective in
319 reducing new cases of COVID-19 registration in the short-term.

320

321 **Conflicts of Interest Statement**

322 The authors declare no conflicts of interest

323

324 **Acknowledgments**

325

326 This study was prepared with support from the Research Group for Biosystems
327 Management, Modeling and Experimentation - GEMABio.

328

329

330

331

332

333 **References**

334

335 Ahmadi, M., Sharifi, A., Dorosti, S., Ghouschi, S. J., Ghanbari, N. Investigation of
336 effective climatology parameters on COVID-19 outbreak in Iran. Science of The Total
337 Environment. 2020. v.729. DOI. <https://doi.org/10.1016/j.scitotenv.2020.138705>.

338

339 Brazil. Ministry of Health. Accumulated COVID-19 cases and deaths by confirmation
340 date. Available at: <https://covid.saude.gov.br/>. Access date: May 9 2020.

341

342 Coccia, M. Factors determining the diffusion of COVID-19 and suggested strategy to
343 prevent future accelerated viral infectivity similar to COVID, **Science of the Total**
344 **Environment**. 2020. <https://doi.org/10.1016/j.scitotenv.2020.138474>.

345

346 Mertz D., Kim T. H., Johnstone J., Lam P. P., Kuster S. P., Fadel S. A. et al.,
347 Populations at risk for severe or complicated influenza illness: systematic review and
348 meta-analysis, *BMJ*. 2013. DOI. <https://doi.org/10.1136/bmj.f5061>.

349

350 BIGS. Brazilian Institute of Geography and Statistics. Populations. Available at:
351 <https://cidades.ibge.gov.br/brasil/pa/panorama>. Date accessed: 10 May. 2020.

352

353 Jahangiri, M., Jahangiri, Milad., Najafgholipour, M. The sensitivity and specificity
354 analyses of ambient temperature and population size on the transmission rate of the
355 novel coronavirus (COVID-19) in different provinces of Iran. 2020. *Science of The*
356 *Total. Environment*. v.728, DOI. <https://doi.org/10.1016/j.scitotenv.2020.138872>.

357

358 Kucharski. A. J., Russell. T. W., Diamond. C., Liu. Y., Edmunds. J., Funk. S.,
359 et al. Early dynamics of transmission and control of COVID-19: a
360 mathematical modelling study. *Lancet Infect Dis.* 2020. DOI.
361 [https://doi.org/10.1016/S1473-3099\(20\)30144-4](https://doi.org/10.1016/S1473-3099(20)30144-4).

362

363 Le. N. K., Parikh. A. V., Brooks. J., Gardellini. J. P., Izurieta. T. R., 2020. Ecological
364 and health infrastructure factors affecting the transmission and mortality of COVID-19
365 *BMC Infect. Dis.* <https://doi.org/10.21203/rs.3.rs-19504/v1>.

366

367 Marson, F. A. L., Ortega. M. M. COVID-19 in Brazil. *Pneumology.* 2020. DOI.
368 <https://doi.org/10.1016/j.pulmoe.2020.04.008>.

369

370 Nikpouraghdam et al. Epidemiological characteristics of coronavirus disease (COVID-
371 19) patients in IRAN: A single center study. 2020. *Journal of Clinical Virology.* 2019.
372 DOI. <https://doi.org/10.1016/j.jcv.2020.104378>.

373

374 SHSP. Secretary of Health of the State of Pará (SHSP). Coronavirus in Pará. Available
375 at: [https://portalarquivos.saude.gov.br/images/pdf/2020/May/09/2020-05-06-BEE15-
376 Boletim-do-COE.pdf](https://portalarquivos.saude.gov.br/images/pdf/2020/May/09/2020-05-06-BEE15-Boletim-do-COE.pdf). Date accessed: 10 May. 2020.

377

378 Prata, D. N., Rodrigues W., Bermejo, P. H. Temperature significantly changes COVID-
379 19 transmission in (sub) tropical cities of Brazil. 2020. *Science of the Total
380 Environment.* v.729. 138862. DOI. <https://doi.org/10.1016/j.scitotenv.2020.138862>.

381

382 Saez, M., Tobias, A., Varga, D., Barceló, M. A. Effectiveness of the measures to
383 flatten the epidemic curve of COVID-19. The case of Spain. *Science of the Total*
384 *Environment*. 2020. v. 727. 138761. DOI.
385 <https://doi.org/10.1016/j.scitotenv.2020.138761>.

386

387 Şahin, M. Impact of weather on COVID-19 pandemic in Turkey. *Science of the Total*
388 *Environment*. 2020. v.720. DOI. <https://doi.org/10.1016/j.scitotenv.2020.138810>.

389

390 See. K. C., Liew, S.M., David C. E. N., et. al., COVID-19: Four Paediatric Cases in
391 Malaysia. 2020. *International Journal of Infectious Diseases*. v. 94, p. 125–127. DOI.
392 <https://doi.org/10.1016/j.ijid.2020.03.049>

393

394 Solé et al. Guidance for the care of neuromuscular patients during the COVID-19
395 pandemic outbreak from the French Rare Health Care for Neuromuscular Diseases
396 Network, *Revue Neurologique*. 2020. DOI.
397 <https://doi.org/10.1016/j.neurol.2020.04.004>.

398

399 Waldecy. R., David. N. P., Wainesten. C. Regional determinants of the expansion of
400 covid-19 in Brazil. 2020, MedRxiv. doi: <https://doi.org/10.1101/2020.04.13.20063925>

401

402 Yang, Y., Peng, F., Wang, R., Guan, K., Jiang, T., Xu, G., Sun, J., Chang, C. The
403 deadly coronaviruses: The 2003 SARS pandemic and the 2020 novel coronavirus
404 epidemic in China. 2020. *J. Autoimmun.* 102434.
405 <https://doi.org/10.1016/j.jaut.2020.102434>

406

407 Wu Y., Jing W., Liu J., Ma Q., Yuan J., Wang Y., Du M., Liu M. Effects of temperature
408 and humidity on the daily new cases and new deaths of COVID-19 in 166 countries.
409 2020. Science of the Total Environment. v.729. 139051. DOI:
410 <https://doi.org/10.1016/j.scitotenv.2020.139051>.

411

412 Zhu Y., Xie J., 2020. Association between ambient temperature and COVID-19
413 infection in 122 cities from China. 2020. Sci. Total Environ. 138201. DOI.
414 <https://doi.org/10.1016/j.scitotenv.2020.138201>.

415

416