

1 **Risk factors and protective factors of mental health**
2 **during COVID-19 outbreak and lockdown in adult**
3 **Indian population- A cross-sectional study**

4

5 Saikarthik Jayakumar¹, Saraswathi Ilango², Siva Thirusangu³

6

7

8 ¹ Assistant Professor of Anatomy, Department of Basic Medical Sciences, Majmaah
9 University, Saudi Arabia

10 ² Assistant Professor of Physiology, Madha Medical College and Research Institute,
11 Chennai, India and PhD Research Scholar, Bharath Institute of Higher Education and
12 Research, Chennai, India

13 ³Tutor in Anatomy, Sri Ramachandra Medical College and Research Centre, India

14

15

16

17 Corresponding Author:

18 Dr Saikarthik Jayakumar

19 Email: s.jaya@mu.edu.sa

20 ¶ All authors contributed equally to this research work.

21 No conflict of Interest

22 Abstract

23 **Background:** The novel Corona virus has derailed the entire world and various
24 steps have been taken by the health authorities to tackle this pandemic. Nationwide
25 lockdown has been imposed to control the spread of COVID-19 outbreak in India,
26 which could have psychological impact on the population. **Aim:** Our study aims to
27 study the effect of the COVID-19 outbreak & subsequent lockdown on mental health
28 status of adult Indian population along with identifying the high-risk groups.
29 **Methodology:** An online survey was conducted during 3rd phase of lockdown
30 gathering details about sociodemographic variables, practice of precautionary
31 measures, awareness and concerns regarding COVID-19 and mental health status
32 of the participants through DASS21 questionnaire from 873 adults. **Results:** The
33 prevalence of depression, anxiety and stress were 18.56%, 25.66%, and 21.99%
34 respectively including higher number of participants with mild depression (15.1%)
35 and stress (14.5%) and moderate anxiety (16.3%). Female gender, age <25 years,
36 unemployment, self-business, employed in private sector, lack of formal education,
37 larger household size, parenthood (>2 kids) were associated with increased
38 likelihood of negative mental health. Confidence in physician's ability to diagnose
39 COVID-19 infection, decreased self-perceived likelihood of contracting COVID-19,
40 lesser frequency of checking for information on COVID-19 and satisfaction of
41 information received were protective against negative mental health. **Conclusion:**
42 This landmark study identified the protective and risk factors of mental health during
43 COVID-19 pandemic, to help authorities and mental health workers to strategize and
44 deliver interventional methods to maintain psychosocial wellbeing of the population.

45 **Keywords:** COVID-19, lockdown, mental health, risk factors, protective factors,
46 depression, anxiety, stress, India

47 **Introduction**

48 COVID-19 outbreak, caused by a novel corona virus, SARS-CoV-2, which originated
49 from China, has spread worldwide, earning the pandemic status by WHO on March
50 11, 2020 [1]. As of the 1st week of June, 2020, India has emerged as the fifth hardest
51 hit country with 247,000 confirmed cases of COVID-19 and 7000 deaths [2].
52 Droplets, contact with immediate environment around infected persons including
53 direct or indirect and airborne in specific circumstances are the main proposed
54 routes of transmission [3].

55 As of now, there are no specific medicines or vaccines available for COVID-19. To
56 tackle this, the Government of India has opted for nationwide lockdown and
57 emphasized on social distancing. Though quarantine and lockdown help containing
58 the spread of infection, it is also accompanied by potential psychological distress in
59 the population. Isolation, fear of contracting the disease, confusion created by
60 rumours, financial strain, apprehension regarding job security, boredom, frustrations,
61 lack of freedom and space due to restrictions, alcohol withdrawal, concerns for the
62 family members that occur during lockdown period could affect the mental health of
63 the population to varying degrees.

64 Studies conducted during earlier epidemics like SARS, equine influenza, Ebola have
65 noted that there was increased psychological distress due to the epidemic and
66 quarantine [4, 5]. Similarly, studies conducted recently in other countries like China,
67 Italy, Iran have noted increased prevalence of mental health disorders like
68 depression, anxiety, stress and sleep disturbances during COVID-19 outbreak [6-8].

69 A recent survey conducted on March 2020 in India, has found that more than three
70 fourth of the study participants had self-perceived need for help for their mental well-
71 being [9].

72 Till date, very few studies have been conducted on the psychological impact of
73 COVID-19 outbreak & lockdown in India, which either focused on specific areas like
74 perceived mental health care need [9] and effect of gender and marital status [10],
75 or conducted on specific population like healthcare workers [11] and pharmacy
76 students [12]. However, there are no studies conducted to assess the impact of
77 COVID-19 outbreak & lockdown on the mental health status of the general
78 population of India with emphasis on the risk factors and protective factors. Our
79 study aims to assess the prevalence of affective components of mental health viz.
80 depression, anxiety and stress along with identifying the high-risk group of
81 population. We believe, our study would help the authorities and mental health
82 professionals in strategizing and delivering mental healthcare to the population
83 targeting on the high-risk group and help maintaining the psychosocial well-being of
84 the Indian population.

85 **Methodology**

86 **Study design and participants**

87 The study was a cross sectional survey conducted through an online survey
88 platform. The invite link to attend the survey was distributed in social networks like
89 Facebook, WhatsApp and Telegram. Only adult (age above 18 years), Indian
90 residents were invited. The objective of the study was explained, and the consented
91 participants filled out the survey and could quit the survey as and when needed. The

92 entire survey was in English and a tentative average time duration needed to fill out
93 the survey was mentioned beforehand (15-20 minutes). Ethical approval was
94 obtained from Institutional Ethics Committee on Fasttrack basis and the anonymity of
95 the participants was maintained.

96 **Survey development and Data collection**

97 The survey included a self-administered questionnaire which was developed after
98 extensive literature survey and included questions pertaining to sociodemographic
99 variables and COVID-19 outbreak & lockdown related variables. Snowball sampling
100 method was used, and the data collected between May 5th to 14th, 2020 during the
101 third phase of lockdown in India was taken for this study. The data collected were
102 sociodemographic variables, COVID-19 outbreak & lockdown related variables and
103 mental health status of the participants.

104 **Sociodemographic variables**

105 Sociodemographic variables included gender, age, educational status, employment
106 status, marital status, monthly income, parental status and household size.

107 **COVID-19 outbreak & lockdown related variables**

108 COVID-19 outbreak & lockdown related variables included (a) practice of personal
109 precautionary measures; wearing masks and gloves in public places, frequency of
110 hand washing with soap or sanitizer per day, The participants were asked if they or
111 any of their peers tested for COVID-19 (b) awareness and knowledge regarding
112 COVID-19 pandemic; route of transmission, means of gathering information,
113 frequency of checking for information and level of satisfaction of the attained
114 information, (c) personal concerns regarding the outbreak; level of confidence in the

115 physician to diagnose COVID-19 infection, self-perceived likelihood of contracting
116 COVID-19 and surviving if contracted with COVID-19 and concerns for family
117 members to contract the infection.

118 **Psychological status of the participants**

119 Affective component of mental health of the participants viz depression, anxiety and
120 stress were assessed using Lovibond and Lovibond's short version of the
121 Depression Anxiety Stress Scale 21 (DASS21). DASS21 is a reliable instrument
122 used in clinical and nonclinical samples which can measure and differentiate
123 between the three negative emotional states [13, 14]. The sub-scores for depression,
124 anxiety and stress were summed up and categorized into "normal", "mild",
125 "moderate", "severe" and "extremely severe". Cut-off score of ≥ 10 for depression, ≥ 8
126 for anxiety and ≥ 15 for stress were considered to be having the aforesaid disorders
127 [15].

128 **Statistical analysis**

129 Analysis was performed using SPSS V.26.0, IBM, New York, USA. Descriptive
130 analysis was performed for all variables. Depression, anxiety and stress scores were
131 expressed as mean and SD. Multicollinearity was checked between independent
132 variables and the variance inflation factor (VIF) was found to be less than 3.

133 To explore potential predictors for depression, anxiety and stress, binomial logistic
134 regression analysis of each independent variable was performed separately, and the
135 results were expressed as crude odds ratio (cOR) with 95% confidence interval (CI)
136 and P value. This was followed by Multivariate binomial logistic regression analysis
137 using 'stepwise forward LR' technique, which included independent variables which

138 were found to be significant ($P < 0.25$, Hosmer-Lemeshow recommendation) by
139 univariate analysis. The regression analysis was performed in two blocks,
140 sociodemographic variables block, and COVID-19 outbreak & lockdown related
141 variables block. The latter block was explored after controlling for the significant
142 sociodemographic factors. The results were expressed as Wald test value, adjusted
143 odds ratio (aOR) with 95% confidence interval and P value ($P < 0.05$ was considered
144 statistically significant).

145 **Results and Discussion**

146 **Descriptive characteristics of the study population**

147 968 responses were received, out of which 64 responses were incomplete, 25
148 respondents were underage, and 6 respondents were of a different nationality and
149 were hence excluded. The final sample size was 873. We had an almost even
150 participation from males (54.1%) and females (45.9%) and majority of the
151 participants were of age group 18 to 45 years (85.1%) with the average age of
152 33.6 ± 12.15 years. The descriptive statistics of the study population is shown in Table
153 1.

154 **Table 1: Descriptive statistics of the study population**

Variables	Subgroups	Number of participants	%
Sociodemographic variables			
Gender	Male	473	54.1
	Female	400	45.9
Age (years)	18 to 25	276	31.6
	26 to 35	225	22.7


	36 to 45	269	30.8
	46 to 55	40	4.5
	56 & above	63	7.2
Educational status	None	28	3.2
	Higher secondary school	261	29.8
	Bachelor's degree	413	47.3
	Master's degree	147	16.8
	Doctorate degree	24	2.7
Marital status	Single	388	44.4
	Married	469	53.7
	Widowed/ separated	16	1.9
Employment status	Student	259	29.7
	Employed – Government	80	9.2
	Employed – private	309	35.4
	Self-business	125	14.3
	Unemployed	100	11.4
Monthly income (INR)	10,000 and less	307	35.1
	10,001-20,000	42	5.4
	20,001-30,000	31	3.5
	30,001-40,000	75	8.5
	40,001-50,000	59	6.7
	50,001-100,000	207	23.3
	100,000 & above	152	17.4
Parental status	No kid	406	46.5
	1 kid	141	16.1
	2 kids	254	29.09
	3 or more kids	72	8.2
Household size	1 member	12	1.3

	2 members	125	14.3
	3-6 members	708	81.2
	More than 6 members	28	3.2
Practice of personal precautionary measures during COVID-19 outbreak and lockdown			
How often do you wear mask and gloves while being outside in public places?	No	103	11.8
	Sometimes	109	12.4
	Yes	661	75.8
How often do you wash your hands with soap or hand sanitizer per day?	less than 5 times	254	29.1
	5 to 10 times	308	35.2
	10 to 15 times	227	26.1
	More than 15 times	84	9.6
Awareness and Knowledge about COVID-19 outbreak			
Route of transmission			
Contact with infected person	No	193	22.1
	Yes	680	77.9
Droplets	No	252	28.9
	Yes	621	71.1
Airborne	No	676	77.4
	Yes	197	22.6
Contact with contaminated objects	No	282	32.3
	Yes	591	67.7
Through food and water	No	816	93.5
	Yes	57	6.5
Pet animals	No	853	97.7
	Yes	20	2.3
Are you aware of the	Yes	843	96.5

increase in number of COVID-19 cases in India?	No	30	3.5
Source of daily information regarding COVID-19 outbreak			
Internet	No	227	26
	Yes	646	74
TV	No	164	18.8
	Yes	709	81.2
Friends and relatives	No	569	65.1
	Yes	304	34.9
MOH messages	No	800	91.6
	Yes	73	8.4
Radio	No	869	99.5
	Yes	4	0.5
Other sources (Newspaper, magazines etc.)	No	821	94
	Yes	52	6
Are you satisfied with the information received daily regarding COVID-19 outbreak?	Highly satisfied	47	5.4
	Satisfied	575	65.9
	Not satisfied	106	12.1
	Highly not satisfied	34	3.9
	I don't know	111	12.7
How often do you check for information about COVID 19 per day?	Less than 5 times	490	56.1
	Less than 10 times	205	23.5
	Less than 20 times	146	16.7
	More than 21 times	32	3.7
Personal concerns regarding COVID-19 outbreak and lockdown			
How confident are you on your physician to diagnose	Highly confident	110	12.6
	Confident	528	60.5
	Not confident	51	5.8

COVID-19 infection?	I don't know	184	21.1
How likely are you to contract COVID-19 during this outbreak?	Highly likely	16	1.8
	Likely	248	28.4
	Not likely	345	39.4
	Highly not likely	89	10.2
	I don't know	175	20
How likely are you to survive, if contracted with COVID-19 infection?	Highly likely	129	14.8
	Likely	456	52.2
	Not likely	55	6.3
	Highly not likely	20	2.3
	I don't know	213	24.4
How concerned are you about your family members to contract COVID-19 infection during this outbreak?	Highly concerned	566	64.8
	Concerned	276	31.6
	Not concerned	11	1.3
	Highly not concerned	12	1.4
	I don't know	8	0.9

155 Average score for the three subscales were, depression: 5.02 ± 4.96 , anxiety:
 156 4.41 ± 3.41 and stress: 7.77 ± 7.42 . 18.56% had symptoms of depression with the
 157 majority having mild depression (15.1%), 25.66% had symptoms of anxiety where,
 158 participants with moderate anxiety predominated (16.3%) and 21.99% had
 159 symptoms of stress with the maximum in mild stress category (14.5%) (Figure 1).
 160 Our results were considerably higher than the prevalence of these negative
 161 components of psychological health assessed by studies in the population, before
 162 the pandemic [16, 17] while being similar to the findings by studies conducted in
 163 other countries during COVID-19 pandemic [6, 8] and SARS pandemic [4].
 164 Figure 1: Prevalence of Depression, Anxiety and Stress in the study population (in
 165 %)



166

167 **Association between sociodemographic, COVID-19**
 168 **outbreak & lockdown related variables and mental health**
 169 **status**

170 Marital status, wearing masks and gloves in public places, awareness of increase in
 171 number of COVID-19 cases, Satisfaction with the information received daily
 172 regarding COVID-19 outbreak (Table 2 and 3) were excluded in the final model for
 173 depression based on the initial binomial logistic regression.

174 Table 2: Bivariate analysis of sociodemographic variables by binary logistic
 175 regression

Independent variables	Depression				Anxiety				Stress			
	P		95% CI		P		95% CI		P		95% CI	
	val	cOR	Lower	Upper	value	cOR	Lower	upper	value	cOR	Lower	upper
Gender												
Male*	-	-	-	-	-	-	-	-	-	-	-	-
Female	.004	1.667	1.182	2.352	.0004	2.624	1.917	3.593	.0005	2.075	1.497	2.876

Age (years)	0.0				0.00				0.00			
	00				0				0			
18 to 25*	-	-	-	-	-	-	-	-	-	-	-	-
-26 to 35	.00	.456	.268	.776	.006	.526	.332	.833	.006	.465	.270	.801
	4											
36 to 45	.04	1.50	1.007	2.260	.087	1.38	.953	2.023	.005	1.78	1.187	2.669
	6	9				9				0		
46 to 55	.79	.893	.374	2.128	.008	2.50	1.267	4.942	.000	5.52	2.759	11.06
	8					3				4		0
56 & above	.36	.701	.326	1.509	.159	1.52	.847	2.762	.002	2.59	1.430	4.723
	4					9				9		
Educational	0.0				0.00				0.00			
status	00				0				0			
None*	-	-	-	-	-	-	-	-	-	-	-	-
Higher	.81	1.16	.333	4.078	.018	.337	.137	.830	.851	.899	.294	2.749
secondary	2	4										
school												
Bachelor's	.10	2.76	.819	9.361	.410	1.42	.613	3.316	.069	2.72	.927	8.015
degree	1	9				6				5		
Master's	.67	1.31	.362	4.753	.106	.464	.182	1.179	.766	.837	.260	2.691
degree	9	2										
Doctorate	.53	1.66	.334	8.324	.117	2.50	.795	7.861	.198	2.47	.624	9.788
degree	4	7				0				1		
Marital status	0.7				0.29				0.00			
	17				8				5			
Single*	-	-	-	-	-	-	-	-	-	-	-	-
Married	.41	.867	.616	1.221	.309	.853	.628	1.159	.252	1.21	.872	1.689
	5									3		
Widowed/ separated	.99	.246	.046	.623	.200	.375	.084	1.678	.001	5.27	1.905	14.62
	8									8		4

Employment	0.0				0.00				0.00			
status	00				0				0			
Student*	-	-	-	-	-	-	-	-	-	-	-	-
Employed -	.07	.373	.128	1.090	.540	1.24	.621	2.482	.997	.042	.009	.103
Government	1					2						
Employed –	.03	1.63	1.027	2.616	.001	2.12	1.368	3.297	.003	1.96	1.252	3.097
private	8	9				4				9		
Self-business	.00	2.86	1.680	4.902	.000	4.26	2.575	7.069	.000	3.85	2.298	6.457
	0	9				7				2		
Unemployed	.00	3.33	1.907	5.844	.000	6.14	3.620	10.444	.000	5.60	3.274	9.589
	0	8				9				3		
Monthly	0.0				0.00				0.00			
income (INR)	00				4				3			
10,000 and	-	-	-	-	-	-	-	-	-	-	-	-
less												
10,001-20,000	.16	.472	.162	1.376	.840	1.07	.517	2.249	.052	.383	.146	1.010
	9					9						
20,001-30,000	.99	.002	.000	.016	.197	1.67	.766	3.647	.998	.007	.000	.015
	8					2						
30,001-40,000	.10	2.37	1.363	4.150	.416	1.26	.720	2.210	.566	1.17	.674	2.059
	2	8				2				8		
40,001-50,000	.99	.004	.000	.023	.997	.018	.004	.048	.997	.017	.000	.031
	7											
50,001-	.00	.052	.358	1.099	.203	.800	.228	1.640	.187	1.29	.881	1.916
100,000	1									9		
100,000 &	.00	.350	.177	.689	.042	.657	.402	1.072	.002	.430	.252	.734
above	2											
Parental	0.0				0.00				0.00			
status	00				0				0			

No kid	.00	.067	.036	.123	.000	.084	.044	.159	.000	.047	.024	.091
0												
1 kid	.00	.002	.000	.016	.000	.013	.005	.034	.000	.003	.001	.013
0												
2 kids	.00	.035	.018	.070	.000	.047	.024	.094	.000	.053	.026	.105
0												
3 or more kids*	-	-	-	-	-	-	-	-	-	-	-	-
Household size	0.0				0.02				0.00			
00					6				0			
1 member	.76	1.25	.292	5.348	.722	.773	.187	3.196	.999	.005	.000	.015
4	0											
2 members	.00	.083	.023	.300	.008	.294	.120	.722	.000	.029	.008	.099
0												
3-6 members	.31	.649	.280	1.504	.147	.563	.259	1.223	.001	.280	.131	.601
4												
More than 6 members*	-	-	-	-								

176 Abbreviations: cOR, crude odds ratio; 95% CI, 95% confidence interval; * Reference
177 group

178 Table 3: Bivariate analysis of COVID-19 outbreak & lockdown related variables using
179 binary logistic regression

Independent variables	Depression				Anxiety				Stress		
	P	95% CI			P	95% CI			P	95% CI	
		value	cOR	Lower Upper		value	cOR	Lower Upper		value	cOR

Wearing	0.06				0.05					0.64		
mask and	6				4					8		
gloves while												
being												
outside in												
public places												
No*	-	-	-	-	-	-	-	-	-	-	-	-
Sometimes	.571	.794	.358	1.763	.017	2.18	1.152	4.127	.518	1.24	.637	2.447
Yes	.176	1.49	.836	2.663	.121	1.51	.896	2.574	.352	1.28	.757	2.186
		2				9				7		
Frequency of	0.00				0.00					.000		
washing	0				0							
hands with												
soap or hand												
sanitizer per												
day												
Less than 5	.000	.091	.052	.161	.000	.108	.062	.188	.000	.084	.047	.149
times												
5 to 10 times	.000	.040	.022	.074	.000	.117	.068	.200	.000	.054	.030	.098
10 to 15 times	.000	.101	.057	.179	.000	.166	.096	.287	.000	.139	.080	.242
More than 15												
times*	-	-	-	-	-	-	-	-	-	-	-	-
Awareness												
of increase in												
number of												
COVID-19												
cases												
Yes	0.99	3.42	1.80	4.30	0.74	0.59	0.041	1.023	0.52	2.51	0.962	2.985
	8											

No	-	-	-	-	-	-	-	-	-	-	-	-
Satisfaction	.212				.000				.040			
with the information received daily regarding COVID-19 outbreak												
Highly satisfied	.023	.235	.067	.820	.023	.276	.091	.837	.004	.161	.047	.556
Satisfied	.411	.814	.498	1.330	.783	.936	.586	1.496	.099	.684	.435	1.074
Not satisfied	.229	.657	.332	1.302	.159	1.52	.847	2.743	.064	.550	.291	1.037
						4						
Highly not satisfied	.812	.892	.347	2.290	.103	3.33	1.501	7.408	.300	.613	.243	1.546
						5						
I don't know	-	-	-	-	-	-	-	-	-	-	-	-
Frequency of checking for information about COVID 19 per day	.000				.000				.000			
Less than 5 times	.043	.419	.180	.974	.759	.879	.384	2.010	.243	2.05	.613	6.909
										8		
Less than 10 times	.121	.494	.203	1.205	.517	.750	.314	1.791	.361	1.78	.514	6.223
										8		
Less than 20 times	.046	2.40	1.014	5.713	.054	2.34	.987	5.557	.000	9.15	2.669	31.37
		7				1				1		6
More than 21 times	-	-	-	-	-	-	-	-	-	-	-	-

Level of confidence on physician to diagnose COVID-19 infection	.000				.000				.000			
Highly confident	.129	1.46	.894	2.407	.883	1.03	.640	1.681	.402	1.23	.756	2.011
		7				7				3		
Confident	.000	.293	.194	.443	.000	.315	.217	.458	.000	.358	.244	.526
Not confident	.002	.143	.043	.478	.808	.924	.487	1.752	.001	.120	.036	.401
I don't know	-	-	-	-	-	-	-	-	-	-	-	-
Likelihood of contracting COVID-19	.000				.000				.000			
Highly likely	.007	4.65	1.526	14.168	.001	6.29	2.157	18.35	.022	3.37	1.191	9.564
		0				2		1		5		
Likely	.011	2.05	1.178	3.589	.014	1.92	1.140	3.235	.823	1.05	.666	1.666
		6				1				4		
Not likely	.334	1.31	.755	2.285	.000	2.56	1.574	4.190	.014	.559	.351	.889
		4				8						
Highly not likely	.000	4.79	2.546	9.015	.000	3.88	2.120	7.136	.003	2.29	1.321	3.978
		1				9				2		
I don't know	-	-	-	-	-	-	-	-	-	-	-	-
Likelihood of surviving, if infected with COVID-19	.000				.302				.000			
Highly likely	.000	.322	.177	.587	.397	.811	.500	1.317	.000	.269	.146	.494
Likely	.000	.413	.280	.608	.029	.668	.465	.959	.001	.538	.373	.774
Not likely	.997	.000	.000	.	.549	.817	.422	1.581	.997	.000	.000	.

Highly not likely	.031	2.78	1.100	7.039	.998	.000	.000	.	.019	3.06	1.198	7.838
I don't know	-	-	-	-	-	-	-	-	-	-	-	-
Level of concern on family members to be infected	.000				.001				.000			
Highly concerned	.080	.286	.071	1.161	.196	.398	.098	1.609	.107	.316	.078	1.282
Concerned	.316	2.66	.391	18.166	.554	1.75	.275	11.15	.554	1.75	.275	11.15
Not concerned	.000	.049	.011	.220	.044	.232	.056	.959	.006	.136	.032	.569
Highly not concerned	.057	11.0	.928	130.32	.999	.000	.000	.	.999	.317	.090	.742
I don't know	-	-	-	-	-	-	-	-	-	-	-	-

180 Abbreviations: cOR, crude odds ratio; 95% CI, 95% confidence interval; * Reference
 181 group
 182 Multiple logistic regression model for depression using stepwise forward LR method
 183 was statistically significant, $\chi^2(52) = 594.77$, $P < 0.000$. The model explained 80.1%
 184 (Nagelkerke R^2) of the variance in depression and correctly classified 94.6% of
 185 cases. Six variables were found to be significant predictors including female gender,
 186 parental status, frequency of hand washing, level of satisfaction with the information
 187 received daily, level of confidence on the physician and likelihood of contracting
 188 COVID-19 (Table 4).

189 Table 4: Multiple logistic regression analysis for depression and its correlates

Variables	Categorie	B	S.E.	Wald	P	aOR	95% CI for aOR
-----------	-----------	---	------	------	---	-----	----------------

	s	value					Lower	Upper
Gender	Male*	-	-	-	-	-	-	-
	Female	1.367	.545	6.300	.012	3.924	1.349	11.412
Parental status				12.381	.006			
	No kid	2.805	4.960	.320	.572	6.529	.001	30.392
	One kid	-5.463	44.34 1	.008	.930	0.396	.053	.461
	Two kids	-1.572	3.955	4.698	.030	0.189	.051	0.440
	Three or more kids*	-	-	-	-	-	-	-
Frequency of washing hands with soap or hand sanitizer per day				10.139	.017			
	Less than 5 times	-1.265	.725	3.047	.081	.282	.068	1.168
	5 to 10 times	-1.877	.644	8.498	.004	.153	.043	.541
	10 to 15 times	-1.615	.753	4.593	.032	.199	.045	.871
	More than 15 times*	-	-	-	-	-	-	-
Satisfaction with the information received daily				20.434	.000			
	Highly satisfied	-3.608	3.813	.895	.344	.127	.089	17.758
	Satisfied	-8.529	1.916	19.818	.000	.405	.159	.740

regarding COVID-19 outbreak	Not satisfied	-2.440	1.594	2.342	.126	.087	.004	1.984
	Highly not satisfied	4.762	9.993	.000	.998	.599	.000	2.105
	I don't know	-	-	-	-	-	-	-
Level of confidence on physician to diagnose COVID-19 infection				18.363	.000			
	Highly confident	.692	.544	1.619	.203	1.998	.688	5.807
	Confident	-1.116	.466	5.738	.017	.328	.132	.816
	Not confident	-.292	.971	.090	.764	.747	.111	5.006
	I don't know	-	-	-	-	-	-	-
Likelihood of contracting COVID-19				15.302	.004			
	Highly likely	-.121	.902	.018	.893	.886	.151	5.193
	Likely	-.988	.562	3.096	.078	.372	.124	1.119
	Not likely	-.097	.503	5.782	.016	.151	.021	.294
	Highly not likely	-1.504	.626	.037	.847	.907	.338	2.432
	I don't know*	-	-	-	-	-	-	-

190 Abbreviations: aOR, adjusted odds ratio; 95% CI, 95% confidence interval; B,
 191 unstandardized beta coefficient; S.E, standard error of the coefficient; * Reference
 192 group

193 According to bivariate logistic regression analysis, among the sociodemographic
 194 variables, marital status (Table 2) and among COVID-19 outbreak & lockdown
 195 related factors, wearing masks and gloves in public places, awareness of increase in
 196 number of COVID-19 cases, Likelihood of surviving, if infected with COVID-19 (Table
 197 3) were excluded from the final model for anxiety.

198 The multiple logistic regression model for anxiety using stepwise forward LR method
 199 showed significant goodness of fit to our observed data, $\chi^2(48) = 455.7$, $P < 0.000$.
 200 The model explained 59.8% (Nagelkerke R^2) of the variance in anxiety and correctly
 201 classified 86.8% of cases. The significant predictors for anxiety were found to be
 202 female gender, educational status, employment status, frequency of checking for
 203 information regarding COVID-19, level of confidence on the physician and likelihood
 204 of contracting COVID-19 (Table 5).

205 Table 5: Multiple logistic regression analysis for anxiety and its correlates

Variables	Categories	B	S.E.	Wald	P value	aOR	95% CI for aOR	
							Lower	Upper
Gender	Male*	-	-	-	-	-	-	-
	Female	1.248	.266	21.947	.000	3.483	2.066	5.870
Educational status				14.839	.005			
	None*	-	-	-	-	-	-	-
	Higher secondary school	-2.069	1.104	3.516	.061	.126	.015	1.098
	Bachelor's degree	-.883	.971	.826	.363	.414	.062	2.775

	Master's degree	-	1.014	4.795	.029	.108	.015	.592
	2.221							
	Doctorate degree	.025	1.239	.000	.984	1.026	.090	11.626
<hr/>								
Employment status				11.472	0.022			
	Student*	-	-	-	-	-	-	-
	Employed Government	-	1.045	.798	1.717	.190	2.845	.596
	13.584							
	Employed – private	-	1.324	.601	4.856	.028	3.757	1.158
	12.191							
	Self-business	1.440	.696	4.281	.039	4.222	1.079	16.523
	Unemployed	1.674	.498	11.293	.001	5.335	2.009	14.164
<hr/>								
Frequency of checking for information about COVID 19 per day				9.651	.022			
	Less than 5 times	-.892	.847	1.109	.029	.410	.078	.657
	10 times	-.167	.740	.051	.821	.846	.198	3.607
	20 times	.875	.939	.869	.351	2.400	.381	15.121
	More than 21 times*	-	-	-	-	-	-	-
<hr/>								
Level of				12.598	.006			

confidence	Highly	.169	.613	.076	.782	1.185	.356	3.942
on	confident							
physician to		-	.520	5.503	.019	.295	.107	.818
diagnose	Confident	1.220						
COVID-19	Not	.389	.946	.169	.681	1.475	.231	9.418
infection	confident							
	I don't	-	-	-	-	-	-	-
	know*							
Likelihood				19.332	.001			
of	Highly likely	.672	.964	.486	.486	1.959	.296	12.965
contracting	Likely	-.970	.563	2.976	.085	.379	.126	1.141
COVID-19	Not likely	-.023	.515	1.002	.014	.777	.356	.981
	Highly not	1.887	.634	8.867	.403	6.599	1.906	22.850
	likely							
	I don't	-	-	-	-	-	-	-
	know*							

206 Abbreviations: aOR, adjusted odds ratio; 95% CI, 95% confidence interval; B,
 207 unstandardized beta coefficient; S.E, standard error of the coefficient; * Reference
 208 group

209 Wearing masks and gloves in public places, awareness of increase in number of
 210 COVID-19 cases were excluded from the final model for stress based on bivariate
 211 logistic regression analysis (Table 3). The multiple logistic regression model for
 212 stress using stepwise forward LR method showed significant goodness of fit to our
 213 observed data, $\chi^2(54) = 621.77$, $P < 0.000$. The model explained 78.2% (Nagelkerke
 214 R^2) of the variance in stress and correctly classified 92.1% of cases. Five variables
 215 were found to be significant predictors including age of the participants, marital

216 status, household size, level of confidence on the physician and likelihood of
 217 contracting COVID-19 (Table 6).

218 Table 6: Multiple logistic regression analysis for stress and its correlates

Variables	Categories	B	S.E.	Wald	P value	aOR	95% CI for aOR	
							Lower	Upper
Age (years)				14.055	.007			
	18 to 25*	-	-	-	-	-	-	-
	26 to 35	-3.543	2.130	2.766	.096	.029	.000	1.882
	36 to 45	-5.531	2.290	5.832	.016	.14	.102	.353
	46 to 55	-4.912	1.832	7.190	.007	.17	.094	.267
	56 & above	-.526	2.557	.042	.837	.591	.004	18.803
Marital status				19.503	.000			
	Single*	-	-	-	-	-	-	-
	Married	7.753	1.756	19.493	.000	2.502	1.701	8.924
	Widowed/ separated	5.100	1.931	6.974	.008	10.411	1.037	14.566
Household size				18.370	.000			
	1 member	-1.996	.927	4.634	.031	.136	.022	.836
	2 members	-5.705	1.104	26.699	.000	.213	.100	.429
	3-6 members	-2.954	.818	13.036	.000	.352	.181	.559

	More than							
	6	-	-	-	-	-	-	-
	members*							
Level of confidence on physician to diagnose COVID-19 infection				16.540	.001			
	Highly confident	2.610	1.174	4.947	.126	3.603	1.363	135.729
	Confident	-5.012	1.648	9.251	.002	.807	.450	.968
	Not confident	-6.395	2.517	6.456	.011	.002	.001	.232
	I don't know*	-	-	-	-	-	-	-
Likelihood of contracting COVID-19				22.887	.000			
	Highly likely	-1.308	.949	1.899	.168	.270	.042	1.737
	Likely	2.161	.754	8.220	.004	1.115	1.026	1.505
	Not likely	-5.219	1.102	22.441	.000	.005	.001	.047
	Highly not likely	-2.510	1.634	2.359	.125	.081	.003	2.000
	I don't know*	-	-	-	-	-	-	-

219 Abbreviations: aOR, adjusted odds ratio; 95% CI, 95% confidence interval; B,
 220 unstandardized beta coefficient; S.E, standard error of the coefficient; * Reference
 221 group

222 We found that, females were 1.7 times more likely to be present with depressive
 223 symptoms (cOR-1.667), 2.6 times as likely to show symptoms of anxiety (cOR-
 224 2.624) and twice as likely to show stress symptoms (cOR-2.075) when compared to
 225 males but when adjusted for other confounding factors, females were quadruple

226 times as likely as males to have depression (aOR-3.924) and 3.5 times as likely as
227 males to have anxiety (aOR-3.483). Our results are similar to previous studies
228 conducted during SARS and COVID-19 pandemic in Italy and China, where the
229 prevalence of psychological disorders were steadily associated with the female
230 gender [4, 6, 8]. This trend was noted in India before the outbreak too where the
231 prevalence in depression and anxiety disorders were more in females than males
232 [18]. There was no significant association between gender and likelihood of stress
233 which is in accordance with studies conducted during earlier epidemic/pandemic
234 situations [5, 19]. Gender specific help and counselling should be offered to mitigate
235 the psychological strain in the population.

236 An interesting finding in our study was the variation in association between age and
237 depression, anxiety and stress. Age of the participants was significantly associated
238 with all three domains, where adult Indians of age 36 to 45 years were 1.5 times
239 more likely to have depression (cOR-1.509) and individuals of 46 to 55 years of age
240 were 2.5 times as likely to have anxiety (cOR-2.503) when compared to young adults
241 of age 18 to 25 years. There was also an increase in likelihood of stress with
242 increase in age (36 to 45 years-cOR-1.708, 46 to 55 years-cOR-5.524) with slight
243 decrease in higher risk of stress in age above 55 years (cOR-2.599). However, when
244 controlled for other factors, individuals above 25 years of age were found to be less
245 likely to exhibit symptoms of stress when compared to younger adults of age below
246 25 years (Table 6). This finding is similar to studies conducted in similar situations
247 where younger age was found to associated with increased likelihood of stress and
248 psychological distress[5, 6]. Young adults may have trouble coping with drastic
249 societal changes and are more active in social media which is swarming with

250 rumours, which may induce fear, anxiety and other psychological effects. Special
251 focus in identifying and providing mental health help for this age group is imperative.

252 With increase in level of educational status there was lower likelihood of anxiety but
253 not depression and stress. Those with higher educational qualification were found to
254 be less likely to show anxiety symptoms when compared to those who had none
255 (higher secondary school - cOR-0.337, master's degree - aOR-0.108). Our finding is
256 similar to the studies conducted both before and in relation to pandemic/epidemic [4,
257 8, 18, 20]. Educational status influences the occupation and income of the individual,
258 which are in turn associated with psychosocial wellbeing. The economic crisis due to
259 lockdown, might put the individuals without formal education at higher risk of
260 developing anxiety. Counselling, guidance or any form of mental health help should
261 include verbal or pictorial representations to aid this group of population.

262 Marital status was significantly associated with stress and not with depression and
263 anxiety. Married individuals were 2.5 times as likely (aOR-2.502) and
264 widowed/separated individuals were 10 times as likely as single ones (aOR-10.411)
265 to have stress symptoms. This is in contrary to the studies conducted in China, Iran
266 and Italy during COVID-19 pandemic, where there was no association between
267 marital status and negative mental status[6-8]. However, in India, being married was
268 found to be strongly associated (6 times as likely as single individuals) with increase
269 in the prevalence of mental health disorders in women unlike other neighbouring
270 countries like China and other high resourced countries [21]. Negative psychological
271 effects were observed in separated and widowed individuals [22]. This trend seems
272 to be continuing during COVID-19 quarantine as well which is evident from our
273 results. Married, separated/widowed individuals should be given additional help in
274 combating psychological distress during this pandemic.

275 Employment status was significantly associated with depression, anxiety and stress
276 where being unemployed and employed as self or in private sector were found to
277 have higher risk of negative mental health when compared to student status. When
278 adjusted for other confounders, unemployed individual were 5 times as likely (aOR-
279 5.335) followed by individuals in self business who were 4 times as likely (aOR-
280 4.222) and private sector employees who were 3.75 times (aOR-3.757) as likely to
281 have anxiety symptoms. This is in contrary to the findings by Wang et al., who found
282 that the student population suffered from higher levels of depression, anxiety and
283 stress during COVID-19 pandemic in China[8]. Uncertainties regarding job security
284 especially in self business and private sector along with the financial strain posed by
285 the quarantine could be the cause for high prevalence of anxiety in the said
286 population.

287 Monthly income was significantly associated with depression, anxiety and stress,
288 where higher income was found to be protective against negative mental health
289 components. Individuals with monthly income above 100,000INR was less likely to
290 have depression (cOR-0.350) and stress (cOR-0.430) when compared to individuals
291 with income less than 10,000INR per month. This is in accordance with previous
292 studies in India where there was an inverse relation between income and common
293 mental disorders [23]. However, when adjusted for the effects of confounders, there
294 was no independent association between monthly income and depression, anxiety
295 and stress. The relative financial stability in the high-income population could be the
296 factor for relatively lesser psychological distress in the said population.

297 Having two kids posed lesser risk of depression when compared to having 3 or more
298 kids (aOR-0.189), while there was no significant association with anxiety and stress.
299 However, in bivariate analysis, not having kids and having 1 or 2 kids were found to

300 be protective against depression and anxiety and having one kid or none were found
301 to be protective against stress when compared to having 3 or more children (Table
302 2). There are varied reports regarding association between parental status and
303 psychological distress. Not having kids was associated with depression during
304 COVID-19 pandemic in Italy [6], having three or more kids were associated with
305 lesser risk of psychological distress during equine influenza epidemic in Australia [5]
306 and having kids was not associated with depression during SARS quarantine in
307 Canada [4]. With the kids being home-schooled, burden on the parents have
308 increased and could be a cause for increase in psychological distress which may
309 increase with the number of kids.

310 Household size of two members was found to be protective against depression and
311 anxiety and household size of 2 members, 3 to 6 members were protective against
312 stress when compared to bigger household size. When controlled for confounders,
313 individuals from smaller family size viz. one member (aOR-0.136), 2 members (aOR-
314 0.213) and 3 to 6 members (aOR-0.352) were found to be less likely to have
315 symptoms of stress when compared to individuals from family size of more than 6
316 members. This is in contrary to a previous study in China where there was no
317 association between household size and psychological distress [8, 24]. Lack of
318 personal space and higher financial strain in larger families could be the possible
319 reasons behind our results. Household size should be taken into consideration by
320 the mental health professionals while offering guidance and counselling.

321 In our study, practice of specific precautionary measures like frequency of washing
322 hands was found to be significantly related to depression, anxiety and stress while
323 wearing masks and gloves in public places had no association when not controlled
324 for confounders. In the final regression model however, lesser frequency of washing

325 hands (less than 15 times) was associated with lesser likelihood of anxiety (Table 5).
326 Previously, anxiety was associated with increase in practice of preventive measures
327 during SARS outbreak [25]. In a survey conducted during 2020 March 22nd -24th in
328 India, 75% of the respondents were found to use gloves and sanitizers, whereas in
329 our study, the number of individuals who do not wear masks and gloves in public
330 places was 11.8% which is higher when compared to China (3.2%) [8, 9]. In our
331 study, 28.9%, 22.1% and 32.3% of participants were not aware that droplets, contact
332 with infected persons and contact with contaminated objects are possible routes of
333 transmission of COVID-19 respectively (Table 1). This lack of awareness could be
334 related to the laxity in practicing personal precautionary measures.

335 Those who checked for information less frequently (<5 times/day) about COVID-19
336 were found to be less likely to show symptoms of anxiety when compared to those
337 who checked for more than 21 times a day (aOR-0.410). 74% of the respondents
338 gathered information about the pandemic through internet including social media
339 which gives many inconsistent and fake news which may cause fear and anxiety.
340 Individuals who were satisfied with the information received were found to be less
341 likely to show depression symptoms (aOR-0.405) which was similar to the findings in
342 China [8]. The population should be urged to follow authentic news provided by
343 reliable sources to avoid psychological distress.

344 Those who felt confident on the physician's ability to diagnose COVID-19 infection
345 were found to be less likely to have symptoms of depression (aOR-0.328), anxiety
346 (aOR-0.295) and stress (aOR-0.807), similar to the study in China [24]. The
347 confidence in physician's ability gives a sense of security to the individuals and
348 hence reduces fear and psychological distress.

349 Individuals who expressed that they were not likely to contract COVID-19 during this
350 outbreak were less likely to have symptoms of depression (aOR-0.151), anxiety
351 (aOR-0.777) and those who thought they were likely to contract COVID-19 were
352 more likely to have symptoms of stress (aOR-1.115). Similarly, the participants who
353 thought that it was highly not likely for them to survive if infected were found to be
354 more likely to have depression (cOR-2.783) and stress symptoms (cOR-3.064) while
355 those who thought it was likely for them to survive had lower levels of anxiety (cOR-
356 0.668) and stress (cOR-0.538). We also found that individuals who were not
357 concerned about their family members contracting COVID-19 were less likely to
358 show symptoms of depression (cOR-0.049), anxiety (cOR-0.232, 95% CI 0.056-
359 0.959) and stress (cOR-0.136). Our findings are similar to previous studies where
360 low levels of perceived likelihood of contracting COVID-19 and surviving the
361 pandemic if infected were protective against depression, anxiety and stress [24].
362 These concerns should be considered by the mental health workers when providing
363 mental health help for the population.

364 **Limitations**

365 Our study is limited by the cross-sectional nature and the non-availability of control
366 group. The longitudinal effects of the pandemic and lockdown is not ascertained. The
367 study participants included only those who had access to internet and those who
368 could respond in English. None of our participants were tested positive for COVID-19
369 and neither had any contact history with known COVID-19 patients. Thus, our
370 findings may not be generalised to the COVID-19 infected patients and their peers.

371 **Conclusion**

372 To our knowledge, our study is the first to assess the mental health status of the
373 adult Indian population during COVID-19 outbreak & lockdown along with identifying
374 the possible risk and protective factors. During the third phase of the lock down, less
375 than one fifth of the adult Indian population suffered from depression, one fourth
376 suffered from anxiety and more than one fifth suffered from stress. Females were
377 more likely to suffer from depression and anxiety when compared to males.
378 Employment in the government sector and higher educational status were protective
379 against anxiety. Age above 25 years, smaller household size and single status were
380 associated with decrease levels of stress. Parents with lesser number of kids (≤ 2) or
381 none were less likely to suffer from depression when compared to parents with more
382 than 2 kids. Increased levels of confidence in physician's ability to diagnose COVID-
383 19 infection, decreased self-perceived likelihood of contracting the infection were
384 associated with decreased levels of depression, anxiety and stress. Less frequency
385 of checking for information on COVID-19 was associated with decreased levels of
386 anxiety and satisfaction of information received about COVID-19 pandemic was
387 associated with decreased levels of depression.

388 The ripple caused by the COVID-19 outbreak & lockdown will be continuing far into
389 the future and providing mental health support to the population, targeting the
390 vulnerable groups is crucial. Our study provides an expansive assessment of risk
391 and protective factors affecting the mental health of the population, which would help
392 to design strategies and interventional methods to address and mitigate the negative
393 impact of COVID-19 outbreak & lockdown on the mental health of the population and
394 help prevent the same.

395 Acknowledgement:

396 The authors would like to thank all the participants in this study. The authors would
397 like to thank the Deanship of Scientific Research, Majmaah University for the support
398 of this project.

399 References

- 400 1. WHO. Rolling updates on coronavirus disease (COVID-19) Geneva2020 [updated 25th May
401 2020. Available from: [https://www.who.int/emergencies/diseases/novel-coronavirus-2019/events-](https://www.who.int/emergencies/diseases/novel-coronavirus-2019/events-as-they-happen)
402 [as-they-happen](https://www.who.int/emergencies/diseases/novel-coronavirus-2019/events-as-they-happen).
- 403 2. Press TC. Asia Today: India sees 10,000 new cases ahead of reopenings [e paper]. New Delhi:
404 The Canadian Press; 2020 [updated 7th June 2020]. Available from:
405 <https://ca.finance.yahoo.com/news/asia-today-india-reports-10-071148089.html>.
- 406 3. WHO. Modes of transmission of virus causing COVID-19:
407 implications for IPC precaution recommendations Scientific brief. Geneva, control C-iap; 2020 29
408 March 2020.
- 409 4. Hawryluck L, Gold WL, Robinson S, Pogorski S, Galea S, Styra R. SARS control and
410 psychological effects of quarantine, Toronto, Canada. *Emerging Infectious Diseases*.
411 2004;10(7):1206.
- 412 5. Taylor MR, Agho KE, Stevens GJ, Raphael B. Factors influencing psychological distress during
413 a disease epidemic: data from Australia's first outbreak of equine influenza. *BMC Public Health*.
414 2008;8:347.
- 415 6. Mazza C, Ricci E, Biondi S, Colasanti M, Ferracuti S, Napoli C, et al. A Nationwide Survey of
416 Psychological Distress among Italian People during the COVID-19 Pandemic: Immediate
417 Psychological Responses and Associated Factors. *International Journal of Environmental Research*
418 *and Public Health*. 2020;17(9):3165.
- 419 7. Moghanibashi-Mansourieh A. Assessing the anxiety level of Iranian general population
420 during COVID-19 outbreak. *Asian J Psychiatr*. 2020;51:102076.
- 421 8. Wang C, Pan R, Wan X, Tan Y, Xu L, Ho CS, et al. Immediate psychological responses and
422 associated factors during the initial stage of the 2019 coronavirus disease (COVID-19) epidemic
423 among the general population in China. *International journal of environmental research and public*
424 *health*. 2020;17(5):1729.
- 425 9. Roy D, Tripathy S, Kar SK, Sharma N, Verma SK, Kaushal V. Study of knowledge, attitude,
426 anxiety & perceived mental healthcare need in Indian population during COVID-19 pandemic. *Asian*
427 *journal of psychiatry*. 2020;51:102083-.
- 428 10. Suseela V. Psychological Issues Based on Gender and Marital Status During Covid-19
429 Lockdown Period. *Tathapi with ISSN 2320-0693 is an UGC CARE Journal*. 2020;19(8):755-64.
- 430 11. Chew NW, Lee GK, Tan BY, Jing M, Goh Y, Ngiam NJ, et al. A multinational, multicentre study
431 on the psychological outcomes and associated physical symptoms amongst healthcare workers
432 during COVID-19 outbreak. *Brain, behavior, and immunity*. 2020.
- 433 12. Suryadevara V, Adusumalli C, Adusumilli PK, Chalasani SH, Radhakrishnan R. Mental Health
434 Status among the South Indian Pharmacy Students during Covid-19 Pandemic Quarantine Period: A
435 Cross-Sectional Study. *medRxiv*. 2020.

- 436 13. Bottesi G, Ghisi M, Altoè G, Conforti E, Melli G, Sica C. The Italian version of the Depression
437 Anxiety Stress Scales-21: Factor structure and psychometric properties on community and clinical
438 samples. *Comprehensive psychiatry*. 2015;60:170-81.
- 439 14. Henry JD, Crawford JR. The short-form version of the Depression Anxiety Stress Scales
440 (DASS-21): Construct validity and normative data in a large non-clinical sample. *British journal of*
441 *clinical psychology*. 2005;44(2):227-39.
- 442 15. Lovibond PF, Lovibond SH. The structure of negative emotional states: Comparison of the
443 Depression Anxiety Stress Scales (DASS) with the Beck Depression and Anxiety Inventories.
444 *Behaviour research and therapy*. 1995;33(3):335-43.
- 445 16. Charlson FJ, Baxter AJ, Cheng HG, Shidhaye R, Whiteford HA. The burden of mental,
446 neurological, and substance use disorders in China and India: a systematic analysis of community
447 representative epidemiological studies. *The Lancet*. 2016;388(10042):376-89.
- 448 17. Pradeep BS, Gururaj G, Varghese M, Benegal V, Rao GN, Sukumar GM, et al. National mental
449 health survey of India, 2016-rationale, design and methods. *PLoS One*. 2018;13(10):e0205096.
- 450 18. Sagar R, Dandona R, Gururaj G, Dhaliwal R, Singh A, Ferrari A, et al. The burden of mental
451 disorders across the states of India: the Global Burden of Disease Study 1990–2017. *The Lancet*
452 *Psychiatry*. 2020;7(2):148-61.
- 453 19. Su TP, Lien TC, Yang CY, Su YL, Wang JH, Tsai SL, et al. Prevalence of psychiatric morbidity
454 and psychological adaptation of the nurses in a structured SARS caring unit during outbreak: a
455 prospective and periodic assessment study in Taiwan. *J Psychiatr Res*. 2007;41(1-2):119-30.
- 456 20. Bjelland I, Krokstad S, Mykletun A, Dahl AA, Tell GS, Tambs K. Does a higher educational level
457 protect against anxiety and depression? The HUNT study. *Social science & medicine*.
458 2008;66(6):1334-45.
- 459 21. Patel V, Kirkwood BR, Pednekar S, Weiss H, Mabey D. Risk factors for common mental
460 disorders in women. Population-based longitudinal study. *Br J Psychiatry*. 2006;189:547-55.
- 461 22. Carr D, Springer KW. Advances in Families and Health Research in the 21st Century. *Journal*
462 *of Marriage and Family*. 2010;72(3):743-61.
- 463 23. Cheng HG, Shidhaye R, Charlson F, Deng F, Lyngdoh T, Chen S, et al. Social correlates of
464 mental, neurological, and substance use disorders in China and India: a review. *The Lancet*
465 *Psychiatry*. 2016;3(9):882-99.
- 466 24. Wang C, Pan R, Wan X, Tan Y, Xu L, McIntyre RS, et al. A longitudinal study on the mental
467 health of general population during the COVID-19 epidemic in China. *Brain, behavior, and immunity*.
468 2020.
- 469 25. Leung GM, Quah S, Ho LM, Ho SY, Hedley AJ, Lee HP, et al. Community psycho-behavioural
470 surveillance and related impact on outbreak control in Hong Kong and Singapore during the SARS
471 epidemic. *Hong Kong Med J*. 2009;15 Suppl 9:30-4.

472